

Index

- Abricta curvica*, 265
 absolute rates
 desirability of, 259–61
Acacia, 287
Acanthiza, 288
Acanthocyclops, 125
 Africa, 48
 central-west, 250
 South, 61
 agria, *see* *Stenocereus gummosus*
Agropyron repens, 231
Alces alces, 86–7
 alcohol
 salicyl, 97
 alga, 100, 340
 red, *see* *Gelidium robustum*
 allelochemical, 79, 86, 94, 99, 324
Allium cepa, 131
Amanita
 muscaria, 130, 131
 pantharina, 130
Amazilia tobaci, 60
 Amazonia, 250
Amblycercus holosericeus, 121
 America
 Central, 48, 121
 North, 48, 106, 110, 118, 149, 165,
 214
 South, 48, 121
 amphibian, 54
 anuran, 2, 37–8, 158
 amphipod
 Caprellidae, 146–7
 Gammaridae, 146–7
 see *Pontoporeia affinis*
 analysis
 detrended correspondence, 270–1
 isoleg, 284
 principal component, 270
 sensitivity, 225
Anolis, 37, 58, 79, 159, 327
 eugenegrahamsi, 338
 evermanni, 74–6
 gundlachi, 74–6
 humilis, 73
 limifrons, 73
 stratulus, 74–5
 vermiculatus, 338
Anoplocephala, 149
 Auostraca 149
 ant, 312
 see *Lasius*, *Conomyrma bicolor*,
 Myrmecocystus kennedyi,
 Conomyrma insana, *Crematogaster*
 emeryana, *Iridomyrmex humilis*
 Antarctica, 48
Anthrocothorax nigricollis, 60
 aphid
 fundatrix, 154
 polymorphism, 154
 summer female, 154
 areography, 4
 Asia, 48, 106
 ASM, *see* model, apparent size
Atherina breviceps, 61
 Australia, 172, 301
 Great Dividing Range, 287
 New South Wales, 272, 288, 303
 Queensland, 288
 south-eastern, 194, 275
 south-western, 275
 Victoria, 9, 272, 287, 288
 Western Australia, 224
 autocorrelation, spatial, 297

424 · Index

- bamboo, *see* *Chusquea*
 bananaquit, *see* *Coereba flaveola*
Banksia, 224
 bat, 334
 insectivorous, 120
 Mollosidae, *see* *Nyctinomops*,
 Molossus
 surface-gleaning, 56
 bathymetry, 50
 bear, grizzly, *see* *Ursus arctos*
 beetle, leaf, *see* *Phratora vitellinae*,
 Galerucella lineola, *Phyllobrotica*
Bellis perennis, 131
Betula pedula, 131
 biogeography, 4
 zone, 275
 bird, 54, 320, 337
 fish-eating, 106
 foliage-gleaning, 172
 frugivorous, 56, 321
 granivorous, 321
 nectarivorous, 67, 126, 321
 rainforest, 250
 body size, of consumer, 315, 330
 Boidae, 37
Boletinus pictus, 130
Boletus rubellus, 130
Bosmina, 125
 burbot, *see* *Lota lota*
 butterfly, 54, 89
 Danainae, 89
 Heliconiinae, 89
 Ithomiinae, 89
 Lycaenidae, *see* *Plebejus argus*
 Nymphalidae, 89
 butterwort, *see* *Pinguicula*
 cacique
 yellow-billed, *see* *Amblycercus*
 holosericeus
 cactus, columnar, 52
Caffrogobius multifasciatus, 61
 Canada, 96
 Alberta, 105
 New Brunswick, 64
 Canadian Shield, 124
 cannibalism, 54, 122, 125
Cantharellus cibarius, 130
 carnivore
 plant, 119
 primary, 110
Carpodacus mexicanus, 277
 carrying capacity, 144, 167, 209, 230,
 295
Catostomus
 catostomus, 105
 commersoni, 105
 cellular automata, 335
Cenococcum
 geophilum, 130
 graniforme, 130
Centrostephanus rogersii, 340
Cervus elaphus, 87
 cestode
 Hymenolepididae, 103
 see *Schistocephalus solidus*
Ch, *see* exploitation, coherent, hard
Chamaelinorops, 37
 chemostat, 360
Chlorella, 133
Chrysolampis mosquitos, 60
Chusquea, 121
 cicada, 292
 see *Abrieta curvicauda*, *Psaltoda*
 moerens, *Henicopsaltria eydouxi*,
 Thopha saccata, *Cystosoma*
 saundersii
 cina, *see* *Stenocereus alamosensis*
Circus cyaneus, 73
 cisco, *see* *Coregonus artedii*
Cittotaenia, 103
 cladistics, 89, 98, 103
 cladocera, 124
Clerodendrum, 98
Clinus spatulatus, 61, 74, 76
Clitopilus prunulus, 130
 clone, 59, 139
 coenocline, 269
 distinct preference, 220
 shared preference, 221
Coereba flaveola, 60, 69
 coevolution, 94
 bilateral, 318
 coexistence, 229

- ecological versatility and, 261–3
- probability of, 262
- Colluricincla harmonica*, 9–10, 265, 273
- Colubridae, 37
- community
 - fungal, 16
 - hammock, 56
 - microbial, 16
 - plant, 16
- competition
 - coefficient, 230–1, 374; equivalence with resource overlap, 222
 - exploitation, 169, 171, 374–5
 - interference, 171
 - interspecific, 15, 27, 213–64, 267, 316, 327–8; actuality of, 214; combining ability, 223; compression, 213, 280–1, 322; context-specificity, 251–3; depressive, 214, 231; ecological versatility and, 220–4; exclusion, 213, 227; intransitivity, 223; manifestations of, 214; matrix, 216; release, 213, 280–1
 - intraspecific, 230, 316, 326
- competitiveness, 236, 242–3
- complementarity, 302
- component
 - environmental, 20
- conformance, 40–1
- Conomyrma*
 - bicolor*, 148
 - insana*, 148
- conservation biology, 203
- constraint, 25
- Contopus latirostris*, 280
- Conus*, 62
 - chaldeus*, 74
- copepod, 76, 106
 - see *Senecella calanoides*, *Epischura*
- Coregonus*
 - artedii*, 105
 - clupeiformis*, 105
- Cortinarius*
 - glaucopus*, 130
 - mucosus*, 130
- cosmology, 1
- cosmopolitanism, 9, 35, 267
- Costa Rica, 56, 96, 121
- Crematogaster emeryana*, 148
- Cs, see exploitation, coherent, soft cue, environmental, 151
- curve, indifference, 23–4
- cyanobacteria, 133
- Cyanophaia bicolor*, 280
- Cyphoderia ampulla*, 36
- Cystosoma saundersii*, 265, 293
- Dacelo novaeguineae*, 265
- Daphnia pulex*, 158
- Darlingtonia haetiana*, 37–8
- DCA, see analysis, detrended correspondence
- Dendroica*
 - castanea*, 118
 - pensylvanica*, 118
 - petechia*, 281
 - plumbea*, 281
- density, of consumer populations, 330
- Deschampsia flexuosa*, 178
- desert
 - Great Basin, 148
 - Mojave, 148
- design, sampling, 329–30, 332–5
- deterrent, chemical, 55, 340
- detoxification, 136, 317
- development, continuous, 159
- Diacyclops*, 125
- Diaphanosoma*, 125
- Diaptomus*, 125
- Dicaeum hirundinaceum*, 301
- digestibility, 87–8
- Dionaea muscipula*, 119
- discrimination, sensory, 323
- disease, 218
- dispersal, 204, 218
 - agents, 49
- displacement, character, 215, 327
- distance, Euclidean, 270
- distribution
 - ideal free, 282–3
 - isotropic resource, 372–3
- disturbance, abiotic, 218
- Dolabella auricularia*, 23, 99

426 · Index

- dormancy, 177, 195
 DRM, *see* model, digestive rate
Drosophila, 52–3, 60, 139, 318
 aldrichi, 53
 arizonensis, 53
 Hawai'ian, 318, 337
 mojavensis, 53
 drought, 54
- earthworm, 59
 echolocation, 120
 ecology, theoretical, 207
 ecophysiology, 266–7
 effect, Allee, 372
 efficiency, digestive, 88
Elaenia martinica, 280
 elaenia, Caribbean, *see* *Elaenia martinica*
 electivity, Ivlev index, 60, 70
Eleutherodactylus, 37–8
 elk, *see* *Cervus elaphus*
Embiotoca lateralis, 146–7, 165
 emigration, 205
Endogone lactiflua, 130
Enhydra lutris, 340
 enslavement, 146, 148, 313
 environment
 predictable versus stochastic, 246–7,
 310, 321
 test, constant, 180–1; stochastic,
 247–8, 254; variable, 180–1
 variability, 245–51, 310–11
Epibolium glandulosum, 131
Epischura, 124
 equilibrium, *see* stability, asymptotic
 Ericales, 130
Esox lucius, 105
 ETE, *see* extinction, expected time to
 ethology, 279
Eucalyptus, 131, 287
 calophylla, 131
 camaldulensis, 9, 131, 273, 287
 dalrympleana, 131
 diversicola, 131
 maculata, 131
 marginata, 131
 microcarpa, 9, 273, 287
 obliqua, 131
 pryoriana, 11, 287
 regnans, 131
 sideroxylon, 288
 sierberi, 131
 st johnii, 131
 tricarpa, 273, 287, 288
 euryphagy, 89
 evolutionary
 biology, 12, 279
 radiation, 12
 EWMA, *see* moving average,
 exponentially weighted
 experiment
 ecological field, 214, 233, 342
 optimal design, 225
 exploitation
 coherent, 162, 179
 hard, 182–7; hard, algorithm, 363–4;
 soft, 182–7; soft, algorithm, 364–5
 efficiency of, 202
 facultative, 142–6, 179
 hard, 175–8
 incoherent, 163, 179
 resource-like, 162, 164, 179; hard,
 187–94; hard, algorithm, 365–7;
 soft, 187–94; soft, algorithm,
 367–8
 soft, 175–8
 specialization, 163, 179; hard, 194–
 201; hard, algorithm, 368–9; soft,
 194–201; soft, algorithm, 369–71
 extinction, 218
 expected time to, 19
 stochastic, 372
- factor, exogenous, 43, 136
Falcunculus frontatus, 304–5
 fantail
 see *Rhipidura*
 grey, *see* *Rhipidura fuliginosa*
 fasting, 177
 feasibility, 216
 fecundity, 51, 146
 finch
 Cocos, *see* *Pinaroloxias inornata*
 Darwin's, *see* *Geospiza*
 Emberizidae xviii, 302

- ground *see* *Geospiza*, small, *see* *Geospiza fuliginosa*
- house, *see* *Carpodacus mexicanus*
- fish, 54, 139
- estuarine, 61
- piscivorous, 157, 160
- planktivorous, 157–8
- fitness, 6–7, 19–20, 333–4
- FIVAR, *see* design, sampling
- fixation
- phenotypic, 151
- floaters, 173
- fluke, blood, *see* *Schistosoma japonicum*
- flux, energy and nutrient, 205–6
- flycatcher
- brown-crested, *see* *Myiarchus tyrannulus (oberi)*
- neotropical (Tyrannidae), 56
- flytrap, Venus, *see* *Dionaea muscipula*
- food, availability
- between-year variation in, 278–9
- seasonal variation in, 276–8
- food web, 29
- theory, 339–40
- forest
- broad-leaf, 118
- coniferous, 118
- deciduous, 68
- montane, rain, 280
- Fragaria*, 131
- chiloensis*, 131
- frog, *see* amphibian, anuran; tadpole
- Gabon, M'Passa Biological Station, 250
- Galeolaria caespitosa*, 194–5
- Galerucella lineola*, 94, 97
- Galium aparine*, 131
- Gasterosteus aculeatus*, 72–3, 106
- gastropod, 57, 340
- Gelidium robustum*, 146–7
- gene flow, 28, 152
- generalization
- costs of, 151
- dietary, 89
- obligatory, 143, 179, 239
- genetic control
- epistatic, 148
- pleiotropic, 148
- genetic diversity, ecological versatility and, 310, 318
- genetics, population, 152
- Geospiza*, 66, 73
- fuliginosa*, 73
- Geum, 131
- germination, 52
- Gerygone*, 288
- gerygone*, *see* *Gerygone*
- Gilchristella aestuarina*, 61, 74, 76
- Glomus*
- caledonius*, 131
- fasciculatus*, 131
- macrocarpus*, var. *geosporus*, 131; var. *macrocarpus*, 131
- monosporus*, 131
- mossae*, 131
- glyph, *see* parameter space, span
- grain, 87
- grass
- silica content of 99
- see* *Agropyron repens*, *Poa pratensis*, *Phleum pratense*
- grassland, 275, 278
- Great Britain, 96, 178, 275
- Kent, 294
- Greya politella*, 129
- gum
- Gippsland manna, *see* *Eucalyptus pryoriana*
- river red, *see* *Eucalyptus camaldulensis*
- Gyrodon meruloides*, 130
- Gyroporus castaneus*, 130
- habitat, 17, 26, 64
- floristics, 35, 270, 286–9
- fragmentation, 203
- patch, 26
- physiognomy, 35, 270, 286–9
- selection, 4, 279; animals, 282–9; density-dependent model, 282–4; plants, 289–90
- sink, 203
- source, 203
- source-sink model 284–6

428 · Index

- habitat (*cont.*)
 stability, 311
 unuseable, 203
 xeric, 22
- hare, snowshoe, *see* *Lepus americanus*
- harrier, northern, *see* *Circus cyaneus*
- hawk, 58
- Heliopsis*, 90
- Henicopsaltria eydouxi*, 265
- herbivory, 324
 generalized, 83–8
 phytophagous insects, 88–99
 small noninsect, 99–100
- hierarchy, social, 146
- Hippodamia convergens*, 93
- honeyeater, 172
 fuscous, *see* *Lichenostomus fuscus*
 regent, *see* *Xanthomyza phrygia*
 white-plumed, *see* *Lichenostomus penicillatus*
 yellow-tufted, *see* *Lichenostomus melanops*
- Hoplosoma*, 98
- host range
 phytophagous insects, 88–100
 taxonomic breadth, 90
- Howardula aaronymphium*, 139
- hummingbird, 68, 176
 Antillean crested, *see* *Orthorhynchus cristatus*
 blue-headed, *see* *Cyanophaea bicolor*
 copper-rumped, *see* *Amazilia tobacis*
 ruby-topaz, *see* *Chrysolampis mosquitus*
- Hutchinsonian series, 215, 327
- idiosyncrasy, 336–9
- immigration, lag, 229, 243–5
- insect, 331
 holometabolous, 158
- interactions
 interspecific, caveats, 80–1
 strength of, 127
- interspecific competition models, *see*
 model, competition, interspecific
- invariance
 ecological, 295
- invasibility, 3, 155, 216–19, 233, 243–5
- invasion, high density, 185–6, 218, 243
- Ipomopsis aggregata*, 69
- Iridomyrmex humilis*, 93
- ironbark
 red, *see* *Eucalyptus tricarpa*
 mugga, *see* *Eucalyptus sideroxylon*
- Islands
 Galapagos 48, 66, 73, Rabida, 73
 Greater Antilles, 338
 Hawaiian, 48, 318
 Hispaniola, 37–8
 Lesser Antilles, 280, 337
 Montserrat, 281
 Santa Cruz, 146
 St Kitts, 281
 Tobago, 60, 67
 Trinidad, 60, 67
 West Indies, 37
- Israel, 285
- Juniperus communis*, var. *siberica*, 131
- kelp, 49, 340
- kingbird, grey, *see* *Tyrannus domenicensis*
- kleptoparasitism, 116
- kookaburra, laughing, *see* *Dacelo novaeguineae*
- kudu, *see* *Tragelaphus strepsiceros*
- lacewing, 110
- Lactarius*
chrysorrheus, 130
deliciosus, 130
helvus, 130
- Lake, Cold (Canada), 105
- Lamia, 98
- landscape, 35, 294
 fractal, 335
- Larix*
decidula, 131
occidentalis, 131
- Lasius*, 128
- latex, 89, 337
- Lavandulaatifolia*, 129
- leaf trenching, 89, 337

- legumes, 133
 - root nodulation, 130
- Lepomis macrochirus*, 157, 165
- Leptoglossus phyllopus*, 90
- Lepus americanus*, 86
- Lichenostomus*
 - fuscus*, 287
 - melanops*, 287, 304–5
 - penicillatus*, 11, 287
- lichen, 133
- life cycle, complex, 2, 106, 156
- limit cycle, 217
 - two-point, 182
- limiting factor, 20, 25, 253
- limpet, 54
- Lithophragma parviflorum*, 129
- lizard, 38, 54, 157
 - see Anolis*
- localized, 9
- Lota lota*, 105
- LVM, *see* model, Lotka–Volterra
- Lycopersicum esculentum*, 131
- Lyophyllum immundum*, 130

- macroecology, 1
- macrohabitat, 17, 26
- Maiathemum dilatatum*, 131
- maintenance, 5
- malentity, 20, 25
- mango, black-throated, *see*
 - Anthrocothorax nigricollis*
- Manorina*
 - melanocephala*, 172
 - melanophrys*, 172, 305
- MDS, *see* scaling, multidimensional
- Megarcys signata*, 137
- Meliphagidae, *see* honeyeater
- metamorphosis
 - disjunctive, 157–9
 - nondisjunctive, 157–9
- metapopulation, 1, 12, 203–7
- Metechinorynchus salmonis*, 104–5
- metric, Canberra dissimilarity, 70
- México, 52
- microhabitat, 17, 26–7, 63–4
- Micronycteris*
 - brachyotis*, 57
 - hirsuta*, 57
 - megalotis*, 57
- Microtus ochrogaster*, 99
- migration, 214
- Mimom crenulatum*, 57
- miner
 - bell, *see* *Manorina melanophrys*
 - noisy, *see* *Manorina melanocephala*
- Mischocyttarus flavitarsus*, 93
- mistletoe, 109, 301, 313
- mistletoebird, *see* *Dicaeum*
 - hirundinaceum*
- mite
 - parasitic, 103–4
 - see* *Unionicola*
- model
 - algorithmic, 168–9
 - apparent size, 157–8
 - community assembly, 217
 - competition, interspecific, 219–20
 - contingency, *see* predation,
 - foraging, optimal
 - core/satellite, 300–1
 - digestive rate, 87–8
 - dominance, 281–2
 - ecological, 207–9, 335–6
 - logistic, 167, 185, 208–9
 - Lotka–Volterra, 219–20, 224,
 - 229–30, 233, 242, 245, 260, 335, 374
 - mathematical, 167
 - nonlinearities, as resources, 246–7
 - partial niche, 142–3
 - patch, 113–15
 - phenomenological, 167
 - prey, 113–15
 - standard conditions, 228
 - statistical, linear, 342
 - ubiquity, algorithm, site-selection,
 - 290–3; contingency, 293–300;
 - fractal landscape, 297–300;
 - hierarchical, 296–7; marginal
 - mosaic, 294–6
 - unpacking of, 18, 260
 - versatility, assumptions of, 371–5;
 - implementation, 375
- Molossus*, 119

430 · Index

- Moniezia*, 103
 monomorphism, 152, 316
 monophagy, 89, 91
 ecological, 89
 monophenism, 151
 moose, *see* *Alces alces*
 morphology, host-specific, 325
Morula marginalba, 194–5
 moth
 Noctuidae, 89
 Notodontidae, 98, 337
 Sphingidae, 89
 Mountains, Andes, 121
 mouse
 deer, *see* *Peromyscus maniculatus*
 pinyon, *see* *Peromyscus truei*
 moving average, exponentially
 weighted, 361–2
 mutation, 153
 mutualism, 126–34, 326
 diffuse, 127
 obligatory bilateral, 128
 outcome variability, 129
 mutualistic associations, 129–33
 mycorrhiza, 130–3
Myiarchus tyrannulus (oberi), 281
Myrmecocystus kennedyi, 148
Mysis relicta, 123–5

 natural enemies, 91, 93
 nectar, 87
 neighbourhood, ecological, 27
 nematode, parasitic, *see* *Howardula aoronymphium*
 niche
 breadth xvii, 4, 8, 40, between-phenotype, 141, 155; statistic, 32–4, 355–7; statistic, conformance, 34, 188–9, 355–7; statistic, nonconformance, 34, 355–7; within-phenotype, 141, 155
 ontogenetic, 141, 156
 overlap, statistic, 32–4, 355–7
 partial, 142–3, 319
 pattern, 269–76
 position, 269, 298
 regeneration, 224

 width, *see* niche, breadth
 numbers, tyranny of small, 339–41
 nutrient, 20
 balancing, 322
Nyctinomops, 119

 Ocean, Pacific, 340
 Oceania, 48
Octolasion tyrtaeum, 59
 OFT, *see*, predation, foraging, optimal
 oligophagy, 55, 89
 omnivory, 122–6
 common-sense or ecological view, 123
Oncorhynchus kisutch, 105
 opportunism, 38
Opuntia willcoxii, 52–3
 organ pipe, *see* *Stenocereus thurberi*
Orthorhynchus cristatus, 280

Pachycephala pectoralis, 273
 Panama, 56, 121
 paradigm
 MacArthurian, 327
 Popperian scientific, 341
 parameter space
 glyph, 226–8
 span, 226–8
 parasite, 214–15, 325
 animal, 102–8, required host (definitive), 105; required host (intermediate), 105; suitable host, 105; unsuitable host, 105
 plant, 108–9
 parasitoid, 101–2, 324–5
 Diptera, Tachinidae, 102, 337
 idiobionts, 101, 313
 koinobionts, 101, 313
 pardalote, striated *see* *Pardalotus striatus*
Pardalotus striatus, 304–5
 particle physics, 1
Paruroctonus mesaensis, 159
Patelloida latistrigata, 194–5
 PCA, *see* analysis, principal component
Penstemon barbatus, 69
 permanence, 3, 219
Peromyscus, 60

- maniculatus*, 66–7, 288
- truei*, 66–7
- persistence, 19
- Peru, Manu National Park, 250
- Petroica*, 288
 - goodenovii*, 265, 273, 287
- pewee, Lesser Antillean, *see Contopus latirostris*
- phenotype, ecological, 146–8, 179, 190
- Philaenus spumarius*, 74, 76
- Phleum pratense*, 131, 231
- phoebe, Say's, *see Sayornis saya*
- phosphorus
 - absorption rate, 178
 - soil concentration, 298
- Phratora vitellinae*, 94, 97
- Phyllobrotica*, 98
- Phyllostomus*
 - discolor*, 57
 - hastatus*, 57
- physiology, thermal, 310, 320
- Physostegia*, 98
- Picea*
 - abies*, 131
 - sichensis*, 131
- pike, northern, *see Esox lucius*
- Pinaroloxias inornata*, 120, 137, 139–40
- pine, ponderosa, *see Pinus ponderosa*
- Pinguicula*, 119
- Pinus*, 131
 - contorta*, 131
 - echinata*, 131
 - monticola*, 131
 - mugo*, 131
 - ponderosa*, 100, 131
 - radiata*, 131
 - strobus*, 130, 131
 - sylvestris*, 130, 131
 - taeda*, 131
 - virginiana*, 131
- pitcher plant, *see Sarracenia*
- Placida dendrittica*, 55, 56, 137, 139–40
- plankton, 157–8
 - phytoplankton, 123–5
- plant
 - calcification, 99
 - chemistry, 100
 - suitability, 324
- Platycercus*
 - eximius* 9–10
 - flaveolus*, 265, 273, 304–5
- platyhelminthes, Monogenea, 103
- Plebejus argus*, 128
- Pleistocene, 266
- Poa pratensis*, 231
- pollen, 87
- pollinator, 49
- polymorphism, 152–5, 316
 - allelic-switch, 152
 - behavioural, 146–8
 - ecological, 146–8
 - protected, 153, 316
 - stable, 152–3, 190–3
- Polymorphus minutus*, 103
- polyphagy, 55, 89
- polyphenism, 316
 - adaptive, 148–52, 263
- Pontoporeia affinis*, 105
- population
 - differentiated, 326–7
 - distinctly staged, 58
 - migrant, 322–3
 - resident, 322–3
 - size-stepped, 58
 - stage-structured, 156–61
- predation, 109–22, 215, 325–6
 - behaviour, 118–21, 314
 - biophysics, 118–21, 312
 - effect on versatility, 110–12
 - foraging, optimal, 87, 112–18, 157, 315
 - intraguild, 54, 122
 - reciprocal, 122
- prickly pear, *see Opuntia willcoxii*
- principle, MacArthur's minimization, 239
- priority effect, 243–5
- problem
 - juvenile bottleneck, 160–1
 - N* populations–*N* resources, 253–9
- process, Wiener, 247
- prokaryote, 129
- Psaltoda moerens*, 265
- Psammogobius krysnaensis*, 61

432 · Index

- Pseudosuga menziesii*, 131
Psophodes olivaceus, 11
 Puerto Rico, 74
Pungitius pungitius, 72–3, 105
- Qs*, *see* *specialization, potential*
- range
 geographic, 4, 9, 267–8, centre, 294–5; periphery, 294–5
- rarity, 275
- recolonization, 218
- recombination, genetic, 154, 263
- recruitment, 284, 327
- refuge, 117, 123
- region, 35, 293
- reproduction, 5
- resin, 89
- resource
 abundance versus availability, 31
 aliasing, 29–30, 32, 49, 124
 availability, 7–8, 160, 314; formula for 360; memory, 361–3; perceived, 361–3; threshold, 83–4; variation in, 204
- complementary, 23–5, 180, 311, 322
 critical, 22
 definition, 20–5
 distribution, clumped, 311, 322
 exclusive access to, 21
 exploitation, 5, 19
 extraneous, 22
 ordering, 29–30
 partitioning, 213, 239
 substitutable, 23–5, 180, 311, 322
 utilization, 5, 19; abstention, 203; antagonism, 309; efficiency of, 223, 310; malleability of, 223; prudence, 242; stage-specific, 263
- response
 functional, 170, 173–5
 mixed, 173–5
 numerical, 170, 173–5
- Rh**, *see* *exploitation, resource-like, hard*
- Rhipidura*, 288
fuliginosa, 273
- Rhodophyllus rhodopolius*, 130
- rhizobia, 133
- Rhyzopogon*
 luteolus, 130
 roseolus, 130
- risk, predation, 116–18
- robin
 red-capped, *see* *Petroica goodenovii*
 see *Petroica*
- rodent, 81
- rosella
 eastern, *see* *Platycercus eximius*
 yellow, *see* *Platycercus flaveolus*
- rotifer, 124
- Rs**, *see* *exploitation, resource-like, soft*
- Rubus spectabilis*, 131
- ruminant, 86
- Russula*
 emetica, 130
 lepida, 130
- salinity, 54, 267
- Salix*
 dasyclados, 94, 97
 fragilis, 94, 97
 viminialis, 94, 97
- salmon, coho, *see* *Oncorhynchus kisutch*
- Salvelinus namaycush*, 105
- Sarracenia*, 119
- satellite, 173
- savanna, 275
- Sayornis saya*, 277
- scale
 ecological, 17
 spatial, 26–9, 274–5
- scaling, multidimensional, 271
- Scaphiopus multiplicatus*, 149, 165
- Schistocephalus solidus*, 106
- Schistosoma japonicum*, 106
- Sciurus aberti*, 99
- Scleroderma aurantium*, 130
- scorpion, *see* *Paruroctonus mesaensis*
- Scutellaria*, 98
- Sea, Caribbean, 48, 60, 267, 327
- sea
 hare, *see* *Dolabella auricularia*
 otter, *see* *Enhydra lutris*

- slug, *see* *Placida dendritica*
 urchin, 340
Senecella calanoides, 124–5
 sequestering, deterrent chemicals, 100,
 136, 315
Sh, *see* exploitation, specialized, hard
 shrew, 176
 shrike-thrush, grey, *see* *Colluricincla*
harmonica
 shrike-tit, crested, *see* *Falcunculus*
frontatus
 shrimp
 fairy, *see* *Anostraca*
 opossum, *see* *Mysis relicta*
 similarity, limiting, 230, 233
 simulation, computer, 168–9
 snake, 37, 320
 water, 157
Solidago, 74–6
 space, ordination, 271
 Spain, 67, 129, 322–3
 sparrow, 278
 western chipping, *see* *Spizella*
 passerina arizonae
 specialization
 dietary, 37–8, 322–3
 ecological, 53
 facultative, 143–5
 obligatory, 143, 320
 physiological, 53
 potential, 196–201
 serial, 159
 stage-specific, 160
 speciation, 27
 specificity
 ecological, 103, 106
 physiological, 106
Sphaerodactylus, 37
Spizella passerina arizonae, 266
 squirrel, Abert's, *see* *Sciurus aberti*
Ss, *see* exploitation, specialized, soft
 stability, asymptotic, 3, 216–19
Stachys, 98
 mexicana, 131
 stagnation, 267
 statistic
 Czekanowski, 189, 359–60
 diagnostic, 358–60
 Hurlbert L, 70
 inverse Simpson diversity, 69
 Levins, 71
 Petraitis W, 70
 proportional similarity, 70, 188, 359
 Shannon–Wiener, 70
Stenocereus
 alamosensis, 52
 gummosus, 52
 thurberi, 52
 stenophagy, 37–8, 88–9, 96, 98, 315
 stickleback, 64, 79
 ninespine, *see* *Pungitius pungitius*
 three-spined, *see* *Gasterosteus*
 aculeatus
Stipa capensis, 285
Stizostedion vitreum, 105
 stonefly, *see* *Megarcys signata*
 strategy
 exploitation, 161–4, hybrid, 262–3;
 tactic, 225, 261
 facultative, 112
 homologous, 182
 sucker
 longnose, *see* *Catostomus catostomus*
 white, *see* *Catostomus commersoni*
Suillus
 bovinus, 130
 flavidus, 130
 flavovirens, 130
 granulatus, 130
 luteus variegatus, 130
 sunfish, bluegill, *see* *Lepomis macrochirus*
 supplementation, 302
 surfperch, striped, *see* *Embiotoca lateralis*
 survey
 climatic zones, 47
 conformance, 59–62
 constraint, 54–5
 differentiated populations, 58–9
 fitness, 51–4
 geographic zones, 47–8
 habitat types, 46–7
 individuals, 55–8
 niche, breadth statistics, 69–70;
 overlap statistics, 69–70

434 · Index

- survey (*cont.*)
 niche statistics, list of, 354–7
 parameters, 41–3
 resource, availability, 59–62
 resource specification, 49–50
 selection criteria, 43–5
 sources, 43–5
 spatial variability, 62–5
 studies, list of, 351–4
 taxonomy, 45–6
 temporal variability, diurnal, 68–9;
 season, 65–8
- survival, 5
 Sweden, 94
Symbiodinium, 133
 symbiosis, 126–7, 312
Syngnathus acus, 61
- tadpole, 149–50
Taxus brevifolius, 131
 temperature, water, 54
 territoriality, 171–2, 279
Tesseropora rosea, 194–5
Thopha saccata, 265
Thuja plicata, 131
 thornbill, *see* *Acanthiza*
 throughput, 86–8
 thyroxine, 149–50
 toad, southern spadefoot, *see*
Scaphiopus multiplicatus
- Tonatia*
bidens, 57
sylvicola, 57
Trachops cirrhosus, 57
Tragelaphus strepsiceros, 85
 transect, Victorian, 272, 287, 292
 trematode, 103–5
Tricholoma
flavobrunneum, 130
flavovirens, 130
imbricatum, 130
pessundatum, 130
saponaceum, 130
vaccinium, 130
Trifolium repens, 131
Trillium ovatum, 131
Tripneustes gratilla, 340
Triticum, 131
aestivum, 131
 trophic continuum, 81, 125
 Tropicodphiidae, 37
 trout, lake, *see* *Salvelinus namaycush*
- Tuber*
albidum, 130
maculatum, 130
Turbo undulata, 340
 turtle, 157
Tyrannus domenicensis, 281
- ubiquity, 8–11, 18, 35
 abiotic effects on, 268
 ecological versatility and, 301–4
 ungulate, 81
 African, 266
Unionicola, 106
 United States, 60, 119
 Arizona, 68, 148, 266, 278
 California, 146, 148
 Colorado, 337
 Florida, 56, 103
 Hawai'i, 337
 Idaho, 60
 Illinois, 60, 69
 North Carolina, 59
 north-eastern, 275
 Tennessee, 59
 Washington, 129
Ursus arctos, 372
- utilization
 pressure, 170–3, 231–2
 rate, marginal, 175; maximum, 171,
 175
- vector, 49
 versatility
 dietary, 87
 ecological, 5–8, 11, 19–20, and
 ubiquity, *see* ubiquity, ecological
 versatility and; density and,
 318–20
 foraging, 22, 303
 viability floor, 371–2
 vole, prairie, *see* *Microtus ochrogaster*
 voracity, 170

- walleye, *see Stizostedion vitreum*
 warbler
 bay-breasted, *see Dendroica castanea*
 chestnut-sided, *see Dendroica pennsylvanica*
 plumbeous, *see Dendroica plumbea*
 wood (Parulidae) xviii, 118–19, 214, 302
 yellow, *see Dendroica petechia*
 wave action, 54
 whipbird, eastern, *see Psophodes olivaceus*
 whistler, golden, *see Pachycephala pectoralis*
 whitefish, lake, *see Coregonus clupeaformis*
 willow, *see Salix fragilis*, *Salix dasyclados*, *Salix viminalis*
 woodland, 272
 evergreen oak, 67
 riparian, 287
Xanthomyza phrygia, 287
Zea mays, 131
Zerna erecta, 178