

## LONDON MATHEMATICAL SOCIETY LECTURE NOTE SERIES

Managing Editor: Professor M. Reid, Mathematics Institute, University of Warwick, Coventry CV4 7AL, United Kingdom

The titles below are available from booksellers, or from Cambridge University Press at <http://www.cambridge.org/mathematics>

- 287 Topics on Riemann surfaces and Fuchsian groups, E. BUJALANCE, A.F. COSTA & E. MARTÍNEZ (eds)
- 288 Surveys in combinatorics, 2001, J.W.P. HIRSCHFELD (ed)
- 289 Aspects of Sobolev-type inequalities, L. SALOFF-COSTE
- 290 Quantum groups and Lie theory, A. PRESSLEY (ed)
- 291 Tits buildings and the model theory of groups, K. TENT (ed)
- 292 A quantum groups primer, S. MAJID
- 293 Second order partial differential equations in Hilbert spaces, G. DA PRATO & J. ZABCZYK
- 294 Introduction to operator space theory, G. PISIER
- 295 Geometry and integrability, L. MASON & Y. NUTKU (eds)
- 296 Lectures on invariant theory, I. DOLGACHEV
- 297 The homotopy category of simply connected 4-manifolds, H.-J. BAUES
- 298 Higher operads, higher categories, T. LEINSTER (ed)
- 299 Kleinian groups and hyperbolic 3-manifolds, Y. KOMORI, V. MARKOVIC & C. SERIES (eds)
- 300 Introduction to Möbius differential geometry, U. HERTRICH-JEROMIN
- 301 Stable modules and the D(2)-problem, F.E.A. JOHNSON
- 302 Discrete and continuous nonlinear Schrödinger systems, M.J. ABLOWITZ, B. PRINARI & A.D. TRUBATCH
- 303 Number theory and algebraic geometry, M. REID & A. SKOROBOGATOV (eds)
- 304 Groups St Andrews 2001 in Oxford I, C.M. CAMPBELL, E.F. ROBERTSON & G.C. SMITH (eds)
- 305 Groups St Andrews 2001 in Oxford II, C.M. CAMPBELL, E.F. ROBERTSON & G.C. SMITH (eds)
- 306 Geometric mechanics and symmetry, J. MONTALDI & T. RATIU (eds)
- 307 Surveys in combinatorics 2003, C.D. WENSLEY (ed.)
- 308 Topology, geometry and quantum field theory, U.L. TILLMANN (ed)
- 309 Corings and comodules, T. BRZEZINSKI & R. WISBAUER
- 310 Topics in dynamics and ergodic theory, S. BEZUGLYI & S. KOLYADA (eds)
- 311 Groups: topological, combinatorial and arithmetic aspects, T.W. MÜLLER (ed)
- 312 Foundations of computational mathematics, Minneapolis 2002, F. CUCKER *et al* (eds)
- 313 Transcendental aspects of algebraic cycles, S. MÜLLER-STACH & C. PETERS (eds)
- 314 Spectral generalizations of line graphs, D. CVETKOVIĆ, P. ROWLINSON & S. SIMIĆ
- 315 Structured ring spectra, A. BAKER & B. RICHTER (eds)
- 316 Linear logic in computer science, T. EHRHARD, P. RUET, J.-Y. GIRARD & P. SCOTT (eds)
- 317 Advances in elliptic curve cryptography, I.F. BLAKE, G. SEROUSSI & N.P. SMART (eds)
- 318 Perturbation of the boundary in boundary-value problems of partial differential equations, D. HENRY
- 319 Double affine Hecke algebras, I. CHEREDNIK
- 320 L-functions and Galois representations, D. BURNS, K. BUZZARD & J. NEKOVÁŘ (eds)
- 321 Surveys in modern mathematics, V. PRASOLOV & Y. ILYASHENKO (eds)
- 322 Recent perspectives in random matrix theory and number theory, F. MEZZADRI & N.C. SNAITH (eds)
- 323 Poisson geometry, deformation quantisation and group representations, S. GUTT *et al* (eds)
- 324 Singularities and computer algebra, C. LOSSEN & G. PFISTER (eds)
- 325 Lectures on the Ricci flow, P. TOPPING
- 326 Modular representations of finite groups of Lie type, J.E. HUMPHREYS
- 327 Surveys in combinatorics 2005, B.S. WEBB (ed)
- 328 Fundamentals of hyperbolic manifolds, R. CANARY, D. EPSTEIN & A. MARDEN (eds)
- 329 Spaces of Kleinian groups, Y. MINSKY, M. SAKUMA & C. SERIES (eds)
- 330 Noncommutative localization in algebra and topology, A. RANICKI (ed)
- 331 Foundations of computational mathematics, Santander 2005, L.M. PARDO, A. PINKUS, E. SÜLI & M.J. TODD (eds)
- 332 Handbook of tilting theory, L. ANGELERI HÜGEL, D. HAPPEL & H. KRAUSE (eds)
- 333 Synthetic differential geometry (2nd Edition), A. KOCK
- 334 The Navier–Stokes equations, N. RILEY & P. DRAZIN
- 335 Lectures on the combinatorics of free probability, A. NICA & R. SPEICHER
- 336 Integral closure of ideals, rings, and modules, I. SWANSON & C. HUNEKE
- 337 Methods in Banach space theory, J.M.F. CASTILLO & W.B. JOHNSON (eds)
- 338 Surveys in geometry and number theory, N. YOUNG (ed)
- 339 Groups St Andrews 2005 I, C.M. CAMPBELL, M.R. QUICK, E.F. ROBERTSON & G.C. SMITH (eds)
- 340 Groups St Andrews 2005 II, C.M. CAMPBELL, M.R. QUICK, E.F. ROBERTSON & G.C. SMITH (eds)
- 341 Ranks of elliptic curves and random matrix theory, J.B. CONREY, D.W. FARMER, F. MEZZADRI & N.C. SNAITH (eds)
- 342 Elliptic cohomology, H.R. MILLER & D.C. RAVENEL (eds)
- 343 Algebraic cycles and motives I, J. NAGEL & C. PETERS (eds)
- 344 Algebraic cycles and motives II, J. NAGEL & C. PETERS (eds)
- 345 Algebraic and analytic geometry, A. NEEMAN
- 346 Surveys in combinatorics 2007, A. HILTON & J. TALBOT (eds)
- 347 Surveys in contemporary mathematics, N. YOUNG & Y. CHOI (eds)

- 348 Transcendental dynamics and complex analysis, P.J. RIPPON & G.M. STALLARD (eds)
- 349 Model theory with applications to algebra and analysis I, Z. CHATZIDAKIS, D. MACPHERSON, A. PILLAY & A. WILKIE (eds)
- 350 Model theory with applications to algebra and analysis II, Z. CHATZIDAKIS, D. MACPHERSON, A. PILLAY & A. WILKIE (eds)
- 351 Finite von Neumann algebras and masas, A.M. SINCLAIR & R.R. SMITH
- 352 Number theory and polynomials, J. MCKEE & C. SMYTH (eds)
- 353 Trends in stochastic analysis, J. BLATH, P. MÖRTERS & M. SCHEUTZOW (eds)
- 354 Groups and analysis, K. TENT (ed)
- 355 Non-equilibrium statistical mechanics and turbulence, J. CARDY, G. FALKOVICH & K. GAWEDZKI
- 356 Elliptic curves and big Galois representations, D. DELBOURGO
- 357 Algebraic theory of differential equations, M.A.H. MACCALLUM & A.V. MIKHAILOV (eds)
- 358 Geometric and cohomological methods in group theory, M.R. BRIDSON, P.H. KROPHOLLER & I.J. LEARY (eds)
- 359 Moduli spaces and vector bundles, L. BRAMBILA-PAZ, S.B. BRADLOW, O. GARCÍA-PRADA & S. RAMANAN (eds)
- 360 Zariski geometries, B. ZILBER
- 361 Words: Notes on verbal width in groups, D. SEGAL
- 362 Differential tensor algebras and their module categories, R. BAUTISTA, L. SALMERÓN & R. ZUAZUA
- 363 Foundations of computational mathematics, Hong Kong 2008, F. CUCKER, A. PINKUS & M.J. TODD (eds)
- 364 Partial differential equations and fluid mechanics, J.C. ROBINSON & J.L. RODRIGO (eds)
- 365 Surveys in combinatorics 2009, S. HUCZYNSKA, J.D. MITCHELL & C.M. RONEY-DOUGAL (eds)
- 366 Highly oscillatory problems, B. ENGQUIST, A. FOKAS, E. HAIRER & A. ISERLES (eds)
- 367 Random matrices: High dimensional phenomena, G. BLOWER
- 368 Geometry of Riemann surfaces, F.P. GARDINER, G. GONZÁLEZ-DIEZ & C. KOUROUNIOTIS (eds)
- 369 Epidemics and rumours in complex networks, M. DRAIEF & L. MASSOULIÉ
- 370 Theory of  $p$ -adic distributions, S. ALBEVERIO, A.YU. KHRENNIKOV & V.M. SHELKOVICH
- 371 Conformal fractals, F. PRZYTYCKI & M. URBAŃSKI
- 372 Moonshine: The first quarter century and beyond, J. LEPOWSKY, J. MCKAY & M.P. TUIITE (eds)
- 373 Smoothness, regularity and complete intersection, J. MAJADAS & A. G. RODICIO
- 374 Geometric analysis of hyperbolic differential equations: An introduction, S. ALINHAC
- 375 Triangulated categories, T. HOLM, P. JØRGENSEN & R. ROUQUIER (eds)
- 376 Permutation patterns, S. LINTON, N. RUŠKUC & V. VATTER (eds)
- 377 An introduction to Galois cohomology and its applications, G. BERHUY
- 378 Probability and mathematical genetics, N. H. BINGHAM & C. M. GOLDIE (eds)
- 379 Finite and algorithmic model theory, J. ESPARZA, C. MICHAUX & C. STEINHORN (eds)
- 380 Real and complex singularities, M. MANOEL, M.C. ROMERO FUSTER & C.T.C. WALL (eds)
- 381 Symmetries and integrability of difference equations, D. LEVI, P. OLVER, Z. THOMOVA & P. WINTERNITZ (eds)
- 382 Forcing with random variables and proof complexity, J. KRAJÍČEK
- 383 Motivic integration and its interactions with model theory and non-Archimedean geometry I, R. CLUCKERS, J. NICAISE & J. SEBAG (eds)
- 384 Motivic integration and its interactions with model theory and non-Archimedean geometry II, R. CLUCKERS, J. NICAISE & J. SEBAG (eds)
- 385 Entropy of hidden Markov processes and connections to dynamical systems, B. MARCUS, K. PETERSEN & T. WEISSMAN (eds)
- 386 Independence-friendly logic, A.L. MANN, G. SANDU & M. SEVENSTER
- 387 Groups St Andrews 2009 in Bath I, C.M. CAMPBELL *et al* (eds)
- 388 Groups St Andrews 2009 in Bath II, C.M. CAMPBELL *et al* (eds)
- 389 Random fields on the sphere, D. MARINUCCI & G. PECCATI
- 390 Localization in periodic potentials, D.E. PELINOVSKY
- 391 Fusion systems in algebra and topology, M. ASCHBACHER, R. KESSAR & B. OLIVER
- 392 Surveys in combinatorics 2011, R. CHAPMAN (ed)
- 393 Non-abelian fundamental groups and Iwasawa theory, J. COATES *et al* (eds)
- 394 Variational problems in differential geometry, R. BIELAWSKI, K. HOUSTON & M. SPEIGHT (eds)
- 395 How groups grow, A. MANN
- 396 Arithmetic differential operators over the  $p$ -adic Integers, C.C. RALPH & S.R. SIMANCA
- 397 Hyperbolic geometry and applications in quantum Chaos and cosmology, J. BOLTE & F. STEINER (eds)
- 398 Mathematical models in contact mechanics, M. SOFONEA & A. MATEI
- 399 Circuit double cover of graphs, C.-Q. ZHANG
- 400 Dense sphere packings: a blueprint for formal proofs, T. HALES
- 401 A double Hall algebra approach to affine quantum Schur-Weyl theory, B. DENG, J. DU & Q. FU
- 402 Mathematical aspects of fluid mechanics, J. ROBINSON, J.L. RODRIGO & W. SADOWSKI (eds)
- 403 Foundations of computational mathematics: Budapest 2011, F. CUCKER, T. KRICK, A. SZANTO & A. PINKUS (eds)
- 404 Operator methods for boundary value problems, S. HASSI, H.S.V. DE SNOO & F.H. SZAFRANIEC (eds)
- 405 Torsors, étale homotopy and applications to rational points, A.N. SKOROBOGATOV (ed)
- 406 Appalchian set theory, J. CUMMINGS & E. SCHIMMERLING (eds)
- 407 The maximal subgroups of the low-dimensional finite classical groups, J.N. BRAY, D.F. HOLT & C.M. RONEY-DOUGAL

Cambridge University Press & Assessment

978-0-521-13860-4 — The Maximal Subgroups of the Low-Dimensional Finite Classical Groups

John N. Bray, Derek F. Holt, Colva M. Roney-Dougal

Frontmatter

[More Information](#)

London Mathematical Society Lecture Note Series: 407

# The Maximal Subgroups of the Low-Dimensional Finite Classical Groups

JOHN N. BRAY

*Queen Mary, University of London*

DEREK F. HOLT

*University of Warwick*

COLVA M. RONEY-DOUGAL

*University of St Andrews*



CAMBRIDGE  
UNIVERSITY PRESS

Cambridge University Press & Assessment  
978-0-521-13860-4 — The Maximal Subgroups of the Low-Dimensional Finite Classical Groups  
John N. Bray, Derek F. Holt, Colva M. Roney-Dougal  
Frontmatter  
[More Information](#)



CAMBRIDGE  
UNIVERSITY PRESS

Shaftesbury Road, Cambridge CB2 8EA, United Kingdom  
One Liberty Plaza, 20th Floor, New York, NY 10006, USA  
477 Williamstown Road, Port Melbourne, VIC 3207, Australia  
314–321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre, New Delhi – 110025, India  
103 Penang Road, #05–06/07, Visioncrest Commercial, Singapore 238467

Cambridge University Press is part of Cambridge University Press & Assessment,  
a department of the University of Cambridge.

We share the University's mission to contribute to society through the pursuit of  
education, learning and research at the highest international levels of excellence.

[www.cambridge.org](http://www.cambridge.org)  
Information on this title: [www.cambridge.org/9780521138604](http://www.cambridge.org/9780521138604)

© J.N. Bray, D.F. Holt and C.M. Roney-Dougal 2013

This publication is in copyright. Subject to statutory exception and to the provisions  
of relevant collective licensing agreements, no reproduction of any part may take  
place without the written permission of Cambridge University Press & Assessment.

First published 2013

*A catalogue record for this publication is available from the British Library*

ISBN 978-0-521-13860-4 Paperback

Additional resources for this publication at [www.cambridge.org/9780521138604](http://www.cambridge.org/9780521138604)

Cambridge University Press & Assessment has no responsibility for the persistence  
or accuracy of URLs for external or third-party internet websites referred to in this  
publication and does not guarantee that any content on such websites is, or will  
remain, accurate or appropriate.

## Contents

	<i>Foreword by Martin Liebeck</i>	<i>page</i> viii
	<i>Preface</i>	x
<b>1</b>	<b>Introduction</b>	1
	1.1 Background	1
	1.2 Notation	4
	1.3 Some basic group theory	6
	1.4 Finite fields and perfect fields	11
	1.5 Classical forms	13
	1.6 The classical groups and their orders	25
	1.7 Outer automorphisms of classical groups	33
	1.8 Representation theory	37
	1.9 Tensor products	43
	1.10 Small dimensions and exceptional isomorphisms	45
	1.11 Representations of simple groups	46
	1.12 The natural representations of the classical groups	48
	1.13 Some results from number theory	52
<b>2</b>	<b>The main theorem and the types of geometric subgroups</b>	55
	2.1 The main theorem	55
	2.2 Introducing the geometric types	59
	2.3 Preliminary arguments concerning maximality	76
<b>3</b>	<b>Geometric maximal subgroups</b>	106
	3.1 Dimension 2	106
	3.2 Dimension 3	110
	3.3 Dimension 4	115
	3.4 Dimension 5	121
	3.5 Dimension 6	122
	3.6 Dimension 7	126
	3.7 Dimension 8	127

3.8	Dimension 9	132
3.9	Dimension 10	135
3.10	Dimension 11	140
3.11	Dimension 12	141
<b>4</b>	<b>Groups in Class <math>\mathcal{S}</math>: cross characteristic</b>	<b>150</b>
4.1	Preamble	150
4.2	Irrationalities	155
4.3	Cross characteristic candidates	159
4.4	The type of the form and the stabilisers in $\Omega$ and $C$	167
4.5	Dimension up to 6: quasisimple and conformal groups	169
4.6	Determining the effects of duality and field automorphisms	187
4.7	Dimension up to 6: graph and field automorphisms	195
4.8	Dimension up to 6: containments	211
4.9	Dimensions greater than 6	220
4.10	Summary of the $\mathcal{S}_1$ -maximal subgroups	253
<b>5</b>	<b>Groups in Class <math>\mathcal{S}</math>: defining characteristic</b>	<b>267</b>
5.1	General theory of $\mathcal{S}_2$ -subgroups	268
5.2	Symmetric and anti-symmetric powers	276
5.3	The groups $\mathrm{SL}_2(q) = \mathrm{Sp}_2(q)$	280
5.4	The groups $\mathrm{SL}_n(q)$ and $\mathrm{SU}_n(q)$ for $n \geq 3$	289
5.5	The groups $\mathrm{Sp}_n(q)$	302
5.6	The groups $\mathrm{O}_n^\epsilon(q)$ , ${}^3\mathrm{D}_4(q)$ , and their covers	304
5.7	The remaining groups and their covers	305
5.8	Summary of $\mathcal{S}_2^*$ -candidates	307
5.9	Determining the effects of duality and field automorphisms	309
5.10	Containments	316
5.11	Summary of the $\mathcal{S}_2^*$ -maximals	318
<b>6</b>	<b>Containments involving <math>\mathcal{S}</math>-subgroups</b>	<b>322</b>
6.1	Introduction	322
6.2	Containments between $\mathcal{S}_1$ - and $\mathcal{S}_2^*$ -maximal subgroups	323
6.3	Containments between geometric and $\mathcal{S}^*$ -maximal subgroups	333
<b>7</b>	<b>Maximal subgroups of exceptional groups</b>	<b>357</b>
7.1	Introduction	357
7.2	The maximal subgroups of $\mathrm{Sp}_4(2^e)$ and extensions	358
7.3	The maximal subgroups of $\mathrm{Sz}(q)$ and extensions	365
7.4	The maximal subgroups of $\mathrm{G}_2(2^e)$ and extensions	368
<b>8</b>	<b>Tables</b>	<b>373</b>
8.1	Description of the tables	373

Cambridge University Press & Assessment

978-0-521-13860-4 — The Maximal Subgroups of the Low-Dimensional Finite Classical Groups

John N. Bray, Derek F. Holt, Colva M. Roney-Dougal

Frontmatter

[More Information](#)

---

	<i>Contents</i>	vii
8.2	The tables	377
	<i>References</i>	429
	<i>Index of Definitions</i>	435

## Foreword

In this book the authors determine the maximal subgroups of all the finite classical groups of dimension 12 or less. This work fills a long-standing gap in the literature. Behind this gap there is a story which I am pleased to have the opportunity to tell.

The completion of the classification of finite simple groups was first announced in the early 1980s. It was clear then (and before) that for many applications of the classification one would need detailed knowledge of the maximal subgroups of the simple groups and of their automorphism groups. Around that time, I gave a Part III course at Cambridge about the classification and its impact. Full of enthusiasm, I set a fearsome exam – I remember giving it to John Conway to check, and him saying that he couldn't do any of the questions, but he thought it was probably OK. The second highest mark was 18%, scored by a rather strong student. The top mark was 97%, scored by Peter Kleidman, a young American.

Soon afterwards, Kleidman started as my first research student. Michael Aschbacher had just published his fundamental theorem on maximal subgroups of the finite classical groups. The time seemed right to attempt to use this to determine all the maximal subgroups of the classical groups of low dimensions (up to 20, say, I thought optimistically). This was Kleidman's initial project. As it turned out, in his thesis he solved many other maximal subgroup problems, and this project occupied just one chapter. Nevertheless it was rather an interesting chapter, consisting of tables of all the maximal subgroups of finite simple classical groups of dimension up to 12. No proofs were given, just an outline of the strategy and a few examples of how the calculations were performed.

After he had graduated, Kleidman and I wrote a book on the subgroups of the finite classical groups, which was an analysis of the structure, conjugacy and maximality of the subgroups arising in Aschbacher's theorem now known as geometric subgroups. For the maximality questions we assumed that the dimen-



sion was more than 12, and it was the intention that Kleidman would extend and write up his thesis work on the low dimensions as a separate book. Indeed, this project was accepted as a volume to appear in the Longman Research Notes series, and has been referred to as such in many articles. Unfortunately, he did not write this book, and left mathematics at the age of about thirty to pursue other interests such as working on Wall Street and producing Hollywood movies.

The non-appearance of Kleidman's book left a yawning gap in the literature for over twenty years. We are fortunate indeed that a number of years ago the authors of this volume took it upon themselves to fill this gap. They have done this in marvellously complete fashion, presenting the material with great clarity and attention to detail. Full proofs and comprehensive background material are given, making the book easily accessible to graduate students. It should also be said that their results go quite a way beyond Kleidman's thesis, in that they handle almost simple classical groups rather than just simple ones, which is important for applications.

It is marvellous to have this volume on the bookshelf where previously there was such an evident space, and I congratulate the authors on their achievement.

Martin Liebeck  
Imperial College London

## Preface

The aim of this book is to classify the maximal subgroups of the almost simple finite classical groups of dimension at most 12. We also include tables describing the maximal subgroups of the almost simple finite exceptional groups that have faithful representations of degree at most 12.

A group  $G$  is *simple* if it has order greater than 1 and has no normal subgroups other than the trivial subgroup and  $G$  itself, and is *almost simple* if  $S \leq G \leq \text{Aut } S$  for some non-abelian simple group  $S$ . A group is *perfect* if it is equal to its derived group. A group  $G$  is called *quasisimple* if  $G$  is perfect and  $G$  modulo its centre is a non-abelian simple group.

The study and classification of the (maximal) subgroups of the finite simple groups and their variations has a long history, and the completion of the classification of the finite simple groups provided further motivation.

The term ‘classical group’ is used frequently in the literature, but it is rarely, if ever, defined precisely. We shall not attempt a formal definition here, and we shall avoid using it in a precise sense. Our general intention is to use it very inclusively. We shall certainly regard all of the named groups (like  $\text{GL}_n(q)$ ,  $\text{O}_n^\epsilon(q)$ ,  $\text{PCSp}_n(q)$ , etc.) in Table 1.2 as classical groups, but we also include among the classical groups arbitrary subgroups between the  $\Omega$ -groups and the  $A$ -groups in the table, and also quotients of the groups in the first of each of the paired rows in the table by arbitrary subgroups of the scalars. Furthermore, we include all almost simple extensions of the simple classical groups.

**Maximal subgroups of classical groups.** In [1], Aschbacher proved a fundamental theorem that describes the subgroups of almost all of the finite almost simple classical groups (the only exceptions are certain extensions of  $S_4(2^e)$  and  $\text{O}_8^+(q)$ ). This theorem divides these subgroups into nine classes. The first eight of these consist roughly of groups that preserve some kind of geometric structure; for example the first class consists (roughly) of the reducible groups, which fix a proper non-zero subspace of the vector space on which the group

acts naturally. Subgroups of classical groups that lie in the first eight classes are of *geometric type*. The ninth class, denoted by  $\mathcal{C}_9$  or  $\mathcal{S}$ , consists (roughly) of those subgroups that are not of geometric type and which, modulo the subgroup of scalar matrices, are almost simple. An alternative proof of Aschbacher's theorem, as a corollary to a version of the theorem for algebraic groups, can be found in [82]. We present a detailed version of Aschbacher's theorem, based on the treatment in [66], in Section 2.2. An interesting version of Aschbacher's theorem is presented in [91], which emphasises the links between the subgroup structure of the finite classical groups and of the algebraic groups of Lie type.

In [66], Kleidman and Liebeck provide an impressively detailed enumeration of the maximal subgroups of geometric type of the almost simple finite classical groups of dimension greater than 12. In this book, we shall extend the work of Kleidman and Liebeck to handle dimensions at most 12, and also classify the maximal subgroups of these groups that are in Class  $\mathcal{S}$ .

With the exception of Kleidman's work [62] on  $\Omega_8^+(q)$ , and classifications of Kleidman [62, 64, 63] and Cooperstein [14] of maximal subgroups of some of the exceptional groups of Lie type, which we simply cite and reproduce in our tables, our approach in this book has been to use these previous classifications for checking purposes only: our proofs make no use of the references below, although we have compared our results with them and, where there are differences, verified that our tables are correct.

The most complete previous work on low-dimensional classical groups is undoubtedly Peter Kleidman's PhD thesis [61], where he presents a classification, without proof, of the maximal subgroups of the simple classical groups in dimensions up to 12. This is a remarkable achievement. Kleidman intended to publish a subsequent book, with the same goal as ours: the classification of maximal subgroups of the almost simple classical groups in dimension up to 12. Unfortunately, this has not been published, and the present work is an attempt to carry out Kleidman's plan.

We base the following historical summary on the surveys by King [60], and Kleidman and Liebeck [65]. The complete description of the subgroup structure of the groups  $L_2(q)$  is usually attributed to Dickson [22], but this topic was also investigated by E. H. Moore [94] and Wiman [115]. The subgroups of  $L_3(q)$  were described by Mitchell for  $q$  odd [92] and then by Hartley for  $q$  even [40]. In both cases the subgroups that lie in  $U_3(\sqrt{q})$  were identified. A more modern treatment of subgroups of  $L_3(q)$  is provided by Bloom in [3]. The maximal subgroups of  $L_4(q)$  for even  $q$  were listed independently by Mwene [95] and Zalesskii [116]. A partial classification for odd  $q$  can be found in [96] and, independently, for  $p > 5$  in [117]. Further results on this case can be found in [59, Section 5]. Mitchell classified the maximal subgroups of  $S_4(q)$  for odd  $q$  in [93]. Flesner [26, 27] partially classified the maximal subgroups of  $S_4(2^e)$ .

The maximal subgroups of  $L_5(q)$  were determined by Di Martino and Wagner for  $q$  odd [23], and independently by Wagner [110] and Zalesskii [116] for  $q$  even. Kondrat'ev classified the quasisimple absolutely irreducible subgroups of  $GL_6(q)$  [71]. There is a brief survey of many of these results in [72].

For higher dimensions, many people have concentrated on the case  $q = 2$ . In 1984, Darafsheh classified the maximal subgroups of  $GL_6(2)$  [21], building on Harada and Yamaki's paper [39], which classified the insoluble irreducible subgroups of  $GL_n(2)$  with  $n \leq 6$ . The subgroups of  $GL_n(2)$  for  $n \leq 10$  were studied extensively by Kondrat'ev: see [67, 68, 69, 70].

In the work of King and others (see [60] and the references therein) a different approach is taken: rather than concentrating on a family of almost simple groups, such as those with socle  $L_6(q)$ , one concentrates on a family of potentially maximal subgroups, such as those of type  $\Gamma L_{n/2}(q^2)$  in  $SL_n(q)$ , and tries to determine maximality. A great deal is known in this direction: see the results cited in [60], together with [15, 16, 17, 99], amongst others. Again, we have not used these works in our proofs, but have compared our results with them: we mention them in the relevant sections of Chapters 2, 3 and 6 of this book.

An approach which is slightly orthogonal to our present purposes, but which has been used as a tool in the proof of many deep theorems, is to classify subgroups of classical groups containing elements of specified orders. We shall not use these results, so will not provide an extensive list of papers, but the interested reader could start by looking at [35, 36] and the references therein.

**Maximal subgroups of non-classical simple groups.** See [83, 114] for an excellent survey and introduction, respectively, on the whole of this field. The ATLAS [12] is also an essential reference in this area.

For the alternating groups, the O'Nan–Scott Theorem (see, for example, Chapter 4 of [8]) provides a subgroup classification similar to, but much simpler than, the Aschbacher classification of matrix groups over finite fields, and results of Liebeck, Praeger and Saxl [78] enable us to determine maximality. As in the case of finite matrix groups, we have a final class of almost simple primitive permutation groups which need to be listed individually by degree, and these lists are currently complete up to degree 4095 [18, 98].

The maximal subgroups of the almost simple exceptional Lie type groups have not yet been fully classified, although a great deal is known about them. A discussion of the overall strategy for their classification, and a brief summary of the main theorems in this area, appear in [91, Chapter 29]; whilst further references to the literature can be found in [66, Table 1.3.B]. In particular, the maximal subgroups of all of the simple, and most of the almost simple groups of this type with representations of degree up to 12 have been classified: see [106, Section 15] for the Suzuki groups  ${}^2B_2(q)$ , [76, 64] for the Ree groups

${}^2G_2(q)$ , [2, 14, 64] for  $G_2(q)$  and [63] for  ${}^3D_4(q)$ . In this book we extend these classifications to the remaining almost simple groups with representations of degree at most 12.

The sporadic groups can be handled on a case-by-case basis. Most of the necessary information is available, including references to the literature, in [12] or, more usefully for computational purposes, in [111]. At the time of writing, all of the maximal subgroups of the almost simple groups with socle a sporadic group are known except for almost simple maximal subgroups of the Monster whose socle is one of a small list of groups. See [111] for a description of the state of play: note that it has recently been shown by R.A. Wilson that  $L_2(41)$  is a maximal subgroup of the Monster, correcting an earlier error in the literature.

**Computational applications.** We were partly motivated to carry out this classification by its applications to computational group theory. In [9], results of Kovács, Aschbacher and Scott dating from the mid 1980s are used to reduce the computation of the maximal subgroups of a general finite group  $G$  to the case when  $G$  is almost simple. Polynomial-time algorithms for constructing the geometric-type maximal subgroups of the classical groups (in all dimensions) are presented in [45] for the linear, unitary and symplectic groups, and in [46] for the orthogonal groups, and they have been implemented in MAGMA [5].

The Class  $\mathcal{S}$  subgroups arising from representations of almost simple groups in their defining characteristic are generally moderately straightforward to construct using standard functionality for computing with modules over groups. Most of the quasisimple Class  $\mathcal{S}$  subgroups that are not in defining characteristic can be constructed by restricting a representation of a quasisimple group in characteristic 0 to the required finite field. The associated almost simple groups can be constructed over the finite field from a knowledge of the relevant group automorphisms and the computation of module isomorphisms.

Various databases of characteristic 0 representations are available either directly on the web, or via computer algebra systems such as GAP [29] and MAGMA. A facility of this type [111] has been under construction and continuous development for several years now. More recently, Steel [103] has constructed almost all of the characteristic 0 representations in [42] of quasisimple groups in dimensions up to 250. We used data from these databases to carry out some of the calculations needed to complete the classification.

Although the bulk of the arguments used in our classification theorems are theoretical, a substantial number of them make use of computer calculations. These calculations require only small amounts of computer time (generally at most a few seconds) and could be easily carried out using existing functionality and databases in either GAP or MAGMA. The MAGMA commands for each of these individual calculations are given in the files on the webpage

<http://www.cambridge.org/9780521138604>, to enable the user to verify them easily. Whenever we use such a calculation in a proof in this book, we refer to it as a “computer calculation” in the text, and direct the user to the individual online file that contains the commands to carry it out. The files have names like `4134d8calc`, which contains calculations with the 8-dimensional characteristic 0 representation of the group  $4'L_3(4)$ . The matrices defining the images of these representations are stored in data files, which are also on the website, and are accessed by the commands that carry out the calculations.

We shall assume that the reader has a general knowledge of group theory and of group representation theory as might be acquired from advanced undergraduate courses on these topics. Some knowledge of the general theory of classical groups over finite fields and of their associated bilinear, sesquilinear and quadratic forms would also be helpful, because we shall only briefly summarise what we need for this book. Good sources are the books by Rob Wilson [114] (which also includes a great deal of information about maximal subgroups of simple groups), Don Taylor [108] or Chapter 2 of [66]. We do not require any familiarity with algebraic groups, but the interested reader should consult [91] for an introduction which is especially well-suited to our current purposes.

Finally, in any classification project of this scale, it is inevitable that some mistakes will have slipped into our tables. At the time of publication, we know of no such errors, but an errata list has been created at

<http://www.cambridge.org/9780521138604>,

and we shall keep this up to date. We would be extremely grateful to be informed of any errata.

**Acknowledgements.** Writing this book has been a long-running project, and a great many people have helped us at various stages of it. We would like to thank Martin Liebeck for kindly agreeing to write the foreword to this volume. Many thanks go to Gunter Malle for his heroic proof-reading of this entire volume, mostly over the Christmas holidays preceding its publication; the book has been vastly improved by his efforts. Special thanks go also to Rob Wilson for proof-reading large portions of the book, and his thoughtful and extremely helpful comments. In addition we would like to thank Jeff Burdges, Peter Cameron, John Cannon, Ian Gent, Steve Linton, Frank Lübeck, Chris Parker and Anna Schroeder. Needless to say, any and all remaining mistakes and infelicities in this volume are entirely the responsibility of the authors! We gratefully acknowledge the financial support that we have received from EPSRC grants GR/S30580/01 and EP/C523229/1, the Mathematisches Forschungsinstitut Oberwolfach, and the seminar budgets of our respective universities. Finally, Derek thanks Cathy for systems support, and Colva thanks Struan for his encouragement, tolerance and sense of humour during the many years it has taken to write this book.