

Cambridge University Press

978-0-521-17457-2 - Galois Groups and Fundamental Groups

Edited by Leila Schneps

Frontmatter

[More information](#)

This book explores recent research underlining the remarkable connections between the algebraic and arithmetic world of Galois theory and the topological and geometric world of fundamental groups. Arising from an MSRI program held in the fall of 1999, it contains ten articles, all of which aim to present new results in a context of expository introductions to theories that are ramifications and extensions of classical Galois theory.

B. H. Matzat and M. van der Put introduce differential Galois theory and solve the differential inverse Galois problem over global fields in positive characteristic; D. Harbater gives a comparative exposition of formal and rigid patching starting from the familiar complex case. S. Mochizuki discusses aspects of Grothendieck's famous anabelian geometry, while the articles by R. Guralnick, A. Tamagawa, and F. Pop and M. Saïdi investigate the structure of the fundamental groups of curves over different kinds of characteristic  $p$  fields. M. Imbert and L. Schneps study the structure of the Hurwitz spaces and moduli spaces of curves, which are of great importance to Galois theory because of the Galois action on their fundamental groups. The first interesting such group is  $SL_2(\mathbb{Z})$ , a family of special subgroups of which is studied by F. Bogomolov and Y. Tschinkel. Finally, R. Hain and M. Matsumoto present their result proving part of a conjecture by Deligne on the structure of the Lie algebra associated to the Galois action on the fundamental group of the thrice-punctured projective plane.

Cambridge University Press  
978-0-521-17457-2 - Galois Groups and Fundamental Groups  
Edited by Leila Schneps  
Frontmatter  
[More information](#)

---

---

**Mathematical Sciences Research Institute  
Publications**

**41**

---

**Galois Groups and Fundamental Groups**

Cambridge University Press

978-0-521-17457-2 - Galois Groups and Fundamental Groups

Edited by Leila Schneps

Frontmatter

[More information](#)


---

## Mathematical Sciences Research Institute Publications

---

- 1 Freed/Uhlenbeck: *Instantons and Four-Manifolds*, second edition
- 2 Chern (ed.): *Seminar on Nonlinear Partial Differential Equations*
- 3 Lepowsky/Mandelstam/Singer (eds.): *Vertex Operators in Mathematics and Physics*
- 4 Kac (ed.): *Infinite Dimensional Groups with Applications*
- 5 Blackadar: *K-Theory for Operator Algebras*, second edition
- 6 Moore (ed.): *Group Representations, Ergodic Theory, Operator Algebras, and Mathematical Physics*
- 7 Chorin/Majda (eds.): *Wave Motion: Theory, Modelling, and Computation*
- 8 Gersten (ed.): *Essays in Group Theory*
- 9 Moore/Schochet: *Global Analysis on Foliated Spaces*
- 10–11 Drasin/Earle/Gehring/Kra/Marden (eds.): *Holomorphic Functions and Moduli*
- 12–13 Ni/Peletier/Serrin (eds.): *Nonlinear Diffusion Equations and Their Equilibrium States*
- 14 Goodman/de la Harpe/Jones: *Coxeter Graphs and Towers of Algebras*
- 15 Hochster/Huneke/Sally (eds.): *Commutative Algebra*
- 16 Ihara/Ribet/Serre (eds.): *Galois Groups over  $\mathbb{Q}$*
- 17 Concus/Finn/Hoffman (eds.): *Geometric Analysis and Computer Graphics*
- 18 Bryant/Chern/Gardner/Goldschmidt/Griffiths: *Exterior Differential Systems*
- 19 Alperin (ed.): *Arboreal Group Theory*
- 20 Dazord/Weinstein (eds.): *Symplectic Geometry, Groupoids, and Integrable Systems*
- 21 Moschovakis (ed.): *Logic from Computer Science*
- 22 Ratiu (ed.): *The Geometry of Hamiltonian Systems*
- 23 Baumslag/Miller (eds.): *Algorithms and Classification in Combinatorial Group Theory*
- 24 Montgomery/Small (eds.): *Noncommutative Rings*
- 25 Akbulut/King: *Topology of Real Algebraic Sets*
- 26 Judah/Just/Woodin (eds.): *Set Theory of the Continuum*
- 27 Carlsson/Cohen/Hsiang/Jones (eds.): *Algebraic Topology and Its Applications*
- 28 Clemens/Kollár (eds.): *Current Topics in Complex Algebraic Geometry*
- 29 Nowakowski (ed.): *Games of No Chance*
- 30 Grove/Petersen (eds.): *Comparison Geometry*
- 31 Levy (ed.): *Flavors of Geometry*
- 32 Cecil/Chern (eds.): *Tight and Taut Submanifolds*
- 33 Axler/McCarthy/Sarason (eds.): *Holomorphic Spaces*
- 34 Ball/Milman (eds.): *Convex Geometric Analysis*
- 35 Levy (ed.): *The Eightfold Way*
- 36 Gavosto/Krantz/McCallum (eds.): *Contemporary Issues in Mathematics Education*
- 37 Schneider/Siu (eds.): *Several Complex Variables*
- 38 Billera/Björner/Green/Simion/Stanley (eds.): *New Perspectives in Geometric Combinatorics*
- 39 Haskell/Pillay/Steinhorn (eds.): *Model Theory, Algebra, and Geometry*
- 40 Bleher/Its (eds.): *Random Matrix Models and Their Applications*
- 41 Schneps (ed.): *Galois Groups and Fundamental Groups*
- 42 Nowakowski (ed.): *More Games of No Chance*
- 43 Montgomery/Schneider (eds.): *New Directions in Hopf Algebras*
- 44 Buhler/Stevenhagen (eds.): *Algorithmic Number Theory*
- 45 Jensen/Ledet/Yui: *Generic Polynomials: Constructive Aspects of the Inverse Galois Problem*

Volumes 1–4 and 6–27 are published by Springer-Verlag

Cambridge University Press  
978-0-521-17457-2 - Galois Groups and Fundamental Groups  
Edited by Leila Schneps  
Frontmatter  
[More information](#)

---

# Galois Groups and Fundamental Groups

*Edited by*

**Leila Schneps**

*Institut de Mathématiques de Jussieu*



**CAMBRIDGE**  
UNIVERSITY PRESS

Cambridge University Press  
 978-0-521-17457-2 - Galois Groups and Fundamental Groups  
 Edited by Leila Schneps  
 Frontmatter  
[More information](#)

*Series Editor*  
 Silvio Levy  
 Mathematical Sciences  
 Research Institute  
 17 Gauss Way  
 Berkeley, CA 94720  
 United States

Leila Schneps  
 Institut de Mathématiques de Jussieu  
 175, rue du Chevaleret  
 75013 Paris  
 France

*MSRI Editorial Committee*  
 Michael Singer (chair)  
 Alexandre Chorin  
 Silvio Levy  
 Jill Mesirov  
 Robert Osserman  
 Peter Sarnak

The Mathematical Sciences Research Institute wishes to acknowledge support by the National Science Foundation. This material is based upon work supported by NSF Grant 9810361.

---

CAMBRIDGE UNIVERSITY PRESS  
 Cambridge, New York, Melbourne, Madrid, Cape Town,  
 Singapore, São Paulo, Delhi, Tokyo, Mexico City

Cambridge University Press  
 The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

[www.cambridge.org](http://www.cambridge.org)  
 Information on this title: [www.cambridge.org/9780521174572](http://www.cambridge.org/9780521174572)

© Mathematical Sciences Research Institute 2003

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2003  
 First paperback edition 2011

*A catalogue record for this publication is available from the British Library*

ISBN 978-0-521-80831-6 Hardback  
 ISBN 978-0-521-17457-2 Paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

## Contents

Introduction	ix
Monodromy Groups of Coverings of Curves	1
ROBERT GURALNICK	
On the Tame Fundamental Groups of Curves over Algebraically Closed Fields of Characteristic $> 0$	47
AKIO TAMAGAWA	
On the Specialization Homomorphism of Fundamental Groups of Curves in Positive Characteristic	107
FLORIAN POP AND MOHAMED SAÏDI	
Topics Surrounding the Anabelian Geometry of Hyperbolic Curves	119
SHINICHI MOCHIZUKI	
Monodromy of Elliptic Surfaces	167
FEDOR BOGOMOLOV AND YURI TSCHINKEL	
Tannakian Fundamental Groups Associated to Galois Groups	183
RICHARD HAIN AND MAKOTO MATSUMOTO	
Special Loci in Moduli Spaces of Curves	217
LEILA SCHNEPS	
Cellulation of Compactified Hurwitz Spaces	277
MICHEL IMBERT	
Patching and Galois Theory	313
DAVID HARBATER	
Constructive Differential Galois Theory	425
B. HEINRICH MATZAT AND MARIUS VAN DER PUT	

Cambridge University Press

978-0-521-17457-2 - Galois Groups and Fundamental Groups

Edited by Leila Schneps

Frontmatter

[More information](#)

Galois Groups and Fundamental Groups  
 MSRI Publications  
 Volume 41, 2003

## Introduction

This volume is the outcome of the MSRI special semester on *Galois Groups and Fundamental Groups*, held in the fall of 1999. Respecting the famous Greek requirements of unity of place, time, and action, the semester was an unforgettable, four-month-long occasion for all mathematicians interested in and responsible for the developments of the connections between Galois theory and the theory of fundamental groups of curves, varieties, schemes and stacks to interact, via a multitude of conferences, lectures and conversations.

Classical Galois theory has developed a number of extensions and ramifications into more specific theories, which combine it with other areas of mathematics or restate its main problems in different situations. Three of the most important of these extensions are *geometric Galois theory*, *differential Galois theory*, and *Lie Galois theory*, all of which have undergone very rapid development in recent years. Each of these theories can be developed in characteristic zero, over the field  $\mathbb{C}$  of complex numbers, over number fields or  $p$ -adic fields, or in characteristic  $p > 0$ ; various versions of the classical and inverse Galois problems can be posed in each situation. The purpose of this introduction is to give a brief overview of these three themes, which form the framework for all the articles contained in this book.

The main focus of study of **geometric Galois theory** is the theory of *curves* and the many objects associated to them: curves with marked points, their fields of moduli and their fundamental groups, covers of curves with their ramification information and their fields of moduli, and the finite quotients of the fundamental group which are the Galois groups of the covers, as well as the moduli spaces and Hurwitz spaces which parametrize all these objects.

To consider a curve  $X$  topologically is tantamount to considering it over the field of complex numbers  $\mathbb{C}$ . As an abstract group, the *topological fundamental group* of the curve depends only on the genus  $g$  and the number  $n$  of marked points chosen on the curve; it can be identified with the group of homotopy classes of loops on the curve based at a fixed (not marked) base point, and is presented by standard generators  $a_1, \dots, a_g, b_1, \dots, b_g, c_1, \dots, c_n$  subject to the unique relation

$$[a_1, b_1] \cdots [a_g, b_g] c_1 \cdots c_n = 1. \quad (*)$$

Cambridge University Press

978-0-521-17457-2 - Galois Groups and Fundamental Groups

Edited by Leila Schneps

Frontmatter

[More information](#)

Note that when  $n \geq 1$ , this group is actually free. The Galois covers of the curve correspond to the finite quotients of this group, which are exactly the finite groups generated by generators  $a_i$ ,  $b_i$  and  $c_j$  satisfying (\*), so they are perfectly understood. The *algebraic fundamental group* of the curve is the Galois group of the compositum of all function fields of finite étale covers of the curve over the function field of the curve itself; in fact, it is exactly the profinite limit of the finite quotients of the topological fundamental group.

This simple situation leads or generalizes very naturally into new regions that contain all kinds of very difficult problems. We sketch some of them:

**1. Fundamental groups in characteristic  $p$ .** When a curve  $X$  is defined over a field in characteristic  $p$ , all relations between its fundamental group and any topological notion of ‘loops’ must be forgotten. Over an algebraically closed field, one defines the algebraic fundamental group directly, exactly as above; it is the profinite limit of the Galois groups (monodromy groups) of finite étale covers of the curve. However, in this situation, for  $(g, n)$  different from  $(0, 0)$  and  $(1, 0)$ , it is extremely difficult to determine the structure of the fundamental group, or even the weaker question of which finite groups can occur as its quotients. Indeed, this is one of the fundamental problems of geometric Galois theory in characteristic  $p$ . In the affine case ( $n \geq 1$ ), the complete solution to the weaker problem was conjectured by Abhyankar; this conjecture was proved over the affine line by M. Raynaud, and the proof extended to all curves by D. Harbater. However, the situation remains completely mysterious in the case of complete curves ( $n = 0$ ,  $g > 1$ ).

Things are better if one considers only the quotients of order prime to  $p$ ; then a result due to Grothendieck states that the groups of order prime to  $p$  which can occur are exactly the finite quotients of order prime to  $p$  of the topological fundamental group of type  $(g, n)$  defined in (\*), and that in fact the prime-to- $p$  quotients of the fundamental groups over  $\mathbb{C}$  and in characteristic  $p$  are isomorphic. In characteristic  $p$ , this group is a quotient of the *tame fundamental group*, which is the largest quotient of the fundamental group having inertia subgroups of order prime to  $p$ ; this group (which is equal to the fundamental group when  $n = 0$ ) is easier to work with than the full group for various purposes. But the structure of the tame fundamental group and the set of its finite quotients are absolutely unknown, except in the non-hyperbolic cases  $(g, n) = (0, 0)$ ,  $(0, 1)$ ,  $(0, 2)$  and  $(1, 0)$ .

The articles by R. Guralnick, A. Tamagawa, and F. Pop and M. Saïdi all work in the situation of curves defined over an algebraically closed field of characteristic  $p$ . Guralnick works on the problem of determining which groups can occur as Galois groups (or their composition factors) of finite separable covers  $f : Y \rightarrow X$ , where  $Y$  is of fixed genus  $g$ , and seeks groups which can specifically be excluded. Tamagawa shows that given the tame fundamental group, it is possible to recover the type  $(g, n)$  of the curve (if  $(g, n) \neq (0, 0)$  or  $(0, 1)$ ).



Results in this and other papers by Tamagawa even tend to imply that in some cases, the tame fundamental group may determine the isomorphism class of the curve completely. This shows how different the characteristic  $p$  case is from the characteristic 0 case, in which as we saw, curves of many different types may have isomorphic fundamental groups (for instance, the fundamental groups of curves of type (2, 2) and (1, 4) are both free of rank 5). Finally, Pop and Saïdi address similar questions, proving, under certain hypotheses on the Jacobians, that at most a finite number of curves can have isomorphic fundamental groups.

**2. Anabelian theory.** We saw above that the isomorphism class of the topological or the algebraic fundamental group is very far from determining even the most basic information about a curve in characteristic 0, such as its type  $(g, n)$ , whereas in characteristic  $p$  it determines much more if not all of the information about the specific curve. However, one can also consider the algebraic fundamental group equipped with its canonical outer Galois action, which should provide more information. Indeed, any variety (scheme, stack) defined over an algebraically closed field can actually be considered as defined over a subfield  $K$ , given by the coefficients of the equations of a defining model, say, and which is finitely generated over the prime field and not algebraically closed. Then there is an exact sequence

$$1 \rightarrow \pi_1(X \otimes \bar{K}) \rightarrow \pi_1(X) \rightarrow \text{Gal}(\bar{K}/K) \rightarrow 1, \quad (**)$$

where  $\pi_1(X \otimes \bar{K})$  denotes the algebraic fundamental group. The *anabelian problem*, which was posed by Grothendieck in his famous letter to G. Faltings, asks which varieties are entirely determined by the group  $\pi_1(X \otimes \bar{K})$  together with the action  $\text{Gal}(\bar{K}/K) \rightarrow \text{Out}(\pi_1(X \otimes \bar{K}))$ . Grothendieck called varieties which are thus determined *anabelian varieties*, and explicitly stated that hyperbolic curves should be anabelian. This is related to the hitherto unproven *section conjecture* for a hyperbolic curve  $X$ , which states that the sections

$$\text{Gal}(\bar{K}/K) \rightarrow \pi_1(X)$$

of (\*\*\*) are in bijection with the rational points of  $X$  if  $X$  is complete, and this set together with the tangential base points if  $X$  is not complete.

S. Mochizuki proved that hyperbolic curves defined over sub- $p$ -adic fields, that is, fields which are subfields of fields finitely generated over the  $p$ -adics, are indeed anabelian. In his article in this volume, he discusses various results related to this theorem, including a partial generalization to characteristic  $p$  and a discussion of the section conjecture over the field of real numbers.

**3. Galois action on fundamental groups.** In his *Esquisse d'un Programme*, completing the letter to Faltings, Grothendieck suggested that not only hyperbolic curves, but also the moduli spaces  $\mathcal{M}_{g,n}$  of curves of type  $(g, n)$  should be examples of anabelian varieties, and that explicitly investigating the Galois action on their fundamental groups should provide information of an entirely new

Cambridge University Press

978-0-521-17457-2 - Galois Groups and Fundamental Groups

Edited by Leila Schneps

Frontmatter

[More information](#)

type about the elements of  $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ ; this is known as *Grothendieck-Teichmüller theory*. The first non-trivial moduli space  $\mathcal{M}_{g,n}$  is the case  $(g,n) = (0,4)$ ; we have  $\mathcal{M}_{0,4} = \mathbb{P}^1 - \{0, 1, \infty\}$ . Following the direction initiated by Grothendieck, the Galois action on the fundamental group of this space (the profinite free group  $\widehat{F}_2$  on two generators) has been studied in a theory known as *dessins d'enfants*. The study has focused on coding the conjugacy classes of finite index subgroups of  $\widehat{\pi}_1(\mathbb{P}^1 - \{0, 1, \infty\})$ , corresponding to the finite étale covers of  $\mathbb{P}^1 - \{0, 1, \infty\}$ , as combinatorial objects (the dessins d'enfants), and using the combinatorics to look for invariants identifying the Galois orbits of these covers.

The only other moduli space of dimension 1 is  $\mathcal{M}_{1,1}$ , the moduli space of elliptic curves (genus one curves with one distinguished point). This space is the quotient of the Poincaré upper half-plane by the proper and discontinuous (but not free) action of  $\text{SL}_2(\mathbb{Z})$ . Finite-index subgroups of  $\text{SL}_2(\mathbb{Z})$  correspond to covers of  $\mathcal{M}_{1,1}$ . As in the case of  $\mathcal{M}_{0,4}$ , many specific families of these subgroups have been studied in detail, most familiarly the modular subgroups  $\Gamma(N)$ . Using graphs in the spirit of the theory of dessins d'enfants, F. Bogomolov and Y. Tschinkel characterize another family of very special finite-index subgroups of  $\text{SL}_2(\mathbb{Z})$ , namely those corresponding to elliptic fibrations.

Before passing from these two curves to what can be said in the case of general moduli spaces  $\mathcal{M}_{g,n}$ , let us make a brief foray out of the geometric situation into the domain of **Lie Galois theory**, a subject that originates in the geometric situation but has been linearized by focusing on graded Lie algebras associated to the profinite fundamental groups rather than the groups themselves. A great deal of work has been done in this subject, mainly by Y. Ihara and his school, but we restrict ourselves here to discussing one conjecture which is a paradigm for the manner in which the problems in the domain arise in geometry, but raise their own interesting arithmetic questions.

Since as above, we have  $\pi_1(\mathbb{P}^1 - \{0, 1, \infty\}) \simeq \widehat{F}_2$ , the exact sequence (\*\*\*) gives a canonical homomorphism

$$G_{\mathbb{Q}} \rightarrow \text{Out}(\widehat{F}_2). \quad (***)$$

As an initial step, the passage from the geometric situation to the Lie situation involves replacing the profinite completions of fundamental groups by their pro- $\ell$  completions, that is, the completions with respect to all finite quotients which are  $\ell$ -groups for a fixed prime  $\ell$ . Denote the pro- $\ell$  completion of  $F_2$  by  $F_2^{(\ell)}$ . This completion is a quotient of the profinite completion by a characteristic subgroup, so that (\*\*\*) yields a homomorphism  $G_{\mathbb{Q}} \rightarrow \text{Out}(F_2^{(\ell)})$ . Following Ihara, define a filtration on  $G_{\mathbb{Q}}$  by setting

$$I^m G_{\mathbb{Q}} = \text{Ker}\{G_{\mathbb{Q}} \rightarrow \text{Out}(F_2^{(\ell)}/L^{m+1})\}$$

where  $L^m$  denotes the  $m$ -th term of the lower central series of  $F_2^{(\ell)}$ , and set

$$\text{Gr}^m G_{\mathbb{Q}} = I^m G_{\mathbb{Q}}/I^{m+1} G_{\mathbb{Q}}.$$

Cambridge University Press

978-0-521-17457-2 - Galois Groups and Fundamental Groups

Edited by Leila Schneps

Frontmatter

[More information](#)

The following conjecture on the structure of the graded Lie algebra associated to the filtration  $I^m G_{\mathbb{Q}}$  was stated (in fuller detail) by Ihara, who attributed it to Deligne.

CONJECTURE. *The Lie algebra*

$$\left[ \bigoplus_{m>0} \mathrm{Gr}^m G_{\mathbb{Q}} \right] \otimes \mathbb{Q}_\ell$$

*is freely generated by generators  $s_3, s_5, s_7, \dots$ , where  $s_m \in \mathrm{Gr}^m G_{\mathbb{Q}}$  is the so-called Soulé element.*

Part of this conjecture, namely the fact that the Lie algebra is actually generated by the  $s_i$ , was proved by Hain and Matsumoto. In their contribution to this volume, they discuss the conjecture and show how to fit it into a motivic framework.

Now let's return to the situation of geometry and consider the (conjecturally anabelian) moduli spaces  $\mathcal{M}_{g,n}$ . The geometry of these spaces has been described by explicitly cutting them into simply connected regions called cells, enumerated by objects known as fatgraphs, which are in fact equivalent to dessins d'enfant. The *Hurwitz spaces* are similar to the moduli spaces, but they parametrize equivalence classes of ramified coverings of Riemann surfaces, where two such coverings  $f : Y \rightarrow X$  and  $f' : Y' \rightarrow X'$  are equivalent if a diagram

$$\begin{array}{ccc} Y & \xrightarrow{f} & X \\ \phi \downarrow & & \downarrow \psi \\ Y' & \xrightarrow{f'} & X' \end{array}$$

commutes for two biholomorphisms  $\phi$  and  $\psi$ . A cellulation of the compactification of these spaces, analogous to that of the moduli spaces, is defined and studied in the article by M. Imbert.

The Hurwitz spaces are closely related to *special loci* in the moduli spaces, namely the loci of points in the moduli spaces corresponding to marked Riemann surfaces admitting a particular group of automorphisms. The article by L. Schneps studies these loci, showing that under certain conditions (which are always fulfilled in genus zero), the special loci are themselves moduli spaces of smaller type. The morphisms mapping these smaller moduli spaces to the special loci of the larger ones are respected by the canonical Galois action on the fundamental groups of the moduli spaces, so that the addition of these morphisms to those previously studied in Grothendieck-Teichmüller theory adds new combinatorial information on the elements of  $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ .

**4. Inverse Galois theory.** One of the most fundamental problems of Galois theory is that of determining which finite groups can occur as Galois groups over a given field  $K$ . This problem has been studied in many different situations and

Cambridge University Press

978-0-521-17457-2 - Galois Groups and Fundamental Groups

Edited by Leila Schneps

Frontmatter

[More information](#)

by many different methods; by various direct methods, by explicit computation and solution of obstructions to embedding problems, and by geometry. The geometry comes in when studying a field of the type  $K(t)$  for an indeterminate  $t$ , and the main tools are curves, since the desired groups are exactly the Galois groups of covers of the projective line over  $K$ .

When  $K = \mathbb{C}$ , the inverse Galois problem is solved by Riemann's existence theorem; every finite group occurs. Although no completely general analog to Riemann's existence theorem exists over arbitrary fields, many partial analogs have been developed in different situations. The key notion is that of *patching*; covers are constructed locally over disks, and the pieces are patched together (agree) on the overlaps. Like the theories described above, the inverse Galois problem exists in characteristic 0 and  $p > 0$ , necessitating the use of different techniques. Patching techniques have been developed using formal schemes (formal patching) and non-archimedean disks (rigid patching) which yield partial or complete solutions to the inverse Galois problem over many different fields (large or algebraically closed fields of any characteristic, fraction fields of complete local rings other than fields, complete fields, henselian fields, and so on). In fact, one can obtain stronger results, such as specific information on the structure of the fundamental group for affine curves; in certain cases, it can even be shown that the fundamental group is free, or that it has the property that every split embedding problem has a proper solution. The contribution by D. Harbater contains a comprehensive account of all these methods and results.

The domain of **differential Galois theory** is an adaptation of the classical inverse Galois problem; instead of considering finite groups as Galois groups of Galois extensions of arbitrary fields, one considers linear algebraic groups as Galois groups of so-called Picard-Vessiot extensions of D-fields, which are fields  $F$  equipped with a derivation  $\partial : F \rightarrow F$ . Over algebraically closed fields of characteristic 0, this problem has been completely solved by the combined results of Ramis, Mitschi, Singer, van der Put and finally Hartmann; any linear algebraic group over such a field is the differential Galois group of a Picard-Vessiot extension. In their contribution to this volume, Matzat and van der Put develop a non-obvious analog of these results in the characteristic  $p$  situation; they introduce iterated differential fields and give a complete formulation and solution to the inverse Galois problem over them.

Leila Schneps

Paris, November 2002