

Index

References to Chapter 7, The Flora, are marked by a tick ('); they are confined to families and genera, making reference to species readily available.

The taxa list in Appendix II is indexed for families only; these entries are marked by a double tick ("').

Major consideration of plant species and Paradox tables is distinguished by bold type.

References to tables and figures are marked with an asterisk (*).

Topics and authors are indexed only for the more important references.

- abbreviations, BIBLIO table, 59*
- MACRO table, 47*, 48*
- PLANT table, 52–53*
- PLANTDS tables, 56–58*
- PLANTHAB table, 54*
- PLSAMPLE table, 45*
- POLLEN table, 45*, 48*
- SITECHAR table, 40–41*
- sites, 62–63*
- TAXA table, 43*
- TAXASORT table, 43*, 44*
- TAXATIME table, 51*
- Abies*, 81', 139
 - alba*, 211
- Acer*, 97'
- Aceraceae, 97', 296"
- Achillea*, 111'
 - millefolium*, 162, 215
- active layer, 231
- Adam, P., 168
- Aghnadarragh, 70
- agricultural revolution, 262
- Agrostis*, 116', 203
- Ajuga*, 108'
- Alaska North Slope, 11, 23, 165
- Alisma*, 112', 139
 - plantago-aquatica*, 213
- Alismataceae, 112', 303"
 - macroscopic remains, 112'
 - pollen, 112'
- Allium*, 115'
 - schoenoprasum*, 206, 207, **218**
- Alnus*, 85'
 - glutinosa*, 211
- Alopecurus*, 116', 203
- alpine species, 186
- Alyssum*, 93'
- analogues, 10–12, 192, 236
- analysis of database, 266
- Andersen, J., 249
- Andersen, S.T., 236
- Andrew, R., 75
- Androsace*, 105'
 - septentrionalis*, **218**
- Anglian, Early, 19
 - late-glacial, 217
 - Middle, 19
 - Stage, 10, 19, 68
 - Late, 19
- annuals, 158
- Anthemis* type, 112'
- Anthoxanthum*, 116', 203
 - odoratum*, 203
- Anthriscus*, 102', 197

- anthropogenic factors, 173
Anthyllis, 95'
Aphanes, 94'
Apium, 102'
 aquatic vegetation, 212
Aquifoliaceae, 97', 296"
Araliaceae, 102', 297"
 arctic, definition, 7
 distributions, origin, 260
 plant communities, 192
 pollen spectra, 192
 rivers, 26
 steppe biome, 201
 term, 249, 265
 insect life, 163
 subarctic categories, 186
 vegetation compared, 162
 arctic-alpine element, 175, 190
 heath, 244
 arctic-subarctic element, 175
Arctostaphylos, 103'
 uva-ursi, 210
 Ardleigh, 187
Arenaria, 90'
 ciliata, 218
Armeria, 24, 106, 188, 216, 217, 218
 macroscopic remains, 105'
 pollen, 105'
 maritima, 106, 133, 206, 207, 212, 216,
 242, 245, 262
Artemisia, 112', 127, 192, 201, 206, 217, 218,
 254
 Grande Pile, 250
 representation, 145, 146*
 ASCII files, 37, 59
 Ashworth, A.C., 255
Aspidiaceae, 286"
 assemblages, dispersal, 153
 biota-dependent factors, 153
 taphonomy-dependent factors, 153
Aster type, 112'
Astragalus, 95'
Athyriaceae, 286"
 Atigun River, 30
 Atkinson, T.C., 247
Atlas of the British Flora, 184
Atriplex, 89'
 hastata, 214, 215
Aurivillius, C., 162
Azolla, 80'
Azollaceae, 80', 286"
 Banks Island, 11
 Banwell-type fauna, 252
 bar-tail pools, 28
 bar-top drapes, 26
Barbarea, 92'
- barrens, 207
Bartsia, 108'
 Bathurst Island, 138*, 199*
 Baventian, 69
 Beestonian, 69, 260
 Beetley, 194, 204, 216
 Behre, K.-E., 251
 Bell, F.G., 6, 75, 180, 236
 elements of the flora, 180, 181*
 Bembridge, 70
 Beringia, 12, 198, 201, 235, 259
 Late Wisconsin vegetation, 206
Betula, 102'
 erecta, 212, 213
Betula, 85', 127, 194, 195, 196, 211, 248
 nana, 133, 188, 195, 209, 211, 219
 nana, distribution, 242
 pubescens, 196
Betulaceae, 85', 287"
 BIBLIO table, 7, 58, 60
 bibliography, 39
Bidens, 111'
 biennials, 158
Bilderdykia, 88'
 convolvulus, 215
 bioclimate, 7
 biological attributes, 155
 biological spectrum, 155
 biome, history, 265
 biome term, 265
 biostratigraphy, 264
 biota, cold stage, 267
 mixed, 257
 biotype, 12, 122
 evolution, 260
 Birks, H.J.B., 75
Bison, 253, 254
Bliss, L.C., 157
 Block Fen, 204, 217
Blysmus, 118'
Boraginaceae, 107', 300"
 boreal distributions, origin, 260
Botrychium, 76, 79'
 Bovey Tracey, 70
 Brandon, 217
 Brimpton, 195
 Brimpton Interstadial, 19, 194, 196, 199
 correlation, 196
 British Flora, origin, 5, 6
 Broome, 210
 Brørup Interstadial, 196
Bruckenthalia, 102', 210
 spiculifolia, 187, 210, 211
Bryophyta, 308"
 bryophytes, 6
 Burn, C.R., 229
Butomaceae, 113', 303"

Index

311

- Butomus*, 113
Buxaceae, 97', 296"
Buxus, 97'
C₃, C₄ species, 204
Calamintha, 108'
Callitrichaceae, 107', 300"
Callitriche, 108', 213
Calluna, 102', 194, 195, 210
 representation, 150, 151*
vulgaris, 210
Caltha, 91', 188
 palustris, 211, 240
Campanula, 111', 262
 rotundifolia, 132, 137, 155, 159, 160, 162,
 206, 208, 216, **219**
Campanulaceae, 111', 301"
Caprifoliaceae, 110', 301"
Capsella, 93'
 bursa-pastoris, 215
carbonate encrustations, 205
Cardamine, 93'
Carduus, 112'
Carex, 119', 137, 188, 216
 flacca, 204
 panicea, 204
Carpinus, 35, 86, 133, 139
Carya, 85'
Caryophyllaceae, 90', 123, 160, 188, 289"
 macroscopic remains, 90'
 pollen, 90'
Centaurea, 112'
 nigra, 132, 206, 215
Centaurium, 106'
Cerastium, 90', 188
 arvense, 206, 215
Ceratophyllaceae, 91', 291"
Ceratophyllum, 91'
 demersum, 213
Chalk, 207
Chamaephyte percentage, 157
channels, abandoned, 31
Chara, 120'
Characeae, 120', 308"
Chelford Interstadial, 10, 19, 103, 194, 195,
 199
Coleoptera, 255
 correlation, 196
Chenopodiaceac, 89', 139, 289"
 macroscopic remains, 89'
 pollen, 89'
Chenopodium, 89'
 rubrum, 214
Chernov, Yu.I., 240, 259
chromosome numbers, 159
CHRONOL table, 37, **42**, 59
chronology, 38
chronostratigraphy, 264
Chrysosplenium, 93'
Cicuta, 102'
 virosa, 213
Cirsium, 112'
 arvense, 215
Cistaceae, 99', 297"
Cladium, 118'
Clementsian view, 190, 252, 265
climate, analysis of distribution, 240
 Arctic chamaephyte, 156
 aridity, 230
 biological evidence, 233
 Chelford Interstadial, 238
 control of distribution, 240
 estimations in NW Europe, 234
 evidence, 227
 evidence from sediments, 230
 evidence from structures, 231
 geological evidence, 230
 indicated by vegetation, 237
 indicated by species, 239, 241, 242
 interpretation, 234
 interpretation of flora and fauna, 247
 interstadial, 238, 249
 Middle Pleniglacial, 235
 plants and animals, 249
 Pleniglacial, 249
 reconstruction, 227
 seasonal, 249, 267
 seasonal differences, 192
 stadial, 235
 temperature indicators, 239
 Upton Warren Interstadial Complex,
 247
 Weichselian, 233
 Weichselian, NW Europe, 238
climatic change, global, 234, 236
 regional, 234
 development, interstadials, interglacials,
 9*, 20
 envelopes, 236, 249
Cochlearia, 92'
Cody, W.J., 229
cold stage, biota, 252
 definition, 17
 Britain, 13–16*
 Ireland, 15–16*
 The Netherlands, 16*
 variation of conditions, 20
Coleoptera, 162
 Late Wisconsin, 255
Coleopteran fauna, 19, 254
 Devensian, 254
Colney Heath, 35, 65, 70, 76
Columella columella, 256
competition, 160, 162

- Compositae, 76, 111', 123, 160, 302"
Liguliflorae, 112', 139
 macroscopic remains, 111'
 pollen, 112'
Tubuliflorae, 112'
 Coniferae, 81', 286"
 continental climate, 168
 shelf, 23
 species, 202
 continentality, 244
 contraction cracks, 232
 Convolvulaceae, 107', 299"
Convolvulus, 107'
 Coope, G.R., 196, 247, 254
Corbicula, 35
Corispermum, 89', 219
 Cornaceae, 101', 297"
Cornus, macroscopic remains, 101'
 pollen, 101'
 sanguinea, 211
Coronopus squamatus, 214
 Corton Sands, 10
 Corylaceae, 86', 288"
Corylus, 35, 86', 133, 139
 representation, 147, 148*
 cover sand, 168, 207, 230
 Coygan-type fauna, 253
 Crassulaceae, 93', 294"
 crevices, 207
Cristatella, 76
 critical species, 121
 Cromer Forest Bed Series, 188
 Cromerian Complex cold stage, 69
 Cruciferae, 92', 123, 139, 160, 188, 292"
 macroscopic remains, 92'
 pollen, 93'
 representation, 142, 145
 cryoturbation, 11, 165
 Cryptogrammaceae, 286"
 Cryptophyte percentage, 157
 Cupressaceae, 83', 287"
Cuscuta, 107'
 cushion plants, 205
 Cyperaceae, 117', 123, 127, 139, 203, 204,
 306"
 pollen, 117', 188, 197, 198, 204
 representation, 142, 143*
Cyperus, 118'
Damasonium, 112'
 alisma, 180, 213, 219
Daphnia, 76
 Darwin, C., 5
 dating, 19
 radiocarbon, 19, 67–68*
 thermoluminescence, 19, 67–68*
Daucus, 102'
 Davis, A.J., 236
 Devensian, Early, 17
 ice sheet, 18
 Late, 17, 23
 late-glacial, 2
 Middle, 17
 records, 187
 Stage, 1, 17, 64
Dianthus, 90, 219
 Dickson, J.H., 6
 Dicotyledones, 287*
Diphasium, 77'
Diplotaxis, 92'
 tenuifolia, 132, 215, 220
 Dipsacaceae, 110', 301"
 Diptera, 162
 dispersal, 160, 162
 macros, 137
 wind, 162
 distribution, climate, 174
 aquatics, 213
 categories, 174, 199
 categories, cold stage floras, 243*
 categories, fells, rocks, etc., 207
 categories, fens, flushes, marshes, 212
 categories, grassland, 207
 categories, meadow, 207
 categories, *Potamogeton*, *Groenlandia*,
 246*
 categories, woodland, scrub, 212
 in time, 187
 limits, north and south, 244
 of taxa, 173
 present, 173
 types of Forbes, 174
 disjunct, 262
 diversity, biota, habitat and climate, 257
 species, 167
Draba, 92', 160, 188
 incana, 207, 208, 220
 drainage, 165
Drosera, 93'
 Droseraceae, 93', 293"
 drought periods, 235
Dryas, 94'
 octopetala, 220, 244, 262
 octopetala, distribution, 242
Dryopteris type, 76, 80'
 dunes, 207
 Earith, 35
 Early Devensian flora, vegetation, 261
 Early Pleistocene pollen diagram, 209*
 Early Pleistocene succession, 15, 17
 Early Pleniglacial, 232
 Early Quaternary, 259
 eastern limit category, 178*

- Ecological Flora Database, 180, 182*, 183*, 190, 240
E/W index, 182
S/N index, 182
 ecotypes, 121
 selection, 160
Elaeagnaceae, 98', 296"
Elatinaceae, 9', 297"
Elatine, 99'
Eleocharis, 118'
palustris, 212, 213
Elymus, 116', 203
Empetraceae, 104', 298"
Empetrum, 194, 210
 macroscopic remains, 104'
nigrum type, 210
 pollen, 104'
 representation, 150, 152*
 entomophilous plants, tundra, 162
 environments, 4
 northern, 233
Ephedra, 83', 254
Ephedraceae, 83', 287"
Epilobium, macroscopic remains, 100'
 pollen, 100'
Equisetaceae, 78', 285"
Equisetum, 78'
 macroscopic remains, 78'
 spores, 79'
Erica, 103', 195
Ericaceae, 103', 209, 210, 298"
 macroscopic remains, 103'
 pollen, 103'
Ericales, 102'
 pollen frequencies, 73
 representation, 147, 149*, 208
Eriophorum, 118', 204
vaginatum, 195, 209, 210
Erodium, 96'
Erophila, 93', 188
Euphorbia, 97'
cyparissias, 206, 220
Euphorbiaceae, 97', 296"
Fagaceae, 86', 288"
Fagus, 87'
 family numbers, 75
 faunas, 201, 202
 Middle Devensian, 253, 254
 Wisconsin, 254
 non-analogue, 257
 fells, 207
 fens, 212
Festuca, 116'
halleri, 203
rubra, 203
Filicales, 285"
- Filipendula*, 94'
ulmaria, 162, 207, 211, 212
 Flandrian Stage, 1, 13, 191
 floatation of macros, 137
 floodplains, 26*, 166
 flora, 77
Flora Europaea, 73, 75, 122, 123, 159, 169
 floras, Middle and Upper Pleistocene, 260
 floristic elements, 174, 175
 mixture, 242, 244
 flushes, 212
 Forbes, E., 5, 188
 forest clearance, Neolithic, 262
 forest limit, 241
 forest-steppe, 239
Frangula, 97'
Fraxinus, 106'
ornus, 244
 freshwater sediments, 31
Fritillaria, 115'
 full-glacial, 8, 20
 future considerations, 268
Galeopsis, 108'
Galium, 107'
 genetic diversity, 262
 resources, 257
Gentiana pneumonanthe, 196
Gentianaceae, 106', 299"
 macroscopic remains, 106'
 pollen, 106'
Gentianella, 106'
 geology, regional, 21
Geraniaceae, 96', 296"
 macroscopic remains, 96'
 pollen, 96'
Geranium, 96'
Geum, 94'
 glaciomarine environment, 33
Glaux, 105', 188
maritima, 212, 214, 215
 Gleasonian view, 190, 252, 262, 265
 Glinde Interstadial, 251
 global warming, 249
Glyceria, 116', 203
 Godwin, H., 5, 6, 75, 77, 175, 188, 261, 262
 Gotland, 262
 Graham, R. W., 254
Gramineae, 116', 123, 127, 139, 188, 202, 203, 305"
 climatic change, 258
 Grande Pile, 250
 macroscopic remains, 116'
 pollen, 116, 188, 197, 198, 201, 202, 203
 representation, 140, 141*
 graminoid productivity, 193

- Grande Pile, 217, 228*
- grassland, 8, 203, 205, 206, 235, 238
- grazing factor, 248
 - fauna, 202
 - population, 169
 - pressure, 202
- Great Ice Age, 2
- Grichuk, M.P. & V.P., 10
- Grime, J.P., 155
- Groenlandia*, 114, **245**
 - densa*, 213, 246
- groundwater discharge, 168
- growing season, 235
- growth, 160
- Guiot, J., 236
- Gulf Stream, 21
- Guttiferae, 98', 296"
- habitats, 164, 169
 - analysis, 169, 170*
 - behaviour 172
 - floodplain, 215
 - heterogeneity, 253
 - light, 172
 - open, 261
 - place, 172
 - preference, 172
 - salinity, 172
 - soil, 172
 - variation, 240
 - vegetation, 169
 - water, 172
- halophytes, 167, 214
- Haloragaceae, 100', 297"
- Hartz, N., 2
- heath taxa, 147
- heathland, 207
 - Early Pleistocene, 208
 - Early Weichselian, 208
- heaths, 207
- Hedera*, 102'
 - helix*, 235
- Hedysarum*, 95'
- Helianthemum*, 75, 127, 139, 188, 216, 218, 262, 263
 - canum*, 132, 159, 179, 188, 206, 218, **220**, **245**
- macroscopic remains, 99'
- pollen, 99'
- Helianthemum-Artemisia* heath, 244
- Hemicryptophyte percentage, 157
- Hengelo Interstadial, 250, 251
- Heracleum*, 102', 197
- herbivores, 203
- Hernaria*, 90, **221**
- Herschel Island, 11, 198, 203
- Highland Line, 18*, 21, 164
- Hill, M., 174
- Hippophae*, 98', 217, **221**
 - rhamnooides*, 217
- Hippuridaceae, 101', 297"
- Hippuris*, 188
 - macroscopic remains, 101'
 - pollen, 101'
 - vulgaris*, 213
- Histon Road, Cambridge, 35, 261
- Historical Northern elements, species, 175, 190, 260, 261
- Tertiary elements, species, 175, 190, 260, 261
- Holocene Epoch, 1, 13, 191
- Holyoak, D.T., 26
- Hordeum*, 203
- humidity, 238
- Huperzia*, 77'
 - selago*, 132, 209
- hybrid taxa, 121
 - zones, 159
- Hydrocharis*, 113'
- Hydrocharitaceae, 113', 303"
- Hydrocotyle*, 102'
 - vulgaris*, 212
- Hypericum*, 98'
- Hypolepidaceae, 286"
- ice limits, 18*
- ice wedges, 228*, 230, 231*
- identification problems, 73
 - macros, 284
- Ilex*, 97'
 - aquifolium*, 235
- insects, anthophilous, in tundra, 162
 - environmental change, 249
- interstadial events, 8
- interstadials, Early Weichselian, 229
 - Early Devensian/Weichselian, 250
 - Middle/Late Devensian/Weichselian, 250
- introduction, 121
- Iridaceae, 115', 304"
- Iris*, 115'
- Isles of Scilly, 36, 70, 76
- Isoetaceae, 78', 285"
- Isoetes*, 78'
- isopolls, 3
- Iversen, J., 234, 235, 239, 265
- Jasione*, 111'
 - montana*, 207
- Jessen, K., 8, 10
- Juglandaceae, 85', 287"
- Juglans*, 85'
- Juncaceae, 115', 305"
- Juncaginaceae, 113', 303"

- Juncus*, 115', 216
biflorus, 162
gerardii, 212, 214
Juniperus, 83', 211
- Kerney, M.P., 256
Koenigia islandica, 87, **221**
Kolstrup, E., 236
- Labiatae, 108', 300"
macroscopic remains, 108'
pollen, 108'
- Lake Hazen, 205
- lakes, 31
- Lambert, C.A., 75
- landscapes, 164, 244
- Late Pleniglacial, 232
- Lathyrus*, 95'
latitudinal differences, 192
- leaf area index, 238
- ledges, 207
- Leguminosae, 95', 139, 295"
macroscopic remains, 95'
pollen, 95'
- Lemna*, 116'
- Lemnaceae, 116', 306"
- Lentibulariaceae, 109', 301"
- Leontodon*, 112'
autumnalis, 207, 215
- lichens, 205
- life form, 52, 155
analysis, 156*, 162, 238
percentages, 156
- life span, 158*
- Liliaceae, 115', 304"
- limestone pavements, 262
- Limonium*, 105'
- Linaceae, 96', 296"
macroscopic remains, 96'
pollen, 96'
- Linaria*, 108'
- Linnean species, 12, 241
- Linum*, 96'
catharticum, 206, 207, 208, 209
perenne, 97, 133, 188, 202, 206
perenne agg., **221**
perenne ssp. *anglica*, 96
- lithology, 20
- Lithospermum*, 107'
- lithostratigraphy, 264
- Littorella*, 110'
- Lloydia*, 115'
- local communities, 258
habitats, 258
- loess, 168, 207, 230
- Loiseluria procumbens*, 210
- Lomatogonium* type, 106'
- Lonicera*, 110'
xylosteum, 196, 194, **245**
- Lotus*, 95'
- Ludham, 209*
- Lupinus*, 95'
- Luzula*, 115', 216
- Lychnis alpina*, 90, 207, **222**
- Lycopodiaceae, 77', 285"
- Lycopodium*, 77'
annnotinum, 132, 209
- Lycopus*, 108'
europaeus, 212
- Lysimachia*, 104'
- Lythraceae, 99', 297"
- Lythrum salicaria*, 99'
- Mackenzie River, 229
- macro taxa, common, 128*
- MACRO table, 17, 37, **46**, 59, 74, 76
- macroscopic remains, abundance, 135*
frequencies in samples, 133, 134*
representation, 131, 132*, 137
- major biome category, 178*
- mammoth steppe, 201, 254
- man-made habitats, 214
- marine environment, 33
- maritime species, 212
- Marks Tey, 210
- marshes, 212
- Matricaria*, 111', 112'
perforata, 215
- Matthews, J.R., 175
- MCSAMPLE table, 37, **44**, 73
- meadows, 205, 207
- Medicago*, 95'
sativa ssp. *falcata*, 206, **222**
- Megaspores (reworked), 120'
- Mentha*, 108'
- Menyanthaceae, 106', 299"
- Menyanthes*, macroscopic remains, 106'
pollen, 107'
trifoliata, 137, 159, 209, 212, 213
- Mercurialis*, 97'
- Middle Pleniglacial, 232
precipitation, 248
- Midlandian, 17, 69
- migration, 253
biotypes, 242
routes, 173
- Milthers, V., 8, 10
- Minuartia rubella*, 90', **222**
verna, 90', 207, **222**
- miscellaneous taxa, 308"
- Moershoofd Interstadial, 250
- Monocotyledones, 303"
- Montia*, 89'
fontana, 139, 213, 215

- MSAMSTAG table, 37, **49**, 74, 187
Munsterian, 17, 69
 Mutual Climatic Range method, 247
Myosotis, 107'
Myosurus, 92'
Myrica, 85'
 Myricaceae, 85', 287"
Myriophyllum, 100'
alterniflorum, 213
spicatum, 213, 234
verticillatum, 213
 Najadaceae, 114', 304"
Najas, 114'
marina, 214
 Nathorst, A.G., 2
 nectar yields, 162
 Nehring, A., 10, 265
Nepeta, 108'
Nitella, 120'
 nomenclature of plants, 73
 non-British species, taxa, 186, 199
 Non-marine mollusc faunas, 256
 Non-marine mollusc faunas, Devensian,
 256
 Northern Drift, 174
Nuphar, 90', 234
 Nymphaeaceae, 90', 291"
 macroscopic remains, 90'
 pollen, 90'
Nymphaea, 90', 188, 234
Nyssa, 119'
 Odderade Interstadial, 196
Oenanthe, 102'
 Oerel Interstadial, 251
 Öland, 244, 262
 Oleaceae, 106', 299"
 Onagraceae, 100', 297"
Onobrychis, 95'
viciifolia, 188, 193, 206, **222**
Ononis, 95'
 oospores, 120'
Ophioglossaceae, 79', 285"
Ophioglossum, 76, 79'
 Orchidaceae, 123
Origanum, 108'
Osmunda, 79'
 macroscopic remains, 79'
 spores, 79'
 Osmundaceae, 79', 285"
Ostrya type, 86'
 overbank deposition, 29
 oxygen isotope curve, 20
 variation, 227, 228*, 229
 Oxygen Isotope Stage 2, 229, 257
 Stage 3, 253, 258
 Stage 4, 254, 257
Oxyria, 88', 192
digyna, 8, **11**, 88, 160
 palaeobioclimatic term, 8
 palaeolithoclimatic term, 10
Papaver, 92', **222**
 Papaveraceae, 92', 292"
 Paradox tables, 37, 38*, 59
 park-steppe, 265
 park-tundra, 265
Parnassia, 94'
palustris, 137, 212
 Parnassiaceae, 94', 294"
Pastinaca, 102'
 patterned ground, 165, 167*
Pedicularis, 109'
palustris, 207, 209
 perennials, 158
 periglacial area, 17, 21
 sediments, 20
 permafrost, 11, 157, 168, 231
Pettitia, 76
 Phanerophyte percentage, 157
 phenotypic plasticity, 160
 physiological plasticity, 160, 173
 variability, 122
 phytogeographical analysis, Godwin and
 Birks & Deacon, 176*, 179
 Matthews, 175, 176*
 Preston & Hill, 177*, 178*, 179
 phytogeographical groups, 174
Flora Europaea, 183, 185*
 phytogeography, history, 5
Picea, 76, 82', 194, 195, 196
abies (cf.) ssp. *obovata*, 194, 195, 196
abies, 137, 138, 187, 196
glauca, 122
 Pinaceae, 81', 286"
Pinus, 76, 82', 127, 139, 194, 195, 196
 representation, 142, 144*
sylvestris, 211
 plant communities, 192
 dispersal, 229
 PLANT table, 37, **51**, 59, 155, 161
 Plantaginaceae, 109', 301"
Plantago, 109', 132, 216, **223**, 263
coronopus, 214
lanceolata, 162, 206
major, 162, 215
majormedia, 139, 216
maritima, 132, 139, 159, 206, 207, 214,
 262
 PLANTDS tables, 174–188, passim
 PLANTDS1 table, 37, **55**, 59
 PLANTDS2 table, 37, **55**, 59
 PLANTHAB table, 37, **53**, 59, 169, 180,
 198–214 passim
Platanaceae, 94', 294"

- Platanus*, 94'
 plateaus, 166, 197
Pleistocene Epoch, 13, 15
pleniglacial, 20
Pleniglacial, The Netherlands, 8
Plio/Pleistocene boundary, 15
Pliocene, 15
PLSAMPLE table, 37, **44**, 73
Plumbaginaceae, 105', 299"
Poa, 116', 203
polar desert, 165, 166, 138*, 199*, 204
 soils, 205
polar steppe, 8
Polemoniaceae, 107', 299"
Polemonium, 107', 139, 207
pollen, deposition rates, 193
 diagram, stadial, 200
 influx data, 202, 203
pollen/spore taxa, common, 129*
pollen, spores, frequencies in samples,
 138
 representation, 132
POLLEN table, 17, 37, **46**, 59, 74
Polunin, N., 7
Polygonaceae, 87', 288"
Polygons, 232
Polygonum, 88', 159
aviculare, 162, 215
aviculare agg., **223**
bistortaliviparum, 216
lapathifolium, 215
 macroscopic remains, 88'
 pollen, 88'
viviparum, 206
polymorphism, 159
Polypodiaceae, 286"
Polypodiaceae s.l., 80'
 macroscopic remains, 80'
 spores, 80'
pools, 31
Populus, 84'
Porsild, A.E., 160, 229
Portulacaceae, 89', 289"
Potamogeton, 114', 123, 132, 245
acutifolius, 213
alpinus, 213
berchtoldii, 213
compressus, 234
crispus, 213
filiformis, 213, 214
gramineus, 213
natans, 213, 234
obtusifolius, 213
pectinatus, 213
polygonifolius, 213
praelongus, 213
pusillus, 213, 214
vaginatus, 187, 213, 214
- Potamogetonaceae*, 113', 303"
 macroscopic remains, 114'
 pollen, 114'
Potentilla, 94'
anserina, 95, 132, 137, 188, 206, 215, **223**
crantzii, 207
erecta, 162, 209, 212
fruticosa, 207, **223**
palustris, 209, 212
- Praetigian*, 15
Pre-Gortian, 69
Pre-Ludhamian, 17
Pre-Pastonian, 69
pre-Quaternary palynomorphs, 120'
precipitation, 11, 239
Preece, R.C., 174, 256
preservation, 121
Preston, C.D., 174
Primula, 104'
Primulaceae, 104', 299"
Printz, H., 206
Prunella, 108'
PSAMSTAG table, 37, **49**, 74, 187
Pteridium aquilinum, 80'
Pteridophyta, 285"
Pterocarya, 85'
Pupilla muscorum, 256
- Quaternary Era*, 1, 13
Quaternary, oxygen isotope record, 227
 stratigraphy, 6, 13, 16*
 subdivisions of the British Isles, 17, 19
Quercus, 87', 133
- Ran, E.T.H.*, 235
Ranunculaceae, 91', 160, 291"
Ranunculus, 91', 188
acris, 162, 216
Batrachium, 127, 137, 216
cf. aquatilis, 216
flammea, 213
hyperboreus, 187, 213, **224**
lingua, 212
 macroscopic remains, 91'
nemorosus, 72
 pollen, 91'
repens, 72, 162, 207, 215
scleratus, 213
- Raunkiaer, C.*, 155
records, in time, 38
 interpretation, 38
 taxa in time, 187
regional vegetation, 258
Reid, C., 2
relict areas, 266
representation, families, 122
 genera, 122
 identified taxa, 127

- representation, families (*cont.*)
 macro and pollen, 150
 species, 122
- reproduction, 160
- reworked taxa, 211
 fossils, 24, 153
 Quaternary spores, 120'
- reworking, 33, 34*–36, 76, 147
- Rhamnaceae, 97', 296"
- Rhamnus*, 97'
- Rhinanthus*, 108'
- Rhodiola*, 93'
- Rhododendron*, 103', 210
- Rhynchospora*, 119'
- rocks, 207
- Rorippa*, 92'
- Rosaceae, 94', 139, 160, 294"
 macroscopic remains, 94'
 pollen, 94'
- Rubiaceae, 107', 123, 299"
- Rubus*, 94'
chamaemorus, 139, 210
fruticosus, 75, 196
- Rumex*, 88'
acetosa, 162, 206, 211
acetosella, 133, 162, 206, 209, 215
 macroscopic remains, 88'
 pollen, 88'
- Sagavanirktok River, 27–29*
- Sagittaria*, 112'
sagittifolia, 213
- Saiga*, 254
- Salicaceae, 84', 287"
- Salicornia*, 89'
- saline conditions, 205
- salinity, 167, 214
- Salix*, 84', 127, 137, 188, 211, 248
herbacea, 133, 207–211, 224
lapporum, 207
 macroscopic remains, 84'
phylicifolia, 211
polaris, 211, 224
 pollen, 84'
repens, 208, 212
reticulata, 211, 224
viminalis, 211
- salt efflorescences, 205
- Salvinia*, 80'
- Salviniacae, 80', 286"
- Sambucus*, 110'
- samples, 38
 numbers in stratigraphical divisions, 65*
- Sanguisorba*, 94'
minor, 206
officinalis, 132, 206
- Saussurea*, 112'
alpina, 207
- Saxifraga*, 93'
oppositifolia, 122, 160, 188, 225, 207
 distribution, 242
- Saxifragaceae, 93', 294"
 macroscopic remains, 93'
 pollen, 94'
- Scabiosa*, 111'
columbaria, 206
- Scheuchzeria*, 113'
- Scheuchzeriaceae, 113', 303"
- Schouw, J.F., 235
- Schweger, C., 198, 206
- Scirpus*, 118'
lacustris, 213
lacustris ssp. *tabernaemontani*, 214
- Scourse, J.D., 36
- scree, 207
- Scrophularia*, 109'
- Scrophulariaceac, 108', 300"
 macroscopic remains, 108'
 pollen, 109'
- scrub, 211
- Scutellaria*, 108'
- sea levels, 23
- sedge fields, 165
- sedimentary regime, 33
 situations, 25*
 situations, number of sites, 66*
- sediments, 26–33, 168
- Sedum*, 93'
- seed banks, 161
- Selaginella*, 78', 139
 megaspores, 78'
 microspores, 78'
selaginoides, 133, 206, 225
- Selaginellaceae, 78', 285"
- Seward, A.C., 5
- Sher, A.V., 250
- shrub tundra, 165
- shrubs, 211
- Silene*, 90'
vulgaris ssp. *maritima*, 214
- SITECHAR table, 37, 39, 60
- SITELOC table, 37, 39, 60
- sites, 38
 Britain, 64
 distribution, 69
 Ireland, 69
 listed, 62*
 map, 61*
 numbers in stratigraphical divisions, 65*
 numbers in cold stages, substages, 64*
 unpublished, 70
- soil classification, 166
 conditions, 22
 moisture, 193

- soils, 165
 arctic, 166, 192
 brunosolic, 166
 cryosolic, 166
 organic, 165
 polar desert, 205
 regosol, 166
 saline, 212
- Solanaceae*, 108*, 300''
Solanum, 108'
Solidago type, 112'
 solifluction, 32, 165
 diamictons, 230
- Somersham, 34, 76, 168, 211, 217, 218
Sonchus oleracea, 215
Sorbus, 94'
Sourlie, 268
 southern element, 181*, 236
- Sparganiaceae, 116', 306"
 macroscopic remains, 116'
 pollen, 117'
- Sparganium*, 116', 117'
erectum, 213, 216
minimum, 213
- Sparks, B.W., 256
- species, in samples, 125*
 numbers, 124*, 131*
 variable, 159, 161
- Sphagnum*, 119', 209, 210
- Spitsbergen flora, 260
- Stachys*, 108'
- stadial events, 8
 flora, extinction, 261
 flora, origin, 260
- stadial/interstadial differentiation, 154
- Stellaria*, 90'
crassifolia, 225
graminea, 162, 211
media, 162, 214, 215
palustris, 212
- steppe, 8, 205, 235
 element, 235, 236, 244, 265
 fauna, 202
 species, 244
 vegetation, 165
- steppe-tundra, 8, 198, 200, 202, 237
 term, 266
- Stratiotes*, 112'
- Stuart, A.J., 253
- Suaeda*, 89'
maritima, 212
- subarctic, definition, 7
- submarine contours, 22*
- subsidence, 32
- subspecies, 160, 161*
- Succisa*, 111', 139
pratensis, 132, 207, 211, 212
- summer warmth, 235
Sympyton, 107'
- tabula rasa* hypothesis, 5, 261
- tall-herb communities, 205
- Tansley, A.G., 21, 164
- taphonomy, 4, 21, 24, 26, 33, 137, 192
- Taraxacum*, 111', 112'
officinale, 159, 162, 207, 215
- taxa, 4, 38
 sample numbers, 127
 frequencies in samples, 133
 temperate stages, 188*
- listed, 72
 numbers, 131*
 records, 38
 represented by macros and pollen, 130*, 133
- TAXA table, 37, 42, 73, 75
- Taxaceae, 83', 287"
- TAXASORT table, 37, 42, 73, 75, 76, 77, 127, 131, 133
- TAXATIME table, 37, 50, 74, 125, 187, 199
- Taxodiaceae, 83', 286"
- taxonomy, problems, uncertainties, 121
- Taxus*, 35, 83, 139, 140
- Tedrow, J.F.C., 166
- Teesdale relict flora, 262
- temperate term, 248, 264
- temperature changes, 228*
 January, 235, 249
 July, 228, 235, 236, 238, 241, 249
 mean annual, 236
- terminology, 264
- Tertiary, 259
- Thalictrum*, 92'
alpinum, 188, 207
flavum, 207, 212
minus, 206, 207, 208
- Thelypteris*, 76
- thermokarst, 32
- Therophyte percentage, 157
- therophytes, 157
- Thurnian, 69
- Thymus serpyllum*, 240
- Tiglian, 15
- Tilia*, 98'
- Tiliaceae, 98', 296"
- Tolypella*, 120'
- Tottenham, 210
- transfer functions, 3, 236, 256
- Trapa*, 100'
- Trapaceae, 100', 297"
- tree limits, 235
- tree/shrub pollen frequencies, 73, 123
- trees, 211
- Trifolium*, 95'

- Triglochin*, 113'
maritima, 214
- Tsuga*, 82'
- Tubantian (Last) Cold Stage, The
 Netherlands, 8
- tundra biome, 259
- tundra, definition, 7
- tundra/steppe, *see* steppe/tundra
- Turner, J., 262
- tussock grasses, 205
- Type X, 119'
- Typha*, 117'
latifolia, 212, 245
- Typhaceae, 117', 306"
 macroscopic remains, 117'
 pollen, 117'
- Ulex*, 95'
- Ulmaceae, 87', 288"
- Ulmus*, 87'
- Umbelliferae, 102', 248, 298"
 macroscopic remains, 102'
 pollen, 102', 197
- uniformitarianism, 265
- Upton Warren, 196
- Interstadial Complex, 10, 19, 194, 196, 199, 247
 - Interstadial Complex, correlation, 251
 - Interstadial Complex, Coleoptera, 247, 255
- Urtica*, 87', 139, 140
dioica, 132, 162, 211, 225
- Urticaceae, 87', 288"
- Utricularia*, 109'
- Vaccinium*, 103', 210
myrtillus, 195, 196, 210
- Valeriana*, 110'
dioica, 207, 212
officinalis, 132, 206
- Valerianaceae, 110', 301"
- Valerianella*, 110'
- Van Huissteden, J., 235
- Van der Hammen, T., 8
- Vandenbergh, J., 248
- variability, 121
- variable species, 161*, 190
- variation, 158
 - biotypic, 173, 174
 - ecotypic, 159, 173
 - taxa number, 125, 126*
- vegetation, 191
 - arctic formations, 265
 - at particular times, 216
 - interstadial, 193
- local, 24
- mosaic, 198
- productivity, 202
- regional, 24
- structure, 193
- types, 164
- types, arctic, 8
- types of Tansley, 192
- Veratrum*, 115'
- Veronica*, 108'
- vertebrate faunas, 253
 - Devensian, 252
 - *Vicia*, 95'
- Villacher Alps, 244
- Viola*, 98, 216, 226
tricolor, 215
- Violaceae, 98', 296"
 macroscopic remains, 98'
 pollen, 99'
- Viscum album*, 235
- Watt, A.S., 252
- weathering of assemblages, 153
- Webb, D.A., 122
- Webb, T., 227
- Weichselian, interstadials, 228*
 - late-glacial climate, 234, 235, 239
 - late-glacial terms, 265
 - late-glacial vegetation, 265
 - The Netherlands, 232
 - vegetation, 235
 - vegetation changes, 228*
- Wesenberg-Lund, C., 247
- widespread species, 175, 184, 190
- Wilmott, A.J., 10
- wind, 235
- Wing, 104, 210
- Wolstonian, 19, 66
 - Early, 19
 - Late, 19
- woodland, 211
- Woodward, F.I., 238
- Wrangel Island, 202, 257
- Wretton, 35, 195, 217, 218, 255, 261
 - pollen diagram, 200
- Yorkshire Wolds grassland, origin, 262
- Young, S.B., 7, 8
- Zagwijn, W.H., 236
- Zannichellia*, 114'
palustris, 213
- Zannichelliaceae, 114', 304"
- zonobiomes, 234