

Cambridge University Press

978-0-521-19154-8 - Design and Analysis of Long-term Ecological Monitoring Studies

Edited by Robert A. Gitzen, Joshua J. Millsaugh, Andrew B. Cooper and Daniel S. Licht

Index

[More information](#)

Index

- abundance and density monitoring. *See also*
 - capture–mark–recapture; indices of abundance
 - counts with perfect detection, 404
 - distance sampling, 404–405
 - N*-mixture models, 405–406
- adaptability and learning. *See also* adaptive resource management
 - adaptive monitoring *sensu* Lindenmayer and Likens (2009), 29–32, 71
 - addressing changes in target population extent, 66–67
 - building from citizen-science monitoring, 464–465
 - improving and optimizing operational monitoring, 49–50, 444, 452–458
 - iterative learning in structural equation modeling, 343
 - modifying and replacing surveys, 69–71, 492
 - planning for flexibility in survey designs, 18
 - planning for potential changes in variance magnitudes, 247
 - supplementing surveillance monitoring after detecting resource degradation, 495
 - unplanned uses of data and unexpected discoveries, 58–59, 468, 504, 505
 - updating models in model-based sampling, 394, 396
- adaptive resource management (ARM), 74, 76, 501–502. *See also* monitoring and management
 - adaptive harvest management of mallards, 93–96
 - competing predictive models, Bayes' theorem, and learning in ARM, 64, 78–79, 88–90, 91, 94–95
 - double-loop learning, 79, 81
 - environmental stochasticity and, 85–87
 - experiments and learning in ARM, 92–93, 96
 - integrating surveillance and ARM-focused monitoring, 96–97
 - monitoring in the ARM context, 57, 79–82
 - partial controllability and, 77–78, 87–88
 - partial observability and, 83–85
 - structural uncertainty, reducing through ARM, 76, 78–79, 80–81, 88–93
 - structured decision making in ARM, 79
- adaptive sampling, 382–384, 387–389
 - adaptive cluster sampling, 383–384
 - adaptive stratified sampling, 385, 389
 - GIS adaptive sampling, 385, 388, 394
 - performance, adaptive vs. non-adaptive designs, 391–394
 - simulations to assess sampling designs, 384, 387, 389–391
 - software for, 389, 395
 - trend detection and adaptive sampling, 383
- allocation of effort and resources. *See also* trade-offs; stratified sampling (allocation of effort)
 - among alternative monitoring programs, 498–499
 - among vs. within sample sites, 46, 402
 - between components of hybrid sampling designs, 368–369
 - between established and new monitoring programs, 505
 - for data analysis, 35, 258
- analysis (data analysis), consideration of during monitoring design, 14, 26, 29, 32, 35, 175–176, 492
- autocorrelation. *See* correlation
- autoregressive model, 171, 330
- Bayes' Theorem, 89, 91
- Bayesian inference, 295. *See also* BUGS; hierarchical modeling; Markov chain Monte Carlo
 - estimation of variance components, 223–224
 - incorporating prior data, 464
 - model selection issues, 440–441
 - structural equation modeling, 338
 - threshold modeling, 279–280, 287–288
 - with missing data, 311–312
- Before–After, Control–Impact (BACI) design, 58, 175

Cambridge University Press

978-0-521-19154-8 - Design and Analysis of Long-term Ecological Monitoring Studies

Edited by Robert A. Gitzen, Joshua J. Millsaugh, Andrew B. Cooper and Daniel S. Licht

Index

[More information](#)

- bias, 15, 25, 30. *See also* frame errors; detectability; measurement error
- bias–variance trade-off, 68–69
- changes in detection ratio, 262–264
- evaluating potential biases affecting trend analyses, 261–265, 469
- identifying and addressing potential sources of during survey planning, 15–17
- sources of bias, 51
- block designs, 192–195, 429
- BUGS, 430, 441
 - capture–mark–recapture estimation, 432, 439, 441–442
 - deviance information criterion (DIC), 440
 - spatio–temporal modeling, 450
 - threshold modeling, 280, 287–288
 - variance component estimation, 213
 - via *R*, 213, 287–288, 432
- calibration of alternative methods, 70
- CAPTURE (software program), 432
- capture–mark–recapture, 406–413
 - assumptions, 406–409
 - closed populations, 406, 409–410
 - Cormack–Jolly–Seber model, 412
 - density estimation, 410, 439
 - heterogeneity, 408–409, 410–411, 431, 435
 - hierarchical modeling, 427–429, 430–433, 436, 438, 439–440
 - joint vs. separate modeling of detectability and abundance, 432, 438–439
 - Lincoln–Petersen method, 193, 409
 - mis-identification and tag loss, 408
 - multi-state (multi-strata) models, 413, 415
 - open populations, 406, 411–413, 415–416
 - recruitment and reproduction estimation, 414–415
 - simulations, comparison of hierarchical trend models, 436
 - spatially explicit models, 410, 413, 439
- chain index analysis method, 301
- changepoint models. *See* threshold modeling
- citizen-science monitoring
 - assessing appropriateness of during program design, 462–464, 503
 - data quality, 463–464, 466–467
 - integrating targeted and citizen-science surveillance monitoring, 476
 - supervision and coordination, 465
 - survey design constraints and trade-offs, 64, 464, 468, 472
 - value
 - for designing subsequent studies, 464–465, 468
 - for large-scale monitoring, 462
 - for public education and engagement, 503
 - of data for exploratory analysis, 467–468
 - volunteer motivation, 464, 465–466
- cluster sampling, 112–114, 274–277
- community-level attributes, monitoring, 400, 401
- conceptual models, 32, 489
- Conditional Autoregressive (CAR) model, 448–449
- conservation and monitoring. *See* management and monitoring; adaptive resource management
- convenience sampling, 9, 105, 401–402
- correlation. *See also* variance components (correlation coefficients)
 - addressing correlations (non-independence) in trend analysis, 471–472
 - autocorrelated data and cross-validation, 474
 - autocorrelation effects on Type I error, 451–452
 - correlation parameters in linear mixed-effects trend model, 158–159
 - design-based inference and spatial correlation, 421
 - non-independence in clustered (grouped) data, 202
 - spatial autocorrelation and occurrence studies, 421–422
 - spatio-temporal modeling of, 448–449, 451–452, 458
- cumulative distribution functions (CDFs), 141–142
- comparison and hypothesis testing, 322–324
- estimation, 313, 316–319
- management thresholds and CDFs, 314
- measurement error and deconvolution, 319–322
- panel designs, CDF estimation with multi-year data, 318
- percentile estimation, 319
- R* software, CDF estimation and testing, 317–319, 322, 323
- data management, 491
 - consideration of during planning, 72
 - in citizen-science monitoring, 465–467
- deconvolution of measurement error, 320–322
- density estimation. *See* abundance and density; capture–mark–recapture: density estimation
- design-based inference, 41, 133, 244. *See also* Horvitz–Thompson estimator
- choosing, vs. model-based inference, 42–43, 423–424
- trend analysis constraints, 153
- detectability, 63–64, 84, 90, 176, 177, 215, 260–261, 263–264, 265, 426–427, 438, 458. *See also* abundance and density; capture–mark–recapture; occurrence; indices of abundance and adaptive sampling, 394–395
- counts with perfect detection, 404
- covariates to account for variation, 465

- hierarchical modeling, 427–429, 430–433, 436, 438, 439–440
- importance of considering, 401, 426–427
- in citizen-science monitoring, 472
- occurrence studies, 416–417
- recruitment and reproduction studies, 414, 415
- within-season heterogeneity, 408–409, 410–411, 431
- Dirichlet tessellation, 142
- distance sampling, 404–405
- domain (subpopulation) analyses, 42–43, 323
- dynamical models, 94–95, 443, 448, 458
- error. *See* total survey error; bias; variance components; sampling error; measurement error
- factor models, 332
- Forest Inventory and Analysis (FIA) Program, US Forest Service, 31–32, 33, 37, 43, 45, 46–47, 49–50
- FRAGSTATS, 365
- Gaussian Markov random field, 448–449
- General Random Samples (GRS), 117–119
- R function, 118, 122–123
- Generalized Random Tessellation Stratified (GRTS)
- design, 101, 116, 117, 118, 131–140, 237, 373–374, 490
- algorithms, 132–139
- analysis in R, 317–319
- and revisit (panel) designs, 147
- evaluating spatial balance, 143–145, 146
- for linear networks and polygons, 138–139
- for point resources, 132–138
- GRTS with stratified sampling, 140
- GRTS with unequal probability sampling, 140
- over sample for adding and replacing sample units, 146, 245
- R implementation, 132, 138, 140, 146, 148, 149–150
- reverse hierarchical ordering, 146
- grid sampling. *See* systematic sampling
- growth-curve trend model, mixed-effects model vs. structural equation modeling, 332
- haphazard sampling, 105
- hierarchical modeling. *See also* variance components
- abundance and detectability, modeling, 427–429, 430–433, 436, 438, 439–440
- data augmentation, 430–431
- experimental treatment effects, modeling, 431
- model selection, 440–441
- random effects variance component model, 201–203
- simulation comparison, mark–recapture and trend modeling, 437
- spatio-temporal trend analysis, 447–452
- survey design optimization, 453–454
- threshold modeling, 296–297
- trend analysis, 302, 331–332, 435, 436
- hockey stick model, 284–285. *See also* threshold modeling
- Horvitz–Thompson estimator, 119, 134, 315–316, 317–319, 389
- extension to continuous population, 316
- hybrid monitoring
- surveillance and targeted monitoring, 96–97
- hybrid survey designs 372
- combining probability and model-based sampling, 362–365
- probability sampling plus purposely chosen sites, 64
- supplementing surveillance monitoring with targeted monitoring, 58
- imputation, 300. *See also* missing data
- inclusion probabilities, 102, 117, 119–120, 315–316. *See also* probability sampling
- second-order, R function, 123–125
- indices of abundance
- analysis methods and missing data, 300–302
- incorporating covariates related to variable detectability, 401, 472
- integrated nested Laplace approximations (INLA), 450–451
- JAGS, 280, 287–288, 450
- judgment sampling, 9, 40–41, 64, 104–105, 128, 491
- Kolmogorov–Smirnov statistic, 141
- learning. *See* adaptability and learning; adaptive resource management
- local neighborhood variance estimator, 133, 160
- machine learning, 471, 474, 475–476
- management and monitoring, 381–382, 383, 399–400, 413, 426, 440. *See also* adaptive resource management
- ecological thresholds and management, 281
- effective partnerships with managers in monitoring, 493
- inadequate monitoring design, management and conservation costs of, 26
- integrating monitoring results into natural resource planning, 494
- management thresholds, 314
- management triggers, 62, 72
- state-dependent decisions, 79–80, 499–501

Cambridge University Press

978-0-521-19154-8 - Design and Analysis of Long-term Ecological Monitoring Studies

Edited by Robert A. Gitzen, Joshua J. Millsbaugh, Andrew B. Cooper and Daniel S. Licht

Index

[More information](#)

- MARK (software program), 432, 438–439, 440
- Markov chain Monte Carlo (MCMC), 450, 453–454
in BUGS, 430, 432, 436
- mark–recapture. *See* capture–mark–recapture
- MaxEnt, software program, 396
- maximum entropy modeling, 386, 396
- measurement error, 260–261, 264. *See also*
detectability
- effects and deconvolution in cumulative
distribution function (CDF) estimation,
319–322
- incorporation into variance expressions, 178–182
- mechanistic models, 94–95, 443, 448, 458
- Microsoft Excel. *See* spreadsheets, analysis with
- missing data, 338–340
and analysis of abundance indices, 300–302
Bayesian analysis, 311–312
choosing among strategies for handling, 310–311,
339–340
complete case method (casewise deletion), 303,
339
imputation methods, 339–340
description, 303–306
simulation comparisons, 306–310
missing at random (MAR), 302, 339
missing completely at random (MCAR), 302,
339
multiple imputation, 306
non-ignorable non-response (missing not
completely at random, MNAR), 302–303,
311, 339
structural equation modeling and, 339–340
- MLPowSim, 219
- MLwiN, 219, 220
- model evaluation and selection, 423
assessing model credibility in adaptive resource
management, 88–90
cross-validation, 473–474
hierarchical model selection, 440–441
in structural equation modeling, 337–338,
341–357
in threshold modeling, 285–287
validation, 473–474
- model-assisted sampling and inference
and hypothesis-driven monitoring, 394,
396
in trend analysis, 153
incorporating predicted species distribution,
384–385, 387–389, 396
iteratively updating models, 394
model inaccuracy, 391–394
- model-based inference, 41–42. *See also*
model-based sampling
choosing, vs. design-based inference, 32, 42–43,
423–424
in demographic monitoring, 422
- model-based sampling, 41, 362, 366
based on graph-theory model of habitat
connectivity, 366–368, 375–377
combining with probability sampling, 362–365
- modeling, statistical. *See* statistical modeling
- models, conceptual, 16. *See* conceptual models
- monitoring
benefits of, 499–504
definitions of, 54, 74
- monitoring programs, general design and
implementation issues. *See also* trade-offs
adaptive resource management as a framework
for monitoring design, 81–82
attributes and measurements, selection of, 36–37,
81–82, 483–487
deciding when to end a monitoring program, 499,
502
effective reporting of results, 492
elements of well-defined objectives, 29
s.m.a.r.t. objectives, 28
importance of clearly defined purposes and
objectives, 14, 20, 28, 56, 397–399,
483–485
institutionalizing a monitoring program,
483
monitoring design and whole-problem planning,
23–26
retrospective vs. prospective focus, 399
statistical thinking and statisticians, key roles of,
5–6, 9–10, 13–14, 20, 23–25
sustainability, 62–63, 507
value and limitations of trend-focused
monitoring, 60–62, 151, 399–400
- Mountford index analysis method, 301
- Mplus, 220, 344, 356
- multiple scales and attributes, monitoring for, 33,
43, 58, 107, 108, 171, 362, 364, 372, 386,
419, 489
hierarchical threshold modeling of multiple
attributes, 296–297
- multi-stage sampling, 108, 109–110, 185, 364
cost functions, 182, 189
variance expressions and estimation, 178–182,
185–189, 192–193, 198
- National Park Service, US, Inventory and
Monitoring Program, 10, 233–234, 370–373,
482–483
commonly selected indicators (attributes) for
monitoring, 488
- nested sampling. *See* multi-stage sampling
- N*-mixture models (spatially replicated counts),
405–406
- nonparametric models, 471
- non-response due to logistical limitations and loss of
access, 64–65, 113, 146

- North American Breeding Bird Survey (BBS), 63, 64, 254, 262–263, 271, 298, 301, 302
as surveillance monitoring, 58, 59
trend examples, 254–259
- objectives. *See* monitoring programs, general design and implementation issues
- observer impacts, 167, 237
- occurrence and occurrence dynamics, 400–401, 416–423, 472
design
sample units, 421
sampling season, 422
target population, 421
effort allocation, 402
integrated monitoring of occurrence and habitat dynamics, 419
monitoring at multiple scales, 419
spatial autocorrelation, issues in occurrence studies, 421–422
- Optimal Design (software program), 219
- optimizing survey design, 444, 452–458
design criteria, 452–453, 454, 458
incorporating costs, 455–457, 458
- paired designs, 192–195
- panel design. *See* revisit design
- percentile estimation, 319
- piecewise linear model, 284–285. *See also* threshold modeling
- pilot studies, importance of, 45–47, 176, 197, 440
- PINT, 219
- population growth rate, estimating, 400, 404, 410, 415–416
- post-stratification, 111
- power, precision, and sample-size examinations, 9, 12–13, 14–15, 17
components of, 48–49
importance of quantitative survey design, 47, 197–199
power for trend detection, 161, 165–167, 195–196
- quantitative sampling objectives, 6, 241
as basis for examinations, 48
consideration of management decision-making, 85, 91, 501
software programs and code for, 49, 169, 196, 219–220, 241
using simulations to assess performance of alternative survey designs, 240–244
- probability (probabilistic) sampling, 41, 104, 129, 315, 362, 402, 490. *See also* model-assisted sampling
advantages and importance of, 102, 104
and design-based inference, 41
and model-based inference, 42
equal probability, 102–104, 132–139
model-assisted sampling, 387–389
non-probability designs, 104. *See also* judgment sampling; convenience sampling
unequal probability, 104, 113, 119, 140, 220, 317, 491. *See also* probability proportional to size sampling
probability proportional to size (PPS) sampling, 381, 385, 388
- process models. *See* mechanistic models
- purpose of monitoring. *See* monitoring programs, general design and implementation issues
- R**
- adaptive cluster sampling, 395
- cumulative distribution functions (CDFs)
estimation and testing, 317–319, 322, 323
- General Random Sample function, 118, 122–123
- Generalized Random Tessellation Stratified (GRTS) design, 132, 138, 140, 148, 149–150, 317–319, 378
- integrated nested Laplace approximations (INLA), 450–451
- local neighborhood variance estimator, 133, 168, 318
- posterior simulations, 292
- second-order inclusion probabilities calculation, 120, 123–125
- simulations, 49, 248, 433
- status estimation, 244, 378. *See also* R: Cumulative Distribution Functions (CDFs)
estimation and testing
- structural equation modeling (SEM), 344
- testing for autocorrelation, 471
- threshold modeling, 287
- variance component estimation, 168
- R packages**
- INLA, 450–451
- R2WinBUGS, 213, 432
- rv, 292
- segmented, 287
- spsurvey, 132, 133, 138, 140, 149–150, 168, 237, 244, 317–319, 322, 323
- WiSP, 49
- random sampling. *See* simple random sampling, probability sampling
- Random Tessellation Stratified (RTS) design, 131, 162
- ratio estimator, 319
- recruitment and reproduction, estimating, 413–415
- representative sampling, 127–132
and analysis of citizen-science data, 469
non-representative sample, 64–65, 68–69, 105, 262–263. *See also* sampling frame: frame error

- representative sampling (*cont.*)
 - representative sample vs. representative sampling process, 129
 - spatial balance and representativeness, 130–132, 141
- response (measurement) design, 45–47
 - index window, 152, 422
 - sample unit size and shape, 45
- revisit design, 44–45, 154–156, 491
 - always revisit (pure panel) design, 154
 - analysis and power examinations for panel designs, linear mixed-effects model framework, 156–160, 165–167, 170
 - and spatial balance, 147
 - augmented serially alternating panel designs, 156
 - balancing trend-detection and status-estimation objectives, 44, 165–167
 - connected vs. unconnected designs, 156
 - cumulative distribution function estimation with panel designs, 318
 - effect of missed visits on power, 170
 - in demographic monitoring, 423
 - never revisit design, 154
 - rotating panel designs, 154
 - effect of temporal correlation on optimal rotation, 182–183
 - serially alternating panel designs, 45, 156
 - split-panel designs, 235, 237
- route regression, 301–302
- sample (sampling) frame, 39, 40, 234. *See also* target population
 - frame errors, 39–40, 66–68, 259–260, 264
 - due to changes in extent and phenology of target population, 40, 44
 - list frame, 368
- SAMPLE (software program), 389
- sample unit, 102
 - defining in occurrence monitoring, 421
- sampling design, 37, 38–44, 101, 401–402. *See also* representative sampling; survey design; and individual sampling methods
 - desired attributes for monitoring designs, 106–107, 114
 - for rare and clustered species, 381, 391, 394. *See also* adaptive sampling
 - inadequate designs, potential causes of, 7–10
 - over sample for adding and replacing sample units, 146
 - scientific (probability sampling) vs. unscientific designs, 102–105
 - spatially explicit designs, 127
- sampling distribution, 25
- sampling error, 51, 278
- SAS
 - GLIMMIX, 213
 - MIXED, 206, 239
 - power calculation, 169
 - trend analysis, 167, 239
 - variance components estimation, 167–169, 213, 239
- segmented regression. *See* piecewise linear model
- selection bias, 128, 260
- semi-parametric models, 469–471
- simple random sampling, 109, 119, 160
 - comparison, vs. stratified and adaptive sampling, 391–394
 - evaluating spatial balance, 143–145, 146
 - over sample for adding and replacing sample units, 146
- simulations
 - assessing performance of alternative survey designs, 240–244
 - assessing statistical performance of hierarchical mark–recapture trend models, 436
 - assessing statistical performance of variance components estimators, 203–204
 - comparing adaptive and conventional sampling designs, 384, 389–391
 - comparison of missing-data imputation methods, 306–310
 - evaluating spatial balance of sampling designs, 143–145, 146
 - simulation approach, estimating variance components on inverse-link (measurement) scale, 224–227
 - system simulations to predict future variability, 234–235, 237–238
 - utility for survey-design planning, 229
- small area estimation, 43
- SMART (multi-attribute weighting) ranking of sampling designs, 391
- software. *See also* R; SAS; and other software program names
 - and accessibility of methods, 277, 441
 - for survey-design simulations, 247–248
 - online data entry system, 467
- spatial balance (coverage), 114. *See also* Generalized Random Tessellation Stratified (GRTS) design
 - and revisit (panel) designs, 147
 - evaluating, 143–145, 146
 - evaluation metrics, 141–143
 - geographic, 148
 - simple random sampling, 143–145, 146
 - through representative sampling, 130–132, 141
 - with dynamic sample-size adjustments/ over sample, 145–146
 - with frame errors, site inaccessibility and uncertain feasible sample size, 116, 120–121
- spatial design. *See* sampling design

- spreadsheets, analysis with, 272–273, 275–277
- Stata, 220
- statistical modeling. *See also* hierarchical modeling;
model evaluation and selection; structural
equation modeling; trends and temporal
changes
- Bayesian estimation, 223–224
- choosing among parametric, semi-parametric, and
nonparametric models, 469–471
- dynamical process vs. descriptive models, 272,
443–444
- generalized additive models, 269–270,
470
- generalized linear and generalized linear mixed
models, 210–211, 217, 268–269, 301–302,
430, 431, 448
- hierarchical vs. two-step analysis, 427–428
- linear mixed-effects model framework for trend
analysis, 156–160
- linear models, 267
- maximum likelihood estimation, 222–223
- mixed effects models, overview, 270–271
- model specification, 265–267, 280
- multi-model inference, 470
- parameter (model) identifiability, 336–337
- scientific model formulation and statistical
inference in threshold modeling, 282–283,
294–295
- status estimation, 152, 313, 315–316
- abundance indices and missing data, 300–302
- cumulative distribution function (CDF), 316–319,
320–322
- step function model, 284. *See also* threshold
modeling
- stratified sampling, 108, 110, 111–112, 113,
129–130, 140, 260, 318, 381, 445–446,
447
- adaptive stratified sampling, 385, 389
- allocation of effort, 111, 130
- and management decision models, 85–86
- comparison, vs. simple random and adaptive
sampling, 391–394
- example analysis, 274–277
- modeling spatial correlation among strata,
448–449, 451
- stratifying based on model predictions, 385,
388–389
- stratifying on non-permanent features, 65–66,
112, 491
- structural equation modeling (SEM)
- assessing model fit, 337–338, 341–342, 346–347,
348, 354
- autoregressive cross-lagged model, 331
- autoregressive latent trajectory model, 335,
354
- autoregressive model, 330, 354
- Bayesian (predictive probabilistic) networks and
SEM, 357
- Bayesian approach, 338
- causal and theory-driven SEM, 325, 329, 341,
343–357
- graphical representation, 326–328
- latent trajectory model, 331, 332–333, 335, 354
- latent variable framework, 328–329
- learning to apply, 341, 357
- maximum likelihood estimation, 338
- missing data, 339–340
- full information maximum likelihood
approach, 340
- model comparison and selection, 341–343
- modeling nonlinear trajectories, 333–334
- observed-variable models, 326–328
- of temporal changes, 329–336, 343–357
- parameter (model) identifiability, 336–337
- software for, 344
- structured decision-making and monitoring, 57. *See
also* adaptive resource management
- subpopulations, estimates for, 42–43
- surveillance monitoring, 97, 364–365, 399–400
- benefits and limitations of, 58–60, 82
- survey design, 38, 489–491. *See also* sampling
design; revisit design; response design
- documenting, 17–18
- evaluating designs with simulations, 384, 387,
389–391
- importance of statistical planning, 37–38
- incorporating costs, 455–457
- optimizing, 444, 452–458
- survey design (sampling) weights, 315, 316, 319
- recommended maximum range, 318
- unequal weights and variance-component
estimation, 220
- survival estimation
- band recovery models, 412
- capture–mark–recapture, 412–413
- systematic sampling, 43, 114–116, 119, 131, 420
- target population (target universe), 38–40, 102, 151,
234, 315
- biological (resource) vs. statistical, 38–39, 421
- changes in extent, location, and phenology, 44,
66–67, 421
- in occurrence studies, 421
- types of spatial populations, 126, 127, 143,
313–315
- targeted monitoring, 57, 82, 394. *See also* adaptive
resource management
- thresholds and threshold (changepoint) modeling
- alternative threshold models, 279
- distinguishing threshold vs. smooth changes, 286
- general model, 283, 285
- hierarchical modeling, 296–297

Cambridge University Press

978-0-521-19154-8 - Design and Analysis of Long-term Ecological Monitoring Studies

Edited by Robert A. Gitzen, Joshua J. Millsaugh, Andrew B. Cooper and Daniel S. Licht

Index

[More information](#)

- thresholds and threshold modeling (*cont.*)
 management applications of threshold modeling, 281
 management thresholds, 314
 maximum likelihood estimation, 287
 model evaluation and selection, 285–287
 nonparametric deviance reduction (CART) approach, 288
 overview of alternative models, 279–280, 285
 total survey error, 46, 51
 trade-offs, 75. *See also* allocation of effort and resources
 balancing trend-detection and status-estimation objectives, 44, 165–167
 competing management objectives, 77
 detectability vs. indices of abundance, 63–64
 investing in monitoring vs. management actions, 502
 long-term vs. effectiveness monitoring, 494–496
 methodological improvements vs. data consistency, 492
 quality vs. quantity, 403, 462, 468
 structured optimization of monitoring effort, costs vs. information value, 93
 with citizen-science monitoring, 462, 464, 468
 trends and temporal changes, 152–153
 autoregressive model, 171, 330
 consideration of during program and survey design, 33–35
 defining and describing trend, 60–62, 254–259
 elapsed years vs. power and precision, 169, 437
 exponential model, 256
 hierarchical modeling, 302, 331–332, 436, 447–452
 incorporating covariates and hypothesized change mechanisms into analyses of, 34, 169, 334–335
 key literature references, 154, 253–254
 linear mixed-effects model framework for trend analysis, 156–160, 171
 log-linear model, 301–302, 435, 448
 modeling interacting trajectories, 331
 modeling nonlinear trajectories, 333–334
 power for trend detection, 161, 165–167, 195–196
 route regression, 301–302
 SAS linear model code, 167
 simulations, hierarchical mark–recapture trend modeling, 436
 structural equation modeling (SEM) of, 329–336, 343–357
 threshold modeling, 296–297
 variability in trends across sites, 157, 159–160, 165, 170–171
 two-stage sampling. *See* multi-stage sampling
 Type I and Type II errors, 30
- unequal probability sampling. *See* probability sampling
- variability, sources of
 defining during survey design, 176–178
 estimates of variance components for stream physical variables in Oregon USA, 162–165
 relative impacts on power for trend detection, 165–167
 variance components, 201, 320
 ANOVA estimation, 167–169, 187, 221–222
 Bayesian estimation, 223–224
 correlation coefficients
 intraclass correlation coefficients, 194, 208
 measurement error and, 184–185
 distributional assumptions and estimation, 168, 208
 estimation of variance components on
 inverse-link (measurement) scale
 latent variable approach, 225
 simulation approach, 224–227
 estimation with small sample sizes, 204, 207
 identification and estimation based on
 fundamental variance expressions, 178–182
 linear model framework for defining, 156–160, 167–169, 176–177, 198, 201–203, 204
 maximum likelihood estimation, 222–223
 restricted (residual) maximum likelihood (REML) estimation, 223, 239
 SAS code for estimation, 167–169
 scientific and management uses of, 200
 sources of information for assessing, 48, 153–154, 197
 statistical properties of estimators
 binomially distributed/categorical data, 211–213, 214–215
 count data, 217–218
 normally distributed/continuous data, 204–209
 uncertainty in estimates of, 16, 191, 207
 using system simulations to predict future variability, 234–235, 237–238
 variance partition components (VPCs), 208–209
 variance expressions
 covariance in stages, 183
 general expression, variance in stages, 178, 179, 180
 log-normal data, 196
 trend parameter in linear model framework, 157, 159, 160
 volunteer monitoring. *See* citizen-science monitoring
- waterfowl, North American monitoring, 61–62, 93–94, 96, 444–446
 WinBUGS. *See* BUGS