

## INDEX

- abscisic acid (ABA), 29, 63, 77, 98,  
 129, 209, 265, 283  
 [<sup>14</sup>C], 170  
 diurnal variation, 171  
 effect on wilted *Pisum* leaves, 171  
 field-grown plants, 172  
 and guard cell ion movements, 66, 67  
 production and leaf turgor, 172, 173  
 responses and CO<sub>2</sub>, 130  
 role in drought, 174–8  
 and root and shoot growth, 177  
 and root water uptake, 175  
 site of synthesis, 168  
 and stomatal behaviour, 168–73  
 stomatal closure, 38, 97, 107, 167, 170  
 and stomatal dependence, 38  
 and water stress, 137, 167–73, 177, 178  
 as water stress substitute, 173, 174
- Acer saccharum*, 141
- Aeonium glutinosum*  
 stomatal opening anomaly, 139
- Aeonium spathulatum*  
 stomata and humidity, 139
- Aeonium urbicum*  
 stomata and humidity, 139
- age and stomatal response, 233
- aerodynamic conductance, 226
- Aichryson dumosum*  
 hairyness, 155, 156  
 stomata and humidity, 139, 155
- Aichryson laxum*  
 hairyness, 156  
 stomata and humidity, 139, 155, 156
- Aichryson villosum*  
 hairyness, 155, 156  
 stomata and humidity, 139, 155
- alanine aminotransferase, 89, 90  
 in guard cells, 94
- Allium cepa*, 22  
 anions in stomata, 76, 77  
 Calvin cycle in epidermis, 95  
 chloride in guard cells, 99  
 chloroplast fluorescence, 110  
 guard cell metabolism, 98  
 guard cell thickening, 3–5  
 guard mother cell division, 2  
 malate accumulation, 77, 99; role of, 99  
 reserve carbohydrates, 76, 98  
 vacuolar fluorescence, 113  
 wall deposition and colchicine, 16
- Allium porrum*, 99
- Alternaria tenuis*  
 toxin and stomatal behaviour, 211, 212
- Amaranthus powelli*, 263
- amphistomatous leaves, 37  
 porometer clamps, 43
- Anemia rotundifolia*  
 cytoplasm appearance and stomatal  
 opening, 80
- anions  
 guard cell cytoplasm and organelles, 80,  
 81  
 light and CO<sub>2</sub> effects on formation, 130–3  
 organic and potassium change, 53  
 in stomatal operation, 71–81
- antitranspirants, *see* ABA and farnesol
- Aphanomyces cochlioides*  
 sugarbeet wilt, 208
- Arachis hypogaea*, 230
- argon, 39
- aspartate, 79  
 epidermal, 93  
 aspartate aminotransferase, 88–90  
 in guard cells, 94
- ATP, guard cell production, 123
- auramine O, cuticle fluorescence, 20, 21
- Avena sativa*, 263, 283  
 ozone response, 188, 189
- benzo-18-crown-6  
 potassium binding, 129
- Betula verrucosa*  
 stomatal response to ABA and water  
 stress, 174, 175
- bromine, [<sup>82</sup>Br], 65, 66
- buffer and guard cell studies, 29
- Calluna vulgaris* microclimate, 155

## 288 INDEX

- Calvin cycle  
 absence and stomatal function, 95, 100  
 and guard cells, 93–7
- canopy conductance, 224–9  
 calculation in millet, 226, 227, 231, 232  
 components, 225, 226  
 crop profiles, 227, 229, 232, 233  
 diurnal variation, temperature and  
 humidity effects, 237  
 and drought, 235, 236  
 factors affecting in wheat, 226–9, 233  
 glasshouse experiments, 235, 239, 240  
 and leaf area, 234, 235  
 radiation relationship, 225, 229–31  
 and shading, 232
- Capsicum annuum*  
 ABA accumulation and water, 172, 173,  
 177
- carbon  
 [<sup>14</sup>C]labelling studies, 89, 90  
 tissue isotope discrimination, 96, 97
- carbon dioxide  
 and ABA, additive effects, 178  
 concentration and stomatal conductance,  
 249, 266–72  
 concentration and stomatal control of  
 photosynthesis, 261–5  
 -free air, stomatal opening, 119, 120, 123,  
 142; potassium effect, 125–30  
 and limits to potassium uptake, 126, 128  
 sink, 248  
 and stomatal opening: effect of humidity  
 and temperature, 143; effect of light,  
 260; effect of water stress, 173, 174  
 stomatal response, 119, 124  
 stomatal sensitivity to, 263, 269; and  
 light, 266–72
- carbon dioxide fixation  
 in disease, 218, 219  
 guard cell, 95; and photosynthesis, 112,  
 113  
 and stomatal movement, 92, 93
- carbonic anhydrase, 94
- Carex* sp., 14  
 guard cell wall fluorescence, 21
- Carex aquatilis*  
 humidity response, 155
- cations in guard cell cytoplasm, 80
- cellulase and cell wall ester release, 19
- cellulose  
 axial deposition, 9  
 pattern of deposition in guard cells, 3, 9
- cellulysin, 104
- chloride  
 entry inhibition, 59  
 and malate accumulation in guard cells,  
 72, 73, 77, 130  
 and potassium balance in guard cells,  
 51–3, 56, 130
- Chlorophytum comosum*  
 low temperature emission spectra, albino,  
 110, 112
- chloroplasts  
 ABA content and water stress, 170  
 ABA synthesis, 168  
 fluorescence of guard cell, 110–13  
 guard cell photosystems, 110, 112
- citrate, 53, 72
- Citrus* sp.  
 stomatal humidity responses, 139
- clamps  
 detachable porometer, 36  
 double hinged, 41  
 leaf attachment, 35, 36  
 and leaf type, 43  
 and mesophyll resistance, 35, 36  
 ventilated leaf disc, 36
- cobalt chloride papers and stomatal  
 conductance, 40
- cobaltinitrite in K<sup>+</sup> analysis, 12, 16, 52
- colchicine, 2  
 effects of, 5  
 and wall thickenings, 15
- Commelina benghalensis*  
 pyruvate P<sub>i</sub> dikinase, 92  
 starch and stomatal opening, 78
- Commelina communis*, 52, 92, 128, 263  
 anions in stomata, 77, 78  
 carbon dioxide sensitivity, 268–70  
 CO<sub>2</sub> fixation in epidermis, 93, 97  
 epidermal ions, 56  
 epidermal malate, 56, 57  
 epidermal membrane potentials, 61  
 epidermal potassium, 55  
 fusicoccin and stomata, 132, 133  
 leakage from epidermis, 98  
 light and stomatal opening, 122, 268  
 malate labelling and stomatal movement,  
 97  
 malate uptake and stomata, 91  
 potassium and stomatal aperture, 64, 65,  
 77, 125–7  
 recent experimental work, 62–7
- Commelina cyanea*, 78  
 anions in stomata, 77, 78  
 aspartate, 89  
 CO<sub>2</sub> fixation and stomatal movement,  
 92  
 epidermal enzymes, 88–90, 92  
 epidermal leakage, 97  
 epidermal malate, 56, 57  
 epidermal potassium, 55
- Commelina diffusa*, 78  
 [<sup>14</sup>C] labelling of epidermis, 89, 90  
 conductance measurements: *see also* canopy  
 conductance, stomatal conductance  
 canopy, 224–9  
 in field experiments, 223–41

- Corylus avellana*, 144  
 p-coumaric acid ester, 19  
 crassulacean acid metabolism, 76, 140  
 and stomatal sensitivity to humidity, 140  
 and stomatal opening, 140  
 cryptochrome, 273  
 C-tumours, 17  
*Cucurbita pepo*, SO<sub>2</sub> effects, 194  
*Cucurbita sativus*, SO<sub>2</sub> effects, 194  
 culture  
 grass leaf-segments, 11–13  
 inadequacy, 13  
 cuticle  
 fluorescence, 20, 21  
 and pathogens, 213  
*Phleum* guard cell, 20, 21  
*Pinus* stoma, 254  
 and vapour loss, 43, 150  
 cutinase, 21  
 cuvettes, 238  
*Cyperus esculentus*  
 field grown stomatal complex, 14, 15  
 guard cell wall fluorescence, 21, 281  
 laboratory stressed, stomatal complex,  
 14–16  
*Cytisus scoparius*  
 windspeed sensitivity and distribution,  
 155  
 cytochalasin B, 2
- darkness and potassium uptake, 126, 128  
*Deschampsia caespitosa*, 155  
*Desmodium uncinatum*: humidity  
 insensitivity, 140, 145  
 diamino-benzidine osmium staining, 19  
 diferulic acid ester, 19  
 diffusive gas flux, 38, 39  
*n*-dodecane, 32  
 driselase, 104  
 drought  
 effects on conductance in tobacco, 235,  
 236  
 slow field development, 238  
*Dupontia fischeri*  
 restricted distribution and humidity  
 response, 155
- electron microprobe analysis, 53, 54, 60  
 electron microscopy  
 scanning, grass stoma, 6; *Vicia* stoma and  
 SO<sub>2</sub>, 195  
 stomatal complex, 6  
 endocytosis, 22  
 energy requiring pump-leak system, 273  
 environment, effects of variability, 229, 230  
 enzymatic cycling techniques, 73  
 epidermal strips, 27–32, 52  
 experimental uses, 28, 29, 62  
 and flux measurements, 62, 72  
 intact, 64  
 isolated, 64, 65, 72  
 and low pH, 29  
 low temperature emission spectra, 110  
 mounting medium, 27, 28  
 osmotic potential and potassium, 65  
 SO<sub>2</sub> effects, 194  
 treatment, 27  
 and turgor pressure, 27, 28
- epidermis  
 carbon dioxide fixation, 92, 93, 97  
 carbon import, 95  
 cell ion measurement, 55, 56  
 cell types, 56  
 enzyme activities, 87–93  
 enzyme inhibitors in extracts, 88  
 leakage, 97, 98  
 malate exit, 97  
 membrane potentials, 61  
*Erwinia amylovora*, 207  
*Erysiphe graminis*  
 and flag leaf stomatal behaviour, 217  
 subsidiary cell invasion, 206  
*Erysiphe pisi*  
 and pea photosynthesis, 216  
 and solute accumulation in peas, 215  
*Eucalyptus pauciflora*, 263, 268
- farnesol, all *trans*, 179  
 and water stress in *Sorghum*, 169  
 fatty acids, free in *Zea*, 180  
 feedback control, 137, 138  
 negative, 265  
 feed forward control, humidity response,  
 137, 140, 154  
 ferulic acid ester, 19  
 fluorescence  
 and electron transfer, 109  
 experimental set-up, 114  
 green, guard cell vacuoles, 113–15  
 and guard cell study, 109–15  
 microscopy and epidermal strip  
 contamination, 110  
 fluorospectrophotometer, 114  
*Fraxinus americana*, 141  
 fructans, 76, 98  
 fusicoccin, 107  
 chemical nature, 211  
 CO<sub>2</sub> response reversal, 131, 132  
 isolation, 211  
 non-specific effects, 212  
 and proton extrusion, 115, 132  
 stomatal responses, 131, 132, 212  
 and vacuolar fluorescence, 115  
*Fusicoccum amygdali*  
 peach canker, 211  
 toxin and stomatal behaviour, 211–13
- gallium-arsenide photomultiplier, 114

## 290 INDEX

- Geum rossii*, humidity response, 155  
 glasshouses, controlled environment  
   and leaf conductance, 235, 239, 240  
 glyceraldehyde-P dehydrogenase, 95  
 glycerate, 72  
 Golgi system in guard cells, 22  
*Gossypium hirsutum*, 33, 263  
   leaf area and transpiration, 234, 235  
 green area index, 220  
*Greenovia aurea*, stomata and humidity,  
 139  
 Gregory and Pearse resistance unit, 35  
 guard cell *see also* wall  
   *Allium* protoplast, vesicles, 22  
   -cell metabolite fluxes, 97, 98  
   cell shape, 3–16  
   coenocytic, 18  
   CO<sub>2</sub> fixation to malate, 78  
   common wall perforations, 17  
   communication, 17, 18  
   Cyperaceae, 14  
   cytoplasm ionic content, 80, 81  
   differentiation in culture, 12, 13, 109  
   emission spectra, 110  
   fluorescence, 109–15  
   grasses, 5–16  
   humidity effects, 137–57  
   intact, 64  
   ionic relations, 51–67, 71–6, 79–81  
   isolated, 64, 65  
   malate levels, 74, 75, 126, 127, 130–3,  
   152  
   membrane potentials, 60, 61  
   metabolism, 87–100, 282, 273; and  
   ABA, 283; and Calvin cycle, 93–7;  
   enzyme activities, 87–93  
   microsurgery, 30  
   model systems, 1  
   osmotic changes and potassium, 63, 64  
   pH effect, 29, 54, 55, 57, 62, 152, 282  
   phenolic esters, 19, 21  
   potassium transport, 29  
   protoplasts, 22, 30, 103–6  
   salt accumulation, 52–60  
   and subsidiary cell damage, 16  
   subsidiary cell interaction, 16–19  
   swelling stage, section 6  
   transient phase, 10, 11  
   turgor control, 126, 127, 165, 172  
   wall deposition, 3–16  
 guard mother cells, 2
- Helianthus annuus*, SO<sub>2</sub> response, 192  
 helium, 39  
*Helminthosporium maydis*  
   toxin effects on stomata, 211  
   heterobaric leaves, 37, 38  
   humidity, 238  
   and photosynthesis control, 258
- plant response and distribution, 154, 155  
   and stomatal responses, 137–52, 249  
 humidity sensors, 42  
 hydrogen, 39  
 hydrogen sulphide, field effects, 198  
 hygrometer, 41  
 hygroscope, 40  
 hypostomatous leaves, 40  
*Hyptis emoryi*, 141
- irradiance and stomatal opening *see also*  
 light, 120, 121
- irrigation  
 and conductance in millet, 226, 231, 232  
 stomatal response, 154
- isopropyl-N-phenylcarbamate  
 and guard cell wall thickening, 15, 16
- isotachophoretic analysis, 54
- leaf  
 area index, 225; and drought, 235, 236  
 growth, stress-induced limitation, 175,  
 176  
 microclimate, factors affecting, 155  
 water potential and transport, 255, 256
- leaf discs  
 clamp, 36  
 sealing, 37
- leaf resistance, hypothetical, factors  
 affecting, 148
- light *see also* quantum flux density and  
 irradiance  
 blue and protoplast response, 107, 108  
 blue and stomatal action, 121–3, 260,  
 261, 267, 268  
 and CO<sub>2</sub>, stomatal response, 262, 266–72  
 and *Erysiphe* effects on pea, 215  
 and leaf structure, 141  
 red and stomatal action, 121–3, 261, 267,  
 268  
 role in stomatal opening, 123, 129; and  
 potassium, 126–30  
 and stomatal humidity response, 140, 141  
 stomatal response, 119–33
- lignin, 254  
 in grass guard cells, 19  
 tests for, 19
- liquid paraffin, 27, 28  
 Lloyd's strips, 27
- Loiseleuria procumbens* microenvironment  
 and form, 155
- Lolium perenne*, humidity response, 145
- Lophodermium pinastre*, guard cell invasion,  
 206
- Lysimachia nummularia*, 145
- lysimetry, 225
- Macroptilium atropurpureum*, 145

- malate, 51, 53, 54, 65, 72, 89, 97, 98, 152  
 epidermal, source, 56, 130  
 epidermal and stomatal aperture, 56, 57,  
 73, 93; effect of light and carbon  
 dioxide, 126, 127  
 in epidermis of *Allium*, 77  
 in epidermis of *Commelina*, 77, 78,  
 88–90, 93, 98  
 guard cell, source in *Vicia*, 73–5  
 pH and valency, 57, 58  
 potassium balance, 65, 72  
 uptake and stomatal stage, 91
- malate dehydrogenase  
 epidermal, 88, 89  
 in guard cells, 94  
 inhibition of, 88
- malic enzyme, 88, 89  
 in guard cell, 94  
 NADP specific in *Commelina*, 90
- Malus pumila*, 263
- mannitol, 107–9
- membrane potential  
 guard-cell, 60, 61; and light, 61
- membrane whorls, myelin-like, 18
- mesophyll  
 epidermal contamination and analysis, 87,  
 88  
 epidermal transfer of carbon, 96  
 porometer resistance, 36, 37, 40; species  
 effects, 37  
 stress messenger, 167
- microelectrodes, ion-sensitive, 55, 77
- microfibrils  
 arrangement and growth conditions, 14, 15  
 composition, 19  
 guard cell alignment, 3–6, 10, 11  
 orientation in transient, 12, 13
- microfilaments, 2
- microspectrophotometry, 19, 110  
 guard cell fluorescence, 112, 113  
 low temperature, 110  
 room temperature, 110
- microsurgery of guard cells, 30, 52, 53
- microtubules, 2  
 in *Allium* guard cell, 4, 5  
 cellulose coorientation, 3  
 deployment, control, 22  
 and division plane, 2  
 membrane cross-bridges, 9, 10  
 pharmacological effects, 5, 15, 16  
 polymerization, 22  
 radial alignment in grasses, 12, 13  
 reorientation in grasses, 9–11
- mitochondria: chloroplast ratio in guard  
 cells, 123
- models  
 gas exchange in leaf, 148, 149, 248  
 parameters used, 237  
 stomatal response in the field, 236–40
- moss sporophyte, guard cell mother division,  
 2
- multivesicular bodies, 18
- Nicotiana rustica*  
 drought responses of two varieties, 236
- Nicotiana tabacum*, 33, 79, 166, 230  
 drought effects on conductance, 235, 236  
 ozone sensitivity, 188, 189  
 SO<sub>2</sub> response, 192
- nitrogen oxides  
 stomatal effects, 197
- nitrous oxide, 39
- nujol, 32
- Opuntia inermis*  
 stomata and humidity, 140
- osmoregulation and ABA, 175
- osmotica  
 guard cell cytoplasmic, 81  
 nature and protoplasts, 107  
 and protoplast isolation, 104
- ozone  
 field effects, 198, 199  
 natural concentrations, 188  
 photosynthesis in oats, 189  
 sinks, 190  
 stomatal response, 188–90; protective  
 effect, 189, 190  
 sulphur dioxide mixture effects, 197–9
- Paphiopedilum* sp  
 mitochondria, 123  
 photosynthesis in guard cells, 114  
 stomata and Calvin cycle, 95
- Paphiopedilum harriseanum*  
 epidermis CO<sub>2</sub> fixation, 96  
 guard cell fluorescence, 113, 114  
 protoplast isolation, 105, 106, 114
- Paphiopedilum venustum*  
 low epidermal CO<sub>2</sub> fixation, 96
- paradermal sections, 26, 27, 104
- pathogens  
 leaf invasion, 213–17  
 motile spores, 207  
 need for water, 206, 207  
 stomatal response, 205–19; and CO<sub>2</sub>  
 exchange, 218, 219; consequences of,  
 217, 219; and water relations, 217, 218  
 toxin production, 210–13  
 and water supply to leaf, 208–10
- Pelargonium zonale*  
 enzyme inhibition, 88  
 no Calvin cycle, 95  
 trichome carbon dioxide fixation, 96
- Pelvetia fastigiata*  
 guard cell cytoplasm, ionic tolerance, 80
- Pennisetum typhoides*  
 canopy conductance, 229

## 292 INDEX

*Pennisetum typhoides* – cont.

leaf area and conductance, 234, 235

perfusion pump, 104

pH, guard cell, 29, 63

and humidity changes, 152, 153

and malate movement, 57, 58

and proton extrusion, 54, 55

and vacuolar fluorescence, 115

phalloidin, 2

*Phaseolus vulgaris*, 166

pollutant mixture effects, 198, 199

SO<sub>2</sub> effects, 194, 198, 199

stomatal conductance and SO<sub>2</sub>, 190, 191, 193

*Phleum pratense*

cuticle development, 20, 21

guard cell development in, 5–9, 15

microtubule alignment, 5, 9

primary leaf culture, 11, 12

stomatal complex, electron microscopy, 5, 6, 17

phosphoenolpyruvate carboxykinase, 89, 92

phosphoenolpyruvate carboxylase

in *Commelina* sp epidermis, 78, 88, 89

in guard cells, 74, 94, 131

ionic strength and inhibition, 80

malate inhibition, 59

in *Tulipa* epidermis, 79

phosphoenolpyruvate synthesis *see* pyruvate P<sub>i</sub> dikinase

phosphoribulokinase, 95

phosphorylase, 94

photosynthesis

cuvette measurement, 238

factors affecting, 224, 225

mesophyll, 262, 265

NO<sub>2</sub> effects, 197

rate of light saturated, 256, 257

SO<sub>2</sub> effects, 192, 193

stomatal control, 256–66, 284, 285; and

carbon dioxide, 261–5; and humidity,

257; and leaf water potential, 258; and

quantum flux density, 259–61; and

temperature, 258, 259

*Phytophthora drechsleri*

and safflower stomata, 208, 209

and wilting, 208, 209

*Phytophthora infestans*, 207

lesion development and stomata, 213

stomata as exits for spores, 208

*Picea engelmannii*

stomata, light and sensitivity to humidity, 141

*Picea sitchensis*, 263

light saturation, 267

stomatal behaviour model, 145

stomatal conductance and xylem water, 258

*Pinus sylvestris*, 263, 267

carbon dioxide sensitivity, 269

stoma cross-section, 254

stomatal conductance, 260

pisatin and stomatal opening, 215, 216

*Pisum* sp, 3

ABA effect on wilted leaves, 171

plasmalemma, proton pumping, 58

plasmodesmata, 18

*Plasmopara viticola*, 207

pollutants *see also* ozone, nitrogen oxides,

sulphur dioxide

damage, 187

inadequate experimentation, 187

stomatal response, 187–200; to mixtures, 197, 199

and stress, 187

*Polypodium vulgare*

epidermal carbon dioxide fixation, 96

hypothetical leaf resistance, 148, 149

stomatal opening and air humidity, 138, 139; effect of temperature, 142, 143;

model, 145–7

porometers

diffusion, 43, 239, 240

diffusive gas flux, 33, 39; Dufour effect,

39; hydrogen, 39; inert gas, 39, 40

mass flow, 33–40; airflow systems, 34,

35; clamp effects, 35–7; and diffusive

flow conductance, 38, 39; hot wire

constant pressure, 35; measurement

systems, 35; uses, 38; Wheatstone

bridge, 35, 38, 42

reliability, 224

responses to, 230

transpiration, 33, 40–4; calibration, 41;

condensation, 42; differential, 41, 42;

humidity sensor, 42; and mass flow

conductance, 42; sensor element, 43,

44

potometry, 29, 33–40

and leaf surface, 231

limitations, 37, 38

mass flow, 33–40

viscous flow, 37

potassium, 29, 51

balance sheet of movement, 59

*Commelina* spp. flame photometry, 55

concentration and stomatal opening,

52–4, 71, 152; organic anions and, 53,

56, 65, 72

iminodiacetate, 54

and light, CO<sub>2</sub> and stomatal response, 125–30

in maize, midtransient guard cells, 12

and osmotic changes in guard cells, 63–6

shuttle, 29, 52, 55, 57

uptake and fusicoccin, 212

proton extrusion, 53–5, 59

and fusicoccin, 131

- proton pump  
*Allium* sp, 77  
 and chloroplast function, 113  
 photoactivation, 61, 108
- protoplasts, 30  
 epidermal, 105, 106  
 guard cell study, 103–15  
 isolated, 105  
 isolation of guard cell, 103–6; organelle  
 isolation, 108, 109; vacuole isolation,  
 108, 109  
 mesophyll, 104, 105  
 microchamber isolation, 104  
 photoresponse, 107, 108  
 swelling, 107, 108  
 test tube method, 105
- Prunus armeniaca*  
 effects on stomatal aperture, 146  
 model of leaf gas exchange, 148, 149, 237
- Pseudomonas tabacci*, 207
- Pseudoperonospora humuli*, 207
- Pseudotsuga menziesii*  
 stomatal behaviour model, 145  
 stomatal conductance and CO<sub>2</sub>, 264
- Psychotria capensis*, 37
- Puccinia graminis*, infection and water  
 stress, 208
- Puccinia hordei*, infection process, 206
- pyruvate and stomatal opening, 91
- pyruvate P<sub>i</sub> dikinase, 89, 94  
 isolation, 92
- quantum flux density *see also* light and  
 conductance  
 and stomatal conductance, effect of CO<sub>2</sub>,  
 260  
 and transpiration, 253–5
- Rhynchosporium secalis*  
 and host affected area, 218  
 and stomatal turgor, 214
- ribulose biphosphate carboxylase, 89; in  
 guard cells, 75, 78, 93, 273, 274
- rubidium  
 [<sup>86</sup>Rb], 65, 66, 74
- salt accumulation, 59
- Scolopia mundii*, 37
- Senecio odoris* stomatal opening, action  
 spectrum, 119, 120, 122
- sensor elements, 43
- Sesamum indicum*, 143
- Sorghum* sp, stomatal closure compound,  
 169
- Spannungspphase*, 25
- starch, 53  
 absence from guard cells, 98, 99  
 source of guard cell, 75  
 and stomatal movement, 75, 76, 78, 92,  
 93, 152–4  
 test kit, 75
- stomata  
 abscisic acid effects, 67, 97, 107, 167,  
 170, 173, 178  
 and bulk leaf water status, 166–74  
 carbon dioxide sensitivity, 263, 268, 269  
 evaporation pathways, 254  
 field behaviour, 223–41  
 and fungal toxins, 210–13  
 humidity sensors, 137  
 isolated, 79  
 number and surface, 38, 231  
 and pathogens, 205–19  
 placement and gas movement, 274  
 pollutant response, 187–200
- stomatal action, hypotheses for, 272–4
- stomatal aperture  
 carbon dioxide effect and water stress,  
 173, 174  
 conductance measurement, 25, 32–44,  
 223–41; gas porometry, 33–40; liquid  
 infiltration, 32, 33, 223; water vapour  
 diffusion, 40–4  
 direct humidity effect, significance,  
 154–6, 252–5  
 factors affecting, 248  
 infiltration scores, 33  
 and ionic state of guard cells, 51–67  
 and light, 146  
 and malate cycling, 58  
 and malate level, 73  
 microscopic measurement, 25–32; effect  
 on aperture size, 26; epidermal strips,  
 27–30; fixed leaves, 27; intact leaves,  
 26; micro-relief methods, 30–2;  
 paradermal sections, 26, 27; ridge  
 dimension changes, 31, 32  
 model describing, 147  
 and osmotic changes, 63  
 and potassium content, 151, 152; in  
*Commelina*, 64, 65; effect of light and  
 CO<sub>2</sub>, 125; and salts, 128, 129  
 and turgor, 27, 28
- stomatal cells  
 and cell division, 2  
 organelles, 22  
 structure and development, 1–22, 281
- stomatal closure, 91  
 and humidity, 151  
 and water deficit, 165
- stomatal conductance  
 definition, 249  
 effects of environment, 229, 230, 249  
 and intercellular space CO<sub>2</sub>, effect of  
 light, 260  
 and irradiance, 226, 238  
 leaf area effects, 234, 235



- stomatal conductance – *cont.*  
 and leaf height, 226–9, 232  
 and leaf surface, 231  
 and leaf turgor, 165, 166  
 and leaf water potential, 165, 166  
 and light, effects of CO<sub>2</sub>, 260  
 modelling, 236–40  
 and plant variability, 231–6  
 response to humidity, 240  
 and stomatal aperture, 25, 26, 32–44  
 and transpiration rate, 248  
 and xylem water potential, 258
- stomatal control  
 of leaf turgor, 166  
 of photosynthesis, 256–66, 284, 285  
 of transpiration, 250–6, 284, 285  
 and water stress, 163–81
- stomatal density and light, 141
- stomatal ledges, 22
- stomatal opening  
 action spectrum, 119–21, 123  
 and air stream humidity, 138, 139  
 and anions, 71–81, 91  
 biochemical pathways and, 94  
 dark, 119, 120, 261  
 extracellular ion exchange, 56  
 factors affecting, 123  
 glucose inhibition, 78  
 and guard cell chloroplasts, 113  
 and guard cell starch, 75, 76, 91  
 hypothesis, 272  
 ion movement and cell interactions, 17  
 irradiance, 120–2  
 and leaf water content, 138  
 measurements needed, 62  
 mechanisms, 73, 124–33; NaCl, KCl  
 effects, 128, 129  
 night in succulents, 119  
 and potassium content, 52–4  
 proton extrusion, 53, 54  
 rhythms, 25, 44, 119  
 role of malate, 57  
 role of potassium, 51, 71, 77  
 starch and, 152, 153
- stomatal resistance (*see also* stomatal  
 aperture and stomatal conductance)  
 and ABA, 168–73
- stomatal response to humidity, 137–57,  
 252–5  
 effect of bulk water potential, 143–5  
 effect of CO<sub>2</sub>, 178  
 effect of CO<sub>2</sub> and temperature, 142, 143  
 effect of light, 140, 141  
 mechanisms, 149–54  
 modelling, 145–9
- stomatal response to light, 266–8  
 effect of CO<sub>2</sub>, 269–72  
 red and blue, 267, 268
- stomatal response to water potential, 165, 166
- subsidiary cells  
 form in grasses, 7  
 function in grasses, 5  
 and guard cell damage, 17  
 and guard cell shape, 16–19
- sucrose, stored and stomatal movement, 93
- sulphur dioxide  
 atmospheric sources, 190  
 and cell turgidity, SEM, 195  
 leaf conductance and cell survival, 194  
 ozone mixture, effects of, 197–9  
 sites of action, 193–6  
 and stomatal responses, 190–6; and  
 humidity, 190–2; hypothesis, 196  
 transpiration, photosynthesis and, 192,  
 193  
 uptake and cucurbit necrosis, 194
- temperature  
 and control of photosynthesis, 258, 259  
 and stomatal conductance, 249
- tentoxin  
 chemistry, 211  
 and potassium uptake, 212  
 and *Vicia* stomatal aperture, 212
- tonoplast  
 fluorescence, 113  
 transport processes, 59, 67
- Tradescantia* sp  
 peristomatal water loss, 150
- Tradescantia albiflora* var. *albo-vittata*  
 epidermal cell ions, 56  
 guard cell membrane potentials, 60, 61  
 potassium movements, guard cell, 59
- Tradescantia virginiana*  
 pore measurement, 31  
 turgor and pore width, 28
- transient in guard cell differentiation, 12–14
- transpiration  
 cuvette measurement, 238  
 factors affecting, 224, 225  
 and leaf area, 234, 235  
 peristomatal, 150, 254  
 ratio, 154  
 SO<sub>2</sub> effects, 192, 193  
 stomatal control, 250–6, 284, 285; and  
 leaf water potential, 255, 256; and  
 quantum flux density, 250, 251; and  
 vapour pressure deficit, 253–5  
 stomatal resistance discrepancy, 42
- trichomes, 96
- Tridax procumbens*, 92
- Trifolium repens*, humidity response of  
 stomata, 145
- Tulipa* sp, chloroplast fluorescence, 110
- Tulipa gesneriana*, 78  
 CO<sub>2</sub> fixation into malate, 78, 79
- tundra, survival in and factors affecting,  
 154, 155



- turgor  
 components of leaf, 165  
 and pathogens, 214
- Uromyces fabae*, stomatal immobilization, 214
- vacuole, guard cell, 67  
 autofluorescence, 22  
 development, 7  
 fluorescence, 113–15  
 isolation, 108, 109  
 pH changes, 58  
 pH and fluorescence, 115  
 and stomatal opening, 80
- Valerianella locusta*  
 starch metabolism, 152, 153  
 stomatal aperture and potassium content, 151–3
- Valonia ventricosa*  
 guard cell cytoplasm ionic tolerance, 80
- vapour pressure deficit *see also* humidity  
 and transpiration control, 253–5
- Verticillium dahliae* and chrysanthemum  
 stomata, 209
- Vicia faba*, 52, 80, 98, 214  
 ABA generation and stress, 168  
 guard cell fluorescence, 110–12  
 guard cell ion content and opening, 52–5, 71–6  
 guard cell protoplasts, 105, 106  
 source of guard cell malate, 74–6  
 starch, 75  
 stomatal aperture, time course, 25  
 stomata and tentoxin, 212  
 sulphur dioxide response, 190, 191  
 vacuolar fluorescence, 113, 115
- virus  
 and stomatal behaviour, 214  
 sugar beet yellows, effects of, 218
- wall, guard cell  
 birefringence, 11–14; loss and digestion, 104  
 composition, 19  
 deposition in grasses, 7–11  
 fluorescence, 18, 19; effect of pH, 19  
 mechanical properties, 3  
 porometer resistance, 40  
 potential, 61  
 secondary, 3  
 thickening; effects of colchicine, 5, 15, 16;  
 species variation, 3–8
- water loss  
 feedback control, 137, 138  
 feed forward control, 137, 140, 154, 255  
 and ABA level, 167, 168  
 and farnesol in *Zea*, 179  
 and lipid metabolism, 180  
 and ozone sensitivity, 188  
 and *Phytophthora* effects on stomata, 213  
 and root and shoot growth, 177  
 and stomatal control, 163–81, 189  
 and *Verticillium* infection, 210  
 wind, stomatal response, 155
- Xanthium strumarium*  
 dark opening of stomata, 120, 263  
 irradiance and stomatal opening, 120, 121  
 stomatal CO<sub>2</sub> sensitivity, 265
- Zea mays* 52, 130, 166, 214, 263  
 epidermal strip, difficulties, 52  
 fatty acids and water stress, 180  
 humidity insensitivity, 140  
 potassium uptake, 79  
 stomatal common wall structure, 18  
 stomatal differentiation sequence, 7, 12  
 water stress and ABA, 169, 170  
 water stress and farnesol, 179  
 water stress and stomata, 163, 164
- Zebrina pendula*  
 pore width and fixation method, 27  
 zwitterions as osmotica, 81