

Author index

-
- Abdullah, N.F. 74, 75
 Acton, A. B. 74
 Adams, M.S. 143
 Alexander, R.D. 144, 166
 Allard, R.W. 74, 75, 128, 129, 130
 Amadon, D. 185
 Assem, J. van den 162

 Babcock, E.B. 65
 Bacci, G. 150
 Baker, H.G. 99, 109, 135, 136, 137
 Baker, R.R. 151
 Banta, A.M. 59
 Bar-Anan, R. 150
 Basrur, V.R. 56
 Bateman, A.J. 169
 Bawa, K.S. 136, 137
 Bell, G. 151
 Bellrose, F.C. 181
 Bengtsson, B.O. 139
 Bergerard, J. 63
 Birky, C.W. 61
 Bodmer, W.F. 19, 24, 78, 86
 Breder, C.M. 182, 183
 Brues, M.A. 24
 Bruning, D.F. 182
 Bull, J.J. 130, 133, 163
 Bulmer, M.G. 141

 Carlisle, T.R. 175
 Carlquist, S. 136
 Carson, H.L. 43
 Catchside, D.G. 77
 Charlesworth, B. 30, 74, 75, 84, 87, 89, 95,
 120, 121, 158
 Charlesworth, D. 84, 87, 120
 Charnov, E.L. 130, 131, 133, 162, 163,
 166
 Chewyreu, I. 162
 Childress, D. 164
 Chinnici, J.P. 74, 75, 76
 Christensen, B. 45
 Clark, R.D. 149

 Clarke, B. 25
 Clarke, C.A. 85
 Clayton, G.A. 11
 Clegg, M.T. 128
 Clutton-Brock, T.H. 179, 184, 186
 Collier, G. 181
 Cox, E.C. 112, 113, 190, 191, 192
 Craig, G.P. 164
 Crosby, J.L. 86
 Crow, J.F. 19, 21, 24
 Cuellar, O. 46, 47, 48, 56, 67

 Darlington, C.D. 23
 Darwin, C. 42, 136, 168, 171, 183, 184
 Davis, G.W.F. 173
 Dawkins, R. 113, 175
 Detlefsen, J.A. 74, 75
 Dewees, A.A. 74, 75
 Dobzhansky, Th. 184
 Drake, J.W. 188, 189, 190
 Drickamer, L.C. 142

 Edwards, A.W.F. 148, 149
 Ehrman, L. 171
 Emerson, A.E. 2
 Ernst, A. 86
 Eshel, I. 14, 82

 Falconer, D.S. 76, 149, 170
 Feldman, M.W. 14, 78, 82
 Felsenstein, J. 10, 15, 22, 32, 33, 73, 78,
 117, 121, 122, 123
 Fisher, R.A. 2, 3, 13, 17, 32, 40, 77, 117,
 119, 129, 146, 147, 157, 159, 160, 171,
 172, 173, 174
 Fossey, D. 142
 Frame, G.W. 152
 Frame, L.H. 142
 Franklin, I. 78, 82
 Freeman, D.C. 162
 Frith, H.J. 182

 Ghiselin, M.T. 133, 135, 138

Cambridge University Press
 978-0-521-29302-0 - The Evolution of Sex
 John Maynard Smith
 Index

[More information](#)

212 *Author index*

- Gibson, T.C. 112, 113, 192
 Gilbert, J.L. 61
 Gilpin, M.E. 2
 Glesener, R.R. 109, 110
 Godley, E.J. 136
 Goldblatt, S.M. 47
 Grant, V. 109
 Greene, P. 144
 Greenwood, P.J. 148
- Haas, R. 172
 Haigh, J. 30, 34, 111, 112
 Haldane, J.B.S. 23, 24
 Hamilton, W.D. 114, 146, 147, 160, 161, 164, 165
 Harcourt, A.H. 142
 Hare, H. 165, 166
 Harper, J.L. 156
 Harper, K.T. 162
 Harrison, B.S. 12
 Harvey, P.H. 141, 179, 184, 186
 Haskins, C.P. 173
 Hebert, P.D.N. 59
 Hickey, W.A. 164
 Hill, J.L. 140
 Hill, W.G. 10, 32, 111, 122, 123
 Howe, H.F. 159
 Hubby, J.L. 24
 Hutchinson, G.E. 60
- Jain, S.K. 130
 Jenni, D.A. 181, 184
 Jinks, J.L. 129
 John, B. 120
- Kahler, A.L. 128
 Karlin, S. 15, 78, 82
 Kaufman, D.W. 130
 Kear, J. 180
 Kerfoot, W.C. 60
 Kidwell, M.G. 74, 75, 76
 Kimball, A.W. 149, 159
 Kimura, M. 19, 21, 24, 78, 188, 189, 194
 King, J.L. 24
 Klikoff, L.G. 162
 Knowlton, N. 155
 Knox, P.B. 62
 Kojima, K. 78, 82
- Lack, D. 179
 Langley, C.H. 84
 Lee, L. 61
 Leigh, E.G. 188, 189, 194
- Lerner, I.M. 129
 Levene, H. 92
 Levin, D.A. 109, 110
 Levins, R. 131
 Levi-Strauss, C. 143
 Lewis, D. 147
 Lewis, K.R. 120
 Lewontin, R.C. 24, 78, 81, 82, 84
 Lokki, J. 57
 Lovell, P.H. 156
- McAlister, W.H. 156
 Macarthur, R.H. 27, 147, 167
 McDonald, I.C. 164
 Marquiss, M. 159
 Maslin, P.T. 46
 Mather, K. 12, 81, 129
 Maynard Smith, J. 2, 3, 14, 15, 17, 20, 22, 24, 26, 28, 29, 30, 31, 90, 93, 96, 101, 104, 106, 111, 112, 126, 127, 130, 132, 147, 148, 168, 173, 174, 175, 176
 Milkman, R.D. 24
 Mitton, J.B. 91, 99
 Mockford, E.L. 53, 54
 Moore, K.G. 156
 Moore, W.S. 69
 Moran, P.A. 17
 Morgenstern, O. 147
 Mukai, T. 85, 189
 Muller, H.J. 4, 18, 32, 33, 36, 117
 Muntzing, A. 65, 66
 Murray, J. 86, 135
- Neaves, W.B. 46
 Neel, J.V. 142, 143
 Newton, I. 159
 Newton, M.E. 164
 Nikolskii, G.V. 156
 Novitski, E. 149, 159
 Nur, U. 43
- O'Donald, P. 172, 173, 184
 Ohta, T. 19
 Olsen, M.W. 45
 Opler, P.A. 136, 137
 Orians, P.H. 180
- Packer, C. 141, 142
 Parker, E.D. 57
 Parker, G.A. 151, 154, 155, 178
 Pijnacker, L.P. 44, 63
 Pimentel, R.A. 156
 Platt, W.J. 156

Author index 213

- Poulsen, D.F. 147
 Preston, F.W. 27
 Price, G.R. 147, 176
 Pusey, A. 142
- Ralls, K. 185
 Reed, T.E. 24
 Rees, H. 120
 Rensch, B. 186
 Richards, A.J. 55
 Roberts, E. 74, 75
 Robertson, A. 10, 11, 32, 111, 122, 123, 150
 Robertson, J.G. 64
 Rosen, D.E. 182, 183
 Rothfels, K.H. 56
 Ruttner-Kulisko, A. 61
- Sabelli, B. 59
 Sakaguki, B. 147
 Saura, A. 57
 Schull, W.J. 142
 Schultz, R.J. 47
 Scudo, F.M. 149
 Seiler, J. 45
 Selander, R.K. 57, 130, 171, 185
 Shaw, D.D. 74, 75, 76
 Shepher, J. 143
 Sheppard, P.M. 85
 Sherman, P.W. 166
 Slatkin, M. 97, 98
 Smith, V.G.F. 151
 Snyder, R.L. 163
 Stebbins, G.L. 23, 45, 49, 51, 54, 55, 65, 109
 Stewart, K.S. 142
 Street, H.E. 51
 Strobeck, C. 30, 91, 111, 116, 120, 121
 Suomalainen, E. 54, 57, 109
 Sved, J.A. 24
- Talmon, Y. 143
 Teitelbaum, M.S. 142, 159
 Thomson, G. 30, 78, 86, 112
 Tilman, D. 109, 110
 Tomlinson, J. 134, 135
 Trivers, R.L. 133, 146, 147, 159, 162, 165, 166, 174, 175, 178, 184
 Turner, J.R. 78, 117
- Uzell, T.M. 47
- Van Valen, L. 26, 54, 110
 Vanzolini, P.E. 48
 Verner, J. 180, 184
 Vessey, S.H. 142
 Von Neumann, J. 147
- Wagoner, D.E. 164
 Wallace, B. 24
 Weir, J.A. 149, 150
 Werner, P.A. 156
 Westergaard, M. 163
 White, M.J.D. 23, 43, 45, 46, 48, 49, 54, 56, 63
 Wiley, R.H. 171, 180, 184
 Willard, D.E. 133, 147, 159, 162
 Williams, G.C. 2, 5, 9, 10, 52, 57, 63, 66, 69, 89, 90, 99, 107, 113
 Willson, M.F. 180, 184
 Wilson, E.O. 27
 Winterbourne, M. 64
 Wood, R.J. 164
 Wynne-Edwards, G.C. 2
- Yamaguchi, O. 85
 Yokoyama, S. 10, 111, 117, 121, 122, 123
 Young, P. 59, 60
- Zaffagnini, F. 59
 Zahavi, A. 173, 174
 Zaret, T.M. 60

Subject index

- Acarini, arrhenotokous families and sub-families of, 53, 160
- Acarophenax* (mite): viviparous, with single male in litter mating with females before birth, 160
- Accipiter nisus* (sparrowhawk), sexual dimorphism and sex ratio in, 159
- Acer negundo* (box elder): sex ratio of, in dry and damp sites, 162
- Actitis macularis* (spotted sandpiper), polyandry in, 181
- Aedes aegypti* (mosquito): sex ratio in, 164
- Aleurodidae (white flies), arrhenotoky in, 53
- Amblystoma* (salamander): thelytokous forms of, 47, 48
- Anas* (river ducks), uniparental care in, 181
- anisogamy, evolution from primitive isogamy to, 39, 146, 151–5
- Anolis garmani* (lizard), large males mate more often, 184
- Anser* (geese), biparental care in, 181
- Antennaria* (Compositae), apomictic species complex of, 55
- aphids, cyclical parthenogenesis in, 62
- apomixis, in thelytokous parthenogenesis, with suppression of meiosis, 43; in animals, 45–6; application of term to plants, 49; facultative, in plants, 55, 56; in plants, commonly associated with polyploidy, 50–1, 55; possible origin of, 48
- arrhenotoky, production of males from unfertilised eggs, 42; combined with inbreeding and excess of females in mites and insects, 160; taxonomical distribution of, 54
- asexual reproduction, in plants: by bulbs, runners, etc., 49; by seeds produced asexually, 50
- Atriplex confertifolia* (shadscale): sex ratio of, in dry and damp sites, 162
- automixis, in thelytokous parthenogenesis, 43–5, 63; artificial selection for, 47; in plants, 50; with pre-meiotic doubling, in *Moraba*, 46, 56
- Avena barbata*, *A. fatua*: partial outcrossing in, 128, 130
- Avena sativa*, plants produced by outcrossing have higher fitness, 129–30
- Aythya* (diving ducks), uniparental care in, 181
- Bacillus rossius* (stick insect): bisexual in part of range, automictic elsewhere, 44, 63
- bacteriophages lambda and T4, mutation rates in, 189
- 'balance' argument: for maintenance of cyclical parthenogenesis, 59–60, 62; for maintenance of recombination, 9, 72; for short-term advantages of sex, 5–6, 38, 57–8, 66–7
- Biomphalaria* (snail): self-compatible, but outcrosses if opportunity occurs, 134
- birds: avoidance of inbreeding in, 140–1; parental care in, 179–83; size dimorphism in, 185
- blood pH: artificial selection for, in mice, with accidental association with sex ratio, 149–50
- Bonasa umbellus* (ruffed grouse), male with central territory obtains most matings, 170
- Bosmina longirostris* (Cladocera): seasonal dimorphism in, related to predation, 60
- Bromus mollis* (grass), partial outcrossing in, 128
- brooding of young: association of hermaphroditism with, 138; resource allocation and, 139

216 *Subject index*

- Calidris alba* (sanderling), parental care in, 181
- Calligrapha* spp. (chrysomelid beetles), facultative parthenogenesis in, 64, 67
- Caryophyllaceae, dioecy in, 135–6
- cattle, sex ratio in, 150
- Cepaea* (snail), self-incompatible hermaphrodite, 125
- Ceriodaphnia cornuta* (Cladocera), dimorphism of, 60
- chiasma formation: genetic variation in frequency and location of, 9, 74; in scheme for origin of sex, 7, 8; selectively favoured? 72–3; *see also* recombination
- chimpanzee, female transfer in, 142
- choice, female, 168–71; of ‘handicaps’ in male, 171–4; self-reinforcing of, 171–2, 173–4
- chromosomes: in scheme for origin of sex, 8; of *Aedes aegypti*, 164
- Clitumnus extradentatus* (stick insect), automictic, 63
- Cnemidophorus uniparens* (whiptail lizard), parthenogenetic, 46, 48, 56
- Cnemidophorus tessellatus*, parthenogenetic: diploid and triploid biotypes of, 57
- Cnephia mutata* (simuliid fly), triploid parthenogenetic biotypes, 56
- Collinsia sparsiflora* (Scrophulariaceae), varies from complete selfing to random mating, 128
- colonisation of new region: parthenogenesis in, 110; by single hermaphrodite individuals, 134, 136, 139; by two populations, and selection for recombination, 15, 96–7
- competition: between *E. coli* wild-type and mutator strains (non-recombinant), 112, 113; between sexual and asexual populations in fluctuating environment, 92–5; for mates, and sex ratio, 160–1
- Crepis* (Compositae), apomictic species complex of, 55
- cyclomorphosis, in Cladocera, 60, 61
- Cygnus* (swans), biparental care in, 181
- Cynipidae (gall wasps), parthenogenesis in, 61
- cytoplasmic ‘genes’, 147
- Daphnia* (Cladocera): cyclical apomixis in, 58–60, 67, 70
- Daphnia pulex*, *D. middendorffiana*: all-female populations of, producing ‘winter eggs’ parthenogenetically, 59
- Dichanthium aristatum* (grass): balance between sex and parthenogenesis in, 62–3, 67, 70
- Dinophilus*, sex determination before fertilisation in, 150
- dioecy, 125; in animals, 133; in plants, 133, 136, 138; and avoidance of inbreeding, 135
- diploidy, possible advantages of, 9
- Diptera: chiasmata absent in meiosis in males of, 9, 73; ‘rare male’ effect in, 140, 171; thelytokous, 56
- dispersion pattern: and avoidance of inbreeding, 140–1; and effect of sib competition on selection for recombination, 108; seeds and, in plants, 49
- Distichlis spicata* (salt grass): sex ratio of, in dry and damp sites, 162
- DNA: recombination and repair of, 7, 193
- Drosophila mangabeirai*, automixis in, 44–5
- Drosophila melanogaster*: linkage disequilibrium in, 84–5; mutation rate in, 189–90; selection for increased or decreased recombination frequency in, 74, 75, 76
- Drosophila mercatorum*, automixis in, 43
- Drosophila* spp.: inversion polymorphisms in, 88; no evidence for genetic variance of sex ratio in, 149; non-random association between alleles determining isozymes and chromosome order in, 87; ‘sex ratio’ condition in, 147
- Drosophila subobscura*: female choice in, 168, 169–70, 175; inbreeding and survival of offspring of, 126, 127, 129
- dry habitats: favour parthenogenesis, 109, and high rates of selfing in hermaphrodites, 128; sex ratio of dioecious plants in, 162–3
- earthworms, parthenogenetic, 46
- ecosystems: convergent and divergent, 27–8; number of species in, as equilibrium between speciation and extinction, 27
- eggs: cleavage of, initiated by sperm with-

Subject index 217

- out contribution of genetic material (*amblystoma*), 47; development of unfertilised, not uncommon, 43, 49; size of mother and size of, 156; winter, of *Daphnia*, 58–9, and rotifers, 6, 61
- enchytraeid oligochaets, thelytokous, 45
- endomitosis (pre-meiotic doubling of chromosomes), in some examples of thelytoky, 46, 48
- environment: evolutionary advance by one species experienced by other species as deterioration of, 26, 110; how rapidly can it change without causing extinction? 21–2; how much more rapidly can a sexual population adapt to, than an asexual one? 22; parthenogenesis tends to occur in biologically simple, 109–10; and sex ratio, 162–3; spatially varying, with migration, selection for sex and recombination in? 89, 96–9; variable or unpredictable, selection for sex and recombination in? 4, 10, 89, 90–6, (evidence from geographical distribution of parthenogenesis) 108–9
- enzymes, polymorphic, 24; frequencies of, as measure of frequency of selfing in plant populations, 128; identification of biotypes by study of, 57
- Ephedra viridis* (mormonitea): sex ratio of, in dry and damp sites, 162
- Escherichia coli*: competition between wild-type and mutator strains of (non-recombinant), 112, 113; mutation rate in, 189, 190–1
- eukaryotes: origin of, 6; scheme for origin of meiosis in, 7, 8
- evolution: potential of parthenogenetic strains for, 51–7; transient and steady-state models of, in asexual population, 30–1, and in sexual population, 28–30
- evolution rate: compared with rate of change in artificial selection experiments, 11–12; depends on balance between exhaustion of variability by directional selection, and its generation by mutation, 12–13; limit on (cost of selection), 23, 24; mutation rate and, 30; in response to continuous change of environment, 19–23; in response to selection based on pre-existing genetic variability, 13–16; in response to sudden change in environment, 16–19; sex and recombination and, 11, 32
- evolutionarily stable strategy (ESS), 147–8, 172; parental care in terms of, 175, 176–9
- extinction of species: balance between parthenogenetic mutations and, 70; lack of evolutionary potential and, 52, 54; law of constant (Van Valen), 26–7; relation between evolution rate, lag load, and, 19–23; simulation of random process of speciation and, 52
- female choice, 168–71; in maintenance of sexual reproduction? 174; of males with 'handicaps', 171–4; and sexual dimorphism, 183–4
- females: natural populations with excess of, 6, 62, 64, 65, 160; species with larger, 185
- fertilisation: internal or external, and male parental care, 178–9
- fish (bony): parental care in, 182–3; parthenogenetic species of, 47
- fitness, genetic: estimate of, for parthenogenetic and sexual forms of *Poecilopsis*, 68–9; 'inclusive', 114, 165; low, of inbred offspring, 124; of males with 'handicaps', 173; mean, of a population, 21, 122–3; additive genetic variance in, 170, 171, 172; variance of, between sexes, and sex ratio, 162–3
- fitness sets: for allocation of resources in hermaphrodites, 130–2; factors favouring convex set, in plants, 132–3
- flukes, digenetic, 61
- forest plants: proportions of hermaphrodite, monoecious, and dioecious among, 136, 137
- frogs, parental care in, 182
- games, theory of, 147, 175, 176
- genes: number of, and mutation rate, 191; for recombination frequency, 76; 'hitch-hiking' and selection of, 112–14; ways in which sex ratio could be influenced by, 146–7, 157
- gonochoristic, use of term, 125
- gorilla, female transfer in, 142
- group selection: argument for maintenance of sex by, 2–5; argument weaker for plants, 51; 'balance' argument against, 5–6; and coexistence of

218 *Subject index*

- sexual and parthenogenetic forms, 68; and cyclical parthenogenesis, 59, 62; fails to explain origin of sex, or maintenance of recombination, 6, 9, 71; share of, in maintaining sex, 6, 37–8, 69–71
- grouse, parental care in, 180
- gynodioecious plants, 42, 147
- gynogenesis (females produce eggs parthenogenetically after mating), 47, 50
- haplo-diploid cycle, scheme for origin of, 7, 8–9
- haplo-diploid systems, 42, 43; sex ratio in, 150
- Hawaiian Islands, dioecy in flora of, 136
- Helix*, self-incompatible hermaphrodite, 125
- Hemidactylus* (geckos), parthenogenesis in, 46
- Hermaphroditism, 125; advantages of, 39–41, 50; in animals, 138–9; models for evolution of, 130–5; as primitive condition in higher plants, 135; resource allocation in, 40, 130–3; selection for self-compatibility in, 125–6; selection for self-sterility in, 126–7; self-sterility in, partial, 127–8; sequential, 135
- heterokaryon: evolution to diploid cell from, 7, 8; selective advantage of, 7
- heterostyly, 86, 125, 136
- Hieracium* (Compositae), apomictic species complex of, 55
- Hill–Robertson effect, 10, 32, 122
- histones: one amino-acid difference in, between animals and plants, 30
- ‘hitch-hiking’ effect, 10; and gene selection, 112–14; in models for evolution, 30; and rate of fixation of mutations, 112, 113, 192–3; and recombination, 114–17, 123
- Hordeum vulgare*, partial outcrossing in, 128
- horse ancestors: rate of change of molar teeth of, 11–12
- Houttuynia cordata* (Saururaceae), apomict with no sexual relatives, 31, 65
- humans: incest taboo in, 142–5; sex ratio in, 158–9, (no evidence for genetic variance of) 148–9, (reproductive condition of mother and) 162
- hybrid vigour, 7, 9; less marked in habitually selfing species? 129
- hybridisation: and origin of sexual dimorphism, 184; association with parthenogenesis, 48, 50–1
- Hymenoptera: arrhenotoky and thelytoky in, 42–3, 52–3; sex ratio in, 150, 165–6, (size of host and, in parasitic species), 162
- Iceryini (Homoptera), arrhenotoky in, 53
- inbreeding: in arrhenotokous species of insects and mites, with large excess of females, 160; avoidance of, in animals, 139–42, and in humans, 142–5; avoided in plants by dioecy and monoecy, 135; and chiasma formation in rye, 120; and fitness of offspring, 124, 125; genetic consequences of protracted, 129–30; in hermaphrodite plants, 160; mechanisms to prevent, 125; and selection for higher recombination, 119–22; and sex ratio, 160–1
- incest taboo, in humans, 142–5
- inversions: and linkage disequilibrium, 87–8; non-random association of, with enzyme loci 84; in thelytokous diptera, 56
- islands: colonists of, 136; number of species on, 27
- isogamy, 38–9; evolution to anisogamy from, 39, 146, 151–5; no advantage of parthenogenesis in, 42
- Jacana spinosa*, polyandry in, 181
- kibbutzim, Israel: marriages of children communally reared in, 143
- Lacerta* (wall lizards), parthenogenesis in, 46
- lag load, *see* load, genetic
- Lagopus lagopus* (red grouse), parental care in, 150
- Leipoa ocellata* (mallee fowl), parental care in, 182
- ‘lek’ species of birds, 170–1
- lima bean, selection for recombination frequency in, 74, 75
- linkage disequilibrium: in natural populations, 73, 83–5, 85–8; and selection for reduced recombination, 80–1; processes acting to maintain, 81–3,

Subject index 219

- randomly generated, favours recombination, 112, 122, 123; and rate of evolution, 14–16
- lizards, parthenogenetic species of, 46
- load, genetic: lag load, 21–2, 26; concept of, irrelevant for intraspecific competition, 26; criticisms of concept of, 23–5; upper critical value of, at which population becomes extinct, 22
- Lolium multiflorum* (grass), partial outcrossing in, 128–9
- Lycaon pictus* (hunting dog), female transfer in, 142
- Macaca mulatta*, male transfer in, 142
- male sterility: in gynodioecious plants, 147; selection for, in parthenogenetic varieties derived from hermaphroditic ancestors, 41–2
- males: female choice of, 140, 171–4; parental care by, 39, 42 (*see also* parental care)
- mammals: with larger females, 185; parental care in, 179
- Marmota monax* (woodchuck), sex ratio in, 163
- mates: hermaphrodites' advantage in finding, 134; local competition for, and sex ratio, 160–1
- meiosis: elimination of male genes during, in *Poeciliopsis* 47; in eukaryotic sex, 8, 9, 39; in male Diptera (chiasmata absent), 9, 73; and sex ratio, 160; suppressed in apomixis, 43, 45; usually prerequisite for development in animals, not in plants, 51, 70
- meiotic drive, 147, 164
- mice, sex ratio in, 149–50
- Micromalthus debilis* (beetle), arrhenotoky in, 53
- migration: into islands, 27, 136; and selection for recombination, 13, 96–9
- monoecy, 125, 135; ensures outcrossing, 138; in forest trees, 137, 138
- monogamy: biparental care common in, 183; in carnivores and primates, 179
- Moraba virgo* (grasshopper), automixis in, 46, 56
- 'Muller's ratchet', 10, 33–6, 119, 122; and mutation rate, 191, 192
- mutations: balance between selection and, 12–13; chain-terminating, 190; diploidy as protection against deleterious, 9; fixation of, 17–19, 28; 'hitchhiking' and rate of fixation of, 30, 112, 113, 192–3; rate of, 188, 192; rate of, in infinite asexual population, 193–4; rate of, and rate of evolution, 30; selection for recombination in presence of, 117–19
- mutator gene in *E. coli*, 112, 113, 192–3
- Neomorpha acutirostris* (huia), sex differences in, 185
- Neurospora crassa*: mutation rate in, 109; mutations in recombination rate of, 77
- Notobranchius guentheri* (cyprinodont fish), 172–3
- Otiorrhynchus scaber* (weevil), parthenogenesis in, 57
- Oxyura* (stiff-tail ducks), uniparental care in, 181
- Papilio memnon* (butterfly), supergene controlling mimicry patterns in, 85–6, 87
- Papio anubis* (baboon), male transfer in, 141
- parental care, 175, (paternal) 39, 42; models of strategies for, 176–9, (applied to vertebrates) 179–83
- parthenogenesis: in animals, 5, 42–9; arrhenotoky and thelytoky in, 42–3; coexistence of genetically different clones in, 31; coexistence of sexual species and, 58, 66–9; cyclical, 58–62, 70; evolutionary potential in, 5, 51–7; facultative, 5, 62–5, 70; geographical distribution of, 26, 57, 108–10; group selection and, 5, 27–8, 68; meiotic, with pre-meiotic doubling of chromosomes, 46, 56; origins of, 47–9; in plants, 49–51; pseudogamy in, 50, 65–6; short-term balance between sexual reproduction and, 62–3; twofold advantage of, 2–4, (factors invalidating advantage) 38–42
- Parus major* (great tit), inbreeding in, 141
- Passer domesticus* (house sparrow), dimorphism in, 185
- peacocks, female choice in, 171, 173
- Peromyscus maniculatus* (deer mouse), inbreeding avoidance in, 140
- Phasmidae (stick insects), facultative parthenogenesis in, 63

220 *Subject index*

- planktonic animals, hermaphroditism in, 138, 139
- Poa* (grass): apomictic species complex of, 55, 70; pseudogamy in, 65
- Poecilia formosa*, parthenogenesis in, 47, 48, 50
- Poeciliopsis*, parthenogenesis in, 47, 48, 68–9
- pollen, fertilisation of endosperm by (pseudogamy), 50, 65–6
- pollination: by insects, in plants with hermaphrodite flowers, 132; by wind or water, in dioecious and monoecious plants, 133, 136, 138
- polyandry, in birds, 181
- polygamy: sexual selection more intense in, 184; size of males, and mating success in, 133
- polygyny: in birds, 180–1; and sexual dimorphism, 186–7
- polyploidy: apomixis associated with, in plants, 50–1, 55; in thelytokous encytraeids, 45
- Potamopyrgus antipodorum* (snail), facultative parthenogenesis in, 64–5, 67
- Potentilla* (Rosaceae): apomictic species complex of, 55, 66, 67, 70; pseudogamy in, 65
- primates: female transfer in, 142; monogamy in, 179; sexual dimorphism in, as function of body size, 186
- Primula*, supergene controlling heterostyly in, 86
- prokaryotes: capacity for genetic recombination in, 7; mutation rates in, 190
- protandry in hermaphrodites: 134; in plants, 125
- protogyny in hermaphrodites, 134
- pseudogamy, 50, 65–6
- psocids, thelytoky among, 53, 54
- Quiscalus quiscula* (grackle), sex ratio in, 159, 171
- recombination, genetic: as adaptation to biologically complex environment, 110; cost of, 85; delays accumulation of harmful alleles, acting as a form of repair, 33–6; and evolution rate, 11, 32; genes affecting, may act locally or generally, 76; genes for high rate of, 112–13; genetic variance for frequency of, 72, 73–5, (nature of variance) 76–7; group selection unable to explain maintenance, of, 6, 9, 71; ‘hitch-hiking’ and, 114–17, 123; origin of, in repair processes for prokaryote DNA, 6, 7; selection against, at equilibrium in uniform environment, 72, 77–83, 89; selection for, in a fluctuating environment, 89, 95–6, in a new region, 15, 96, in presence of recurrent mutations, 117–19, and in spatially varying environment, 89, 96–9; short-term advantages of, 10; short-term disadvantages of, 9, 73; selfing, and selection for higher rate of, 112, 119–22; *see also* sexual reproduction
- red deer, fighting of males of, 169
- Red Queen hypothesis, 26, 54, 110
- resource allocation: to brooding of young, 139; to male and female functions in hermaphrodites, 40, 130–3
- Rhea americana*, parental care in, 182
- rotifers: bdelloid, without males, 53, 69; of order Monogononta, 53, 60; with overwintering sexual and immediately hatching asexual eggs, 6, 61
- Rubus* subgenus *Eubatus* (blackberries), apomictic species complex of, 55
- Rumina* (snail), self-compatible hermaphrodite, with occasional outcrossing, 130
- rye: chiasma frequency in, 120
- Salmonella typhimurium*, mutation rate in, 187
- Schistocerca gregaria*, selection for recombination frequency in, 74, 75, 76
- seeds, size of, 156
- selection: artificial, for automixis, 47, for blood pH in mice, 149–50, for number of abdominal bristles in flies, 11, 12, and for recombination frequency, 74, 75; cost of (Haldane), 23, 24; group and individual, 1–2 (*see also* group selection); ‘hard’ and ‘soft’, 24–5, 25–6; kin, 66; models for evolution with intermittent, 28–9, 30, and with long-continued, 29, 30; ‘second order’, on genes altering mutation and recombination rates, 112–13; sexual, 10, 168–87; threshold, 24, 25
- self-fertility: advantageous if a mate is not available, 124; selection for, in hermaphrodites, 125–6

Subject index 221

- self-sterility, in plants, 125, 126–8
- selfing species: selection for self-incompatibility in, 125–9; in tropical environments, 26; selection for higher recombination in, 112, 119–22
- sessile habit: association of hermaphroditism with, in animals, 138, 139; diminishing returns from increased production of seeds and pollen in, 133
- sex, mechanisms for determining, 164
- sex ratio, 40, 64, 146–8; genetic variance in, 148–50; with local competition for mates, 160–1; with random mating, 157–60; in social Hymenoptera, 165–6; stable, 157, 166–7; when value of male and female varies with circumstances, 161–3
- sexual dimorphism: barrier to hybridisation in early stages of speciation, and production of, 184; ecological differentiation and, 184–5; polygamy and, 184, 186; sexual selection in production of, 183–4; in size, as function of body size in primates, 186; in size, with larger females, 185
- sexual reproduction: as adaptation to biologically complex environment, 90, 109–10; and evolution rate, 11, 32; female choice in maintenance of? 174; long-term advantages of, 4, 37; is it maintained by group selection? 37–8; origin of, 6, 7–9, 71; and rate of evolution, 22–3; share of group selection in maintaining, 6, 69–70, 70–1; short-term advantage of, 5, ('balance' argument for) 5–6, 38, 57–8, 66–7, (with sib competition) 89–90, 99–108, (in fluctuating environment?) 4, 10, 89, 90–6, (in spatially varying environment?) 89, 96–9; short-term balance between parthenogenesis and, 62–3; *see also* recombination, genetic
- sexual selection, 10, 168; female choice in, 168–74; monogamy, polygamy, and sexual dimorphism in, 183–7; parental care and, 175–83
- sib competition: between offspring of sessile organisms, 133; effect on, of many loci concerned with a single environmental feature, 107–8; patterns of dispersal and, 108; short-term advantages for sex and recombination, 10, 99–100, 123; simulation of, 100–7
- Sipylodea sipylus* (stick insect): parthenogenesis in, 63, 67
- skin grafts, showing genetic identity of different populations of *Cnemidophorus*, 56
- skua, Arctic: sexual selection in, 184
- snails, supergenes controlling shell colour and pattern in, 86
- Solenobia triquetrella* (moth), parthenogenesis in, 57
- Solidago* (golden rod); size of seeds in, 156
- species abundance distribution (Preston), 27
- Spergula arvensis* (corn spurrey), genetic polymorphism of seeds of, 156
- stickleback, polygyny in, 183
- supergenes, promote linkage disequilibrium, 85–7
- syngamy, in eukaryotic sex, 8, 9, 39
- Taraxacum* (dandelions): asexual clones of, produce functionless pollen, 41; species complex of, 54–5, 67
- Thalictrum fendleri* (meadow rue): sex ratio of, in dry and damp sites, 162, 163
- thelytoky (production of females from unfertilised eggs), 43; geographical distribution of, 54; with normal meiosis and subsequent fusion of pronuclei or early cleavage nuclei (automixis), 43–5, 47–8; with pre-meiotic doubling of chromosomes, 46, 48; with suppression of meiosis (apomixis), 43, 48, (in animals) 45–6, (in plants) 49, 50–1, 54, 55; taxonomic distribution of, 53, 54
- Thysanoptera (thrips), arrhenotoky in, 53
- trees: hermaphrodite, monoecious, and dioecious, 136, 137; pollination of, 136, 138
- Tribolium castaneum*, selection for recombination frequency in, 74, 75
- tropics: forest trees in, 136, 137; parthenogenesis rare in, 109
- turkeys, hatching of unfertilised eggs of, 45
- variability, genetic: generation of, by chiasma formation, 8; release of, by recombination, 11; and response to selection, 11–13; in populations with and without sex, 13–16

Cambridge University Press
978-0-521-29302-0 - The Evolution of Sex
John Maynard Smith
Index
[More information](#)

222 *Subject index*

- variance, genetic: in fitness (additive), 170, 171, 172; in frequency of recombination, 9, 72, 73–7; in sex ratio, 148–50
variation, selectively neutral, 24
Volvocidae: size of colony, and degree of anisogamy in, 155
weevils, apomictic, 45–6; geographical distribution of, 54
Xyleborus (bark beetles), arrhenotoky in, 53