

Index

The number in *italics* refers to the figure on the page indicated.

The suffix ‘t’ refers to the table on the page indicated.

- abundance: absolute, of fish, 344, 355–6; algal cover, 185; benthic fauna, sampling methods for, 106–18, 120–5; fish egg, 358; meifaunal, 142ft; plant, measurement of, 53–62 (qualitative estimation, 55–6; quantitative estimation, 56–9); relative, of fish, 344, 350, 355; scales(s), 169–71, 170t, 174, 177–9, 216–17, 217; species, 201 (on rocky shores, 174, 183); zooplankton, 359
- Acartia* sp., 296, 300
- Acartia clausii*, 294
- accretion, measurement of 66–7
- acridine orange dye, for bacteria, 246, 248
- Actinia equina*, 180, 184
- adenosine triphosphate (ATP) determination, 249–50
- adenylate energy charge measurement, for estimation of bacterial activity, 254
- Agassiz trawl, 121
- age determination, in fish, 361–3
- Agropyron pungens*, 14t
- Agrostis stolonifera*, as indicator species, 70t 71
- air–water interface, 241
- Alaria esculenta*, as indicator species, 164
- alcohol, as preservative, 135
- Alcyonium digitatum*, 216t
- algae/algal: abundance scales for, 170t; blue-green, 293, 300 (identification of, 407; as pollution indicator, 89); cover estimation, 180, 181, 185; identification of, 407–8; microscopic, sampling in salt marshes, 60; morphology variation, 189–91; symbiotic, in plankton, 299; *see also* macroalgae; microalgae
- amphipod(s), 297, 300; identification of, 412
- angiosperms, zonation in salt marshes, 38
- Anglesey, 168, 168, 179, 180, 234
- Annelida, identification of, 410
- Antedon bifida*, 216t
- Anurida* sp., 167
- Arctica islandica*, 136
- Arenariomyces trifurcatus*, 265
- Arenicola marina*, 83, 88
- Armeria maritima*, 59, 69, 70t
- arrow worms, *see* chaetognath(s)
- arthropods, rapid survey of, 73–5
- ascomycetes, 258; marine, on driftwood, 263t
- Ascomycotina, 262, 265, 267
- Ascophyllum* sp., 164, 172
- Ascophyllum nodosum*, 166
- Aster tripolium*, 14t, 69
- Atlantic Bight, 299
- Atriplex hastata*, 62
- auks, 383, 385, 395, 396; population, 387; *see also* Guillemot, Puffin, Razorbill
- autoradiography, for estimation of bacterial activity, 251
- Baccaris halimifolia*, as indicator species, 41
- bacteria(l), activity, 245, 251–4; marine, 238–58, 312 (biomass estimation of, 248–50; distribution, 238–9; environmental effects of, 239–40; identification of, 406–7; numerical estimation of, 245–8); role in food webs, 297–9; sampler for, 243
- baited hooks, for fish collection, 345t, 358

INDEX

441

- baited techniques, for collection of marine fungi, 261–8 (blending and plating, 264–5; sea-water, 267; sand, 268; sub-sampling, 265)
- Balanus perforatus*, abundance scale for, 170t
- Baltic Sea, 108, 286, 300; coast, birds on, 375; salinity, 166
- barnacle(s), 172, 180, 181, 183, 185; abundance scale for, 170t; identification, 187, 411; larva, 294; *see also Balanus; Chthamalus; Semibalanus*
- basidiomycete, marine, 259
- bass, 344
- beached-bird surveys, 390–9, 396; corpse recovery in, 393; oiling of corpses in, 391–2, 394, 394, 395–8, 397
- Beatrice Oilfield, *Scoloplos armiger* distribution in, 22
- behaviour, fish, 343, 344–6, 348
- benthic organisms, distribution, 87–8 (sampling strategies, 106–18, 120–5)
- biomass: of fish, 344; measurement of, 59, 61, 62, 94, 137 (bacteria, 245, 248–50, 251; macrofauna, 96, 109; marine fungi, 270–1; meiofauna, 151; phytoplankton, 90; phytoplankton, 288–9; plankton, 303, 310, 312–16; samples for, 94, 122); of meiofauna, 141–3, 142t, 145; of plankton, 298, 299, 300, 308, 309; zooplankton, 302; *see also plant performance*
- birds, 374–400; identification of, 413
- Blakeney (England), 67
- bloom, phytoplankton, 391, 300–2, 301, 319
- boats: for fish observation, 347; for sampling, 118, 316 (safety procedures for, 430–1)
- Bogorov tray, 311, 315
- Bolinopsis* sp., 296
- Botany Bay, 111, 112
- Brent Goose (*Branta bernicla*), 378; Dark-bellied, 380, 381
- Bridgwater Bay, 45, 66
- Bristol Channel, 285t, 299–300; meiofauna in, 142t
- brittle-star larva, 294
- bryozoa, identification of, 413
- Buergenerula spartinae*, 258
- bulk-staining, 136
- caesium, radioactive, 281 (Cs-137, 67)
- Cairnhead, 165
- Calanus* sp., 295–6
- calorimeter, microbomb, 312
- capture techniques, for fish, 346, 350–9; damage to fish by, 346
- carbon, particulate, analysis of, 312
- Carcinus maenas*, 69, 83, 180
- Caryophyllia smithii*, 216t, 220
- catch, recording procedure for, 359
- catchability coefficient, 355–6
- Cefni salt marsh (Wales), 70
- Cellepora pumicosa*, 216t
- central-limit theorem, 87–8
- Cerastoderma edule*, 83
- Ceriosporopsis capillacea*, 259
- chaetognath(s) (arrow worms), 294, 306
- Chlorophyceae, on tidal flats, 95
- chlorophyll: concentrations, 322–3, 322, 359; determination/measurement, 93–4, 313–4, 316, 324, 328–24 (variability 320t); extraction, 329 (for estimation of microbenthic algal biomass, 90–1, 92–3; of phytoplankton biomass, 312–14; from sediment cores, 90, 92–3); fluorescence of, 311–12; –salinity diagrams, 287; sea water profiles, 282
- choanoflagellate(s), 297, 298
- Chthamalus* sp., 165
- Chthamalus montagui*, 179
- Chthamalus stellatus*, 179
- ciliate(s), 297–9, 299, 300, 301, 310, 311; tintinnid, 298, 298, 302, 311
- Cirripedia, identification of, 411
- Cladophora* sp., identification of, 408
- classification: data, 13–16, 14t, 15t, 20, 404; floristic, 63–4; plant community, 63–5; salt marsh, 63–5
- Clavatospora stellatacula*, 266
- cliff(s): -nesting seabirds, 383–8; safety procedures for, 430
- clothing, for field trip, 426
- Clupea harengus*, 358
- CLUSTAN program, 65
- cluster analysis, 20, 65
- coccolithophorids, 308, 310, 311
- Coccilithus* sp., 293
- cockle-counting apparatus, 99, 99
- Coelenterata, identification of, 409
- cohort analysis, 363
- coliform bacteria, 256–8
- colony, breeding, of seabirds, 383–90, 387t; bird mortality near, 394; mapping, 389–90
- community(ies): analysis, 21 (rocky shore, 185; salt marsh, 63–5); on subtidal wreck, 218–19; stone/boulder, 166, 189
- computer(s): for data handling, 16, 64–5, 175, 181, 182, 322; for interpretation of digital remote sensing data, 27, 32–3, 35–6; for multivariate analysis, 20; program for sigma-T (seawater density) calculation, 326
- conservation, 1, 375; value, 73–5, 107, 159, 198 (criteria, 71–2; of mud flats, 82)

442 INDEX

- contaminants: collection of samples for analysis of, 121; of sediments, 106–7, 108
- continental shelf, 281–2
- cooling system intake screens, retention of fish by, 355
- Coot (*Fulica atra*), 380
- copepod(s), 233, 285t, 294, 300, 306; calanoid, 295–6; harpacticoid, 141, 142t, 143–4, 145, 149, 151; identification of, 411
- core(s), sediment, microalgae in 90–3.
- core sampling: for fish, 345t; for meiofauna, 117, 148–9; sediment, 96–7 (for bacteria, 244; for fauna, 229–30); soil, for salt marsh fauna, 69
- corer(s), 96–7, 101, 121–2; Craib, 148–9; Reineck (box), 122
- Corollospora maritima*, 260, 265
- Corophium* sp., 83
- Corophium volutator*, 132
- correlation tests, 21
- Corynactis viridis*, 216t
- Coulter counter, 312; for particle size distribution, 126
- counting techniques: for bacteria, 246–8 (total, 246; viable, 247–8); for estimation of phytoplankton populations, 90, 91–2
- counting units: for seabirds, 384–5; for terns, 389
- cover-abundance scales, 56, 58
- cross-staff levelling, 176–7, 176, 178
- ctenophore(s), 294, 296
- culture techniques, 239; for bacteria, 247, 255
- Cumacea, identification of, 411
- Curlew (*Numenius arquata*), 377
- currents, measurement of, 289–90; *see also* tidal currents
- cyanobacteria, 293, 312; *see also* algae, blue-green
- cycles: seasonal, for plankton, 300–2, 301; tidal effect of on sampling salt marsh, 68, 69–70
- Cyprinus carpio*, 358
- Czekanowski's Quantitative Index, 20
- DAFOR scale, 55–6, 55
- dangers, potential, assessment of, 424
- data: analysis, 17–21; 320 (Fourier series, 322; ordination, 20; (*see also* rarefaction)); classification, 13–16, 14t, 15t, 20; comparability, 207, 208; fish (length, 360; weight, 360); handling, 16–23 (microcomputers for, 16); presentation, 22–3, 320–3; recording methods, 16–17 (for Diptera, 74)
- datum levels/points, 42, 49–51, 65, 160, 177; for accretion measurement, 67
- decapod(s), 297; identification of, 412
- Decca Navigator, 119, 213
- decompression, of divers, 200
- Dee estuary, Knot population in, 379
- Delta area (Netherlands), 41; distribution of *Orchestia* sp. in, 84
- density of sea water, 282; calculation of, 326; profile, 282, 289
- Deuteromycotina, 265
- diatom(s), 291, 292, 300, 301, 302; benthic, 293, 308, 309; identification of, 407
- Dictyocha* sp., 292, 293
- Diet determination, in fish, 361
- Digitatipsora marina*, 259
- Dinobryon* sp., 293
- dinoflagellates, 291, 292, 293, 298, 299, 300, 302, 308, 309, 310, 311; identification of, 407; production of toxic substances by, 291
- Diplomastigomycotina, 267
- Diptera, rapid survey of, 73–4
- distress signals, international, 427
- distribution, species: diurnal, 167; effect of biological interaction on, 169; seasonal, 167–8
- Diver(s) (*Gaviidae*), 380, 381
- diver(-operated) sampling techniques, 98–9, 121–2, 209–34, 203t, 358
- diversity indices, 18–19, 404
- diving, 198–234; dry-suits for, 200; for fish observation, 348; team, for surveys, 199 (safety procedures for, 432–6)
- dominance index, Simpson's, 19
- dominant species, definition, 19–20
- dredges, 120–1, 345t
- drogues, 290, 316, 324
- dry weight determination, 151; zooplankton, 314
- duck(s), 378, 380; Long-tailed (*Clangula hyemalis*), 381
- Dunlin (*Calidris alpina*), 377
- Echinocardium cordatum*, 115
- Echinodermata, identification of, 413
- Echinus* sp., 221
- Echiura, identification of, 410
- echo sounder(s), 350; for sampling station relocation, 213
- egg counting, fish, 360
- Eider (*Somateria mollissima*), 381, 397
- Ekofisk oilfield, 144
- electrofishing techniques, 352–3; safety procedures for, 436–8
- electron transport system (ETS) method for estimation of bacterial activity, 253–4
- elutriation, for separation of macrofauna, 132–3; for separation of meiofauna, 149–50

INDEX

443

- Ems estuary, 98, 142t
 English Channel, 350, 395
Ensis sp., 122
Enteromorpha sp., 41, 167; identification of, 408
 environment, chemical, effect of on bacteria, 241
 environmental factors, measurement of, 102
 epifluorescence, for counting plankton, 311–2
 equipment, field, 426
Escherichia coli, as indicator, 240, 256–8
 estuary(ies): bird distribution in, 374–5;
 fish in, 344; flushing time, 284–6, 285t,
 289; hydrography of, 282–90; mixing in,
 288, 289, plankton sampling in, 317
 euphausiid(s), 285t, 296, 305; identification of, 411
Eurytemora sp., 296, 300
Eutrepia sp., 293, 300
 eutrophication, 319; *see also* bloom
 excursion, field: preparation for, 425;
 procedure following, 425–6
 exposure: of rocky shores, 172 (assessment of, 161–5; biological indicators of, 163–5; grade, 190, 191); variation in algal morphology with, 189–91
- fall-out, as sediment marker, 67
 Faroe, 164
 fecundity, fish, 360
Festuca rubra, 58; as indicator species, 71
 fetch, 161, 162, 163
 field cards: for intertidal sediments, 84–7,
 85; for rocky shores, 171–3, 172; *see also* habitat recording card
 filtration: of plankton, 310–11, 312; for spore collection of marine fungi, 266
 first aid, 427–8; for electric shock, 437–8
 Firth of Clyde, 284–6
 fish, 342–65; identification of, 413; parent stock size, calculation of, 358–9;
 population, 359; recorders, electronic, 347; sex determination of, 360; stock, 359; tagging, 356–8, 357
 fished area, estimation of, 356
 fishery statistics, 364–5
 fishing gear, 345t, 350–5, 351
 fjord(s), 284–6, 296, 299, 300, 302
 flagellates, 293, 292, 297, 298, 300, 308,
 309; identification of, 408
Flagellospora sp., 266
 flounder (*Pleuronectes flesus*), 344
 flowmeter, for plankton net, 306–7
 fluorometer calibration, 333
 fluorometry, for chlorophyll determination, 313–14, 324, 330–1, 334
 flushing, estuarine, 299, 302 (time-scale estimation, 327–8); time, 284–6, 285t,
 289, 324
 foam samples, for marine fungi, 269
 food web/chain, 297–300, 361, 363;
 detritus, 297, 299, 300, 302
 Foraminifera, 299; identification of, 408;
 sampling, 69
 formalin/formaldehyde, as preservative, 92, 135, 137, 149, 233, 244, 308–9, 310,
 311, 259; safety procedures, 438
 Fouling Organisms, Marine, Catalogue of Main, 406
 Fourier series analysis, 322
 frequency: sampling, 151, 152t; species, measurement of, 57, 59, 202, 225;
 survey, 114–15, 151
 freshwater–brackish interface, 300
Fucus sp., identification, 186
Fucus vesiculosus, 62, 165, 189–91, 190
 Fulmar (*Fulmarus glacialis*), 383–6;
 population, 387
 Fungi: imperfecti, 258; marine, 238–40,
 258–71, 259t (environmental effects, 239–40); identification of, 407; isolation of, 266
- gas–liquid chromatography, 250
 gastrotrichs, 149
 goby (*Gobius* sp.), 344
 Godwits (*Limosa* sp.), 377
 gonad ripeness, in fish, 360
Gonyaulax sp., 291, 292
 grab(s), 120–5, 123, 149, 245t;
 hand-operated, 98, 98, 121, 230;
 samples, 116, 120, 122, 124–5;
 Campbell, 122; Day, 122–4, 123;
 Smith-McIntyre, 124, 230; van Veen, 117, 124–5
 gradient(s): environmental, 147–8, 149,
 151, 158; salinity, 289; species frequency, 202; temperature, 289
Grantia compressa, 216t
 Grebes (*Podicipitidae*), 380
 Grevelingen estuary (The Netherlands), 82–3, 83
 grids: for mapping, 82–3; sampling, 88,
 147, 212
 growth rate estimation in fish, 363
 Guillemot (*Uria aalge*), 383–7, 387t, 395;
 population, 388
 gulls, 374, 383–3, 388; roosting, 382
Gyptis rosea, 133
 habitat recording sheet, 204–5, 205; *see also* field card
Halimione portulacoides, 14t, 60, 62, 70t
Halosphaeria hamata, 258
Haptomastigomycotina, 267

444 INDEX

- Heterocapsa* sp., 300
Heteromastus filiformis, 83
 heterotrophic plankton, 297–9, 300, 301
Hyale sp., 167
 hydrography, 281–90, 316, 321; effect of,
 on plankton distribution, 289
 hydromedusae, 296; identification of, 409
 hydroplane/hydrofoil, for diver towing,
 211–12, 211, 215, 348
 hyphomycetes, 266
- incubation of marine fungi, 262–4
 index/indices: diversity, 18–19;
 Czekanowski's Quantitative, 20;
 dominance, 18–19; morpho-edaphic,
 359; Shannon–Wiener, 18–19; similarity,
 20; Simpson's, 19; vulnerability, 13, 15t
 indicator species: of environmental
 conditions, 151–3; of exposure, 163–5;
 for salt marsh, 41, 70–1, 70t
 infrared radiation recording, 125
 Insecta, identification of, 412
 intercalibration exercises, 10
 International Biological Programme, 54,
 60, 71
 International Council for Exploration of
 the Seas (ICES), 108
 international distress code, 427
 International Waterfowl Research Bureau
 (IWRB), 378–9
 Irish Sea, 185, 281, 393, 395
 Isle of Man, 163, 184
 Isopoda, identification of, 411
Iva frutescens, as indicator species, 41
- jellyfish, 296, 306
- Kattegat, 395
 kelp(s), 216t; forest, 224–5; holdfast,
 232–4
 Kittiwake (*Rissa tridactyla*), 383–6;
 population, 387
 Knot (*Calidris canutus*), 377, 379
- Labyrinthulales, identification of, 407
 lagoons, 286
Laminaria digitata, 189
Laminaria hyperborea, 201, 232, 234
Lanice sp., 122
Lanice conchilega, 83
 LANDSAT satellite, 32–5, 33t, 34t
 Lapwing (*Vanellus vanellus*), 377
 larval recruitment, 114–15, 296
 lateral thinking, for survey planning, 4
 length-frequency analysis, for age
 determination in fish, 361–2
 levels, measurement of, 49–51, 174, 176–7,
 177
- lichens: abundance scale for, 170t; cover
 estimation, 180, 181; identification of,
 408
Limonium sp., 14t, 70t
Limonium vulgare, 69, 70t
 limpet(s), 172; abundance scale for, 170t;
 distribution, 158, 159; identification, 186
 see also *Patella* sp.
 line(s), fishing, 345t, 350, 353
 linescan imagery, 125
Lithothamnion sp., 172; abundance scale
 for, 170t
Littorina sp., 166, 167; identification, 187
Littorina irrora, 68
Littorina littorea, 83, 164
Littorina mariae, 179, 180
Littorina obtusata, 12t, 179, 180
 Loch Ardbhair, 285t
 Loch Creran, 285t, 287, 301
 Loch Eil, 115
 Loch Ine, 206
 Loch Spelve, 321, 323–5
 Loch Striven, 322
 Lofoten Islands (Norway), 393
 log normal distribution, 404; for
 community estimation, 21
 Ludox-flotation method, 150
 Lugol's iodine, acidic, 308, 309
Lulworthia sp., 258
 Lundy Island (Bristol Channel), 206,
 218
Lutraria sp., 122
 Lynher estuary (Cornwall), 145
- mackerel (*Scomber scombrus*), 350, 359
 macroalgae, 95, 238; identification of, 408;
 tidal flat distribution, 95
 macrofauna, 95–9; definition, 95, 106;
 sampling methods for, 95–9, 106–18,
 120–5; separation from sediment, 131–3;
 sorting methods for, 136–7
 Malta, 206
 Mammalia, identification of, 414
 Mann–Whitney U-test, 74
 map(s): for data presentation, 320;
 preparation of, 45–52 (see also
 mapping); for sampling station location,
 88, 290
 Maplin Sands, 13, 14t, 40
 mapping: bird abundance and distribution,
 380; breeding colonies, 389–90;
 instrument, large area, 48–51;
 phytoplankton distribution, 314; rock
 profiles, 214–15, 214; salt marsh
 vegetation, 51–2, 63, 64; small area,
 45–8; species distribution, 3t; tidal flats,
 82–3, using grids, 82; using transects,
 82–3, 220

INDEX

445

- markers: for accretion measurement, 66; for fish, 356–8; plant, 62; for relocation of sampling sites, 42–4, 46, 160, 213; on sea-bed, 212–13; transect (for rocky shores, 181–2; for tidal flats, 100)
- Mastigomycotina, identification of, 407
- Mediterranean Sea, 206, 286
- Meganyctiphanes* sp., 296
- meiofauna(l), 140–53; anaesthesia of, 150; biomass, 141–3, 142t; definition of, 140; density, 141–3; distribution, 141; dry weight determination, 151; generation times, 141; preparation of, 150–1; sampling method for, 117; separation of, from sediment, 149–50
- Menai Straits (Wales), 69
- mesh size, 133–3; for plankton nets, 305–6; for retention of macrofauna, 95, 106, 117, 133; for retention of meiofauna, 140, 150
- Mesodinium* sp., 299, 311
- Michaelis-Menten kinetics, 252–3
- microalgae, 89–94; epipelagic, 89, 91; epipsammic, 89, 91; estimation of, 89–94; sampling methods for, 89–91 (benthic, 89; salt marsh, 60)
- microhabitat, on rocky shores, 158–9
- microp plankton, 291
- microscope, inverted, for plankton counting, 309–10
- migration: bird, 374–5; fish, 344, 347; vertical, of zooplankton, 296
- Milford Haven, 41, 110–11, 319
- mineral salts, 281
- Mnemiopsis* sp., 296
- Mollusca, identification of, 412
- monitoring, biological, 2
- Monostroma* sp., identification of, 408
- Moray Firth, Scotland, 181, 182
- Morecambe Bay, 41, 65; Knot population in, 379; quadrat for, 53
- morpho-edaphic index, 359
- mortality: bird, 390–8 (winter, 394–5); estimation in fish, 363
- mosaic, photographic, for mapping, 48
- most probable number (MPN) method for bacteria, 248
- moulting grounds, for birds, 375
- mounting techniques for marine fungi, 262–4
- mud flats, 81–102; global extent of, 81; *see also* tidal flats
- mud-sledge, 101, 102, 429
- mullet (*Mugil* sp.), 344, 347
- muramic acid determination, 250
- mussel(s), 172; abundance scale for, 170t; farm, 323; spat, 323–4; *see also Mytilus edulis*
- Mya* sp., 83
- mysid(s), 297, 300; identification of, 411
- Mytilus edulis*, 83, 167, 216t
- narcosis: of ciliates, 311; of fish, 352
- navigation techniques, 118–20
- nekton, 294
- Nematoda/nematodes, 142t, 143–4, 145, 147, 149, 151, 152–3, 233; identification of, 409–10
- Nemertini, identification of, 409
- nets: drift, 353; fishing, 345t, 350, 351, 353; hand, 345t, 352; plankton, 304, 305; push, 345t 351, 354–5; seine, 345t, 350, 351, 354, 355, 358
- neuston screen, for collection of foam, 269
- New York Bight, 116
- nitrate, in sea water, 282, 317, 324
- North Sea, 108, 114–15, 116, 119, 281, 285t, 393; oilfields, 111–14, 113
- Nucella emarginata*, 164
- Nucella lapillus*, 164, 167, 180, 184
- nutrient(s), 238, 283–4, 297, 317–18, 319, 322; bacterial requirements, 241–2; inorganic, 317; mineral, 300
- observation techniques, for fish surveys, 346–50
- oil pollution: bird vulnerability to, 396–8; effects, 106–7, 111, 113, 124, 168, 383, 390–9; incidents, 393, 397, 398, 399
- Oligochaeta/oligochaetes, 142t, 143, 151; identification of, 410
- oligotrich(s), 298, 301
- Orbimyces spectabilis*, 261
- Orchestia cavimana*, 84
- Orchestia gammarella*, 84
- ordination analysis, 20, 404
- organic material, dissolved, 317
- oribatid mites, sampling of, 69
- Orkney, 183
- Oscillatoria* sp., 293
- Ostracoda/ostracods, 143, 150, 151; identification of, 411
- otoliths, for age determination, 361–3, 362
- oxygen, dissolved, determination, 252
- Oystercatcher (*Haematopus ostralegus*), 377
- Palmaria palmata*, 168
- Patella* sp., 164, 167, 179, 180
- Patella aspera*, identification, 186
- Patella vulgata*, 184
- Pelvetia* sp., 172
- Pentapora foliacea*, 216t
- Phaeocystis* sp., 293
- Phaeosphaeria typharum*, 258
- phosphorus, in sea water, 317

446 INDEX

- photographic equipment: for small area mapping, 46–7; underwater, 209, 222–3, 222, 224, 349
- photographic film: false colour, 28–9, 30, 31; full colour, 28–9, 30, 31; infrared, 28–30, 31; panchromatic, 289–9, 32
- photography: aerial, 27–32, 87, 171, 347; for mapping, 46–8, 82, 125; for relocation of sampling sites, 42, 260, 181–2; of sampling sites, 87, 94, 174; stereoscopic, 198, 223, 349; for surveying 94, 110, 125, 182, 203, 221–4; underwater, 198, 209, 221–4, 349
- photosynthetic rate, measurement of, 94
- phytoplankton, 281, 284, 286, 291–3; biomass, 288–9, 312–14, 319, 323, 324
- picoplankton, 293, 297, 312
- pigment extraction, for microbenthic algal population estimation, 90–1, 92–4; *see also* chlorophyll extraction
- pitfall trap, 68, 73, 99, 189
- plaice (*Pleuronectes platessa*), 344
- plankton, 280–334; distribution, 291, 293, 296–7, 299–300, 318–20, 323, 324; identification, 314; measurement of 303–16; nets, 304, 305; permanent, 295–6, sub-sampling, 315–16; temporary, 296
- plant: cover estimation, 57; density estimation, 54, 56–7; frequency estimation, 57; performance (estimation in salt marsh 60–2)
- Plantago coronopus*, 70t
- Plantago maritima*, 59, 70t
- plating, marine fungi, 264–5, 266, 269
- platyhelminthes, identification of, 409
- Pleospora* sp., 258
- Pleurobrachia* sp., 296
- Plover: Golden (*Pluvialis apricaria*), 377; Ringed (*Charadrius hiaticula*), 377
- poison station, for fish sampling, 348, 352
- pollutants, 126, 238, 242, 281; analysis for, 126–7 (electronic techniques, 127); *see also* contaminants
- pollution, 163, 233, 234, 395; effects, assessment of, 2, 106–7, 151; indicators of, 151–3; monitoring, 21, 107–15, 124, 133–4, 145, 158, 319, 390; oil, 15t, 106–7, 111, 113, 124, 168, 383
- Polydora ciliata*, 179, 216t
- Pomatoceros* sp., abundance scale for, 170t
- Pomatoceros lamarckii*, 179
- Pomatoceros triquetus*, 179
- population: dynamics for seabirds, 383–4; estimation, benthic microalgae, 90–4
- Porifera, identification of, 409
- position fixing, 99–101, 118–20
- Potamogeton* sp., 94
- production: bacterial, 251; fish, 343, 359, 363 (egg, estimation of, 358–9); meiofaunal, 145; net aerial, 61; phyto benthic, 90; phytoplanktonic, 299–300, 319; primary, 94, 140, 299, 359; *see also* plant performance; standing crop
- protozoa, 297; identification of, 480–9
- Puccinellia* sp., 14t
- Puccinellia distans*, 45
- Puccinellia maritima*, 39, 39, 41, 58, 59, 61, 70t
- Puccinellia rupestris*, 45
- Puffin (*Fratercula arctica*), 383, 385–6, 395; population, 387, 388
- pumps, plankton, 307–8, 316, 325; for fish collection, 345t
- pycnocline, 283, 284, 318
- Pycnogonida, identification of, 411
- pyrenomycetes, marine, 268
- quadrat(s), 5, 9, 45, 53–4, 54, 57, 59, 65, 96, 178, 203t, 206; fixed, 167; box, 231; permanent, 42, 89, 159, 160–1, 168, 181–2, 183–5, 184; point, 94; random 183; reference, 217, 220, 227; shape, 53; size, 53, 183, 184–5, 220; on stone/boulder shore, 189
- Quinaldine, 352
- radar, for position fixing, 100, 119
- radiochemical methods for estimation of bacterial activity, 252
- radiolarians, 299
- radiotelemetry, for fish surveys, 347
- rarefaction, 19
- Razorbill (*Alca torda*), 383–6, 387t, 395; population, 388
- redox potential discontinuity layer, 153
- Redshank (*Tringa totanus*), 377
- reference collection, 137–8, 405
- remote sensing, 27–36, 171; digital, 27–8, 32–6
- research vessels, safety procedures, 431–2
- results, presentation of, 22–3; (proportional circles, calculation of, 23)
- Ribble estuary, Knot population in, 379
- ringing, bird, 375; recoveries from, 390
- river flow/salinity method for flushing time calculation, 327–8
- rock: intertidal, 157–89 (safety procedures for, 430; type, 165); exposure of, 161 (biological indicators of, 163–5); pools, 166, 172, 347; profiles, 214–15, 214; sublittoral, 206 (collection of, 226–7, 226)
- rose bengal stain, 92, 135, 149, 315
- Rotenone, 352
- Ruppia* sp., 94

- Sabatieria pulchra*, 153
 safety, 424–38
Sagitta sp., 296, 299
Salicornia sp., 14t, 63
Salicornia europaea, 62
 salinity, 166–7, 241, 283–90, 287, 322, 344; effect of, on plankton distribution, 299; limits for marine fungi, 258–9; measurement of, 289, 317, 324; profiles 282–3, 282
 salinometer, 289
 salmonids, 344, 361
 salt marsh(es), 14t, 34, 38–72, 238; classification of, 63–5; conservation value of 3t, 71–2; in estuary systems, 41–2; fauna sampling of, 67–70; limits of, 40–1, 46; safety procedures for 428–30; scientific value criteria of, 72; topography, 51; transition, 61–2; vulnerability to oil spill damage, 15t
 sample(s): collection of, by divers, 225–34; containers, 135–6; dilution of bacterial, 245–6; elutriation of, 132–3; preservation, 135–6, 137–8, 359 (bacteria, 244; meiofauna, 149; plankton, 303, 308–9); random location of, 88, 109; replication, 115–17, 147–8; size, 9–10, 87–8, 90, 108, 117, 206–7; vessel for bacteria, 240–1; washing, 131–2
 sampler(s): cockle, 99, 99; diver-operated, 98–9, 121–2, 198, 227–32; flushing, 97; Johnson-ZoBell, 242, 243; for macrofauna, 96, 117–18, 120–5, 123; Niskin, 242; plankton, 304, 305–8, 306 (Bongo net, 307, 316; Clarke-Bumpus sampler, 307; Hardy-Longhurst sampler, 307; pumps, 307–8, 316, 325; Working Party 2 net, 307); suction, 98, 121–2, 198, 203t, 228, 228, 229, 230–1, 348 (with reference cylinder, 231–2, 231)
 sampling: area, 9–10; deck, 359; depth, 202; equipment, diver's, 209; errors in bird surveys, 376–8, 386–7, 387t; frequency, 10, 88, 108, 115, 290, 317, 318–19, 386, 392; mixed age salt marsh, 60; point, 58–9, 223; pump, 242, 305, 307–8; qualitative, for plankton, 305; quantitative (from bedrock, 227–8, 227; of sediments, 229–32; zooplankton, 305–8 *see also* survey, quantitative); strategy, 7–13, 87–8, 108–17 (temporal considerations, 12–13); stratified, 116; units, 201–2
 sampling methods, 5, 67–70, 203t; for bacteria, 240–4; for marine fungi, 259–70; for intertidal sediments, 81–102; for macrofauna, 95–9, 108–18, 120–5; for meiofauna, 117, 148–9; for microalgae, 89–91; for plankton, 281, 305–12; standardization of, 10–12, 108–9
 sampling site(s)/station(s); damage to, 44–5, 61; distribution of, 109–14, 111, 112, 113, 241; evaluation of, 71–2; grid(s), 87–8, 109, 110–11, 111, 112, 113, 114; location of, 6–7, 46, 100, 118–20, 147–8, 174, 202–6, 213, 290, 318; relocation of, 13, 42–4, 100, 118–20, 200, 212, 213–14
 sand grain method, for sporulation of marine fungi, 267
Sanderling (*Calidris alba*), 377
 Sandpiper, Purple (*Calidris maritima*), 377
 satellite(s): imagery, 125, 347; for navigation, 119–20, 213; for remote sensing, 27, 32–5, 35t
 SatNav satellite, 119–20
 scales, fish, for age determination, 361–2, 362
Scirpus maritimus, 70t
Scolelepis sp., 83
Scoloplos sp., 83
Scoloplos armiger, distribution in Beatrice Oilfield, 22
Scomber scombrus, 359; *see also* mackerel
 Scoters (*Melanitta*), 381, 297
 SCUBA diving, 198, safety procedures for, 432–6; for fish observation, 348
 Sedgewick-Rafter chamber, 311
 sediment(s): chemistry, 241–2; effect of tidal pressure on, 239; extraction of fauna from, 131–6, 149; intertidal (biological sampling of, 81–102; meiofauna in, 142t); particle size analysis, 126; salt marsh, faunal sampling of, 68–9; samples for bacteria, 242–3 (preparation of, 254–5); samples for marine fungi, 270; subtidal, faunal sampling of, 106
 sedimentation: chamber, for plankton, 309–10; in salt marshes, 38 (measurement of, 66–7)
 Sellafield, 281
Semibalanus balanoides, 182, 184, 294
Serratia indica, 257
 Severn Barrage, 107
 Severn Estuary, 164; salinity, 166
 Shannon-Wiener Index, 18–19, 147
 Shearer method, for sporulation of marine fungi, 267
 Shearwater, Manx (*Puffinus puffinus*), 383
 Shelduck (*Tadorna tadorna*), 381
 Shetland, 59, 162, 163, 164, 189–91, 190; accretion measurement in, 67; quadrat for, 53; *see also* Sullom Voe
 sieve(s), 131–5; types, 134–5, 134, *see also* mesh size
 sigma-T (sea water density), 326

448 INDEX

- silicate, in sea water, 282, 317
 Simpson's Index of dominance, 19
Sipuncula, identification of, 410
 Skallingen marsh (Denmark), 62
Skeletonema sp., 291, 300, 301, 302
Skeletonema costatum, 313
 Skomer Island (Wales), 206
 sledge: mud 101, 102, 429; underwater, 211
 Snipe (*Gallinago gallinago*), 377
 sonar, for fish surveys, 347
 Southampton Water, 111, 112, 126; distribution of *Spartina anglica*, 55
Spartina sp., 14t, 65, 258
Spartina alterniflora, 58, 61, 258
Spartina anglica, 41, 55, 61, 62
Spartina cynosuroides, 58
Spartina maritima, 70t
 specific growth rate, 327
 spectrophotometry, for chlorophyll measurement, 93–4, 313, 330, 334
Spergularia marina, 45, 70t
 Spermatophyta, identification of, 408
 spore collection, for marine fungi, 265–6
 sporulation of marine fungi, 262, 264, 265, 267 (sand grain method, 267; Shearer method, 267)
 SPOT satellite, 32–4, 33t, 34t
 stains: acridine orange (for bacteria), 246, 248; crystal violet, 246; eosin, 135; rose bengal, 92, 135, 149, 315; safranin, 246
 standard error, 17–18
 standardisation of methods/procedures, 10–12, 108–9, 343
 standing crop, 300
 starvation, in birds, 394, 395
 statistical analysis, 17–23, 73–4; for sample size, 207; *see also* cluster analysis
 stomach content(s): analysis, 361; of fish predators, 345t, 359
 stratification, of water column, 282–6; salinity, 284
 stress: in birds, 393; osmotic, in fish, 344
Strongylocentrotus droebachiensis, 164
 substrata, artificial, for rocky shore species, 185–6
 suction sampler, 98, 121–2, 198, 203t, 228, 228, 229, 230–1, 348 (with reference cylinder, 231–2, 231)
 Sullom Voe, Shetland, salt marshes, 39–40, 39, 54
 surveillance, biological, 2, 145
 survey(s): aerial, for birds, 382; area, limits of, on rocky shores, 158, 174; beached-bird, 390–9; detailed extensive, 173–5; frequency, *see* sampling frequency; meiofauna, 144–6; microbial, 239; presence-absence, 215–16; primary, 171; qualitative, 83–7, 146, 147, 148, 202; quantitative, 87–99, 146–8, 181–3, 202, 220–1; reasons for carrying out, 1–2, 144–6; semi-quantitative, 202, 216–19 (results, 218–19); transect, for rocky shores, 175–83
 survey methods/techniques: for birds, 376, 384, 389, 391; for fish, 343, 346–59, 345t (egg and larval, 358–9); for rocky shores, 188t
 survey planning, 1–23, 146–8; bird, 374; criteria, 3–13; for collecting bacteria, 240–2; for collecting fungi, 258–60; for fish, 342–6; for plankton collection, 318–20; for rocky shores, 157–61; for salt marshes, 39–42; using divers, 199–209
 sweep net, 68
Synechococcus sp., 293
 TABORD program, 64
 tagging, fish, 356–8, 357
 Tanaidacea, identification of, 411
 tape recorder, underwater, 209–10, 210, 215, 219, 220
 taxonomic group, species identification by, 405–6
 Tees estuary, salinity, 166
 telephone, diver-to-surface, 210, 211, 215, 219
Tellina fabula, 122
 temperature: of sea water, 282–3 (measurement of, 288, 289, 316–17, 324); profile, 282; –salinity diagrams, 286–8, 287, 322; –salinity probes, 289
 tern(s), 388–90; Arctic (*Sterna paradisaea*), 388, 390; Caspian (*Hydroprogne caspia*), 388; Common (*Sterna hirundo*), 388, 390; Gull-billed (*Gelochelidon nilotica*), 388; Little (*Sterna albifrons*), 338, 390; population, 390; Roseate (*Sterna dougallii*), 388; Sandwich (*Sterna sandvicensis*), 388–90
 tetrazolium salt reduction for estimation of bacterial activity, 253–4
Thalassiosira sp., 291, 292
 thermocline, 241
 Thistle oilfield, North Sea, 113, 114
 Thraustochytriales, 267, 270; estimation of, 267–8, 269; identification of, 407
Thysanoessa sp., 296
 thymidine, radio-labelled, for measurement of bacterial production, 251
 tidal cycle, effects of: on fish, 344; on sampling salt marsh, 68, 69–70, 429
 tidal currents, 281–2
 tidal exchange method, for flushing time calculation, 327, 328

Cambridge University Press

978-0-521-31191-5 - Biological Surveys of Estuaries and Coasts

Edited by J. M. Baker and W. J. Wolff

Index

[More information](#)

INDEX

449

- tidal flats, 81–102; delimitation of, 81; safety procedures for, 428–30; sampling methods for, 89–91, 92, 94, 95–9; sieving samples on, 132
- tidal pressure, effect on sediments, 239
- tides, and field work, 428–30
- time-series analysis, 322
- tow nets, for plankton, 304, 305–7, 324; size, 305
- towing speeds, for plankton collection, 307
- towing systems, for divers, 211–12
- trampling, effect of: on salt marshes, 44–5; rocky shores, 160–1
- transect(s), 65–6, 109, 110, 111, 113, 114, 147, 157–8, 159, 160, 169–70, 177–83, 382; continuous belt, 179–81; interrupted belt, 177–9, 178; for mapping, 82–3, 220; marking (on rocky shores, 181–2; on tidal flats, 100)
- transponder(s), for sampling site relocation, 213–14
- transport, on tidal flats, 101–2
- traps, 73–5; fish, 345t, 350, 351, 353, 355, 358; pitfall, 68, 73
- trawl(s), 345t, 350, 354, 355; Agassiz, 121; beam, 354, 355–6; otter, 351, 354, 356
- Trichodesmium* sp., 293
- trichomycetes, 270
- Triglochin maritima*, 60, 707t
- Triglochin palustris*, 70t
- tugs, underwater, 212, 215, 348
- Tunicata, identification of, 413
- Turbellaria/turbellarians, 141, 142t, 143, 149, 151; identification of, 409
- turbidity, 233, 299–300, 302, 319; maximum, 300
- turbulence, 281–4
- Turnstone (*Arenaria interpres*), 377
- TV cameras, for fish observation, 349
- ultrasonic communication systems, 210
- Ulva* sp., identification of, 408
- underwater communications systems, 209–11, 210
- underwater tugs, 212
- UV-fluorescence technique, 246, 249
- variance, 17–18
- Vaucheria* sp., identification of, 408
- Verrucaria maura*, 165; identification, 186
- vulnerability index, shore line types to oil damage, 13, 15t
- Wadden Sea, 82, 88, 90, 285t, 286
- waders, 374, 376–7, 380; arctic, 375; counts, 378
- Wales, 164
- Wash, the, 66–7; Knot population in, 379
- water: column, 282–4; movement of, 281–2, 319
- water-bottles: for sampling bacteria, 243; for plankton sampling, 305, 306, 307
- water quality, 166–7; for bathing, 256–7, 256t; for shellfish cultivation, 257
- wave action: effects of on rocky shores, 161; measurement of, 162–3
- weather: effect of, on beached-bird survey, 393; forecast, before field trips, 429
- Wentworth scale, 173
- Westernport Bay, Australia, 116
- Wigeon (*Anas penelope*), 381
- wildfowl, 374, 376; counts, 378; population, 380
- winches, on sampling boats, 118, 316
- wind, effects of, on water movements, 281–2
- winkles, abundance scale for, 170t; *see also Littorina* sp.
- wood, decaying, for collection of marine fungi, 268–9
- yield models, for fish, 363–4
- zinc sulphate, for crab sorting, 133
- zonation: of marine fungi, 259; rocky shore, 158; 159; salt marsh, 38, 65
- zone: black lichen, 165; euphotic, 295
- zooplankton, 281, 286, 294–9, 302; biomass, 302, 314–16; identification of, 411; vertical migration, 283–4, 294–6
- Zostera* sp., 14t, 40, 41, 83, 83, 94; identification of, 408
- Zostera marina*, 62, 201