

*The production of new potato  
varieties:  
technological advances*


*Edited by*

**G. J. JELLIS**

Principal Scientific Officer  
Plant Breeding Institute, Cambridge

**D. E. RICHARDSON**

National Institute of Agricultural Botany



**CAMBRIDGE UNIVERSITY PRESS**

*Cambridge*

*New York New Rochelle*

*Melbourne Sydney*

Published by the Press Syndicate of the University of Cambridge  
The Pitt Building, Trumpington Street, Cambridge CB2 1RP  
32 East 57th Street, New York, NY 10022, USA  
10 Stamford Road, Oakleigh, Melbourne 3166, Australia

© Cambridge University Press 1987

First published in 1987

Printed in Great Britain at the University Press, Cambridge

*British Library cataloguing in publication data*

The production of new potato varieties:  
technological advances.

1. Potatoes – Europe – Varieties

I. Jellis, G. J. II. Richardson, D. E.

633'.4917 SB211.P8

*Library of Congress cataloguing in publication data available*

ISBN 0 521 324580

## CONTENTS

List of contributors	x
Preface	xvi
Editors' note and acknowledgements	xvii
Introduction	xviii
J.G.Th. HERMSEN and K.M. SWIEZYNSKI	
The development of potato varieties in Europe	1
G.J. JELLIS and D.E. RICHARDSON	
<b>Genetic Resources</b>	9
Genetic resources: their preservation and utilization	10
N.E. FOLDØ	
Gene pooling of modern potato varieties	28
D.R. GLENDINNING	
<b>Breeding Strategies</b>	31
Potato breeding strategy in the German Democratic Republic	32
M. SCHOLTZ	
Potato breeding strategy in the Federal Republic of Germany	38
M. MUNZERT	
Potato breeding strategy in the Netherlands	45
J.P. VAN LOON	
Potato breeding strategy in Poland	55
K.M. SWIEZYNSKI	
Potato breeding strategy in the United Kingdom	60
G.R. MACKAY	
Private potato breeding in the United Kingdom	68
J.M. DUNNETT	
<b>Selection and Screening Methods</b>	71
The efficiency of early generation selection	72
J. BROWN	
Problems associated with early generation selection of potato clones in West Siberia	75
B. DOROZHKIN and B. CADYCHEGOV	

Influence of weight of seed tubers on selection of first year clones: preliminary results K.M. LOUWES and A.E.F. NEALE	78
A joint cyst nematode/late blight test for early generation screening of potato clones C.N.D. LACEY, G.J. JELLIS, N.C. STARLING and S.B. CURRELL	81
Screening for resistance to diseases in a potato breeding programme G.J. JELLIS, R.E. BOULTON, N.C. STARLING and A.M. SQUIRE	84
Breeding for resistance to potato viruses with special reference to cDNA probes R.E. BOULTON, G.J. JELLIS and A.M. SQUIRE	86
Screening for resistance to diseases and pests G.R. MACKAY	88
Breeding for resistance to and tolerance of potato cyst nematode M.F.B. DALE	91
Breeding multi-resistant potato germplasm M.W. MARTIN	94
Resistance to storage diseases in breeding stocks A. PAWLAK, J.J. PAVEK and D.L. CORSINI	96
<u>Phytophthora</u> research at the Foundation for Agricultural Plant Breeding (SVP), The Netherlands A.G.B. BEEKMAN	99
Effect of time interval between inoculation and assessment on relative content of potato virus Y <sup>N</sup> in leaves of potato plants XU PEI WEN	102
Selection and evaluation of potatoes for improved tolerance of environmental stresses D. LEVY	105
<b>Variety Assessment</b>	109
Potato variety assessment in the Federal Republic of Germany W. BÄTZ	110
Potato variety assessment in France P. PERENNEC	114
Variety assessment in The Netherlands K. VAN DER WOUDE	119
Potato variety assessment in Poland J. BORYS	131
Potato variety assessment in the UK D.E. RICHARDSON	135
Use of common origin seed for potato trials S.P. KERR	140

<i>Contents</i>	vii
Micropropagation - an aid in the production of new varieties P. WOOSTER and T.J. DIXON	142
Testing potato varieties for response to drought and irrigation S.J. FLACK	146
Testing varieties for resistance to and tolerance of <u>Globodera pallida</u> G.M. GURR and S.P. KERR	148
Testing for glycoalkaloids J. WHITE	151
Methods for calculating 1-9 values to express the resistance of potato varieties to diseases P.T. GANS and P. WOOSTER	153
Establishing standards in variety assessment M. TALBOT	157
Consumer quality requirements in the United Kingdom R.M. STOREY and D.P. HAMPSON	163
The effects of fertilizer treatments on a range of old and new early-maturing potato varieties A.J. THOMSON	165
Variety trials in Egypt, with special reference to dormancy M.K. IMAM	168
<b>Semi-conventional Breeding Methods</b>	171
Efficient utilization of wild and primitive species in potato breeding J.G.Th. HERMSEN	172
Advances and limitations in the utilization of Neotuberosum in potato breeding R.L. PLAISTED	186
Breeding at the $2x$ level and sexual polyploidization S.A. HERMUNDSTAD and S.J. PELOQUIN	197
Haploids extracted from four European potato varieties L. FRUSCIANTE and S.J. PELOQUIN	211
Heterosis for tuber yields and total solids content in $4x \times 2x$ FDR-CO crosses M.F. MASSON and S.J. PELOQUIN	213
Variability of $F_1$ progeny derived from inter-ploidy ( $4x \times 2x$ ) <sup>1</sup> crossing J. JAKUBIEC	218
Unreduced gametes in the breeding of potatoes at the diploid level E. ZIMNOCH-GUZOWSKA and I. WASILEWICZ	222

Desynapsis and FDR 2n-egg formation in potato: its significance to the experimental induction of diplosporic apomixis in potato E. JONGEDIJK	225
Utilizing wild potato species via <u>Solanum phureja</u> crosses R.N. ESTRADA	229
The use of diploid <u>Solanum phureja</u> germplasm C.P. CARROLL	231
Advances in population breeding and its potential impact on the efficiency of breeding potatoes for developing countries H.A. MENDOZA	235
<b>True Potato Seed</b>	247
Breeding strategies for true potato seed M.T. JACKSON	248
Evaluation of yield and other agronomic characteristics of true potato seed families and advanced clones from different breeding schemes L. CONCILIO and S.J. PELOQUIN	262
Evaluation of true potato seed families obtained from different breeding schemes in the south of Italy L. FRUSCIANTE, S.J. PELOQUIN and A. LEONE	264
Potato production from true potato seed in Italy L. MARTINETTI	266
Field seeding of true potato seed in a breeding programme M.W. MARTIN	269
<b>Unconventional Breeding Methods</b>	271
Recent progress in molecular biology and its possible impact on potato breeding: an overview R.B. FLAVELL	272
Combined application of classical and unconventional techniques in breeding for disease resistant potatoes G. WENZEL, S.C. DEBNATH, R. SCHUCHMANN and B. FOROUGH-WEHR	277
Auther culture of dihaploid <u>Solanum tuberosum</u> H <sub>3</sub> 703 N.P. BATTY and J.M. DUNWELL	289
Genetic manipulation in potato using <u>Agrobacterium</u> G. OOMS	293

Prospects of using tumour-inducing plasmid-mediated gene transfer for the improvement of potato varieties A. BLAU, P. ECKES, J. LOGEMANN, S. ROSAHL, R. SANCHEZ-SERRANO, R. SCHMIDT, J. SCHELL and L. WILLMITZER	309
Use of protoplast fusion and somaclonal variation in potato breeding M.G.K. JONES	315
The potential value of somaclonal variants in potato improvement A.J. THOMSON	327
Use of <u>in vitro</u> culture of <u>Solanum tuberosum</u> in potato breeding B. COLIN, F. LAMMIN and Y. DATTEE	331
<b>Commentary</b> A practical breeder's view of the current state of potato breeding and evaluation A.J. THOMSON	335 336
Index	347

## THE DEVELOPMENT OF POTATO VARIETIES IN EUROPE

G.J. Jellis and D.E. Richardson

### HISTORY

The potato was introduced into Europe from South America some-time between 1565 and 1573. It was first grown in Spain but by the early part of the seventeenth century it was found in the botanical gardens of many European states. This was largely due to the efforts of the botanist Charles d'Ecluse, or Clusius, who received two tubers and a fruit from the Prefect of Mons, Philippe de Sivry in 1588, multiplied the tubers and distributed them to a number of friends. The first botanical description of the potato was published by the Swiss botanist, Caspar Bauhin in 1596. Bauhin also gave the potato its Latin binomial, Solanum tuberosum, although he later added esculentum. A second description was published in England, where the potato was probably introduced between 1588 and 1593, by Gerard in his 1597 "Herball".

The original introductions were almost certainly from the Andean regions of Peru or Columbia and were of the subspecies andigena (hereafter called Andigena potatoes), with a short-day photoperiodic response. Under European conditions they would have produced many stolons but poor yields of late-developing tubers. During the following two centuries, seedlings from these original introductions were selected for yield and earliness, to give rise to clones well adapted to longer summer days. This selection may have been made in hybrids and selfs of only two original introductions (Salaman 1926). By the middle of the eighteenth century the potato had become a universally cultivated field crop in Europe. A major problem was "Curl", a degenerating condition which meant that varieties had a limited life. To overcome this, seedlings were raised from self-set potato fruit and so there were large numbers of varieties. Towards the end of the century a number of distinct varieties came into use, some of which were specified for human consumption and others for animals.

Two events of major importance in the history of potato breeding



occurred in the middle of the nineteenth century. The late blight (Phytophthora infestans) epidemics of 1845-46 demonstrated the lack of blight resistance in contemporary varieties. This was not surprising as previous selection had been done in the absence of blight. New varieties were developed with some degree of resistance. These included Paterson's Victoria, Champion and Magnum Bonum, all of which feature in the ancestry of many modern varieties (Figure 1).

Figure 1. Three UK potato varieties grown extensively in the nineteenth century; Champion (top left, introduced 1876), Magnum Bonum (top right, introduced 1876) and Lumper (bottom, introduced prior to 1810 and widely grown in Ireland until 1846, when its failure due to late blight caused the famine).



The second event was the introduction of Rough Purple Chili by the Rev. Chauncey Goodrich of New York in 1851. He obtained it as one of a small collection of tubers via the US Consul in Panama. Goodrich assumed that it was a Chilean variety and since it was well adapted to the long days of New York, this may have been the case (Hawkes 1967). From Rough Purple Chili Goodrich raised a number of selfed seedlings including Garnet Chili, which was widely grown in the USA for many years. Among the varieties derived from Goodrich's stocks were Beauty of Hebron, Russet Burbank and Early Rose. The last named variety, in particular, has been very important in European breeding programmes.

Another source of breeding material in the last century was an Andigena clone known as Daber which was introduced into Germany in about 1830 and gave rise to many new varieties, including President.

The next major stimulus to potato breeding was the spread of wart disease (Synchytrium endobioticum) in the early part of this century. The recognition of field immunity in a few varieties, including Snowflake, led to the breeding of a whole range of wart-immune varieties. Good blight resistance was still being sought, and we can imagine Salaman's delight when he found field immunity, first in S. edinense and later in S. demissum (Salaman 1985). By 1914 he had produced a series of families segregating for resistance, and a programme which was to occupy the minds of many potato breeders, particularly in the UK, Germany and the USSR, for the next half a century had been initiated. The first signs that this major gene resistance was not durable came in 1932 but it was another 30 years before breeding for this type of resistance was abandoned. Nowadays, breeding programmes worldwide are aimed at introducing race non-specific resistance into new varieties. This is not proving to be an easy task; the resistance is controlled by a number of genes and is often associated with undesirable traits such as late maturity.

Degeneration of potato stocks, or "Curl", has already been mentioned. During the early part of this century it was discovered that this phenomenon was caused by a number of viruses. Sources of resistance were identified and breeding programmes initiated. Virus resistance is now a major objective in many breeding programmes, using many different Solanum species as sources of resistance.

Another objective common to many present-day breeding programmes is resistance to potato cyst nematodes (Globodera spp.). Breeding work was made possible by Ellenby (1952) who found resistance to G. rostochiensis in

five accessions of *Andigena* and also in *S. vernei*. Populations of cyst nematodes which multiplied on clones incorporating this resistance were found in many European countries. This led to the recognition of a range of pathotypes and eventually to two separate species being described, *G. rostochiensis* and *G. pallida*, each with pathotypes. Resistance to pathotypes of *G. pallida* has been found in accessions of *Andigena* and *S. vernei* and also in other *Solanum* species. An international scheme for identifying and classifying potato cyst nematodes was proposed by Kort *et al.* (1977) and this is presently being modified.

Potatoes have many uses and programmes have been designed specifically to breed for the requirements of the processing and starch industries. Some European countries have a large export market for seed potatoes and have breeding programmes geared to producing varieties suitable for foreign markets.

As will quickly be appreciated in the following chapters, potato breeding technology is rapidly developing, providing new opportunities for crop improvement. Early generation screening techniques are leading to more rapid identification of the superior genotypes. Breeding at the diploid level and exploiting unreduced gametes are providing ways of introducing new sources of variation and transferring them efficiently (Hermundstad & Peloquin, this volume). Just around the corner is the promise of directed variation by genetic manipulation and protoplast fusion. Already we are reaping some of the benefits of molecular biology in the form of efficient virus and viroid detection techniques (Boulton *et al.*, this volume; Flavell, this volume). Tissue culture has resulted in the rapid propagation of breeding material, and the production of variation through regenerating calluses has led to the exploitation of somaclonal variation (Jones, this volume; Thomson, this volume).

Finally, there is the whole new concept of breeding true potato seed (TPS). The best way of producing such seed is still being investigated (Jackson, this volume) but already TPS is having a major impact in some developing countries.

The "humble potato" has come a long way in the past 400 years, but its future looks just as exciting as its past.

For more detailed and fascinating accounts of the origins of the potato in Europe, the reader is referred to Salaman's book "The History and Social Influence of the Potato", recently reprinted (Salaman 1985), and Hawkes' 1966 Royal Horticultural Society Masters Memorial Lecture "The History

of the Potato" (Hawkes 1967).

#### PLANT BREEDERS' RIGHTS

In Britain during the early part of this century, there was a multiplicity of "new" varieties - bogus selections of existing varieties masquerading under new names. Stocks with the new names usually fetched a higher price than those with the original name. This led the National Institute of Agricultural Botany to set up its Potato Synonym Committee in 1919, when over 200 synonyms of one variety, Up to Date, were found. This Committee continued in operation until the introduction of the distinctness regulations for the Plant Breeders' Rights (PBR) in the 1960s.

The concept of PBR is an extension of the ownership of an invention protected by patent legislation. The aim is to provide a fair return on the long term investment required to breed and assess a new variety; this normally takes about 12 years for potatoes. Protection on all propagating material is given for a minimum period of 15 years from the date of registration; this has been extended for potatoes to 30 years in some countries. Royalties are collected on the basis of the area of potatoes grown for seed certification.

The owner of a variety may transfer the proprietary rights on his variety to individuals or companies for the purpose of growing and selling seed material. He has the legal right to exclude others from reproducing, selling, exporting or importing seed of this variety. There are, however, no restrictions on anyone growing a variety exclusively for their own use, or for selling it for consumption, either fresh or processed.

Plant variety production is internationally controlled by the Union pour la Protection des Obtentions Vegetales (UPOV), which was established in 1961. Membership extends to most European countries, Australia, New Zealand, Israel, Japan, South Africa and the United States. Its aim is to promote progress by stimulating plant breeding and facilitating the interchange of varieties between countries. The benefit of UPOV membership is the availability of reciprocal rights, which avoids duplication of effort by both breeders and government authorities. UPOV has produced test guidelines including 107 taxonomic characters which may be used to establish distinctness in potato varieties (Anon. 1974).

Until now PBR has provided protection only for the variety, which is the end product of plant breeding. In future, with the introduction of genetic engineering there is justification for some form of patent protection

of certain technological or microbiological processes used to produce a new variety. The legal implications of this concept have been discussed by Lange (1985).

The general consensus of opinion is that PBR have been advantageous in encouraging investment in private breeding and in increasing the numbers of improved varieties. It has been suggested, however, that PBR are responsible for the recent interest of large multinational chemical and oil companies in taking over seed and breeding establishments, and that such companies may restrict access to varieties with improved disease and pest resistance, but this remains unproven.

It has also been suggested that PBR may cause a reduction of government support for state breeding, which may reduce cooperation and exchange of information between the state and private breeders. In most European countries this is not a problem since the state breeders tend to concentrate on doing basic research and providing the private breeders with basic breeding material.

#### COOPERATION

There is a good spirit of international cooperation in both potato breeding and variety assessment. This is fostered by such organisations as the European Association for Potato Research (EAPR), European Association for Research in Plant Breeding (EUCARPIA), Potato Association of America (PAA), International Potato Center (CIP) and the International Board for Plant Genetic Resources (IBPGR).

In order to help with the collection, conservation, documentation, evaluation, exchange and use of germplasm, guidelines on descriptors have been produced (IBPGR 1977). These include over 100 "passport", taxonomic, agronomic, disease and quality characteristics. More recently a shorter minimum list of descriptors has been produced by the Commission of the European Communities Agricultural Research Programme on Plant Productivity in collaboration with IBPGR (CEC 1985).

#### MERIT TESTING

Within the European Economic Community (EEC) a new variety can only be marketed if it is included on the Common Catalogue. This is a list of all varieties included on the National List of all member states. A derogation order can be obtained by any member state to prevent a variety being grown in their country if it presents a particular disease or pest hazard.

In order to comply with the EEC Directive for the inclusion of a variety on a National List it must be distinct, uniform and stable (DUS) as for PBR, and have value for cultivation and use (VCU). The testing systems in several countries are described in this volume.

The need for VCU testing to be a statutory requirement has sometimes been challenged, but few agree that the success or failure of a variety should be dictated by market forces alone. Some form of independent merit testing is generally considered necessary for advisory purposes and for protection against exaggerated or misleading claims. However, potato experts agree that it is difficult to predict from trials the future of some potato varieties and that commercial experience may also be required.

Fifty years ago the Scottish potato breeder of the famous "Arran" varieties, Donald MacKelvie (1937), wrote "No one can foretell what will be the verdict of the practical grower on a new variety of potato, or its future in the market, even if it is successful in trials". To a lesser extent this is still true today, and few varieties get beyond the "promising" stage.

#### REFERENCES

- Anon. (1984). Test Guidelines for Potato (TG/23/2). Geneva: International Union for the Protection of New Varieties of Plants.
- Ellenby, C. (1952). Resistance to the potato root eelworm, Heterodera rostochiensis Wollenweber. Nature, 170, 1016.
- Hawkes, J.G. (1967). The history of the potato. Masters Memorial Lecture, 1966. J.R. Hort. Soc., 92, 207-24 and 249-302.
- Huaman, Z., Williams, J.T., Salhuana, W. and Vincent, L. (Eds.) (1977). Descriptors for the Cultivated Potato. Rome: International Board for Plant Genetic Resources.
- Kort, J., Ross, H., Rumpfenhorst, H.J. and Stone, A.R. (1977). An international scheme for identifying and classifying pathotypes of potato cyst-nematodes Globodera rostochiensis and G. pallida. Nematologica, 23, 333-9.
- Lange, P. (1985). The nature of Plant Breeders' Rights (plant variety protection law) and their demarcation from patentable inventions. Pl. Var. Prot., 44, 17-27.
- MacKay, G.R., Hijink, M.J. & Mix, G. (Eds.) (1985). Minimum List of Characters of European Communities. Rome: International Board for Plant Genetic Resources.
- Salaman, R.N. (1926). Potato Varieties. Cambridge: Cambridge University Press.
- Salaman, R.N. (1985). The History and Social Influence of the Potato. Revised Impression ed. by J.G. Hawkes. Cambridge: Cambridge University Press.