

Cambridge University Press

0521 392802 - Vitamin Receptors: Vitamins as Ligands in Cell Communication

Edited by Krishnamurti Dakshinamurti

Frontmatter

[More information](#)

Vitamins are essential micronutrients available to animal organisms through the diet. This book takes a fresh approach to vitamin-binding proteins, with emphasis on the nature of the binding of the vitamin ligand to a protein and its sequela. The role of vitamin-binding proteins as initiators of the metabolic response is evaluated. Experts in the field from around the world present a state-of-the-art account of their work on the interaction of vitamins with specific intracellular systems through the appropriate binding proteins and how this interaction results in the biological action of vitamins. This is the first comprehensive book dealing with the subject. The book will be of interest to research workers and postgraduate students in the fields of biochemistry and nutrition.

Cambridge University Press

0521392802 - Vitamin Receptors: Vitamins as Ligands in Cell Communication

Edited by Krishnamurti Dakshinamurti

Frontmatter

[More information](#)

Intercellular and intracellular communication

General editor: Professor B. Cinader

Vitamin receptors

Vitamins as ligands in cell communication

Cambridge University Press

0521 392802 - Vitamin Receptors: Vitamins as Ligands in Cell Communication

Edited by Krishnamurti Dakshinamurti

Frontmatter

[More information](#)

Intercellular and intracellular communication

- 1 Hormones, receptors and cellular interactions in plants
Edited by C.M. Chadwick and D.R. Garrod
- 2 Receptors in tumour biology
Edited by C.M. Chadwick
- 3 Receptors and ligands in psychiatry
Edited by A.K. Sen and T. Lee
- 4 Receptors and ligands in neurological disorders
Edited by A.K. Sen and T. Lee
- 5 Intracellular trafficking of proteins
Edited by C.J. Steer and J.A. Hanover

Cambridge University Press

0521 392802 - Vitamin Receptors: Vitamins as Ligands in Cell Communication

Edited by Krishnamurti Dakshinamurti

Frontmatter

[More information](#)

Vitamin receptors: *Vitamins as ligands in cell communication*

EDITED BY

KRISHNAMURTI DAKSHINAMURTI

*Professor of Biochemistry and Molecular Biology, Faculty of Medicine,
University of Manitoba, Winnipeg, Canada*



CAMBRIDGE
UNIVERSITY PRESS

Cambridge University Press
0521 392802 - Vitamin Receptors: Vitamins as Ligands in Cell Communication
Edited by Krishnamurti Dakshinamurti
Frontmatter
[More information](#)

Published by the Press Syndicate of the University of Cambridge
The Pitt Building, Trumpington Street, Cambridge CB2 1RP
40 West 20th Street, New York, NY 10011-4211, USA
10 Stamford Road, Oakleigh, Melbourne 3166, Australia

© Cambridge University Press 1994

First published 1994

A catalogue record of this book is available from the British Library

Library of Congress cataloguing in publication data

Vitamin receptors: vitamins as ligands in cell communication/edited
by Krishnamurti Dakshinamurti.

p. cm.

Includes index.

ISBN 0 521 39280 2 (hardback)

1. Vitamins—Receptors. 2. Carrier proteins. I. Dakshinamurti,
Krishnamurti, 1928–
QP771.V5734 1994

574.19'26—dc20 93-4680 CIP

ISBN 0 521 39280 2 hardback

Transferred to digital printing 2004

CONTENTS

<i>List of contributors</i>	ix
<i>Preface</i>	xi
<i>Introduction</i>	xv
1 Serum and cellular retinoid-binding proteins <i>Dianne R. Soprano</i>	1
2 Retinoic acid receptors <i>Vincent Giguère</i>	28
3 Vitamin D receptors and the mechanism of action of 1,25-dihydroxyvitamin D ₃ <i>J. Wesley Pike</i>	59
4 Cobalamin binding proteins and their receptors <i>Bellur Seetharam and David H. Alpers</i>	78
5 Folate binding proteins <i>Steven Weitman, Richard G.W. Anderson and Barton A. Kamen</i>	106
6 Riboflavin carrier protein in reproduction <i>P. Radhakantha Adiga</i>	137
7 Binding proteins for α -tocopherol, L-ascorbic acid, thiamine and vitamin B ₆ <i>Krishnamurti Dakshinamurti and Jasbir Chauhan</i>	177
8 Biotin-binding proteins <i>Krishnamurti Dakshinamurti and Jasbir Chauhan</i>	200
<i>List of abbreviations</i>	250
<i>Index</i>	253

CONTRIBUTORS

- P. Radhakantha Adiga *Center for Reproductive Biology and Molecular Endocrinology and Department of Biochemistry, Indian Institute of Science, Bangalore 560 012, India*
- David H. Alpers *Division of Gastroenterology, Washington University School of Medicine, St. Louis, MO 63110, U.S.A.*
- Richard G.W. Anderson *Departments of Pediatrics, Pharmacology and Cell Biology, University of Texas Southwestern Medical Center, Southwestern Medical School, Dallas, TX 75235, U.S.A.*
- Jasbir Chauhan *Department of Biochemistry and Molecular Biology, Faculty of Medicine, University of Manitoba, Winnipeg, Canada R3E0W3.*
- Krishnamurti Dakshinamurti *Department of Biochemistry and Molecular Biology, Faculty of Medicine, University of Manitoba, Winnipeg, Canada R3E0W3.*
- Vincent Giguère *Division of Endocrinology, Hospital for Sick Children, and Department of Molecular and Medical Genetics, Faculty of Medicine, University of Toronto, Toronto, Canada M5G 1X8.*
- Barton A. Kamen *Departments of Pediatrics, Pharmacology and Cell Biology, University of Texas Southwestern Medical Center, Southwestern Medical School, Dallas, TX 75235, U.S.A.*
- J. Wesley Pike Sr. *Division of Biochemistry, Ligand Pharmaceutical, 9393 Towne City Drive, San Diego, CA 92121-3016, U.S.A.*
- Bellur Seetharam *Division of Gastroenterology, Medical College of Wisconsin, Milwaukee, WI, U.S.A.*
- Dianne Robert Soprano *Department of Biochemistry and Fels Institute for Cancer Research and Molecular Biology, Temple University School of Medicine, Philadelphia, PA 19140, U.S.A.*
- Steven Weitman *Departments of Pediatrics, Pharmacology and Cell Biology, University of Texas Southwestern Medical Center, Southwestern Medical School, Dallas, TX 75235, U.S.A.*

PREFACE

Contemporary research has allowed us to discern a linguistic structure in biological interactions in which molecules in solution and in or on membranes react with one another, sequentially, and thus form sentences of a language. Interlocking ligand–receptor systems constitute an intracellular and intercellular communication system in which molecules, such as cytokines, hormones, metabolic products, vitamins and foreign macromolecules can serve as words and convey signals – messages – through combination with membrane structures, i.e. receptors. These signals can initiate synthesis of other factors which can combine with receptors and form the sentences of the molecular language. Thus, molecular language consists of sequential interactions between receptors and ligands, which are either cell-bound or secreted by one cell and taken up by membrane structures of another cell. A succession of these interactions at the membranes of cells and organelles coordinate cell metabolism within the same and between different cells and organs.

The analogy between language and molecular communication can also be detected in the evolutionary diversification of words, i.e. in the existence of superfamilies of molecules, having a common two-chain structure and involved in recognition of different ligands. Members of these families show homologies in a variety of cell types and among cells of different animals, from invertebrates to vertebrates.

Within cells, some types of organelles acquire macromolecules from the cytoplasm. The site-specificity of these acquisitions depends on receptors, which recognize appropriate proteins by signals that may consist of amino acid sequences or of post-translational modifications.

In communication, occurring between different types of cells, receptors can be activated through soluble factors, and hence at a distance. Receptor–ligand interaction can also occur between membranes of

xii *Preface*

different cell types, i.e. via adhesion molecules that play a role in structural development of organs, exemplified by neural cell adhesion and embryological development under the influence of 'master' cells.

Once a ligand has combined with the membrane receptor of a cell, a change in membrane configuration can ensue and activate a series of metabolic events. Alternatively, ligands and receptors can enter the cell through pathways which start in 'coated pits', and can then be delivered to lysosomes for degradation or be recycled to the surface. This process of endocytosis mediates and regulates the entrance and half-life of various types of molecules, such as growth factors, viruses, toxins and nutrients.

Cell communication is regulated by activation events and by processes which limit the period during which a given stimulus can affect biochemical reactions that are initiated via a particular receptor. This limitation is achieved by processes such as endocytosis, recycling, and affinity changes in receptors, and through disassociation of macromolecular complexes with which the ligand binding site is associated.

Function of each cell type involves internal exchange between cytoplasm and various membrane-bound compartments, i.e. organelles. The function of each organ involves communication between different cell types; that of the organism as a whole involves communication between different organs.

The vast majority of molecules involved in biological sentences, whether receptors or soluble ligands, are autologous. However, the impact of external waves of energy can trigger the sentences of the neuronal system and external molecules can trigger sentences in the immune and nutritional systems.

Past volumes of this series have dealt with various aspects of molecular language, reviewing hormone receptors, cellular interactions in plants, tumor biology, neurology and psychiatry. The fifth volume of our series examined the steps by which molecules can enter the cell interior and be transported with and without receptor to the locations within the cell in which they interact or in which they are being disposed of. It thus dealt with traffic within the cell and with the interrelation between pathways of endocytosis and exocytosis. The present (sixth) volume of this series considers vitamins in terms of the processes by which they enter the system and the receptors by which they affect and activate the system. It deals with nutrients, i.e. with molecules, synthesized in the external world, and hence with an instance of communication initiated by encounters with externally generated molecules. This particular feature is common to the nutritional and the immune systems.

Vitamin receptors deals with chemically diverse molecules, water- and

Preface

xiii

lipid-soluble vitamins, taken up in the gastrointestinal tract through specific saturable membrane components. It surveys binding mechanisms, intracellular transport and, in some instances, transport to the nucleus and the interaction with sections of the genome, resulting in transcription of specific proteins. So far, there is little indication of a common ancestral receptor molecule from which vitamin receptors have descended; they may be derived from different lines of descent. There are indications that some vitamin receptors may have common molecular antecedents with receptors of the endocrine system. The identification of receptor structures remains an important challenge for the future development of analysis of the language of communication in which vitamins are involved.

Bernhard Cinader

INTRODUCTION

This volume attempts to assemble the current information on the ligand (vitamin) – receptor (binding protein) interaction as it applies to each of the vitamins, from the site of enteric absorption to the site of vitamin function. For the absorption and transport of most vitamins, specific proteins, which have a recognition factor incorporated into them, are involved. At the plasma membrane of the cell, a second recognition factor has to function before internalization of the vitamin. Intracellular vitamin-binding proteins and their interaction with cellular systems resulting in the biological action of the vitamins provides the last step in the successive stages of communication. Thus, we have the concept of a ligand, an extraorganismic molecule, reacting with a specific protein entity, the receptor, facilitating absorption and transport of the ligand to its site of action with the resultant sequence of physiological events to which we refer as ‘vitamin function’. The term ‘receptor’, in the context of vitamins, is used to refer to proteins with a recognition function and not to receptors in the traditional pharmacological sense. It includes proteins which bind to the vitamin ligand and function as extracellular or intracellular transporters as well as the smaller number of vitamin-binding nuclear proteins. In the latter case, vitamins might act as small molecular modulators of protein–DNA interaction, a situation analogous to that of some hormones. The designation ‘receptors’ for these categories of vitamin-binding proteins appears to be appropriate as it connotes a function, more than ligand binding.

Vitamins are required by all protozoans and metazoans that have been studied. They do not conform to any single chemical classification. Some vitamins, such as retinol (vitamin A), cholecalciferol (vitamin D), α -tocopherol (vitamin E) and phyloquinone (vitamin K), are lipid-soluble; others, such as thiamine (vitamin B₁), riboflavin (vitamin B₂), niacin,

xvi *Introduction*

pyridoxine (vitamin B₆), panthothenic acid, lipoic acid, biotin, folic acid, cyanocobalamin (vitamin B₁₂) and ascorbic acid (vitamin C), are water-soluble. While the unlimited availability of vitamins may have permitted the evolutionary loss of the ability of animals to synthesize vitamins, the ability to absorb vitamins from the environment or from digested food is essential. The modes of uptake, intra- and intercellular transport of these two broad categories of vitamins are different, based on their solubility characteristics. Lipoprotein binders are implicated in the transport of the lipid-soluble vitamins. Among the water soluble-vitamins the bindings of cyanocobalamin and folate, respectively, to specific proteins were recognized early. The intestinal absorption of other water-soluble vitamins was assumed to be through simple diffusion. Increasing evidence has accumulated to indicate that the absorption of many water-soluble vitamins in the gastrointestinal tract is through saturable mechanisms. In addition, the concentrations of many water-soluble vitamins in the brain are much higher than in plasma. This is true of placental transport as well, implicating specific transporters for individual vitamins across the blood-brain and placental barriers, respectively.

Absorption of vitamins by most cells occurs through membrane-associated transport systems which are present in low concentrations. These receptor-like proteins may consist of an extracellular vitamin-binding domain linked to a transmembrane anchor. The fertilized egg and embryo obtain vitamins from the external environment using cell-surface vitamin receptor/transporter proteins. The vitamins are used directly without storage. In contrast, the eggs of birds and reptiles contain both the vitamins and their respective vitamin-binding proteins deposited in the yolk for transport to the oocyte.

The physiological effects of vitamins follow their entry into the cell, metabolic transformation and association with specific apoenzymes, resulting in accentuation of a metabolic pathway. In this respect vitamins are metabolic initiators. Where the vitamin function is other than as a coenzyme or prosthetic group of an enzyme its effect is through regulation of cellular protein synthesis. The mechanisms of action of lipid-soluble vitamins are similar to those of the steroid hormones. Water-soluble vitamins such as biotin and folic acid also seem to have a similar function.

The transcriptional patterns of selected genes are often altered by external signals that the cell receives. Thus changes in nutritional supplies, interaction with hormones, viral infection or exposure to environmental factors such as metals, light, high temperature or radiation can either induce or repress the transcription of specific genes or gene sets. In

Introduction

xvii

many cases the regulation of transcription (up or down) is mediated by alterations in the availability of essential transcription factors or by change in the factors' ability to bind to specific regulatory sequence motifs.

The nuclear vitamin D receptor has been characterized as belonging to the steroid family of receptors. Retinol and retinoic acid and their respective binding proteins may serve unique vital functions within different tissues and organs at different times during development. The large number of retinoic acid receptor isoforms expressed in a subtype-specific fashion during development and in differentiated tissues is consistent with the diversity of retinoid action. Folate binding proteins might be involved in transmission of messages for cell division as the accumulation of folate is essential for purine and thymidine synthesis. Biotin has been shown to mimic the actions of insulin in inducing glucokinase mRNA and repressing phosphoenolpyruvate carboxykinase mRNA. The effect of biotin is seen in the starved as well as the diabetic animal in a way that does not involve the synthesis or secretion of insulin. The presence of nuclear binding proteins for biotin and pyridoxal phosphate is indicated, although their functions are not yet known.

We are still at a preliminary stage in our exploration of the groups of vitamin binding proteins. This volume aims to focus attention on the protein-bound vitamin function. The role of some vitamins as the prosthetic group of an enzyme or as a coenzyme participating in enzymatic catalysis is well recognized. This traditional aspect of vitamin function is not considered here. The emphasis is rather on the nature of the binding of the vitamin ligand to a protein, and its sequela.

Krishnamurti Dakshinamurti