

## Index

ASP prompts, menu selections, and output labels appear in **Boldface Type**. File names and ASP menus are CAPITALIZED.

- Accuracy of ASP, 102*n*
- Across-groups, 171
- Add Or Delete Variable Or Case**, 15
- Adding variables to data matrix, 57–60
  - transformed variables, 45
- Adjacent values, 66
- ALT COMMANDS MENU, 8
  - truncated, 44
- Analysis, regression, 79–83, 189, 196–199
- Analysis of variance
  - F* test, 171
  - one-way, 170–174
  - repeated measures, 179–181
  - two-way, 174–179
    - fixed-effects, 174–176
    - mixed-effects, 177–179
    - random-effects, 176–177, 178–179
- ANALYSIS OF VARIANCE MENU, 170
- ANOVA (*see* Analysis of variance)
- Antilog, 88
- ASCII, 2
  - text editor, 26
- .ASF, 13
- ASP, accuracy of, 102*n*
- Association, tests of, 211–219
  
- Basic quantities (computational, in **Summary Statistics** output, 39)
- Biased estimator (of variance), 123
- Binomial probability, 93–97
  - ASP notational conventions, 93
  - cumulative, 95–97
  - normal approximation to, 102, 104
- Bivariate regression, 197
- Bounce bar, 4
- Box-and-whisker plots, 66–68
- Break Down By Third Variable**, 72
- Breaking down variables, 72,
- Buffers, 2
  
- Calculated Value**
  - of *t*, 149
  - of *Z*, 98, 145
- Calculating class limits, 24, 28, 29
- Calculator (*see* Pop-up calculator)
- Case(s),
  - analysis of variance, 168
  - columns in transposed matrix, 108
  - Case(s) (*cont.*)
    - deleting, 15–16, 63
    - labels
      - changing, 15, 106
      - entering, 17
    - Pearson's *chi*-square tests, 211–212, 214
    - rows in ASP data matrix, 4
    - selecting
      - by case label, 63
      - directly by row number, 16, 152–153
      - for deletion, 15–16, 63
      - for samples, 152–153,
- CASE TRANSFORMATION MENU, 110–111, 123–124
- Case transformations
  - mean, 110
  - sample variance, 123–124
  - variance, 111
- Central Limit Theorem, 113
- Change Names/Labels/Cases/Vars.**, 15
- Change Sign**, 90
- Changing case labels, 15, 106
- Changing number of digits displayed, 162
- Changing sign (of entry in pop-up calculator), 90, 100
- CHI-SQUARE**
  - goodness-of-fit test, 201
  - hypothesis test about  $\sigma^2$ , 162
  - tests of association, 213
- Chi*-square
  - distribution, 135
  - tests (*see* Pearson's *chi*-square tests)
- CHOOSING RANDOM INTEGERS**, 106
- Class limits, calculating, 24, 28, 29
- Coding variables (for ANOVA), 168, 180
- Coefficient of
  - correlation, 73, 190
  - determination (**R SQ.**), 73, 81, 196
  - kurtosis, 40
  - skewness, 40
  - variation, 40
- Combinations, 89–90
- Confidence intervals, 126
  - mean
    - using normal distribution, 126–128
    - using Student's *t*, 132–135
  - proportion, 129–132
  - variance, 135–136

- Confidence intervals (*cont.*)  
 menu options, 130
- Consistency, 115  
 of mean, 115–116  
 of sample proportion, 116–122  
 of variance ( $S^2$ ), 123
- Constant, scaling, 47
- Contingency tables, 211–214
- Copying  
 disks, 6  
 output, 26
- Correlation  
 coefficient of, 73, 190  
 different values using **Correlation Matrix** and **PROD. M. COR(XY)**  
 options, 192–193  
 matrix, 76–78  
 ratio, 196  
 testing hypotheses about one, 189–193  
 testing hypotheses about two, 193–194  
 using **Correlation Matrix**, 76–77
- Correlation Matrix**, 76–78, 189
- Covariation, linear vs. nonlinear, 196
- Crosstab/Contingency**, 203, 211, 212, 215, 219
- CROSSTABULATE Variables Or Enter TABLE Directly**, 212, 215, 217
- Crosstabulation in tests of association, 214–219
- Cumulative probability  
 binomial, 95–97  
 normal, 99–100
- Cursor, 2
- D.F.** (*see* Degrees of freedom)
- Data  
 directory, 9–10  
 file, 4  
 changing, 14  
 creating, 10–14  
 matrix, 4  
 creating, 11–12  
 editing, 14  
 getting, 19–20  
 listing, 13  
 saving, 12  
 transposing, 42*n*, 108  
 missing, 42*n*, 57, 192–193
- Decile, 49
- Default, 2
- Degrees of freedom (**D.F.**)  
*chi-square*  
 goodness-of-fit tests, 206  
 tests of association, 213  
 tests of hypotheses about  $\sigma^2$ , 161
- F* test  
 analysis of variance, 171, 175, 177, 179  
 hypotheses about two variances, 164  
 Student's *t*, 134, 146, 149, 156
- Deleting  
 cases, 15–16, 63  
 OUTPUT file, 27  
 variables, 107
- Dependent variable  
 in ANOVA, 167  
 in regression, 72, 197  
 forecasting (predicting), 82–83
- Determination, coefficient of (**R SQ.**), 73, 81, 196
- Deviations from linear regression, 198
- Digits, changing number in display, 85
- Directory, 3  
 data, 9–10  
 root, 3
- Disk, 2  
 floppy, 2  
 formatted, 4  
 hard (fixed), 2
- Disk drive, 2
- DISKCOPY (DOS command), 6
- Dispersion  
 captured by Pearson's *chi-square* tests, 203  
 statistics, 39–40
- Display, changing number of digits in, 85
- Distribution  
 binomial, 94–97  
*chi-square*, 161  
 Fisher's *F*  
 hypotheses about two variances, 163  
 one-way analysis of variance, 171  
 two-way analysis of variance, 176, 178–179  
 normal, 98–100  
 properties captured by Pearson's *chi-square* tests, 203  
 sampling, 105, 110–114  
 of mean, 113–114  
 shape of (skew and kurtosis), 40  
 Student's *t*, 132, 134, 146–149, 158  
 uniform, 106
- DOS (*Disk Operating System*), 3  
 commands from ASP, 10
- Drawing random samples, 105–108
- Drive, disk, 2

- E** (ASP exponential notation), 21*n*, 38  
**Edit Or Create Data Matrix**, 11, 14  
 Editing a data file, 14  
 Effects (in analysis of variance), 171  
   fixed, 174, 176, 177  
   main, 176  
   mixed, 176, 177–179  
   random, 176–177, 178–179  
 Efficiency, 115  
 Entering case labels, 17  
 Equation, regression (linear or least squares), 80, 81, 196, 197–198  
 Error  
   as departure from linearity, 198  
   ASP output label (**PURE ERROR**), 198  
   sums and means of squares in ANOVA, 176  
   ASP output label (**ERROR**), 171  
 Error variance, pooled estimate of, 178–179  
 ESC (Escape key), 16  
 Estimate  
   interval (*see* Confidence intervals)  
   pooled, of error variance, 178–179  
   standard error of, 82  
 Estimation  
   point, 115–116  
 Estimator (*see also* Consistency)  
   point, 115  
   unbiased, of variance, 123  
   *See also* sample variance  
 Exiting  
   ASP, 16  
   pop-up calculator, 96  
 Expected frequencies and proportions, 203  
  
**F** distribution (*see* Distribution, Fisher's *F*)  
**F(D.F)**, 175  
*F* test (*F* ratio)  
   for hypotheses about two variances, 163  
   for one-way analysis of variance, 171  
   for two-way analysis of variance, 176–177, 177–178, 178–179  
**F Test For LINEAR Fit**, 198  
**F1=TRUNCATED MENU**, 44  
**F2=CALCULATOR**, 74  
**F3=TOGGLE PRINT**, 27  
 Factor, 167  
 Factorial, log, 87–89  
 Factorials, 85–86  
 Failure, in binomial experiment, 94  
 Fences, inner and outer, 66  
  
 File, 3  
   data (*see* Data file)  
   **DATA** entry for hypothesis tests, 139  
   label, 12, 14–15  
 Fisher's *F* distribution (*see* Distribution, Fisher's *F*)  
 Fisher's *r*-to-*z*, 191–192, 193–194  
 Fit, lack of, 198  
 Fixed effects, 174, 176, 177  
 Floppy disk, 2  
**FORECAST Dependent Variable**, 82  
 Formatted disk, 4  
 Freedom, degrees of (*see* Degrees of freedom)  
 Frequencies  
   crosstabulating, 212, 214–219  
   expected and observed, 203  
 Frequency distribution(s), 20–30  
   grouped, 24–25  
   ungrouped, 21–24  
**FREQUENCY PLOT**, 20  
 Function keys, 4  
  
**Get Data Matrix From ASP File**, 19–20  
**GOODNESS OF FIT TEST**, 203, 205  
 Goodness of fit tests, 203–208  
   for normality, 207–208  
   with data, 204–206  
   with numbers, 207  
 Grand median, 217  
 Grouped frequency distributions, 24–25  
 Groups, across and within, 171  
  
 Hard disk, 2  
 Highlight, 4  
 Homogeneity, Pearson *chi*-square test of, 211–214  
 Hypothesis tests for  
   entire distributions (*see* Pearson's *chi*-square tests)  
   many means (*see* Analysis of variance)  
   one correlation, 189–193  
   one mean, 144–149  
   one proportion, 139–143  
   one variance, 161–163  
   regression, 196–199  
   two correlations, 193–194  
   two means, 155–158  
   two proportions, 151–154  
   two variances, 163–164  
**HYPOTHESIS TESTS MENU** (*see* Menus)  
  
 Independence, Pearson *chi*-square test of, 214–217

- INDEPENDENT Samples Or MATCHED pairs**, 156, 160–161 (Exercise 7.5.9)
- Independent variable(s)  
analysis of variance, 167  
regression, 72, 197
- Indicator random variables, 220
- Inner fences (in box-and-whisker plots), 66
- INSTRUCTIONS**, 8
- Integers, choosing random, 106
- Interaction, 176  
mean squares as denominator in  $F$  test for random- and mixed-effects analyses, 176–177  
sum of squares pooled with sum of squares error, 178–179
- Interquartile range (IQR), 65
- Interval estimates (*see* Confidence intervals)
- Intervals, confidence (*see* Confidence intervals)
- Keyboard **DATA** entry for hypothesis tests, 139
- Kurtosis  
captured by Pearson's *chi*-square tests, 203  
coefficient of, 40
- Label(s)  
case (*see* Case labels)  
file, 12, 14–15
- Lack of fit, 198
- Large samples, 129
- LARGE Or SMALL Population**, 133
- Line, (least squares) regression, 81, 198
- Linear covariation, 196
- Linear regression  
deviation from, 198  
equation, 80
- List Data Matrix**, 13
- Location  
captured by Pearson's *chi*-square tests, 203  
statistics, 39
- Log factorial, 87–89
- Logarithm, natural (*see* Natural logarithm)
- m** (missing value code), 42
- M** (store in memory of pop-up calculator), 75, 103
- Main effects, 176
- MAIN MENU**, 7
- Major outlier(s), 33, 66
- Matched pairs, 156, 160–161 (Exercise 7.5.9)
- Matrix, 4  
correlation, 76–78, 189, 192  
data (*see* Data matrix)  
transposing, 42*n*, 108
- Mean, 39  
case transformation, 110  
confidence intervals for, 126–128, 132–135  
frequency plot of, 112–113  
sampling distribution of, 113–114  
**Summary Statistics** output, 39  
testing hypotheses about, 144–149, 155–158
- Mean SQ.**, 175
- Mean squares across treatments, 180
- Mean squares residual, 180
- Measures, repeated (ANOVA), 179–181
- Median  
ASP method of calculation, 39  
grand, 217  
**Summary Statistics** output, 39
- Median test, 217–219
- Menus  
**ALT COMMANDS MENU**, 8  
truncated, 44  
**ANALYSIS OF VARIANCE MENU**, 170  
**CASE TRANSFORMATION MENU**, 110  
**CROSSTAB**, 213  
**HYPOTHESIS TESTS MENU**  
**GOODNESS OF FIT TEST**, 203  
**MEAN X = MEAN Y**, 156  
**MEAN X = x**, 147  
**PROD M. COR(XY) = x**, 189  
**PROP. X = PROP. Y**, 152  
**PROPORTION X = x**, 140  
**VAR(X) = VAR(Y)**, 163  
**VAR(X) = x**, 161  
**x% CONF. INTERVAL**, 130  
**MAIN MENU**, 7  
**MISCELLANEOUS PLOTS MENU**, 20  
**BOX AND WHISKER PLOT**, 66–68  
**FREQUENCY PLOT**, 20  
**QUANTILE PLOT**, 49–52  
**SCATTER PLOT**, 71–75  
**STEM AND LEAF PLOT**, 30–31  
**ORDINARY LEAST SQUARES**, 82  
**PROBABILITY DISTRIBUTIONS MENU**  
**BINOMIAL**, 94–97  
 $F$ , 176, 177, 178–179  
**NORMAL**, 98–100, 103, 127, 142  
 $P = \text{COMB. } (a/b)$ , 89  
 $P = \text{FACTORIAL } a$ , 86  
 $P = \text{Ln FACTORIAL } a$ , 88  
 $P = \text{PERM. } (a/b)$ , 86  
 $t$ , 134, 149, 158  
**REGRESSION MENU**, 79  
**SELECTING CASES MENU**, 15–16, 150–151 (Exercise 7.3.6), 152–153

- Menus (*cont.*)  
 VARIABLE TRANSFORMATION MENU,  
 42–46, 56–57, 117–118  
 operators and relations, 43
- Minor outlier(s), 66
- Miscellaneous Plots Menu**, 20
- Missing scores, 42*n*, 57, 192–193
- Mixed effects, 177–179
- Mode, 39
- MODEL**, 80
- Moment, product, 77
- Monitor, 1, 5
- Multiple regression analysis, 197
- Natural logarithm, 87–89  
 in Fisher's *r*-to-*z*, 191, 193–194
- Nonlinear covariation, 196
- Normal approximation to binomial, 102, 104
- Normal probability, 98–100
- Normality, goodness-of-fit test for, 207–208
- Notation, 5, 21*n*, 38, 93
- Observed frequencies and proportions, 203
- On-off switch, 1
- One-way analysis of variance, 170–174
- .OUT, 26
- Outer fences (in box-and-whisker plots), 66
- Outlier(s), 62–68, 225  
 major, 33, 66  
 minor, 66
- Output, 26
- p*-value (*see* Significance probability)
- P-VALUE**, 142, 149, 162, 162*n*, 164, 171, 176*n*,  
 192, 199
- P-VALUE/2**, 140*n*, 143, 149, 164, 192
- Pairs, matched, 156, 160–161 (Exercise 7.5.9)
- Paull's Rule, 178
- Pearson's *chi*-square tests  
 association, 211–219  
 using contingency tables, 211–214, 217  
 using crosstabulations, 214–219  
 goodness of fit, 204–208  
 for normality, 207–208  
 with data, 204–206  
 with numbers, 207  
 median test, 217–219  
 properties of distributions captured by, 203  
 sample sizes required for, 214
- Percentile(s), 49–52
- Percentile rank, 50, 51–52
- Permutations, 86–86
- Plots (*see* Menus, MISCELLANEOUS PLOTS MENU,  
 and individual entries, e.g., Scatter plot)
- Point estimation, 115–116
- Pooled estimate of error variance, 178–179
- Pop-up calculator, 74–75, 90, 96–97, 99–100, 103,  
 127, 135, 142, 144–145, 148, 176–179, 193–194
- Population coefficient of determination, 196
- Population correlation ratio, 196
- Population proportion (*p*)  
 confidence intervals for, 129–132  
 testing hypotheses about, 139–143, 151–154
- Population variance, 123
- Printer buffer, 2
- Printing, 27
- Probability  
 binomial, 93–97  
 normal, 98–100  
 significance (*see* Significance probability)
- PROBABILITY DISTRIBUTIONS MENU** (*see* Menus)
- Probability table (binomial, as ASP output), 94
- PROBABILITY Or TABLE Value**, 98, 103, 127,  
 145, 149
- PROD M. COR(XY) = x**, 189
- Product moment, 77
- Prompts, 5
- PROP. X = PROP. Y**, 152
- Proportion  
 confidence intervals for, 129–132  
 population, 139  
 sample, 117  
 testing hypotheses about, 139–143, 151–154
- PROPORTION X = x**, 140
- Proportions, expected and observed, 203
- Quantities, basic computational in **Summary  
 Statistics** output, 39
- QUANTILE PLOT**, 49–52
- Quantiles, 49, 50
- Quartile(s), 49, 65
- R** (recall from pop-up calculator memory), 75
- R. SQ.**, 73, 81
- r*-to-*z*, 191–192, 193–194
- Raise to a Power** (on pop-up calculator), 74, 99
- RANDOM BLOCK/REPEATED MEASURES**, 180
- Random effects, 176–177, 178–179
- Random Integers**, 106
- Random samples, drawing 105–108
- Random variable(s)  
 indicator, 116–117  
 standard normal, 98
- Range, 39

- Rank, percentile 50, 51–52
- Ratio, correlation, 196
- Raw score, synthesizing missing, 62
- Reciprocal, obtaining with pop-up calculator, 90
- REGRESSION**, 73, 81
- Regression, 72, 79–84  
hypotheses about, 196–199
- Regression Analysis**, 79, 189, 196–199
- Regression analysis, multiple, 197
- Regression equation, linear (*or* least squares), 80, 81, 196, 197–198
- Regression line (least squares), 81, 198
- REGRESSION MENU**, 79
- Repeated measures (ANOVA), 179–181
- Replacement,  
sampling with, 106, 112*n*  
sampling without, 133*n*
- Replacing variables in data matrix, 45, 118
- Residual, mean squares, 180
- Root directory, 3
- Sample mean, 39  
case transformation, 110  
frequency plot of, 112–113  
sampling distribution of, 113–114  
**Summary Statistics** output, 39
- Sample proportion, 117
- SAMPLE PROPORTION OF X**, 143
- SAMPLE SIZE OF X**, 143
- Sample standard deviation, 40
- Sample variance, 40  
case transformation, 123–124  
**Summary Statistics** output, 40
- Samples  
drawing random, 105–108  
independent *vs.* correlated groups, 156
- Sampling  
distribution (*see* Distribution, sampling)  
with replacement, 106, 112*n*  
without replacement, 133*n*
- Saving  
data matrix, 12  
results, 26
- Scaling constant, 47
- Scatter plot (*and* **SCATTER PLOT**), 71–75
- Schematic plot, 67
- Score(s)  
missing, 42*n*, 57, 192–193  
standard, 54–56  
synthetic, 62
- Scr./Data Dir./Prt.Port Dflts**, 9
- Screen buffer, 2
- SD. ER. EST.**, 73, 81
- Select Cases By CASE Label**, 63, 151  
(Exercise 7.3.6e)
- Select Cases DIRECTLY By Row Number**, 16, 152–153
- Select Cases If: X (<,>=,etc.) a**, 150–151 (Exercise 7.3.6b)
- SELECTING CASES MENU**, 15–16, 150–151  
(Exercise 7.3.6), 152–153
- Shape of distribution (skew and kurtosis), 40
- Significance level ( $\alpha$ ), 144, 163
- Significance probability (*p*-value)  
ANOVA, 171, 176, 176*n*, 177, 178, 179  
correlation, 192, 194  
Pearson's *chi*-square tests, 213, 219  
regression, 199  
tests of means, 144, 145  
tests of proportions, 139, 142–143, 143*n*, 154  
tests of variances, 161, 163, 164  
*See also* **P-VALUE** *and* **P-VALUE/2**
- Skew (*and* skewness)  
captured by Pearson's *chi*-square tests, 203  
coefficient of, 40
- Sort Or Transpose Data Matrix**, 42*n*, 108
- Space bar, 5
- Standard deviation, 40  
sample, 40  
**Summary Statistics** output, 40
- Standard error of estimate, 82
- Standard normal random variable, 98
- Standard scores, 54–56
- Statistics  
dispersion, 39–40  
location, 39  
summary (**Summary Statistics**), 37–40  
suppress (**Suppress Statistics**), 23, 24
- Status line, 5
- Stem-and-leaf plots, 30–36  
stretched, 34–35  
twigs, 34
- Student's *t* (and Student's *t* distribution), 132, 134, 146–149, 158
- Subdirectory, 3
- Success, in binomial experiment, 94
- Sum**, 39
- SUM OF-SQ'S** (in ANOVA output), 175
- Sum Of Squares** (basic computational quantity), 39
- Sum Of Squared Dev.**, 39
- Summary Statistics**, 37–40
- Suppress Statistics**, 23, 24
- Synthesizing raw scores, 62
- Synthetic (raw) score, 62

- t** (*see* Menus, PROBABILITY DISTRIBUTIONS MENU)
- t** (*and t* distribution), 132, 134, 146–149, 158
- T-scores**, 56–57
- Table(s)**  
contingency, 211–214, 217  
probability (binomial, as ASP output), 94
- Table Value**, 100, 127
- Test(s)** (*see* Hypothesis tests *or* specific entries, e.g., Goodness of fit tests)
- Text editor**, ASCII, 26
- Transformation(s)**, 42–47  
case (*see* Case transformations *and* Menus, CASE TRANSFORMATION MENU)  
variable (*see* Menus, VARIABLE TRANSFORMATION MENU *and* Variable transformation)  
*See also* Standard scores *and* T-scores
- Transposing data matrices**, 42*n*, 108
- Treatment(s)**  
groups, 167–168  
mean squares across (in repeated measures ANOVA), 180
- Truncated ALT COMMANDS MENU**, 44
- Two-way analysis of variance**, 174–179
- Unbiased estimator (of variance), 123
- Unbiasedness, 115
- Ungrouped frequency distributions, 21–24
- Uniform distribution, 106
- V=View**, 26
- Value**  
calculated (*see* Calculated Value)  
table (*see* PROBABILITY Or TABLE Value)
- Values**  
adjacent, 66
- Values (cont.)**  
missing, 42*n*, 57, 192–193
- VAR(X) = VAR(Y)**, 163
- Variable(s)**  
adding to matrix, 45, 57–60  
breaking down, 72  
coding as levels in ANOVA, 168, 180  
columns in ASP data matrix, 4  
crosstabulation of, 212, 214–219  
deleting from a matrix, 107  
dependent (*see* Dependent variable)  
independent (*see* Independent variable)  
predictor, 72  
replacing in data matrix, 45, 118  
rows in transposed matrix, 108  
selecting samples from, 156  
transformation, 42, 54, 117–120 (*see also* Menus, VARIABLE TRANSFORMATION MENU, Standard scores, *and* T-scores)
- Variance**, 40  
analysis of (*see* Analysis of variance)  
case transformation, 111  
confidence intervals for, 135–136  
population, 123  
sample (*see* Sample variance)  
**Summary Statistics** output, 40  
testing hypotheses about 161–164  
unbiased estimate of, 123  
*See also* Sample variance
- Variance of estimate**, 82
- Variation**, coefficient of, 40
- V = View/Rename/Delete OUTPUT File**, 26
- Window**, 5
- Within-groups**, 171
- x% CONFD. INTERVAL**, 130