

Index

Page numbers for volume 2 are indicated by (2).

- absolute value 1–2
 - archimedean absolute value 1
 - non-archimedean absolute value 1
 - trivial absolute value 1
- action
 - by differential operator (2) 37, (2) 40, 153, 160, 172
 - by right translation (2) 36, (2) 40, 153, 160, 172
 - of Maass operators on Whittaker functions 92
 - of \mathbb{Q} on the adèles and ideles 8
 - of the Hecke algebra 399
 - on the Kirillov space 224
- additive character 15
 - adelic 18
- adele 7
 - finite adèles 8
 - group 107–108
- adelic
 - automorphic form 118, 119, (2) 26
 - automorphic representation 152–161
 - automorphic representation for $GL(n, \mathbb{A}_{\mathbb{Q}})$ (2) 36–41
 - Bruhat-Schwartz function 20
 - Bruhat-Schwartz space 46
 - Cartan decomposition 111
 - cuspidal form 119, 311, (2) 29
 - Whittaker coefficient of (2) 33
 - space of 160, 311, (2) 40
 - Fourier transform 18–22
 - integral 20
 - Iwasawa decomposition 110
 - lift 119–126, 136–141
 - of a classical automorphic form 136–141
 - of a vector valued automorphic function (2) 25
 - of Maass form 119–126, 166
 - matrix 419
 - Poisson summation formula 30–31
- admissible
 - $(\mathfrak{g}, K_{\infty}) \times GL(2, \mathbb{A}_{\text{finite}})$ -module 173–178, 403–406
 - $(\mathfrak{g}, K_{\infty}) \times GL(n, \mathbb{A}_{\text{finite}})$ -module (2) 38
 - $(\mathfrak{g}, K_{\infty})$ -module 259–260, (2) 82
 - dual (2) 106
- admissible representations
 - of $GL(2, \mathbb{A}_{\text{finite}})$ 404
 - of $GL(2, \mathbb{Q}_p)$ 183–192, 252–253, 404
 - of $GL(n, \mathbb{Q}_p)$ 2(52)
- algebra
 - of differential operators 114
 - simple 418
- algebraic direct sum of vector spaces 265
- archimedean 1
- associative algebra 112
- asymptotic
 - behavior 84
 - of matrix coefficient 323, 330–343
- Atkin-Lehner theory 97, (2) 46
- automorphic
 - cuspidal representation 160, 172–173, 177, 372, (2) 41
 - form 40, 76–77, 118–119, (2) 26
 - adelic 40, 118, 119, (2) 26
 - classical (2) 11, *see also* Maass form
 - classical associated to an adelic automorphic form 175
 - for $GL(n, \mathbb{A}_{\mathbb{Q}})$ (2) 26
 - K_{∞} -fixed (2) 1
 - \mathcal{L}^2 - 83–84, 92–94, 119, 126, 369, 374, (2) 97, (2) 119
 - non-holomorphic 414, 418
 - function (2) 3–13
 - for $\Gamma_0(N)$ 79
 - of integral weight 78–80
 - vector space of 82
 - vector valued (2) 6
 - with multiplier ψ 76
 - relation 82, 115, 117, (2) 6
 - representation 65–68, 152–178, (2) 36–41
 - adelic 152–161
 - with central character 160, (2) 40
- automorphy relation 82, 115, 117, (2) 6

- basic types of
 - generic irreducible unitary representations of $GL(n, \mathbb{Q}_p)$ (2) 101
 - generic irreducible unitary representation of $GL(n, \mathbb{R})$ (2) 98
 - p -adic irreducible unitary representations of general linear groups (2) 100
 - unitary representations of real general linear groups (2) 96
- basis 207, 265
- Bernstein-Zelevinsky classification (2) 75
 - for $GL(n, \mathbb{Q}_p)$ (2) 70–75
 - of discrete series representations of $GL(n, \mathbb{Q}_p)$ (2) 77
 - of tempered representations of $GL(n, \mathbb{Q}_p)$ (2) 77
- Bessel function 85, 261
- bi K -invariant function 396
- bilinear form
 - canonical 280, 294
 - invariant 279, 280, 293, 294, 315, 331
 - nondegenerate (2) 55
 - on $S(\mathbb{Q}_p^s)$ 285
 - on the vector space of adelic cusp forms 311
- Borel
 - set 424
 - subgroup (2) 17, (2) 21
 - of $GL(n, \mathbb{R})$ (2) 17
 - of $GL(n, \mathbb{Q}_p)$ (2) 21
 - subset 309
- bottom of the spectrum 92–94
- bound on parameters, unitary representation 361, (2) 79
- boundedness of matrix coefficients, unitary representation 375, (2) 78
- Bruhat decomposition
 - for $GL(2, \mathbb{Q}_p)$ 319
 - for $GL(2, \mathbb{R})$ 299
 - for $GL(n, \mathbb{Q}_p)$ (2) 22
 - for $GL(n, \mathbb{R})$ (2) 17
- Bruhat-Schwartz function/space
 - for $\mathbb{A}_{\mathbb{Q}}$ 20, 46
 - for $GL(2, \mathbb{Q}_p)$ 188
 - for $\text{Mat}(2, \mathbb{A}_{\mathbb{Q}})$ 419, 430
 - for $\text{Mat}(n, \mathbb{A}_{\mathbb{Q}})$ (2) 125
 - for $\text{Mat}(2, \mathbb{Q}_p)$ 433
 - for $\text{Mat}(n, \mathbb{Q}_p)$ (2) 125
 - for \mathbb{Q}_p^* 215
 - for \mathbb{R} 15
- Cartan decomposition
 - for $GL(n, \mathbb{Q}_p)$ (2) 20
 - for $GL(2, \mathbb{R})$ 104
 - for $GL(n, \mathbb{R})$ (2) 16
 - p -adic 107
- Casimir 164
- Cauchy-Riemann condition 82
- Cauchy-Schwartz inequality 360
- cell
 - big 319
 - little 319
- center of the universal enveloping algebra 117, 269, (2) 8
- central character 118, 119, 187, (2) 54
 - of an adelic lift 140
- character
 - additive 15
 - central 118, 119, 187, (2) 54
 - Dirichlet 41
 - Hecke 40
 - normalized unitary character 71, 201, 263
 - ramified 44
 - sign 263
 - spherical Hecke 402, (2) 103
 - unramified 43
- characterization
 - of irreducible tempered $(\mathfrak{g}, K_{\infty})$ -modules 368
 - of irreducible unitary principal series representations 363
 - of representations of $GL(2, \mathbb{Q}_p)$ via the growth of matrix co-efficients 326
 - of the finite dimensional irreducible representation of $GL(n, \mathbb{Q}_p)$ (2) 54
 - of the irreducible tempered representations of $GL(2, \mathbb{Q}_p)$ 363
 - of the Kirillov space 225
 - of unitary special representations of $GL(2, \mathbb{Q}_p)$ 365
- Chinese Remainder Theorem 9
- class field theory 51
- classical automorphic form 175
- classical L-function 60–65
- classification
 - Bernstein-Zelevinsky (2) 70–75
 - Langlands (2) 77
 - of irreducible admissible $(\mathfrak{g}, K_{\infty})$ -modules 269–274
 - of irreducible unitary representations of $GL(n, \mathbb{R})$ (2) 97
 - of linear algebraic groups (2) 106
 - of smooth irreducible representations of $GL(n, \mathbb{Q}_p)$ via the growth of matrix coefficients (2) 75–78
 - of supercuspidal representations (2) 67
 - of the generic irreducible unitary representations of $GL(n, \mathbb{Q}_p)$ (2) 101
 - of the generic irreducible unitary representations of $GL(n, \mathbb{R})$ (2) 98
 - of the irreducible, generic, unramified unitary representations of $GL(n, \mathbb{Q}_p)$ (2) 102–105
 - of the smooth irreducible representations (2) 75–78
 - of the smooth irreducible unitary representations of $GL(n, \mathbb{Q}_p)$ (2) 101

- Rough classification of the irreducible admissible representations of $GL(n, \mathbb{Q}_p)$ (2) 69
- cocycle
 j-cocycle 76
 classical (2) 5
 one-cocycle (2) 3, (2) 5
 relation (2) 4, (2) 5
- commutative ring containing “1” (2) 27, (2) 28
- commutator subgroup 254
- compact-open topology (2) 105
- compactly supported
 function 70
 modulo the center 287
- complementary series representation 367, (2) 98
 of $GL(2, \mathbb{Q}_p)$ 363
 of $GL(2, \mathbb{R})$ 367, (2) 98
 of $GL(n, \mathbb{Q}_p)$ (2) 80
- complex parameter 361, (2) 79
- complexified universal enveloping algebra 115, (2) 81
- composition factor (2) 73
- conductor 44
 of a locally constant compactly supported function 70
- congruence subgroup 77, (2) 2–3
 non-congruence subgroup 79
 principal congruence subgroup 77
- conjecture
 Ramanujan conjecture 413–414
 Ramanujan-Petersson conjecture 414
 Selberg conjecture 413–414
- constant term
 of a classical automorphic form (2) 11
 of an adelic automorphic form (2) 29
- continuation
 analytic 48, 54, 199, 206, 419
 meromorphic 47, 60
- continuous homomorphism 40
- continuous representation 254
- contragredient
 of a (\mathfrak{g}, K_∞) -module 290, (2) 82
 of a $(\mathfrak{g}, K_\infty) \times GL(2, \mathbb{A}_{\text{finite}})$ -module 303–306, 369
 of a generic representation (2) 58, (2) 87
 of a parabolically induced representation (2) 65
 of a principal series representation 283
 of a smooth representation of $GL(n, \mathbb{Q}_p)$ (2) 54
 of a special representation 283–285
 of a supercuspidal representation 285–288
 of a unitary (\mathfrak{g}, K_∞) -module 366
 of a unitary representation 358, (2) 78
 of an irreducible admissible cuspidal automorphic representation 315
 of irreducible (\mathfrak{g}, K_∞) -modules 298
 contragredient representation 277–281, 289–294
 of a cuspidal automorphic representation 311–316
 of a principal series representation 281–283, 294–303
- convolution algebra 396
- coset representative 80
- cuspidal cusp 78
 form 370
 holomorphic 414
 space of cusp forms 160, (2) 40
 inequivalent cusps 80
 parameter 87
- decomposition
 Bruhat 299, 319, (2) 17, (2) 22
 Cartan 104, 107, 111, (2) 16, (2) 20
 Iwasawa 103, 105, 110, (2) 2, (2) 18
- decomposition of representations 388–396
- decomposition theorem 389
- degree 37
- dense 108, (2) 23
- diagonal embedding 8
- diagonal matrix 110
- differential form 77
- differential operator 112, 161, 162
- Dirichlet character 41
- Dirichlet L-function 45
- discrete series 268, (2) 77
 essentially 268
 limit of 268
 of $GL(n, \mathbb{Q}_p)$ (2) 77
 representation 372 (2) 77
- discrete series representation 372
 of $GL(2, \mathbb{R})$ 471–474, (2) 99
 of $GL(n, \mathbb{Q}_p)$ (2) 77
 of $GL(n, \mathbb{R})$ (2) 99
- division ring 418
- divisor 37
- Dixmier’s lemma 186
- double coset 203
- dual (2) 105–106
 admissible (2) 106
 Langlands (2) 106
 of a (\mathfrak{g}, K_∞) -module 289
 of a $(\mathfrak{g}, K_\infty) \times GL(2, \mathbb{A}_{\text{finite}})$ -module 303
 Pontryagin (2) 105
 representation 277
 spherical (2) 106
 unitary (2) 105–106
- ϵ -factor 451, *see also* root number
- eigenfunction 370
- eigenvalue 174
- Eisenstein series 89–90
- elliptic curve 52
- embedding 168
 diagonal 8, 108
- equivalence of absolute values 1
- equivalent conditions for supercuspidality (2) 67

- Euler product 378–379
 - factorization 378
 - exact 350
 - explicit computation of $\gamma(s, \omega)$ 57
- factor of automorphy 76
- factorizable function 20, 39, 430, (2) 125
- factorizability for representations 67, 68, 378–413, (2) 41–43
- factorization
 - Euler 378
 - initial factorization of a $(\mathfrak{g}, K_\infty) \times GL(2, \mathbb{A}_{\text{finite}})$ -module 404
 - Iwahori 343
 - of adelic integral 21
 - of \mathfrak{h}^n (2) 10
 - of Haar measure 230, 425, (2) 132
 - of integral 295
 - of the global Whittaker function 133
 - of the global zeta integral 430–432, (2) 124–125
 - of unramified irreducible admissible cuspidal automorphic representation 383–388
 - unique factorization of adèles 111
- field of formal Laurent series 34
- finite, finiteness
 - finite adèles 8
 - finiteness property of a (\mathfrak{g}, k_∞) -module 156
 - K_∞ (2) 26
 - right K 118
 - $Z(U(\mathfrak{g}))$ 118, (2) 17
- finite dimensional representation
 - continuous of $GL(n, \mathbb{Q}_p)$ 106
 - of K_∞ (2) 4, (2) 82
- finite length (2) 52, (2) 65
- finitely generated representation 191, (2) 52
 - smooth 191, (2) 53
- fixed vector
 - $GL(2, \mathbb{Z}_p)$ fixed/ k -fixed vector 192, 400
 - $K_{0,1}(p^m)_p$ (2) 44, (2) 45
 - K_∞ 260
- flat section 206
- Fourier coefficient 96
- Fourier expansion 23–30, 80–81
 - of an adelic automorphic form 126–128, (2) 31–36
 - of an adelic cusp form (2) 32
- Fourier inversion 453
 - on \mathbb{Q}_p 17
 - on \mathbb{R} 15
 - on the adèles 22
- Fourier transform
 - adelic 18–22
 - on \mathbb{R} 15–18
 - p -adic 15–18
- Fourier-Whittaker expansion 87–89
- Frobenius reciprocity 239
- function
 - adelic 19
 - automorphic (2) 3–13
 - Bruhat-Schwartz 20, *see* Bruhat-Schwartz function/space
 - Bessel 85, 261
 - characteristic 17
 - compactly supported 12, 70
 - entire 49
 - Euler’s 71, 172
 - factorizable 20
 - holomorphic 80–81
 - K-Bessel 85, 261
 - K -invariant 396
 - locally constant 12
 - locally integrable 307
 - periodic adelic 23
 - periodized Bruhat-Schwartz 23
 - radially symmetric 398
 - restriction 221
 - Schwartz 340
 - smooth vector valued (2) 6
 - spherical 398
 - test 47, 438, 440, 441, 463
 - vector valued (2) 8
 - vector valued automorphic (2) 6
- functional 13
- functional equation
 - Dirichlet L-function 45
 - for the induced representation (2) 137
 - global 62
 - local 56, 432–434, (2) 125–128
 - of Jacquet’s Whittaker function 262
 - of the global zeta integral 62, 426, (2) 120
- fundamental domain 9–11, 78, 109, 111
- fundamental identity for local integrals of unitary supercuspidal representations of $GL(2, \mathbb{Q}_p)$ 454
- (\mathfrak{g}, K_∞) -module 156, 259, 289, (2) 37
- (\mathfrak{g}, K_∞) -module of $GL(n, \mathbb{R})$ (2) 81
- $(\mathfrak{g}, K_\infty) \times GL(n, \mathbb{A}_{\text{finite}})$ -module (2) 38
- $(\mathfrak{g}, K_\infty) \times GL(2, \mathbb{A}_{\text{finite}})$ -module 157, 368–374
- function field 33
- G -module 191, (2) 53
- G -submodule 192, (2) 53
- Gauss sum 58
- generic (2) 31, (2) 58
- generic representation (2) 56–60
 - irreducible unitary (2) 98, (2) 101
 - of $GL(n, \mathbb{Q}_p)$ (2) 56–60
 - of $GL(n, \mathbb{R})$ (2) 85–88
- global
 - conductor 54
 - functional equation 62
 - matrix coefficients 374, (2) 119
 - new vector (2) 45
 - Whittaker function 128–134, 141–147
 - zeta integral 425–430
 - factorization of 430–432, (2) 124–125

- for $GL(2, \mathbb{A}_{\mathbb{Q}})$ 425–430
 - for $GL(n, \mathbb{A}_{\mathbb{Q}})$ (2) 118
 - Tate's 61
- Godement-Jacquet L-function 418–419
- Godement-Jacquet method 419
- growth
 - condition 76
 - moderate 40, 79, 118, (2) 7, (2) 26
 - of matrix coefficients 316–330, (2) 75–78
 - polynomial 81
 - property 311, 316–330
- Haar measure 423–425
 - additive 13, 455
 - existence and uniqueness of 229, 424
 - factorization of 230, 425
 - left and right 424, 425
 - multiplicative 14, 426
 - on $GL(2, \mathbb{Q}_p)$ 228–232, 318, 319
 - on $GL(n, \mathbb{Q}_p)$ (2) 69
 - on $GL(2, \mathbb{R})$ 299
 - on \mathbb{Q}_p 13
- harmonic analysis 52
- Hecke
 - algebra 396–403, 408, (2) 103
 - character 66
 - newform 172
 - operator 68, 95, 378, 396
- Hecke algebra
 - spherical 396–403, (2) 103
 - S -spherical 408
- Hecke character 66
 - unitary 40
- Hermitian form 358, (2) 91
 - invariant 369
 - on adelic cusp form 369
- Hilbert space (2) 92
- idele, ideles 7
- idelic
 - absolute value 47
 - differential 47
 - lift 41, (2) 25
 - $GL(n)$ idelic lift of a Dirichlet character (2) 25
 - of a Dirichlet character 41
- index 77
- induced
 - map 349
 - non-normalized representation (2) 94
 - representation 238–240, 257
 - space (2) 63
- inner product 372
- integrable
 - locally integrable function 307
- Integration
 - Integral on $Z(\mathbb{A}_{\mathbb{Q}})\backslash GL(2, \mathbb{A}_{\mathbb{Q}})$ 307
 - Integral on $Z(\mathbb{A}_{\mathbb{Q}})GL(2, \mathbb{Q})\backslash GL(2, \mathbb{A}_{\mathbb{Q}})$ 309
 - on $GL(2, \mathbb{A}_{\mathbb{Q}})$ 306–311
 - on $GL(2, \mathbb{Q}_p)$ 228–232
 - on $GL(2, \mathbb{R})$ 299
- intertwining map 66
 - injective (2) 107
 - of $(\mathfrak{g}, K_{\infty})$ -modules 158, (2) 39
 - of $(\mathfrak{g}, K_{\infty}) \times GL(n, \mathbb{A}_{\text{finite}})$ -modules 159, (2) 39
- invariance of the integral 310
- invariant
 - bilinear form 315, 331
 - on principal series of $GL(2, \mathbb{Q}_p)$ 282
 - on principal series of $GL(2, \mathbb{R})$ 296, 300
 - differential operator 155, (2) 8
 - Hermitian form on cusp forms 369
 - linear functional 234
 - pairing 280, 294, 306
- inverse limit 6
- involution 398
- irreducible 188, 279
 - $(\mathfrak{g}, K_{\infty}) \times GL(2, \mathbb{A}_{\text{finite}})$ -module 157
 - $(\mathfrak{g}, K_{\infty}) \times GL(n, \mathbb{A}_{\text{finite}})$ -module (2) 38
 - quotient, (2) 53
 - representation 66, 184
 - admissible 252–253
 - automorphic 174
 - cuspidal automorphic 177
 - discrete series 471–474
 - special 364–365
 - spherical 435
 - square integrable 329
 - supercuspidal 364–365, 440
 - tempered 363, 368
 - unitary Hilbert space (2) 106
 - subquotient (2) 74
- Iwahori
 - factorization of K_n 343
 - subgroup 134
- Iwasawa decomposition 103, 105–107
 - adelic 110
 - for $GL(n, \mathbb{Q}_p)$ (2) 26
 - generalized (2) 50
 - p -adic 105
- Jacquet functor 350
- Jacquet integral 205
 - analytic continuation of 206
- Jacquet module 236–238
 - for $GL(n, \mathbb{Q}_p)$ (2) 68
 - of a principal series representation 347
 - of a special representation 351
- Jacquet's Whittaker function 86
 - local 195–200, 260–263
 - local K_{∞} -invariant 261
- Jordan-Hölder series (2) 73

Cambridge University Press

978-0-521-47423-8 - Automorphic Representations and L-Functions for the General Linear Group, Volume I

Dorian Goldfeld and Joseph Hundley

Index

[More information](#)

546

Index

- K_n -symmetrization 344
- K -invariance 436
- Kirillov
 - model 214–221, 243–252
 - of a supercuspidal representation 252
 - of the principal series representation 221–228
 - representation 214
 - space 221, 225
 - uniqueness of 243–252
- \mathcal{L}^2
 - automorphic form 83–84, 92–94, 119, 126, 369, 374, (2) 97, (2) 119
 - equivalence (2) 95
 - modulo the center (2) 119
 - space 83, 84, 374
- Langlands parameter 439
- Langlands program 51
- Laplace operator (Laplacian), weight K 82
 - relation with casimir 176–177
- left and right translation invariant measure 424
- level 84, (2) 6
- Levi
 - factor (2) 28
 - subgroup (2) 28
- L-functions 45–55, 60–65
 - analytic theory of 418–419
 - associated to automorphic representation 62
 - classical Dirichlet 48
 - Dirichlet 45
 - Godement-Jacquet 418–419
 - Hasse-Weil 52
 - local 55–60, 64, 434–440, (2) 143
 - of a discrete series representation (2) 142
 - of a supercuspidal representation (2) 128
 - of an automorphic form on $GL(2)$ 378
- L-group (2) 106
- Lie
 - algebra 116, (2) 80–85
 - bracket 112, (2) 80–85
 - group 77
- lift
 - adelic 119–126, (2) 24–25
 - central character of 140
 - from the upper half-plane \mathfrak{h} to $GL(2, \mathbb{R})^+$ 136
 - idelic 41, (2) 25
 - of an automorphic function (2) 7
- limits of discrete series 268
- linear fractional transformation 103
- local
 - component at p 444
 - functional equation 56
 - for $GL(1)$ 56
 - for parabolically induced representation (2) 137
 - for principal series representations 446–450, 467–471
 - for tensor products (2) 128–130
 - for the supercuspidal representations 452–462, (2) 128
 - for unitary special representations 450–452
- L-function 55–60, 64, 434–440, (2) 126–144
 - as a common divisor 64
 - associated to an irreducible discrete series representation 471–474
 - for the unitary special representation 444–446
 - of an irreducible principal series representation 441–444, 463–466
 - of an irreducible supercuspidal representation 440
 - of an unramified irreducible admissible representation 439
- matrix coefficient 426
- new vector (2) 44
- representation 193–195, 223, 226, (2) 85
 - from a Maass form 193–195, 223, 226
- root number 62
- unitary character 55
- Whittaker function 195, 260
- zeta integral 55
 - associated to a supercuspidal representation 440, 452
 - evaluation of 438, 440, 441, 463
 - for a parabolically induced representation (2) 130–138
 - for discrete series (2) 138–143
 - for irreducible unitary generic representation of $GL(n, \mathbb{R})$ (2) 143–151
 - meromorphic continuation of 60
 - operator valued 455
- locally compact group 388–396
 - abelian (2) 105
 - totally disconnected 389
- locally constant 12
- Maass cusp form 89
 - non-holomorphic 396
 - vector valued (2) 12
- Maass form 81–84, 98
 - cusp 89
 - even 119
 - odd 126
- Maass raising and lowering operators 90–92, 162
- map
 - induced 349
 - intertwining 66
 - linear 277, 399
- matrix coefficient 281, 301, (2) 84
 - factorizable 374, 430, (2) 124
 - for (\mathfrak{g}, K_∞) -module 299, 302
 - for $GL(2, \mathbb{Q}_p)$ 281, 343–353
 - for $GL(n, \mathbb{Q}_p)$ (2) 55
 - global (2) 119
 - growth of 316–353, (2) 75–78

- of a cuspidal automorphic representation 374, (2) 119
- of a principal series representation 318
- of a special representation 316, 328
- of a supercuspidal representation 288, 326, 440, 461, 462
- of a unitary representation 359, (2) 78
- of an automorphic representation 374, 425–426, 430–431, (2) 119–120, (2) 124–125
- of the contragredient representation 426
- maximal compact subgroup 105
- maximal standard parabolic subgroup (2) 30
- maximal unipotent subgroup (2) 31
- measure
 - Haar 423–425
 - on \mathbb{Q}_p 12
 - product 425
- Mellin inversion 336
 - p -adic 71, 70
- Mellin transform 336
 - p -adic 70–72
- minimal parabolic (2) 141
- mirabolic subgroup (2) 33
- model 213
 - consisting of matrix coefficients 317
 - Kirillov 214–221, 243–252
 - of a representation 213
 - Whittaker 213, 223, (2) 56, (2) 57, (2) 85, (2) 86
- moderate growth 40, 79, 118, (2) 7, (2) 26
- modular form
 - Fourier expansion of a holomorphic 80–81
 - holomorphic 77
- modular group 77
- modular quasicharacter (2) 61, (2) 91
- modular relation 418
- module
 - (\mathfrak{g}, K_∞) - 156, 259, 289
 - $(\mathfrak{g}, K_\infty) \times GL(2, \mathbb{A}_{\text{finite}})$ 157, 368–374
 - $(\mathfrak{g}, K_\infty) \times GL(n, \mathbb{A}_{\text{finite}})$ (2) 38
 - Jacquet 236–238
- multilinear algebra (2) 66
- multiplicative Haar measure
 - on \mathbb{Q}_p^\times 14
 - on $GL(2, \mathbb{Q}_v)$ 426
- multiplicativity of the Fourier coefficients 96
- multiplicity one
 - for $GL(n, \mathbb{Q}_p)$ (2) 59
 - for $GL(n, \mathbb{R})$ (2) 87
- natural projection (2) 60, (2) 88
- newform 97, 169
 - classical 97, (2) 45
 - for $GL(n)$ (2) 43–47
 - Hecke 172
 - holomorphic Hecke 172
 - local new vector (2) 44
- non-archimedean 1
- non-normalized induction (2) 63
- nondegenerate (2) 55
- nonunitary representation (2) 145
- norm-square 316, 329
- normalized
 - Haar measure 230
 - Haar measure on $GL(n, \mathbb{Q}_p)$ (2) 69
 - parabolic induction for $GL(n, \mathbb{Q}_p)$ (2) 62
 - parabolic induction for $GL(n, \mathbb{R})$ (2) 93
 - unitary character 71
 - of \mathbb{Q}_p^\times 201
 - of \mathbb{R}^\times 263
- normalizer (2) 29
- odd 126, 337
- oldform 97, 169
- one-dimensional 400
- operator
 - Casimir 164, 176–177, (2) 7–8
 - classical Hecke 95
 - differential 90–94, 112–117, 153–155, 160–166, 174, 176–177, 469, 497, (2) 8–10, (2) 37, (2) 40, (2) 81–82
 - Hecke 68, 95, 396, 378, 435–437
 - intertwining 66, 159, (2) 40
 - Laplace 82
 - lowering 162
 - Maass lowering 90, 162
 - Maass raising 90, 162
 - projection 390
 - raising 162
 - slash (2) 6, 83
- ordered partition (2) 27
- orthogonality relation 72, 177, 220, 461, (2) 149
- p -adic 2
 - Cartan decomposition 107
 - field 3
 - Fourier transform 15–18
 - integer 6
 - integration 12–15
 - Iwasawa decomposition 105
 - Mellin inversion 71, 70
 - Mellin transform 70–72
 - number 2
 - representation 414
- pairing, *see also* bilinear form
 - Bruhat 322
 - canonical bilinear (2) 107, 280, 294
 - invariant 280
 - nondegenerate bilinear (2) 110, 306
 - tensor product (2) 110
 - uniqueness of invariant 294, 280
- parabolic
 - induction 238–240
 - for $GL(n, \mathbb{Q}_p)$ (2) 60–66
 - for $GL(n, \mathbb{R})$ (2) 88–100

Cambridge University Press

978-0-521-47423-8 - Automorphic Representations and L-Functions for the General Linear Group, Volume I

Dorian Goldfeld and Joseph Hundley

Index

[More information](#)

548

normalized (2) 62, (2) 93
 subgroup (2) 27, (2) 28
 parabolically induced
 Hilbert space (2) 92
 positive definite Hermitian form (2) 91
 pre-Hilbert space (2) 91
 representation (2) 65
 parity of an irreducible admissible
 (\mathfrak{g}, K_∞) -module 260
 partition (2) 27
 periodic adelic function 23
 periodized Bruhat-Schwartz function 23, 24
 Petersson inner product 83
 Poisson summation formula 30–31
 for $GL(2, \mathbb{A}_\mathbb{Q})$ 419–423
 for $GL(n, \mathbb{A}_\mathbb{Q})$ (2) 114–118
 Pontryagin dual of a locally compact abelian
 group (2) 105
 positive definite Hermitian form 358
 parabolically induced (2) 91
 positive definite norm on \mathbb{C}^r (2) 10
 pre-Hilbert space (2) 91
 prime 2
 infinite 2
 principal series representation 200–205, 226,
 263–268
 Kirillov model of the 221–228
 of $GL(n, \mathbb{R})$ 296, (2) 95
 the vector space $\mathcal{V}_\infty(s, \omega)$ 263
 the vector space $\mathcal{V}_p(s, \omega)$ 201
 product
 associative 112
 convolution 396
 direct 425
 infinite 62
 product formula 2
 products of general linear groups (2) 61
 projection operator 191, 278, 291, 344, 390,
 (2) 62, (2) 150
 pure tensor 67

 quasicharacter (2) 61, (2) 91
 quotient (2) 53
 of a representation (2) 56–60
 space (2) 53
 unique irreducible (2) 142

 Ramanujan conjecture 439, *see also* Selberg
 conjecture
 at ∞ 466
 for $GL(2, \mathbb{A}_\mathbb{Q})$ 413–414
 for $GL(n, \mathbb{A}_\mathbb{Q})$ (2) 106
 Petersson 414
 ramification 54
 ramified 44
 (\mathfrak{g}, K_∞) -module for $GL(n, \mathbb{R})$ 260,
 (2) 82
 at p 406, (2) 42
 character 44

Index

representation 192
 representation of $GL(n, \mathbb{Q}_p)$ (2) 53
 rank of a matrix 421, (2) 117
 Rankin-Selberg method 69
 rapid decay 311, (2) 119
 rational function field 33
 reducibility
 of a representation induced from
 supercuspidal (2) 72
 of principal series 267
 reducible (2) 70
 parabolically induced representation (2) 127
 representation (2) 127
 region of absolute convergence 445, (2) 127,
 (2) 128, (2) 139
 representation
 automorphic 65–68, 152–178
 automorphic cuspidal 172–173, 372
 complementary series 363, 367, (2) 80
 essentially square integrable 330
 group 65
 induced 238–240, 257
 induced from supercuspidals (2) 72
 irreducible 66, 184, 252–253
 isomorphic 66
 Kirillov 214
 nonunitary (2) 145
 of finite length (2) 52
 p -adic 414
 parabolically induced (2) 65
 parabolically induced from supercuspidal
 representations (2) 71
 principal series 200–205, 226, 263–268
 quotient of a (2) 56–60
 ramified 192
 restricted 239
 smooth 183, 184, 404
 special 232–236, 364–365
 spherical 402
 square integrable (2) 76, 329
 supercuspidal 240–243, 364–365
 tempered 363, 368, (2) 76, (2) 80
 unitarizable 358
 unitary group (2) 83
 unramified 192
 residues of Eisenstein series 83
 restricted
 direct product 7, 67, 107, (2) 22
 representation 239
 tensor product of local representations 382,
 (2) 41
 tensor product of vector spaces 381
 Riemann hypothesis for curves 414
 Riemann-Roch formula 38
 Riemann zeta function 46, 50
 ring of Borel sets 424
 root number 54–55, 62–65, 434, 450–452, 468,
 (2) 126–128, (2) 131, (2) 144
 local 62

- scaling matrix 89
- Schur orthogonality (2) 149
for unitary supercuspidal representations 461
- Schur's lemma
for irreducible admissible (\mathfrak{g}, K_∞) -module 292
for irreducible and admissible $(\mathfrak{g}, K_\infty) \times GL(2, \mathbb{A}_{\text{finite}})$ -modules 304
for irreducible smooth representations 186
- Schwartz function 340
- Selberg conjecture 413–414
- Selberg spectral decomposition 83
- self-adjoint 370
extension 93
- slash operator 83, (2) 6
- smooth, smoothness 109, 188, (2) 25
adelic function 24
 $(\mathfrak{g}, K_\infty) \times GL(2, \mathbb{A}_{\text{finite}})$ -module 157
 $(\mathfrak{g}, K_\infty) \times GL(n, \mathbb{A}_{\text{finite}})$ -module (2) 38
 H -module 236
linear map 277
representation 183, 184, 404
vector valued function (2) 6
- special representation 232–236, 364–365
- spectrum
bottom of the 92–94
continuous 89
- spherical, *see also* unramified
dual (2) 106
function 398
Hecke algebra 396–403, (2) 103
Hecke character 402, (2) 103
representation 402
- square integrable, *see also* direct series
essentially 330
modulo the center 326
representation 329, (2) 76
- stability subgroup 84
- stabilizer
open (2) 52
- standard
Levi subgroup (2) 28
maximal parabolic subgroup (2) 49
parabolic subgroup (2) 27
representative (2) 76
- strong approximation 109
for adèles 10
for congruence subgroups 134–135
for $GL(n)$ (2) 22–24
for ideles 11
for $K_0(N)$ (2) 23
for prime power level subgroups 134
- subquotient 159, (2) 40, (2) 53
- subrepresentation (2) 53
- supercuspidal
equivalent conditions for supercuspidality (2) 67
representation 240–243, 364–365
representation of $GL(n, \mathbb{Q}_p)$ (2) 66–70, (2) 71
support (2) 74
- Tate series 69
- Tate's thesis 51
- tempered representation 363, 368, (2) 78, (2) 80, (2) 100
- tensor product
infinite 381–383
of a (\mathfrak{g}, K_∞) -module and a representation of $GL(2, \mathbb{A}_{\text{finite}})$ 404
of (\mathfrak{g}, K_∞) -modules and representations 379–380
of local representations 380, 381–383
of 2 group representations 379
of unitary representations (2) 89
pairing (2) 110
theorem 406–413
theorem for $GL(n)$ (2) 41–43
- topological group 8, 389
- topological space 7, 389
- topological vector space 66
- torus 238
quadratic 188
- totally disconnected group 388, 389
- translation
additive 13
right 153, 65
- trivial extension (2) 62
- twist
of a $GL(2, \mathbb{Q}_p)$ representation by a character 330
of a representation of $GL(n, \mathbb{Q}_p)$ (2) 72
of a representation with unitary central character 330
of a tensor product representation of $M_k(\mathbb{R})$ (2) 90
- type
 K_∞ -type 270, 272
- uniformity of behavior of matrix coefficients 317
- unimodular group 425
- unipotent radical (2) 28
- uniqueness of factors (tensor product theorem) 412, (2) 43
- unitarizable representation 358
- unitary
central character 330
dual (2) 105–106
 (\mathfrak{g}, K_∞) -module 365–368, (2) 83
Hecke character 40
principal series representation 360–364
representation 358–365, (2) 78–83
universal enveloping algebra 112–117, (2) 8
complexified 115, (2) 81
- unordered partition (2) 74

Cambridge University Press

978-0-521-47423-8 - Automorphic Representations and L-Functions for the General Linear Group, Volume I

Dorian Goldfeld and Joseph Hundley

Index

[More information](#)

550

unramified 260
 at p 406, (2) 42
 character 43
 local character 43
 representation 192, 202, (2) 53
 twists (2) 127
 unramified representation 192, 202,
 (2) 53
 admissible irreducible 400
 of $GL(n, \mathbb{Q}_p)$ and $GL(n, \mathbb{R})$ (2) 97
 upper half plane 78
 extended 78
 generalized (2) 1

vector valued automorphic function (2) 6

weak approximation 10
 weight k Laplacian 82, 176, (2) 7

Weyl
 element (2) 47, (2) 109
 group of $GL(n)$ (2) 17

Whittaker coefficient of an adelic cusp form
 (2) 33

Index

Whittaker function
 classical 84–87, 92
 global 128–134
 Jacquet's 86, 141–147, 195–200, 260–263
 local 195, 260
 relative to ψ (2) 57, (2) 86

Whittaker model
 of a local representation of $GL(n, \mathbb{R})$
 (2) 85–86
 of a local representation of $GL(n, \mathbb{Q}_p)$
 56–57
 of a principal series representation 213, 223

Whittaker's differential equation 84, 88

$Z(U(\mathfrak{g}))$ -finite 118

Zariski
 closed (2) 50
 open (2) 50

zeta function 418
 Riemann 46, 50

zeta integral, *see also* local zeta integral
 global 419–423, (2) 118

Zorn's lemma 192