

Index

- abundance, diversity, 114
 - and distribution, extinction probability, 119–20
 - and size, patterns, 77
 - variations in, 122
- accidents
 - causes of extinction, 112–17, 127
 - and interactions, 128
- Achatinella*, 89
- adaptations, key, 110, 312–16
- adaptive differences, 177–8
- adaptive radiation, driven by competitive speciation, 109–10
- Africa
 - large mammals, 137, 349
 - rainforests, species diversity, 279
- Alcelaphini*, extinction and speciation rates, 137
- algae
 - disturbance experiment, 37–8
- Allmon, Warren, 366
- Alocasta macrorrhiza*, 154–5
- alpha diversity, 33, 380
- Amazon, river basin
 - equal-area samples, 294
 - fish species, 265, 277
- Ambrosia*, ragweed, 167
- American chestnut, *Castanea dentata*, 123–5, 173
- amphibians
 - predator–victim ratios, 82–6
 - salamander growth, 313
- amphipods, old lakes, 180
- Anas*, ducks, 152–3
- Anatidorcas*, extinction and speciation rates, 137
- angiosperms
 - extinction rates, 138
 - spread, 313–16
 - see also plants
- Anolis sagrei*, lizard, 185
- Antarctic brachiopods, productivity, 45
- antbird
 - Gymnophithys*, 155
 - Hylophayx*, 155
 - Phaenostictus*, 155
- Antillean vertebrates, species–area curves, 15
- ants
 - army ant, *Eciton burchelli*, 155
 - Barro Colorado Island, Panama, 119–20
 - cherapachyine, 17–18
 - Connecticut ants, 20
 - New Guinea, 17–18, 260
 - Panama, 119–20
 - ponerine, 17–18
 - species diversity and latitudinal gradients, 291
- Apyceros*, extinction and speciation rates, 137
- archipelagos
 - geographical speciation, 89
 - isolation effects, 240
 - species–area curves, 14–22
 - z-values, 240–1
 - see also islands
- Arctic tundra
 - crustacea, 351, 352
 - copepods, 188
 - Daphnia*, 61, 187–8
 - floras, polyploid species, 338
 - Norway, 325
 - Wrangel Island, Russia, mammoths, 146
- area
 - competition, hares and rabbits, 234
 - and disturbance rates, 38–9, 344
 - energy–area hypothesis, 331–2
 - and geographical barriers, 288
 - hypothesis and evaluation, 369–71
 - insect diversity, 61
 - large-scale effects, 288–96
 - latitudinal gradients, 286–96
 - species diversity dynamics, 376–7
 - latitudinal gradients, 284–5, 286–96
 - Mercator projection, 284–5
 - population interactions on islands, 233–4
 - as primary regulator of species diversity, 381
 - and productivity, 369–70
 - scales, 378
 - species diversity, 376
 - size, lower extinction rate curves, 257–8
 - tropics vs northern lands, 284–5
 - see also species–area curves
- Arizona
 - deserts, rodents, 355
 - hummingbirds, 361
 - short grass patch, intermediate stage, 365
- Arrhenius equation, 12, 66, 67, 380
- point diversities, 380

- Artamus*, white-breasted woodswallow, 185
Artemisia monosperma, 172
 artifacts
 habitat diversity, 307–10
 sampling, 191
 Asilidae, robberflies, 155
 Atlantic Ocean
 biomes, 288
 megafaunas, 348
 molluscs, 292
 ostracodes, 58
Atriplex, 97–8, 154–5
 Audubon, J.J.
 on passenger pigeon, 122–3
 on writing, *quoted*, ii
 Australia
 Broken Hill, NSW, species–area curves, 33
 equal-area samples, 291
 Green Island, 245
 heathlands, nutrient enrichment, 347
 islands
 birds, 184, 356
 plants, 245, 250–2
 species–area curves, 15
 Lake Eyre, brine shrimps, 174
 mammals, 204–6
 Norfolk Island flora, 278
 Queensland, rainforest understory plants, 155
 South-Western Botanical Province kwongan, 29, 176, 187
 Tasmania, eucalypts, 47–8
 tropical aquatic species, seasonality, 71–2
 turtles, 192
 Western Australia
 birds, 184
 islands, plant extinctions and immigration, 250–2
 kwongan, 29
 wheatbelt, 204–9, 347
 Azores
 birds, 234
 ferns, 235–6
 Bahamas
 birds, 235
 energy–area hypothesis, 331–2
 orb-web spiders, 129
 snails, 217
 spider predation, 233
Balanus balanoides, barnacle, 152
 bamboo
 Pharus latifolius, 153–4
 Streptochaeta sodiroana, 153–4
Banksia hookeriana, 187
 barnacle
 Balanus balanoides, 152
 Chthamalus stellatus, 152
 disturbance experiment, 37–8
 barriers, *see* geographical speciation
 Barro Colorado Island, Panama, ants, 119–20
 bats
 North America, 293
 North and Central America, 26
 species–area curves and z-values, 264–5
 bees, Santa Catalina mountains, 360
 beetles, 89
 Bornean tropics, 74
 Lake Mälaren, 226
 St Helena, 215
 total numbers, 1–3
 benthos, *see* marine benthos
 beta diversity, 33
 biogeographical provinces, 191–2
 biological province, defined, 212, 264
 biomass, and productivity, 44–5
 biotas, types, *see* patterns
 biotic revolutions, 110
 birch forest, Finland, 326
 birds
 chaparral, compared, 16
 Darwin's finches, 89
 habitat and species diversity, 2, 35
 hummingbirds, 235, 361
 islands
 extinction rates, 139–40
 off Panama, 255–7
 see also named islands and archipelagos
 of:
 Åland, Finland, 218–24
 Azores and Madeiras, 234–5
 Bahamas, 235
 Britain, 69, 139, 294
 Guam, 231–2
 Hawaii, 230
 lowland tropics, 369
 Neotropics, species–area curves, 13, 14
 New Guinea, 185
 Ohio, 64, 69, 71
 Pacific archipelagos, 240–1
 Panama, 180–1
 Patagonia, 182–3
 Pennsylvania, species–area curve, 23
 Peru, 32
 Sweden, 193–5
 tropical rainforests, 356
 West Indies, 15
 Páramo, species–area curve, 20
 passenger pigeon, 122–3, 125–6
 seasonality, 69–72
 blue palo verde, *Cercidium*, 172, 173–4
 body size, 73–7
 and optimal foraging, 302
 trade-off principle, 155
Boiga irregularis, brown snake of Guam, 231–3
Bolitoglossa, salamander, 313
 Borneo
 beetles, 74
 species diversity, 2–3
 brachiopods
 Paleozoic, 59

- productivity, 45
- bracken fern, *see* ferns
- Bretsky, Ordovician benthos, 52, 307
- brine shrimp, *Parartemia*, 174
- Britain, birds
 - breeding, 69
 - island populations, 139
 - and Lepidoptera, equal-area samples, 294
- Britain, plants
 - species–area curves
 - Hertfordshire, 10
 - Surrey, 9
 - trees and insects, 267–8
- Brookhaven, NY, oak–pine forest, 63
- bryophytes, productivity, 43
- bryozoans, budding, 312
- Burns, R., *To a Mouse*, vii
- California
 - birds, 16
 - Ellwood Beach, 37
 - plants, 17
- Calligonum comosum*, 170
- Cambrian, 50
- Canada
 - Arctic tundra, crustacea, 351–2
 - islands
 - Devon Island, 326
 - ice bridges, 237
 - lakes
 - glacial lakes, 335
 - zooplankton, 42
- canonical log-normal distribution, 271, 379
- capture–recapture estimation, 141–3, 301
- fossils, 53–4
- carrying capacity, 100, 102
- Castanea dentata*, American chestnut, 123–5, 173
- catastrophes, *see* mass extinctions
- Caucasus Mountains, 48
- cave-dwelling species, 266–7
- censuses, repeated, maximum likelihood estimation, 138–41
- centrifugal speciation, 94, 288, 297
- Cercidium*, blue palo verde, 172, 173–4
- Challenger, data, 43
- Channel Island plants
 - ferns, 235–6
 - z-values, 17, 18
- chaos, 6, 320
 - defined, 305
 - and dynamics of diversity, 305
- chaparral, birds, species–area curves, 16
- characteristic equation, eigenvalues, 318
- chemical defense, 360
- Chenopodium album*, 173
- chestnut, American, 123–5
 - developmental switch, 173
- chestnut blight, *Endothia parasitica*, 123–5
- China
 - chromosomal variants, mammals, 46–7
 - Père David deer, 112
- chipmunk, 360
- Chondrus crispus*, alga, 342–3
- chromosomal variants, mammals, seismic activity, 46–7
- chronoseres, succession, 62–3
- Chthamalus stellatus*, barnacle, 152
- cichlids, old lakes, 180
- clades, 298
 - extinction rates, 313
- cladocerans, 61, 187–8
- clonal plants, 166–7
- Cocos island, 89, 109
- coevolution, model, 311
- competition
 - apparent, 358
 - assembly rules, 229
 - contest, 360
 - equations, 374
 - and habitat selection, 162–6
 - interference, hypothesis and evaluation, 359
 - and morphology, 230
 - optimal habitat selection, 368
 - positive/negative feedback, 303
 - scramble, 360
- competitive displacement, 346
- competitive exclusion, 126, 230
- competitive extinction, 229
- competitive speciation, 97–105, 109–13
 - evidence for, 105–7, 109–13
 - models, 97–105
- competitive structure
 - changes, hypothesis and evaluation, 359–61
 - and productivity, 359–61
- competitive subordination, 346
- competitor species
 - addition, 162–6
 - intertaxon, and productivity, 358–9
- Congo, river basin, fish species, 265, 277
- Connechaetus*, extinction and speciation rates, 137
- Connecticut ants, species–area curve, 20
- conservation ecology, 353
- continental drift
 - effect on species–area curves, 279–84
 - fossil evidence, 280–4
 - interprovincial curves, 282–4
- copepods
 - Arctic tundra, 188
 - old lakes, 180
- Cordyline rubra*, 154–5
- Cormack method, 140
- corvids, diversity and body size, 75–7
- Costus scaber*, 121
- covariance in population densities, 367–8
 - hypothesis, 370–1
- crayfish, *Orconectes*, 156
- Cretaceous–Tertiary mass extinction, 37, 110, 147
- crustacea

- Arctic tundra, 61, 187–8, 351–2
- productivity, 44–5
- cumaceans, productivity, 44
- Cuscuta*, dodder, 167–8
- Damaliscus*, extinction and speciation rates, 137
- Daphnia*, Arctic tundra, 61, 187–8
- Darwin's finches, 89
- De Candolle, 375
- decapods
 - productivity, 44–5
 - unimodal pattern of diversity, 44
- deer, China, 112
- density-dependent fitness, 99–100, 115, 160–3
- density-dependent optimal foraging, 367–8
- Déscartes, *quoted*, 300
- deserts
 - polar, 327
 - rodents, 348–9, 361
 - Arizona, 355
 - Israel, 355
 - trophic levels, 321–2
- detritivory, 335–6
- Dicoelostia*, 59
- differential equations, 374–6
 - immigration and extinction, 375
 - species diversity dynamics, 374–6
 - species–area curves, 376
- dimensionality hypothesis, 330
- dimensionless patterns, 5, 73–86
 - body size, 73–7
 - food chains and webs, 77–86
- Dipodomys microps*, 302
- diseases, measles, 228
- dispersal ability, and immigration rate, 234–8
- disruptive selection, 101
- distributions
 - ideal free, 160–2
 - see also* log-normal distribution
- disturbance rates, 36–9
 - experiment, 37–8
 - hypothesis and evaluation, 362–7
 - collateral predictions, 364–5
 - intermediate disturbance frequencies, and
 - species diversity, 341–4
 - intermediate disturbance levels, 36–9
 - marine and terrestrial productivity, 363
 - and non-equilibrium systems, 341–4
 - and productivity, 362–7
 - scale, 37
 - species–area curves, 344
 - z-values, 344
- divergence rates, and isolate formation, 92–9
- dodder, *Cuscuta*, 167–8
- doomed species, extinction, 145–6
- dragonflies, dimensionless patterns, 73
- Drosophila*, 89, 105–6
- ducks, *Anas*, 152–3
- dynamic game theory, 166
- dynamic hypothesis, 327, 335
- dynamic instability
 - delicate balance, 319
 - hypothesis and evaluation, 357
 - and productivity, 357
- dynamic stability, 317–20
 - defined, 317–18
 - linearized stability analysis, 318
 - Pimm and Lawton's hypothesis, 317
 - qualitative concept, 319
- earthquakes, 46
- echinoderms, productivity, 44–5
- Ecton burchelli*, army ant, 155
- ecological opportunity, 100, 101
- ecological techniques, palaeobiological patterns,
 - 55–72
- ecological time, 62–72
- ecological vicars, 192
- Ectopistes migratorius*, passenger pigeon, 122–6
- ectoprocts, Pacific, 290
- eigenvalues, 318, 319, 320
- Einstein, and Newton, 374–5
- El Niño droughts, 120
- Elephas*, extinction and speciation rates, 137
- Ellobius talpinus*, 46–7
- Ellobius tancrei*, 46–7
- Elton–Hutchinson theory, thermodynamics,
 - 375
- Endler's clinal model, founder effects, 107
- Endothia parasitica*, chestnut blight, 123–5
- energy, second law of thermodynamics, 320–1
- energy flow and diversity, 345–72
 - global phase, 370–1
 - productivity, experimental increase, 345–7
 - prospects, 371–2
 - unimodal pattern
 - intertaxonomic, 350–70
 - regions, 348–50
- energy–area hypothesis, 331–2
- Enteromorpha*, 342–3
- environmental heterogeneity, and productivity,
 - 354–7
- environmental perturbations, 120
- environmental variance, effects on habitat diversity, 380
- Eocene mammals
 - diversity estimates, 56, 142
 - extinction rates, 142
 - and speciation, 301
- Eocene molluscs, recovery from mass extinction,
 - 149–50
- ephemerals, 170
- epochal steady states, 52–60
- equal-area samples
 - Amazon basin, 294
 - Australia, 291
 - birds and Lepidoptera, Britain, 294
- equitability, 201
- Eriophyllum ambiguum*, 171
- eucalypts, *Eucalyptus* spp.

- hybrid zones, Tasmania, 47–8
- jarrah, 173
- eurytopism, defined, 99
- evapotranspiration, 364
- evolution
 - habitat diversity, 175–89
 - new genes, 94
- evolutionary compromise, 311
- evolutionary deadlock, 311
- evolutionary mutualists, 303
- Evolutionary Stable Strategy, trade-off, 311
- evolutionary time, 50–62
 - species–area curves, 264–96
 - time scale, 138
- excess penetrance, 104–5
- exchange of species, 71–2
- experimental increase of productivity, 345–7
- explanations, 6–7
- extinction, 112–50
 - of abundant, widespread species, 122–6
 - causes
 - accidents, 112–17
 - competitive extinction, 229
 - population interactions, 113, 229–34
 - population size, 114–17, 217–28
 - by predation, 126–7, 231–2
 - doomed species, 145–6
 - and geographical range, 114–21, 118–22
 - logarithmic process, 135–7
 - population size, 317
 - probabilities, species–island combinations, 115
 - rates, 299
 - abundance and distribution, 118–20
 - balance with speciation, 300–1
 - capture–recapture estimation, 141–3
 - curvature, 129, 250
 - effect of number of provinces, 279–84
 - equilibrium, 214
 - formula, 136–8
 - and immigration, 213
 - vs immigration rate, 138–41
 - islands, 130–5, 143–5, 250–2
 - latitudinal gradients, 339
 - lower, larger areas, 257–8
 - maximum likelihood estimation, repeated censuses, 138–41
 - measurement, 135–45
 - size of islands, 131–5
 - and species diversity, 128–35
 - and variation in abundance, 122
 - widespread, abundant species, 122–5
 - reptiles on islands of known age, 143–5
 - see also* mass extinction
- Falco concolor*, 174
- feedback, *xiii*, 263, 297, 299, 303, 375–6
- ferns
 - Azores and Channel Islands, 235–6
 - constraints in growth, 314
 - fern-eating arthropods, 236–7, 261, 266
 - productivity, 43–4
 - Pteridium* (bracken fern), 237
- Fiji, birds, 185
- Filago californica*, 171
- Finland
 - Åland birds, 218–24
 - birch forest, 326
- fire, 172
- fish species
 - Amazon and Congo river basins, 265
 - coastal fishes, latitudinal gradients, 28
 - numbers, 109
 - old lakes, 180
 - productivity, 44–5
- Fisher's α
 - effect of abundance, 202
 - equitability, 201
 - log-series distribution, 193
 - QBASIC program, 194
 - sample size, 202, 218
 - species diversity, 193–8, 203
- fitness
 - density-dependent, 99–100, 163
 - frequency-dependent, 99
 - and optimization theory, 157–8
 - tolerance, 163
 - Wrightian surface, 100
- flat laxity, 330
- flies, 113
- Florida
 - Keys, invertebrates, 426
 - mangrove islands, 245
- food chains and webs
 - deserts, 320–3, 327
 - insects, 350
 - length, 320–32
 - and second law of thermodynamics, 320–7
 - and dynamical instability, 318–20, 327
 - return times, 328, 334
 - testing hypotheses, 380
 - testing major hypotheses, 380
 - total energy, 331–2
 - and trophic levels, 77–81, 325–38
- omnivores, 81–2, 332–6
- patterns, 77–86
 - diversity and higher trophic levels, 77–81
 - omnivory, 81–2
 - predator–victim ratios, 82–6, 336–8
- population dynamics, 317–44
- see also* trophic levels
- Foraminifera, marine, latitudinal gradient, 29
- fossils
 - biases in record, 307–10
 - capture–recapture estimation, 53–4
 - 'jack-knifing' technique, 54–5
 - see also* paleobiological patterns
- founder effects
 - bottlenecks, isolates, 93
 - stasipatric speciation, 107
- Fouquieria*, ocotillo, 173

- fractal landscape, 189
- France, Montpellierais region, 63
- freshwater species
 - predator–victim ratios, 84
 - seasonality, 71–2
- frogs, *see* amphibians
- fungi
 - diversity, 2
 - eucalypt hybrid zones, 46–7
- fynbos, South Africa, 29, 176–8, 289
 - species diversity, 31
- Γ statistic (R_N/R_{MAX}), 272
 - range of values, 273
- Galapagos archipelago, 89, 109
 - isolation, 279
- gall formers, 78
- game theory, dynamic, 166
- gamma diversity, 33
- gastropods
 - extinction and length of life, 119
 - Lake Tanganyika, 179
 - old lakes, 180
 - productivity, 44
 - species flocks, 178
 - versatility, 312
- Gatun Lake, Panama, 255–7
- Gause's principle, niches, 127
- genetic drift, 113, 115
- geographical barriers, 90–2
- geographical range, 90–2
 - and extinction, 114–21, 118–22
 - and speciation, 90–2
- geographical speciation, 87–96, 108–9, 297–305
 - evidence for, 108–9
 - within more than one trophic level, 301–5
 - within one trophic level, 297–301
- Geospizinae (Darwin's finches), 89
- Gerbillus andersoni*, 158–9, 165
- Gerbillus pyramidum*, 158–9, 165
- germination, 168–73
 - activity, adaptations in plants, 169–74
- Giraffa*, extinction and speciation rates, 137
- giving up density (GUD), 165
- glacial lakes, Canada, 335
- Gleason semi-logarithmic plots, 11, 198, 209
 - and equation, 64–8
- Glechoma*, mint, 167
- Gobi desert, Mongolia, rodents, 348–9
- golden whistler, *Pachycephala*, 185
- Goldilocks contest, 105
- Gough Island, 185
- granivores, 359
- Greenland, polyploidy, 108
- Guam
 - birds, 231–2
 - brown snake, 231–3
- GUD (giving up density), 165
- guided multiple linear regression, 377
- guild, defined, 161
- Gymnophithys*, antbird, 155
- gymnosperms, 314
- habitat
 - predictability, 171
 - specialization, 383
 - use, 157–75
 - variance, vs phenotypic variance, 101
- habitat diversity, 32–6
 - colonization of new habitats, 306–7
 - effect on species–area curves, 190–210
 - environmental variance effects, 380
 - evolution, 175–89
 - coevolution with species diversity, 151–89
 - hypothesis and evaluation, 354–7
 - increase, paleobiological patterns, 306–16
 - information theory, 34
 - new habitats, 306–7
 - niche space exploration, 310–16
 - and population size, 217–18
 - two-dimensional vs three-dimensional, 330–1
- Haila's islands, 256–7
- Halcyon*, kingfisher, 185
- Haldane, J.B.S., *quoted*, 1
- hare
 - competition and area, 233–4
 - extinctions, 258
- Hawaii
 - adaptive radiations, 89, 178
 - bird introductions, 230
 - snails, 178
- heathlands
 - nutrient enrichment, 347
 - Western Australia, 29
 - see also* fynbos; kwongan
- helminths, species diversities, 61–2
- Hesperodaptomus arcticus*, copepod, 188
- high-risk species, 122
- Homoptera, seasonality, 70
- honeycreepers, 89
- host age relationship, 60–2
- host race formation, 98
- hummingbirds
 - and flower mites, 120–1
 - Arizona, 361
 - West Indies, 235
- hybrid zones, species diversity, 47–8
- Hylophyax*, antbird, 155
- Hypomocoma*, 89
- ice bridges, colonization of islands, 215, 237
- ideal free distributions, 160–2
 - assumptions, 161
 - samples, 160
- immigration, supertramps, 215
- immigration rate
 - and dispersal ability, 234–8
 - and extinction rate, 214
 - vs extinction rate, 138–41
 - islands of W. Australia, 250–2
 - and isolation, 234

- and speciation, 110–11
 - taxon-specific, 215
- inactivation process, mutation rates, 94–6
- independent variables, reduction, 88
- indices of species diversity, 199–200
 - see also* Fisher's α ; Simpson's index
- individuals curve, defined, 270
- Indonesia, Sunda Islands
 - mammals, 257–8
 - relict species, 130–5
- information theory, habitat variety, 34
- insecticides, knockdown, 2, 32
- insects
 - British trees, 267–8
 - diversity, 2–3
 - indices of diversity, 200
 - standing diversity, 377
 - eucalypt hybrid zones, 47–8
 - fern-eating, 236–7, 261, 266
 - food chains and webs, 350
 - host age relationship, 60–2
 - tropical, seasonality, 70
- intensive variables, 6
- interactions, and accidents, 128
- intercept value, 12
- interference competition, 359–60
- interprovincial curves
 - continental drift, 282–4
 - species–area curves, 276
 - time scales, 277
 - z-values, 282
- intertaxonomic competition, 371–2
 - hypothesis and evaluation, 358–9
- intertaxonomic patterns, productivity–diversity, 350–1
- islands
 - biogeographical island, defined, 19, 211
 - continental, 130
 - defined, 19
 - distance effect, 234–43
 - effect on habitat use, 180–6
 - endemism, 109
 - equilibria, 243–4
 - away from equilibrium, 248–50
 - extinction rates, 130–5
 - curves, 250
 - and immigration, 250–2, 257
 - nine hypotheses, 133
 - reptiles, 143–5
 - immigration, 211, 250–2, 257
 - islands theory, 212–63
 - isolation *see* islands, distance effect
 - land-bridge, 130
 - large/small, net extinction rate, 254–5
 - mountaintop, 130–1
 - non-equilibrial, 248–50
 - non-interactive dynamics, 214
 - patterns, 210–12
 - species–area curves, 210–63
 - population interactions
 - competition, 229–31
 - interactions and area, 233–4
 - predation, 231–3
 - population size, and habitat diversity, 217–28
 - rabbit introductions, 233–4
 - relaxation time, 426
 - saturation, 239
 - size, and extinction rate, 131–5
 - species diversity
 - compared with mainland, 180–6
 - and net turnover, 252–3
 - self-regulation, 214
 - steady state, 246–8
 - summary, 263
 - theory of island biogeography, 128–35, 212–17
 - 'contradictions' in real turnover rates, 250–8
 - 'contradictions' in z-values, 241, 262–3
 - turnover on islands, 243–6
 - contradictions, 250–8
 - z-values, 258–63
 - time scale, 259
 - see also* archipelagos
- isoclines, and steady state, 303–4
- isolates
 - centrifugal speciation, 94
 - formation, 90–2
 - and divergence rates, 92–9
 - founder-effect bottleneck, 93
 - rates, 88
- isolation, *see* islands; distance effect
- isoleg theory, 163–6
- Israel
 - deserts, rodents, 355
 - Wilderness of Paran, 321
- Jack pine, 172
- 'jack-knifing' technique, fossils, 54
- jarrah, *Eucalyptus*, 173
- Johnson *et al.*, model of competitive speciation, 103–5
- Jolly–Seber model, populations, 56, 142
- karyotypes, 46–9, 96–7, 107–8, 338–41
- Kaspari, El Niño droughts, 120
- Kemp, C.D., SI as a statistic, 199
- key adaptations, 110, 312–16
- kingfisher, 91, 185
- knife barriers, geographical speciation, 90–1
- knockdown insecticides, 2, 32
- Kobus*, extinction and speciation rates, 137
- kwongan, South Western Australia, 29, 176–8
- Lake Baikal, 179–80
 - isolation, 279
- Lake Gatun, Panama, 255–7
- Lake Huron, colonization of islands, 215, 237
- Lake Mälaren, snails, 226
- Lake Michigan, ice bridges, 237

- Lake Tanganyika, 179–80
 lakes, species flocks, 179–80
Lates niloticus, Nile perch, 126
 latitudinal gradients, *ii*, 25–32
 body size and diversity, 73–7
 defined, 4
 large-scale effects of area, 286–96
 and polyploid species, 48–9, 338–9
 productivity, 379
 scale, 294
 speciation, 294, 339
 Terborgh's pattern, 284–6
 Law of Large Numbers, theory of log-normal distribution, 268
 leafhoppers
 extinction risk, 118
 seasonality, 70
 Lepidoptera
 Britain, 294
 cumulative diversity, 65–6
 diversity, cycle, 65, 70
 moths, species diversity, 64–7
 Rothamsted, 193, 195–7
Lepus, hare, 233–4, 258
 Lethbridge, moths, 64–5, 67
 Levings and Wheeler, El Niño droughts, 120
 Liapunov, 318
 life history evolution, 157–8
 lighthouse keeper's cat, 128, 146, 381
 limiting similarity, niches, 6, 127–8, 186–9
 limpet, *Lottia alveus*, 121
Littorina littorea, sea snail, 342
 lizard
 Anolis sagrei, 185
 and orb-web spiders, 129
 see also reptiles
 log-normal distribution
 abundances, 352
 canonical, 271, 379
 theory, Law of Large Numbers, 268–9
 log-series distribution, Fisher's α , 193
 long-term increase in diversity, 306–16
 paleobiological artifacts, 307–19
 lorikeet, *Vini vidivici*, 233
Lottia alveus, limpet, 121
Loxodonta, extinction and speciation rates, 137
 Lyell, fossils, 112

M (index of diversity), 200
 MacArthur, R.H.
 (dedication), *iv*
 on models, 6, 32–5, 73–4, 128, 157, 181–2, 186, 212–17, 243, 247, 263, 348, 375
 MacArthur–Wilson hypothesis, islands, extinction rates, 133
 Madeira Islands, birds, 234–5
 Maine, mammals, 426
 mainlands, defined, 212
 maize, selection, 105
 Malaysia, Sunda Islands, 130–5, 257–8

 Malesian rainforest, 30–2
 mammals
 chromosomal variants, seismic activity, 46–7
 diversity, 2
 Eocene, 56
 herbivorous, Africa, 349
 Indonesia, Sunda Islands, 257–8
 mountaintops, 249–50
 North America, 294
 fossils, large, 57
 quadrupedal, latitudinal gradients, North and Central America, 26
 southeast Australia, 204–6
 teeth, 313
 z-values, 280, 283
 mammoths
 extinction, 145
 Wrangel Island, Russia, 146
 mangrove islands, 227, 245
 marine benthos
 Antarctic brachiopods, productivity, 45
 megafaunas, 348
 Ordovician, 52
 Paleozoic, 362
 productivity, 43–4
 biomass, 44–5
 sediment type, and ocean depth, 362
 marine invertebrates, 36, 51, 309
 epifaunal, 36
 Florida Keys, 426
 Phanerozoic time, 51, 309
 marine species, ranges, 91
 mass effect, 259
 species, 376
 mass extinctions, 146–50
 causes, 149–50
 Cretaceous–Tertiary mass extinction, 37, 110, 147
 effect of number of provinces, 279–84
 intervals between, 147, 305
 Maastrichtian, 150
 periodicity, 147, 305
 probabilities, 147
 see also extinction
 mating
 mating preference genes, 103
 random mating, 101–2
 Mauritius, species diversity measures, 196–7
 measles, diseases, 228
 Mercator projection, 284–5
 mesquite, *Prosopis*, 174
 metabolism, time retreat, 175
Metapeira datona, spider, 233
Metrosideros polymorpha, 89
 Mexico, Sea of Cortez, reptile extinction rates, 143, 378
 Michigan plants, species–area curve, 11
 migratory species, 174–5
 minimum-spanning tree, 230, 232
 Minnesota, old fields, 62–3

- mint, *Glechoma*, 167
- Miocene, mammal extinctions, 145–6
- mites
 - flower, hummingbirds, 120–1
 - soil, species diversity, 35
 - species diversity and latitudinal gradients, 291–2
- models, simplicity, 6
- molluscs
 - Eocene, 56, 142
 - extinction rates, 118, 138
 - freshwater, species diversity, 34
 - mass extinction, 149–50
 - Pacific, 290
- Mongolia, Gobi desert, rodents, 348–9
- mongooses
 - diversity and body size, 75–6
 - lizard predation, 232
 - predation on lizards, 129
- monocotyledons, extinction rates, 138
- Monte Carlo computer simulation, 81–2
- moths
 - diversity, 65
 - Lethbridge, 64–5, 67
- ‘mountaintop islands’, 130
- mountaintops
 - mammals, 249–50
 - plants, 250
- mouse
 - Mus musculus*, 47
 - Peromyscus*, 93, 93–4, 288
- mouse, addressed by Robert Burns, vii
- multiple linear fishing expedition, 377
- multiple linear regression, non-independent variables, 208–10
- mutation rates
 - inactivation process, 94–6
 - population size, 95
- mutualism, evolutionary mutualists, 303, 304
- Namib Desert, trophic levels, 322
- Neotoma, Ohio, birds, 64, 69
- nested design, species–area curves, 10
- Netherlands, Wadden Sea islands, 234, 258
- New Guinea
 - ants, 17–18, 260
 - birds, 185
 - island colonizations, 237
 - z-values, 240–1
- new habitats, 306–7
- New Mexico, pinyon–juniper woodland, 209–10
- New Zealand
 - archipelagos, 279
 - land snails, 178–9
- Newton, and Einstein, 374–5
- niches
 - breadth, coevolution, 157–66
 - disappearance, 120–1
 - exploration, 310–16
 - Gause’s principle, 127
 - limiting similarity, 6, 127–8
 - overlap, isolog theory, 164
 - species closeness, 186–9
- Nicolet River valley, Quebec, Ordovician invertebrates, 52, 68–9, 109–11, 306–7
 - increase in diversity, 306–7
 - ‘jack-knifing’ technique, 54–5
- Nile perch, *Lates niloticus*, 126
- non-independent variables, multiple linear regression, 208–10
- non-linear regression technique, 12
- non-linearity, species–area relationship, 30
- Norfolk Island, 278
- North American mammals, 294
 - large fossils, 57
 - mass extinctions, 145–6
- North Atlantic megafaunas, 348
- North Atlantic ostracodes, 58
- North Pacific Gyre, 281
- Norway, Arctic, 325
- Nothofagus*, southern beech, 182–4
- nutrient enrichment, experimental, 39, 346
- nutrients, and productivity, 39–40
- oak, leaf miners, 61
- oak–pine forest, Brookhaven, NY, 63
- Oceania, ants, 17–18
- oceans
 - depth, and sediment type, 362
 - provinces, 281
 - see also marine benthos; named oceans
- ocotillo, *Fouqueria*, 173
- Odynerus*, 89
- Ohio, birds, breeding, 71
- old fields, Minnesota, 62–3
- omnivory, 81–2, 332–6
 - life-history omnivory, 335
- optimal foraging, 158–9, 302, 375
 - density-dependent, 367–8
- optimal habitat use, 157–75
 - effect of islands on, 180–6
 - ideal free distribution, 160–2
 - in the presence of competitors, 162–6
 - by plants, 166–74
 - seasonality, 174–5
 - seed germination, 169–73
- optimization
 - migratory species, 174–5
 - optimal behavior, 160
 - theory, and fitness, 157–8
- orb-web spiders, 116, 122, 129
- Orconectes*, crayfish, 156
- Ordovician invertebrates, 138, 306
 - increase, 310
 - ‘jack-knifing’ technique, 54–5
 - Nicolet River valley, Quebec, 52–5, 68–9, 109–11, 306–7
- origination rate, 376
- Oryctolagus cuniculus*, rabbit, 233–4, 258
- ostracodes

- North Atlantic, 58
- old lakes, 180
- trade-off principle, 152
- Pachycephala*, golden whistler, 185
- Pacific archipelagos
 - birds, 240–1
 - non-linear regression technique, 12
- Pacific islands, lizard predation, 232
- Pacific Ocean
 - biomes, 287–8
 - North Pacific Gyre, 281
- paleobiological patterns, 50–60, 297–316
 - epochal steady state, 50–60, 297–305
 - extinction and range size, 118–20
 - increase of diversity, 306–16
 - artifacts, 307–10
 - new habitats, 306–7
 - niche space, 310–16
 - pull of recent, 308–10
 - number of provinces, 279–84
 - productivity–diversity, 28–9, 363, 365–6
 - trends in speciation, 109–11
- Paleozoic
 - brachiopods, 59
 - marine benthos, 362
- Pamiro-Alay Mountains, mammal chromosomal variants, 46–7
- Panama
 - ants, 119–20
 - birds, 180–1, 356
 - island birds, 255–7
 - isthmus as land link, 280
- Pangaea, and Permo–Triassic catastrophe, 281, 283
- Paramecium*, 113
- Páramo islands, Andes
 - birds, 241–3
 - species–area curve, 20
 - defined, 241–2
 - z-values, 242
- Paran, Wilderness of, Israel, 321
- Parartemia*, brine shrimp, 174
- parasitoid systems, 334
- Partulina*, 89
- passenger pigeon, *Ectopistes migratorius*, 122–6
- Patagonia, birds, 182–3
- patterns
 - defined, 4–5
 - dimensionless, 73–86
 - food chains and webs, 77–86
 - four types:
 - among biotas, 24–5
 - archipelagos, 14–22
 - large pieces of a biota, 9–14
 - tiny pieces of a biota, 22–3
 - islands, 210–12
 - paleobiological, 297–316
 - population dynamics and food chains and webs, 317–44
 - Preston's temporal pattern, 63–8
 - processes and explanations, 5–7
 - regional, 40–5
 - small-scale, 39–40
 - species–area curves
 - island patterns, 210–63
 - mainland patterns, 190–210
 - in time, 50–72
- Pectocarya*, 171
- pelagic species, 290
- Pennsylvania, birds, species–area curve, 23
- Père David's deer, 112
- Permo–Triassic catastrophe, Pangaea, 281, 283
- Peromyscus* (mice), 93–4, 288
- perturbation analysis, 149, 318
- Peru, birds, 32
- Phacochoerus*, extinction and speciation rates, 137
- Phaenostictus*, antbird, 155
- Phanerozoic time, 50–2
 - marine invertebrates and plants, 51
- Pharus latifolius*, bamboo, 153–4
- phenotype cartel breakup, speciation, 156–7
- phenotypic dissociations, 312
- phenotypic variance
 - vs habitat variance, 101
 - and random mating, 101–2
- photosynthetic capacity, and efficiency, trade-off, 154
- Piedmont, diversity, 63
- Pimm–Lawton hypothesis
 - dynamical stability, 317, 327–33, 375
 - omnivores, 332–3
 - parasitoid systems, 335
- pine
 - Jack pine, 172
 - Rhus ovata*, 172
- pinyon–juniper woodland, Utah and New Mexico, 209–10
- Plagithmysus*, 89
- plankton, 118
 - Canadian lakes, 42
- Plantago patagonica*, 171
- plants
 - angiosperms and gymnosperms, 314
 - Broken Hill, species–area curve, 23
 - Channel Islands, z-values, 17, 18
 - clonal, ramets, 166–7
 - desert, germination, 169–74
 - extinction rates, 138
 - habitat selection, 166–74
 - islands
 - W. Australia, 250–2
 - z-values, 242–3
 - land, first appearance, 313–16
 - latitudinal gradients, 25–32, 48–9
 - mountaintops, 250
 - opportunism, 172–4
 - Phanerozoic species diversity, 51
 - phenotype variation, 167

- pollination strategies, 168–9
- polyploid species, 48–9
- rootless stages, 168–9
- seeds, 168–73
 - germination activity, 169–74, 314–16
 - number and size, 314–16
- sex strategy, 167
- species diversity, and productivity, 40–4
- spread of angiosperms, 313–16
- time specialization, examples, 168–74
- trade-off principle, 153
- various, species–area curves, 24
- vine habit, 167–8
- see also* trees
- Pleistocene, marine benthos, 362
- Pliocene, mammal extinctions, 145–6
- point diversities, 33
 - measuring, 380
- pollen dating, trees, 60
- pollination, 168
- Pollock's robust design, 142
- polychaetes, productivity, 44
- polyploid species, 338–41
 - allopolyploidy, 96
 - and altitude, 48–9
 - Arctic floras, 338
 - autopolyploidy, 96
 - evidence for, 96–7, 108
 - higher orders, 97, 339–40
 - latitudinal gradients, 48–9, 338–9
 - prevalence, 108
 - rates of polyploidy, 299
 - relative species diversity, 48–9
- ponds, 187–8
- population densities
 - covariance in, 367–8
 - and productivity, 367–8
 - minimum viable, 321, 352
- populations
 - closed [*footnote*], 53
 - dynamics, and food chains and webs, 317–44
 - interactions, 113
 - causes of extinction, 113, 229–34
 - role, 125–35
 - Jolly–Seber model, 56
 - size
 - causes of extinction, 114–17, 217–28
 - and habitat diversity, 217–28
- predation
 - puzzles, 374
 - theories, 127
- predator–victim ratios, 82–6, 336–8
 - changes
 - hypothesis and evaluation, 357–8
 - and productivity, 357–8
 - constancy, 82–6
 - destabilization, 346
 - feasible community, 336
 - freshwater species, 84
 - mammals, 85
 - vertebrate/insect systems, 333
- prediction and testing in science, 334–5
- Prestonian theory
 - canonical log-normal analyses of species
 - abundance, *ii*
 - extension of theory, 352
 - temporal pattern, species–time curve, 63–8
 - veil line, 352
- productivity
 - and area, 369–70
 - and competitive structure, 359–61
 - defined, 39
 - and disturbance, 362–7
 - and environmental heterogeneity, 354–7
 - experimental increase, 345–7
 - high, consequences, 330
 - indexes of, 39
 - of:
 - competitor species, intertaxon, 358–9
 - decapods, 44–5
 - ferns, 43–4
 - gastropods, 44
 - protobranchs, 44
- patterns, 4–5
 - decrease phase hypotheses, 354–70
 - dynamical instability, 357
 - increase phase, 351–3
 - latitudinal gradients, 379
 - regional, 40–5
 - small-scale, 39–40
 - unimodal in regions, 348–70
- productive space hypothesis, 331
- reduction in covariance of population densities, 367–8
- species diversity, 348
 - decline, 379
 - plants, 40–4
 - prospects, 371–2
 - and time, 361–2
- productivity–diversity
 - biogeographical provinces, 370–1
 - decrease phase, 354–70
 - global scale, 370–1
 - increase phase, 351–3
 - intertaxonomic patterns, 350–1
 - regional patterns, 348–50
- profits from loss, 166
- propagules
 - as bottlenecks, 93
 - defined, 88, 212–13
 - mammals, 429
 - sizes, 216
- Prosopis*, mesquite, 174
- Proterrhinus*, 89
- protobranchs, productivity, 44
- province
 - defined, 212, 264
 - number and diversity, 279–84
 - see also* biogeographical provinces
- Pteridium* (bracken fern), 237

- puzzles, 373–84
- pyramid of numbers, 114
- qualitative stability, 318
- quantum speciation, 110
- Quebec, *see* Nicolet River valley, Quebec
- rabbits
 - extinctions, 258
 - introduction to islands, 233–4
- ragweed, *Ambrosia*, 167
- Rahbek
 - species–area curves, 13, 14
 - productivity–diversity in Neotropical birds, 369–70
- random mating, and phenotypic variance, 101–2
- rare species, 382–3
- refraction, 192–3
 - Hart and Horowitz, 194
- rat, Polynesia, 233
- Red Queen principle, defined, 311
- relict species, Indonesia, Sunda Islands, 130–5
- reptiles
 - extinction rates
 - islands of known age, 143–5
 - Mexico, Sea of Cortez, 143, 378
 - latitudinal gradients, 27
 - West Indies, 15
- rescue effect, defined, 259
- reserves, subdivisions, 383
- resource–density assumption, IFD, 161
- retreat, metabolic, 175
- return times, length, in food chains and webs, 328, 334, 357
- Rhagoletis* (fruit flies), 106
- Rhinoseius*, hummingbird flower mites, 120–1
- Rhus ovata*, 172
- robberflies, Asilidae, 155, 321
- rodents
 - chromosomal variation, seismic activity, 46–7
 - deserts, 361
 - Arizona, 355
 - Gobi desert, Mongolia, 348–9
 - Texas, 41
- Rothamsted, Lepidoptera, 193, 195–7
 - cumulative diversity, 65–6
- Rowettia*, Gough Island bunting, 185–6
- Russia, Wrangel Island, mammoths, 146
- St Helena, beetles, 215
- St Lawrence River, islands, 237
- salamander, *Bolitoglossa*, 313
- salinity gradient, ponds, 188
- sample size
 - equal-area samples, 378
 - Fisher's α , 202, 218
 - Simpson's index, 199, 218
 - and species diversity, 190–2
 - species–area curves, shape generated by larger and samples, 202–3
- scale(s)
 - dependence of z and c , 21
 - disturbance rates, 37
 - species–area curves, 5, 268
- Schismus barbatus*, 171
- Schizoporella floridana*, 312
- Sea of Cortez, reptile extinction rates, 143, 378
- seasonal variation, 68–72
- seasonality, 174–5
- second law of thermodynamics, 320–1, 320–2
- second species effects, 165
- secondary loss, 384
- seeds and germination, 168–73, 314–16
 - see also* plants
- seismically active regions, chromosomal variants in mammals, 46–7
- self-regulation, species diversity in islands, 214
- sex strategy, plants, 167
- Shannon–Weiner index of diversity, 59, 145
- shared preferences, tolerance differences, 368
- sharks, in food chains and webs, 79
- Sierola*, 89
- Silurian brachiopods, 59
- Silurian terrestrial life, 306
 - plants, 51, 313
- Simpson's index
 - sample size, 218
 - species diversity, 198–202
- simulations
 - model of competitive speciation, 103–5
 - Monte Carlo, 81–2
- sink species, 211, 260, 376, 381
- size of body, 73–7, 155, 302
- Skokholm Island, Wales, 426
- SLOSS controversy, 382
- Smallwood, guild studies, 161–2
- snails, 178–9
 - Bahamas, 217
 - Hawaii, 178
 - Lake Mälaren, 226
 - New Zealand, 178
 - sea, *Littorina littorea*, 342–3
- snakes
 - bolyerine, 313
 - brown snake, *Boiga irregularis*, 231–3
 - pit vipers, 312–13
 - see also* reptiles
- source species, 211, 260
- South Africa
 - Breede River fynbos, 176
 - Cape Floristic Region, 176
 - fynbos, 29, 176–8
 - rainforest, species–area relationship, 29–30
- South American birds
 - biomes, species–area curves, 13
 - Páramo islands, 241–3
- southern beech forest, 182–4
- speciation, 87–111
 - centrifugal speciation, 94
 - cladistic methods of modeling, 94

- competitive speciation, 97–108, 298
 - and adaptive radiation, 109–10
- divergence after isolate formation, 92–6
- ecological opportunity, 298
- geographical speciation, 87–96, 108–9, 297–305
 - and immigration, 110–11
 - latitudinal gradients, 284, 339
 - modes, importance, 107–10
 - phenotype cartel breakup, 156–7
 - polyploidy, 96–7
 - rate, 298
 - and productivity, 375
 - stasipatric, 107
 - sympatric, predatory insects, 106–7
 - without geographical barriers, 103–5
 - see also* polyploidy
- species, total numbers, 1–3
- species–area curves, 8–25, 190–263
 - among biotas, 23–4, 204, 264–8, 277–8, 297–305
 - among islands of one archipelago, 14, 210–63
 - among large pieces of biota, 9, 190–210
 - among tiny pieces of a biota, 22, 192–203, 279
 - differential equation, 376
 - disturbance pattern, 344
 - in ecological time, 190–263
 - effect of continental drift, 279–84
 - evolutionary time, 264–96
 - first, 9
 - four patterns, 5, 8, 190–2
 - graph, combined, 276–9
 - habitat hypothesis, 191, 204–5
 - how to build, 9–25
 - intercept value, 13
 - interprovincial patterns, 264–8
 - island patterns, 210–63
 - theory, 212–17
 - W. Australia, 250–2
 - z-values, 258–63
 - linearity, and their slopes, 268–76
 - lizards, islands of known age, 143–5
 - logarithmic base, 21–2
 - mainland patterns, 190–210
 - effect of biogeographical provinces, 191–2
 - effect of habitats, 190–210
 - effect of individuals, 190
 - mechanisms, 258–63
 - nested design, 10
 - nested scales, *i*
 - non-linear regression technique, 12
 - provincial effects, 204
 - regional pattern, 9–14, 278
 - scale, 268
 - small-scale pattern, 22–3, 279
 - southeast Asia, 131
 - synthesis
 - several scales of place and time, 276–9
 - several scales of space and time, 276–68
 - theory, re-doing, 379
 - three scales, 277
 - z-values, 15–22, 203
 - islands, 258–63
 - single islands, 238–42
 - various taxa, West Indian Islands, 237
 - see also* area
- species–area effects
 - equation, scale dependence of *z* and *c*, 21–2
 - and species diversity, 376–7
 - and species–time relationships, 66
 - variable, global scale, 370
- species curve of log-normal distribution,
 - defined, 270
- species diversity
 - alpha, beta, gamma diversities, 33
 - biogeographical islands, compared with mainland, 180–6
 - coevolution with habitat diversity, 151–89
 - dynamics, 299–301, 373–84
 - area effects, 376–7
 - differential equations, 374–6
 - future exploration, 377–8
 - hierarchical puzzle, 373–84
 - problems, 378–81
 - SLOSS controversy, 382
 - and energy flow, 345–72
 - hybrid zones, 47–8
 - increase of diversity, paleobiological patterns,
 - artifacts, 307–10
 - indices, 199–200
 - Fisher's α , 193–8, 203
 - Shannon–Weiner index of diversity, 59
 - Simpson's index, 198–202
 - Yule's index, 200–1
 - see also* Fisher's α ; Simpson's index
 - intermediate disturbance frequencies, 341–4
 - intermediate disturbance levels, 36–9
 - new habitats, 306–7
 - predator–victim ratios, 84–6
 - and productivity, 40–4
 - prospects, 371–2
 - relative, polyploid species, 48–9
 - sample size, 190–2
 - standing diversity, 377
 - steady state, 339–40
 - trophic levels, food chains and webs, 77–81
- species–energy hypothesis, 331, 352
- species flocks, 176–80, 380
- species richness [*footnote*], 201
- species–energy hypothesis, 331, 352
- species–time curves, 377, 379
- species–time relationships, 66–7
- spiders
 - orb-web, 116, 122
 - Bahamas, 129, 233
- stability *see* dynamical stability
- standing crop, 44
- standing diversity, insects, 377
- stasipatric speciation, 98, 103, 107
- steady states
 - epochal, 52–60, 297–305

- islands, 246–8
 - and isoclines, 303–4
 - steady state diversity, 339–40
- stenotopy, defined, 99
- Stephen Island wren, *Traversia lyalli*, 128
- stochasticity, 115, 127–8
- Streptochaeta sodoirana*, bamboo, 153–4
- succession, 62–3
- Sunda Islands, Indonesia
 - mammals, 257–8
 - relict species, 130–5
- super-hero phenotype, 151
- supertramps, 215
- Surrey
 - first species–area curve, 9
 - Lepidoptera, cycle of diversity, 65, 70
- surrogate variables, 208
- swallow, *Artamus*, 185
- Sweden
 - birds, 193–5
 - Lake Hjälmaren plants, 198, 224–5
- sympatric speciation, 98, 105–7
- Syncerus*, extinction and speciation rates, 137
- Tadzhikistan, rodent karyomorphs, 46
- Tasmania, eucalypt hybrid zones, 47–8
- Taurotragus*, extinction and speciation rates, 137
- Taylor series, 318
- teeth, mammals, 313
- temporal habitats, optimality, 174
- temporal patterns, 50–72
- Terborgh's pattern
 - latitudinal gradients, 284–6
 - tropical rainforest birds, 32
- termites, latitudinal gradients, 27
- territoriality, 359
- Texas, rodent diversities, 40
- Texas, Big Bend National Park, IFD in a guild
 - of species, 161–2
- thermodynamics
 - Elton–Hutchinson theory, 375
 - hypothesis, 327
 - second law, 320–2
- top ten current problems, 378–81
- Thuja* swamp forest, 270
- Timbuktu, polyploid species, 48
- time
 - dividing (plants), 169–74
 - ecological time, 62–72
 - evolutionary time, 50–62
 - habitats, optimality, 174
 - hypothesis and evaluation, 361–2
 - plant specialization, examples, 168–74
 - and productivity, 361–2
 - temporal patterns of diversity, 50–72
 - see also* species–time curves
- time travel, 361
- tolerance differences, shared preferences, 368
- trade-off principle, 151–6
 - costs, 166
- Evolutionary Stable Strategy, 311
 - giving up density (GUD), 165
- Tragelaphos*, extinction and speciation rates, 137
- trajectories, stability, 320
- Traversia lyalli*, Stephen Island wren, 128
- trees
 - birch forest, Finland, 326
 - deciduous habit, 68
 - host age relationship, 60–2
 - insects, 267–8
 - leaf miners, 61
 - oak–pine forest, 63
 - pollen dating, 60
 - southern beech forest, 182–4
 - Thuja* swamp forest, 270
- triclads, old lakes, 180
- trophic levels
 - defined, 80
 - food chains and webs, 77–81
 - higher, and diversity, 77–81
 - maximum, error, 324
 - Namib Desert, 322
 - population dynamics, 317–18
 - steady state, 301–5
 - see also* food chains and webs
- tropical rainforest, species diversity, 31
- tropical rainforests
 - Bahia, Brazil, 29
 - Malaysia, 30–2
 - plants, 265
 - species diversity, 2–3, 29, 31
 - understory plants, photosynthetic efficiency, 154
- tropics
 - latitudinal gradients, 284
 - lowland
 - birds, 369
 - species diversity, 380–1
 - ranges, spillover into non-tropical zones, 296
- Turdus*, thrush, 185
- turtles, North America and Australia, 192, 293
- Ukraine, chromosomal variants, mammals, 46–7
- Utah, and New Mexico, pinyon–juniper woodland, 209–10
- variation, in abundance, 122
- veil line, 352
- versatility, 312–15
- vertebrate/insect systems, predator–victim
 - ratios, 333
- vine habit, 167–8
- Vini vidivici*, conquered lorikeet, 233
- Viverridae, *see* mongooses
- Wadden Sea islands, Netherlands, 234, 258
- Wales, Skokholm Island, 426
- Wallace, island endemism, 109
- Washington, Puget Sound, colonization experiments, 227–8

- wasps, 113
- weasel, disturbance effect, 37
- weevils, 113
- West Indian Islands
 - hummingbirds, 235
 - z-values, various taxa, 237
- West Virginia, Greenbriar Valley, cave-dwelling species, 266–7
- wheatbelts, Australian vertebrates, 207–9
- Whittaker’s principle, 301
- Wilderness of Paran, Israel, 321
- Wisconsin
 - American chestnut, 124
 - fish, 204
- world, area, tropics vs northern lands, 284–5
- Wrangel Island, Russia, mammoths, 146
- wren, *Traversia lyalli*, 128
- Wrightian surface, 100
- writing style, *ii*
- Yule’s index, 200–1
- z-values, 273
 - actual values, 273
 - between-province, 264
 - on islands, 258–63
 - single islands, 238–42
 - mammals, 280, 283
 - reduced, higher species diversity, 276
 - small provinces, 383
 - species–area curves, 15–22, 258–63
 - islands, 258–63
 - see also* species–area curves
- zooplankton, Canadian lakes, 42
- Zostera marina*, 121