

Index

- absorption
 - large polaron, 75–77
 - large-polaron low-frequency Drude, 78–79
 - small polaron, 77–78, 83
 - small-(bi)polaron low-frequency polarization
 - conductivity, 79–80
 - softening bipolaron, 63, 78
- adiabatic
 - formalism, 21–23
 - limit, definition, 20, 29
 - small-polaron hopping, 101
- bandwidth, polaron
 - strong-coupling, 34
 - strong-coupling, large polaron, 34, 42
 - strong-coupling, small polaron, 35–36, 42
 - weak-coupling, 18
- barrier to self-trapping
 - free-carrier small-polaron dichotomy, 30
 - large-polaron small-polaron dichotomy, 32
 - persistent photo-conductivity, 31
 - small-polaron formation, 30
 - time delay for self-trapping, 30
- bipolaron
 - attraction between large bipolarons, 164–165
 - definition, 3, 54
 - electron correlation, 57–58
 - energy barrier to formation, 57
 - ferroelectric materials, 56
 - inert-pair effect, 59
 - Jahn–Teller, 55, 63
 - large, 54, 55–58
 - large four-lobed, 58
 - negative U , 59
 - pair-breaking in semiclassical hopping, 121–123
 - small, 54, 58–59, 96
 - softening, 55, 59–64, 78
 - softening, boron carbides, 59–60, 63–64
- bipolarons
 - mutual interactions, 164–165
- Born–Oppenheimer method. *See* adiabatic formalism
- boron carbides
 - neutron detector, 133
 - properties, 59–60
 - Seebeck coefficient, 132
 - softening bipolaron, 63–64
- carrier mobilities
 - conductivity mobility, 135
 - drift mobility, 135
 - Hall mobility, 135
 - standard, 135
 - time-of-flight mobility, 135
- carrier-induced atomic changes
 - altered atomic equilibrium positions, 1
 - altered atomic vibration frequencies, 1
- carrier-induced softening
 - atomic desorption, 186–190
 - causes, 129–130
 - definition, 25
 - electronic polarizability, 26, 92
 - enhanced Seebeck coefficient, 130–133
 - Holstein’s molecular-crystal model, 27, 52
 - increased thermal resistivity, 130
 - large-polaron scattering, 92
 - molecular polaron, 51
 - phonon localization, 130
 - Seebeck coefficient enhancement, 64
 - softening bipolaron, 59–64
 - strong-coupling polaron, 28–29, 33–34
 - vibrations’ free-energy, 28–29, 33–34
 - weak-coupling polaron, 13–20
- chalcogenide glasses
 - Hall effect sign anomalies, 158
 - optical properties, 84–85
 - small polarons, 185
 - threshold switching, 184
- coherent transport
 - definition, 86
 - Hall mobility, 136–138
 - large polaron, 88
 - minimum mobility, 87
- desorption
 - carrier-induced softening, 187–189
 - delayed desorption, 187, 190

desorption (cont.)
 immediate desorption, 187, 190
 surface electronic localization, 186–187
dielectric constants
 covalent materials, 10
 ferroelectric materials, 10
 ionic materials, 10
 large-bipolaron formation, 56
 optical, 10
 static, 10
effective mass, polaron
 at high frequencies, 89
 large-polaron, atomic tunneling regime, 89–91
 large-polaron, classical treatment, 88–89
 strong coupling, 3
 strong coupling, large polaron, 34, 88–91
 weak coupling, 14
electronic polarizability, 26
electronic transfer energy, 100–101
electronic velocity operator, 103
electron–phonon interactions, 9–12
 deformation-potential short-range interaction, 2, 11, 28, 30, 31
 Fröhlich coupling constant, 14
 Fröhlich long-range interaction, 2, 9, 28, 30
exciton
 Frenkel, 82
 self-trapped, 82
 Wannier, 82
Frenkel–Poole effect, 118
 for polar molecular solids, 51
geminate recombination
 definition, 81
 Onsager, 81
 small polaron, 81–82
Hall effect
 adiabatic Hamiltonian, 23
 adiabatic small-polaron hopping, 147–155
 Hall coefficient, definition, 135
 Hall mobility, Boltzmann solution, 136–138
 Hall mobility, definition, 135
 LaMnO, 47–48
 magnetic phase factors, 139
 MnO, 158
 non-adiabatic small-polaron hopping, 141–147
 sign anomalies, 3, 47–48, 137, 155–159
 sign anomalies in amorphous covalent semiconductors, 157–158
 sign anomalies in chalcogenides glasses, 158
 small-polaron hopping, square lattice, 143–146
 small-polaron hopping, triangular lattice, 141–143, 147–155
Holstein’s molecular-crystal model, 32, 51–52
 for molecular polarons, 51–52
 for transition-metal oxides, 27, 52
 hopping activation energy, 106

 pseudo-molecules, 27, 52
hopping. *See* phonon-assisted hopping
incoherent transport
 definition, 86
 small bipolaron, 96
 small polaron, 95–96, 138
large bipolaron
 Drude absorption, 169
 ghost modes, 169
 ordering, 170
 photo-absorption, 168
 photo-emission, 168
 positron annihilation, 169
 superconductivity. *See* superconductivity
 suppression of Seebeck’s spin entropy in cuprates, 171
large polaron
 absorption, 75–77
 definition, 3
 Drude absorption, 78–79
 Hall mobility, 138
large-polaron formation
 adiabatic limit, scaling analysis, 30
 quasi-one-dimensional, 32
large-polaron motion
 coherent, 3, 88
 Drude tail, 94
 effective mass, 34, 88–91
 Einstein-phonon scattering limit, 94
 Hall mobility, 138
 mean-free-path, 93
 mobility, 93, 94
 phonon scattering, 3, 91–94
 phonons’ thermal diffusion, 94
light-atom diffusion
 Condon approximation breakdown, 192–193
 high-temperature hopping, 191–193
 hydrogen diffusion in bcc metals, 193
 low-temperature isotope dependence, 196–197
 semiclassical isotope dependence, 193–196
 versus classical diffusion, 195
 versus small-polaron hopping, 195
luminescence
 exciton, 82
 optical cooling, 83–84
 self-trapped exciton, 82
 small polaron, 82–83
 Stokes shift heating, 83
magnetic field
 affect on atoms’ motion, 23
magnetic polaron
 bound magnetic polaron BMP, 68, 69–72
 colossal magnetoresistance CMR, 65, 71
 definition, 65
 EuO, 65, 71–72
 EuS, 71
 EuSe, 71
 EuTe, 65, 68, 71

- ferromagnetism, 65
- magnetic susceptibility, 65, 68
- small-polaron collapse, 68
- small-polaron collapse, thermally induced, 68–72
- stability, 67–68
- magnetic semiconductors
 - small-polaron hopping, 123–124
- Meyer–Neldel compensation effect, 116
- molecular polaron
 - definition, 49
 - Frenkel–Poole behavior, 51
 - Meyer–Neldel compensation effect, 51
 - scaling analysis, 49–51
 - Seebeck coefficient enhancement, 51
- Peltier heat
 - definition. *See* Seebeck coefficient
- persistent photo-conductivity. *See* barrier to self-trapping
- phonon-assisted hopping
 - activation energy, 106, 115
 - adiabatic semiclassical small-polaron, 110–117
 - adiabatic small-polaron, 98
 - adiabatic versus non-adiabatic regimes, 116–117
 - bipolaron pair breaking, 121–123
 - detailed balance, 108
 - Frenkel–Poole effect, 117–118
 - Hall effect sign anomalies, 155–159
 - inverted regime, 108
 - jump distance dependence, 117
 - magnetic semiconductors, 123–124
 - negative differential conductivity, 108
 - non-adiabatic impurity conduction, 98
 - non-adiabatic semiclassical small polaron, 108–110
 - non-adiabatic small polaron, 102–108
 - non-Ohmic at high fields, 180–184
 - non-Ohmic at low temperatures, 175–180
 - non-Ohmic hopping formalism, 173–175
 - shallow impurity, 97
 - small polaron, 97–102
 - threshold switching, 180–185
- polaron
 - absorption, 75–78
 - solvated electron, 3
 - strong-coupling, definition, 1
 - weak-coupling, definition, 1
- polaron radius
 - quasi-one-dimensional polaron, 32
 - strong-coupling, large polaron, 30, 34, 42
 - strong-coupling, small polaron, 31, 42
 - weak-coupling, 13
- rare-earth chalcogens
 - EuO, 65, 71–72, 133
 - EuS, 71
 - EuSe, 71
 - EuTe, 65, 68, 71
- recombination
 - geminate, 81–82
 - luminescence, 82–84
 - non-radiative, 83–84
- scaling analysis. *See* self-trapping: adiabatic limit
- Seebeck coefficient
 - carrier-induced softening’s enhancement, 53, 130–133
 - definition, 125
 - hopping carrier, 127–129
 - LaMnO, 47
 - low-temperature limit, 127
 - magnetic semiconductors, 133–134
 - magnetic-field dependence, 126
 - narrow-band limit, 125–126
 - polaron hopping, 128
 - singlet bipolaron, 126
 - spin degeneracy, 126
 - trap-modulated conduction, 129
 - with temperature-dependent energy levels, 126–127
- self-trapping
 - adiabatic approach, 1, 23–29
 - adiabatic limit, scaling analysis, 29–32, 189–190
 - barrier to self-trapping, 30, 190
 - beyond adiabatic limit, 33–39
 - definition, 1, 42
 - desorption, 186–187
 - electronic dimensionality effect, 29, 31–32, 189–190
 - electronically constrained, 2
 - nonlinear eigenvalue equation, 27
 - nonlinearity, 2
- small polaron
 - absorption, 77–78
 - definition, 3
- small polaron, defect-related
 - from breaking a trapped electron pair, 45
 - from trapping a large polaron, 44
 - impurity conduction. *See* small-polaron motion
 - trapped small polaron, 44
- small-polaron formation
 - adiabatic limit, scaling analysis, 30
 - band narrowing, 36
 - bandwidth, 35
 - barrier to self-trapping, 30, 32
 - beyond adiabatic limit, 33–39
- small-polaron formation, coexistence. *See* barrier to self-trapping
 - with free carriers, 30, 31, 36–39
 - with large polarons, 32
- small-polaron formation, induced by, 42
 - disorder, 41
 - electronic localization, 39–40
 - long-range electron–phonon interaction, 32
 - magnetic disorder, 5, 68–72
- small-polaron motion
 - adiabatic hopping Hall Effect, 147–155
 - adiabatic semiclassical hopping, 110–117
 - adiabatic versus non-adiabatic hopping, 116–117
 - bipolaron pair breaking, 121–123
 - coherent with atomic tunneling, 35–36
 - coherent-motion’s suppression, 36, 96, 138
 - Frenkel–Poole effect, 117–118
 - generalized for light-atom’s hopping, 191–193
 - Hall-effect sign anomalies, 3, 45–48, 155–159
 - hopping, 3, 36, 98–102

- small-polaron motion (cont.)
 - hopping activation energy, 106, 115
 - hopping Hall effect. *See* Hall effect
 - impurity conduction, 45–48
 - incoherent, 3, 95–96
 - incoherent for classical atoms, 35, 95–96
 - jump-distance dependence, 117
 - low-frequency polarization conduction, 79–80
 - magnetic semiconductors, 123–124
 - non-adiabatic Hall effect, 141–143
 - non-adiabatic hopping, 102–108
 - non-adiabatic semiclassical hopping, 108–110
 - non-Ohmic hopping, 180–185
 - photo-conductivity, 81
 - pinned Fermi level, 45
 - Seebeck coefficient, 127–129
 - Seebeck coefficient, magnetic semiconductors, 133–134
 - temporally correlated hopping, 118–119
 - thermally induced conductivity transition, 39, 42
 - threshold switching model, 180–185
 - time-of-flight mobility, 119–120
- strong-coupling polaron
 - effective mass, 3, 34
 - Fröhlich polaron energy, 14
 - large. *See* large polaron
 - optical absorption, 2
 - small. *See* small polaron
- superconductivity
 - BCS condensate, 163, 165
 - compared with superfluidity, 163
 - large bipolarons, 164
 - large-bipolaron liquid, 165–166
 - large-bipolaron properties, 168–172
 - large-bipolarons' excitation spectrum, 167–168
 - large-bipolarons' proximity effect, 172
 - mutual attraction between large bipolarons, 164–165
 - small- versus large-bipolaronic superconductivity, 163
 - superfluid helium's condensate, 163
 - suppression by large-bipolaron ordering, 171
- thermoelectric figure-of-merit, 132
- threshold switching, 185
 - chalcogenide glasses, 185
 - small-polaron model, 180–185
- time delay for self-trapping. *See* barrier to self-trapping
- time-of-flight mobility, 119–120
- transition-metal oxides
 - La₂CuO₄, 170, 171
 - LaMnO₃, 45–48
 - LiNbO₃, 85
 - MnO, 43, 52, 133, 158
 - NiO, 44
 - SrTiO₃, 44, 170
- vibrational dispersion
 - adiabatic small-polaron Hall effect, 150
 - adiabatic small-polaron hopping, 112–116
 - correlated hopping, 118–120
 - large-polaron scattering, 93–94
 - phonon-assisted hopping, 98
 - semiclassical small-polaron hopping, 109
 - small-polaron hopping, 105
- Wannier–Stark localization, 86–87, 103
- weak-coupling polaron
 - carrier-induced softening, 15–20
 - deformation-potential short-range electron–phonon interaction, 14
 - effective mass, 14
 - Fröhlich long-range electron–phonon interaction, 13
 - Fröhlich polaron energy, 14