

Index

Page numbers in **bold** refer to tables and in *italic* to figures

- Abies alba* (silver fir), 414
 abundance, variation across species' range, 315, 317
Accipiter gentilis (goshawk), 57, 58
Acer, 17
Acer saccharum, 10, 11
 adaptation, 440
 aggressive interactions avoidance, 140
Aglais, 8
Agropyron, 405
Alauda arvensis (skylark), 143, 144
 albatross, 83
 alder (*Alnus glutinosa*), 413, 414
 Allee effect, 308, 354, 355, 356, 359
 integral kernel-based models, 367, 368
 integrodifference model, 369
 stratified diffusion model, 360, 362
 allele frequency data, 89–107
 advantages/disadvantages, 90
 dispersal pattern differences
 detection in natural population, 98, 100, 101
 testing difference, 101–2
 F_{ST}
 derivation, 91–2
 dispersal estimation, 90, 439
 estimation, 92–3
 $N_e m$ (number of migrants) relationship, 94–6, 95, 96
 pairwise estimates for detecting restricted dispersal, 96–8, 99
 temporal variation studies, 106–7
 testing, 93–4
 $GUST_w$, 92
 mutation mechanisms, 104–6
 ϕ_{ST} , 104–6
 θ (co-ancestry), 92
 allozyme markers
 dispersal pattern studies, 101
 insect dispersal studies, 38
 seabird studies, 79
Alnus glutinosa (alder), 413, 414
 altitudinal movements, 41
 amphibian larvae, artificial markers, 52
 amplified fragment length polymorphisms (AFLPS)
 insect dispersal studies, 38
 yellow rust virulence mutations, 399, 400, 401
Androctonus australis toxin gene construct, 190
Andropogon gerardii, 166
 annuals
 insect resource patches, 113
 migration rate, 4
Anser caerulescens (snow goose), 147
Anser fabalis (bean goose), 140
 Antarctic fur seal, 81
 antennal sensilla, 125, 126
 ants
 dispersal swarms, 26
 seed dispersal, 5, 7, 16, 417
 aphids, 26, 113, 116–18, 117, 129, 437
 antennal sensilla, 125, 126
 chemosensory abilities, 125–6, 127
 condition-dependent dispersal, 146
 dispersal distances, 121, 122
 dispersal event sequence, 118–28, 119
 directed movement towards hosts, 124–5
 long-distance flight termination, 124
 sailing with wind, 120–4
 take-off, 120
 dispersal-related mortality, 124
 flight speeds, 114, 117
 monitoring networks, 29–30
 ocelli, 124
Aphis fabae, 121, 122, 125
Aphytis, 381
 arctic fox, 418
 ARGOS system, 53, 76
Aricia agestis (brown argus), 272, 273, 432
 mark–release–recapture studies, 259–61, 260, 262–3
Aricia artaxerxes (northern brown argus), 432
 artificial markings, land vertebrates, 51–2
Arvicola terrestris (water vole), 414

Cambridge University Press

978-0-521-54931-8 - Dispersal Ecology

Edited by James M. Bullock, Robert E. Kenward and Rosie S. Hails

Index

[More information](#)

446 INDEX

- Asarum canadense*, 416
 Asteraceae, seed fat storage, 154, 157
 Atlantic cod (*Gadus morhua*), 146–7
 Atlantic salmon (*Salmo salar*), 106
 auditory stimuli, 239
 Audouin's gull (*Larus audouinii*), 79
Autographa californica, 175, 176, 189, 190
Avena sativa (oat), 304
 azimuthal bias in seed dispersal, 5
- bacteria, 219–34
 adaptive change, 225
 role of plasmids, 226
 cosmopolitan populations, 220
 endemic populations, 220, 221
 horizontal gene transfer, 220, 221–2, 224, 226
 leaf-colonizing, 223, 224
 dispersal by phytophagous caterpillars, 229–33, 231, 231, 232
 levels of dispersal, 221, 222, 222–3
 long-distance spore dispersal, 354
 transposons, 224–5
- bacteriophages, 220, 221, 222
- baculoviruses, 173–5
 chitinase/cathepsin gene products, 175
 destruction of host, 175–7, 176
 lytic baculovirus gene products, 175–6
 dispersal within insect populations, 175
 genetic modifications for pest control, 174, 175, 177–8
 AaIT scorpion-toxin construct, 189, 190, 190–1
 fitness trade-offs, 178
 models of competition with wild-type strains, 186, 187–91, 189, 190
 population dynamics, 183
 risk assessment, 181, 184, 187
 host behaviour modification, 177, 187
 insect–pathogen interaction models, 178–85, 182
 diffusive instability, 184, 185
 dispersal, 183–4, 185, 186–7
 occlusion bodies, 174, 175, 177, 178
- badger (*Meles meles*), 52
- ballistic seed dispersal, 5, 7, 16
- bands (rings), 52, 53, 59, 60
 microelectronic tags attachment, 76, 77
 seabird/seal dispersal measurement, 74, 78
- barnacle goose (*Branta leucopsis*), 142, 143, 147
- bean goose (*Anser fabalis*), 140
- bear (*Ursos arctos*), 413, 414
- beaver (*Castor fiber*), 335, 336, 343
- beech (*Fagus sylvatica*), 413, 414
- behavioural approaches
 butterfly dispersal, 272, 273, 274
 landscape-level dispersal, 336
- behavioural observations, insect dispersal studies, 27, 29
 pest invasions forecasting, 30
- Belding's ground squirrel (*Spermophilus beldingi*), 146
- Bemisia tabaci*, 120
- Bertholletia*, 7
- Beta vulgaris* spp. *maritima* (sea beet), 101, 101, 103, 103–4
- Betula*, 17
- bias in dispersal detection, 60, 61, 66
 distance-weighted sampling bias, 259, 261, 268
 binary-coded wire tags, 53
- biodiversity
 consequences of climate change, 418–19, 420
 threats from exotic introductions, 374, 375
- biogeography, 303–21
 impact of ecology, 313–14
 impact on ecology, 314–19
 range
 dynamics, 305–13
 structure, 304–5
- biotic resistance, 379–80
- biotrophic fungal pathogens, 395–408
 asexually produced spores, 397, 408
 cereal–parasite coevolution, 405–7
 emergence of virulent genotypes, 399, 407–8
 epidemiology, 402–5
 host crop populations distribution, 397–8
 host specificity, 395
 gene-for-gene relationship, 395, 396, 397, 399
 long-distance spore dispersal, 398, 404–5, 407, 408
 spread of virulence mutations, 399–401, 400, 401
 obligate biotrophy, 395
 population genetics, 398–401
 sex as adaptation to host diversity, 407–8
 sexual structures, 407, 408
 special forms (host specialization), 395, 405, 407
- birds
 artificial markers, 52
 dispersal initiation, 64
 exotic species establishment on islands, 381, 383, 384, 384, 385
 metal rings (leg bands), 52
 philopatry, 145
 seed dispersal, 296, 417
 by caching, 5–6, 16, 19
 by defecation, 5, 7, 8, 16, 19
 determining seed source, 16
 epizoochory, 19
 long-distance, 5–6, 7, 8, 19
 mechanistic models, 15
 success-mediated dispersal, 145–6
 see also seabirds
- black-browed albatross, 77
- black fly (*Simulium vittatum*), 142
- black guillemot (*Cephus grylle*), 81
- black-tailed godwit (*Limosa limosa*), 140, 145, 147
- blackbird (*Turdus merula*), 63
- bleached markers, land vertebrates, 52
- blue butterfly (*Cyaniris semiargus*), 267
- blue tit (*Parus caeruleus*), 143, 144
- Blumeria graminis*
 evolution, 405–6, 407
 phylogeny, 406–7
 special forms, 395, 405
- Blumeria graminis* f.sp. *avenae*, 395, 406
- Blumeria graminis* f.sp. *hordei*, 395, 396, 397
 long-distance dispersal of virulent clones, 401
- Blumeria graminis* f.sp. *secalis*, 395, 406

- Blumeria graminis* f.sp. *tritici*, 406
epidemiological study in China, 402, 402–5, 404
- bottlenecks
insect mitochondrial DNA studies, 40
metapopulation dynamics, 205
seabird/seal genetic studies, 79
- Bracon*, 381
- Branta b. bernicla* (dark-bellied brent goose), 140
- Branta b. hrota* (light-bellied brent goose), 140
- Branta leucopsis* (barnacle goose), 142, 143, 147
- Brassicaceae, seed fat storage, 154
- breeding dispersal, 54, 55, 55, 56, 437
colonial breeding seabirds/seals, 80–3
depletion models, 140
interference models, 140
pre-/postnuptial movements, 56
territoriality, 141
- breeding failure, 82
- breeding site, seabird/seals, 73
population surveys, 74
- Briza media*, 163, 163
- Bromus*, 405
- Bromus tectorum* (cheat grass), 352, 361
- brown argus (*Aricia agestis*), 272, 273, 432
mark–release–recapture studies, 259–61, 260, 262–3
- brown bear, 331
- brown trout (*Salmo trutta*), isolation by distance, 98, 100, 101
- Brownian movements, 308
- Brünnich's guillemot (*Uria lomvia*), 81
- bryozoans, freshwater, 197–8
statoblasts, 197, 198
temporal gene flow, 211
- buffer effect, 135, 136–8, 139
observational studies, 142, 143, 144
- bumble bee, harmonic radar studies, 31, 33, 34, 35
- Buteo buteo* (buzzard), 57, 58, 59, 60, 62, 63, 336, 344
- butterfly dispersal, 257–75, 437
distances, 258
geographical barriers, 41
habitat management programmes, 266, 267
harmonic radar studies, 31, 33
long-distance, 262, 263, 264, 267–71
consequences, 269, 271
indirect measurement methods, 267, 268
mark–release–recapture studies, 259–66, 260, 265
distance-weighted sampling bias, 259, 261, 268
finite area adjusted (FAA) model, 261, 262
metapopulation biology, 258
pattern-based study approaches, 271–2
relationship to individual behaviour, 272, 273, 274
population dynamics, 266–7
radio-tagging, 259
study methods, 258
swarms, 26
synchrony in population fluctuations, 266
- buzzard (*Buteo buteo*), 57, 58, 59, 60, 62, 63, 336, 344
- cabbage moth (*Mamestra brassicae* L.), 177, 230, 231
- caddis fly, 31
- CAIC, 155
- Cakile*, 437
- Calathea ovandensis*, 287, 293
- Calidris canutus* (knot), 142, 143
- California gnatcatcher (*Poliophtila c. californica*), 336
- California sea otter, 352
- Callorhinus ursinus* (northern fur seal), 81
- Calluna vulgaris*, 167, 287, 288
capture bias, 60
- carabids, harmonic radar studies, 31
- carrion crow (*Corvus corone*), 142
- carrying capacity, population models, 332, 333
- Castor fiber* (beaver), 335, 336, 343
- cathepsin, baculovirus gene product, 175
- causes of dispersal, 64, 336, 438
seabirds/seals, 81–3
- cellular occupancy, 63–4
- Centaurea nigra*, 285
- Cephus grylle* (black guillemot), 81
- Cerastoderma edule* (cockle), 145
- Chamerion angustifolium*, 167
- Charadrius hiaticula* (ringed plover), 141
- cheat grass (*Bromus tectorum*), 352, 361
- checkerspot (*Euphydryas editha*), 267
- chitinase, baculovirus gene product, 175
- Chorthippus parallelus* (grasshopper), 413, 414
- Chorthippus vagans* (heath grasshopper), 422
- cladocerans, 206, 207
resting egg banks, 210
- climate change, 314, 410–24, 440
biodiversity consequences, 418–19, 420
conservation issues, 421
translocation projects, 422–3
dispersal response, 413
since last glaciation, 413–18
evidence for warming, 411–12
future greenhouse gas emissions, 412
impact on organisms, 412–13
advanced phenology, 418
recent responses, 418
invasions, 423–4
past climate reconstructions, 411
past migration rates, 415–17, 416
past migration routes, 413, 414, 415
- climate envelope analysis, 418–19
- clipping, vertebrate markers, 51–2
- clips, fish fins, 52
- cluster analysis, insect molecular marker studies, 39
- co-ancestry (θ), 92
- cockle (*Cerastoderma edule*), 145
- coevolution, powdery mildews with cereals/grasses, 405–7
- Coleoptera
flight polymorphism, 26
harmonic radar studies, 31
- collar tags, 53
- collared dove (*Streptopelia decaocto*), 308, 309, 352
- colonization, 64, 65
butterfly dispersal studies, 258
competition trade-off, 161–7, 163
effectiveness measures in seeds, 153
leaf-colonizing bacterial dispersal by phytophagous caterpillars, 229–33, 231, 231, 232

- colonization *cont.*
 local species richness, 196
 metapopulation dynamics, 195
 plants, 291, 292, 292
 plant recruitment, 279
 gap dynamics, 281–4, 282, 283
 invasion of new areas, 279, 284–90
 recolonization following extinction, 279
 range margins expansion, 307–8
 seal population dynamics, 79
 spatial patterns, 269
 zooplankton, 207, 209
- coloured markers, land vertebrates, 52
- common guillemot (*Uria aalge*), 81
- community structure
 composition changes, 318, 319
 consequences of climate change, 418–19
- competition, 438
 colonization trade-off, 161–7, 163
 definition, 163
 exotic organisms establishment in disturbed habitats, 385
 seed-limited abundance, 164–6, 165
- Compsohalieu harrisi* (Galapagos cormorant), 81
- condition-dependent dispersal, 146
- conidia, 397
 wind-borne dispersal, 403, 404
- Conopodium majus*, 164
- conservation, climate change issues, 421
- conspecific attraction, 240
- copepods, 206, 207
 resting egg banks, 210
- Corvus corone* (carrion crow), 142
- Crataegus monogyna* (hawthorn), 423
- crested newt (*Triturus cristatus*), 413, 414
- crested tit (*Parus cristatus*), 146
- Cristatella mucedo*
 metapopulation dynamics, 205, 211–12
 microsatellite studies, 198, 200–2, 202, 203, 205
 mitochondrial DNA sequence data, 202, 203, 204–5
 population genetic studies, 198–207, 199
 statoblasts, 198, 198, 210
- Cyaniris semiargus* (blue butterfly), 267
- Cytisus scoparius*, 286
- Dactylis*, 405
- Danaus plexippus* L., 29
- Daphnia*, 207, 209
- Daphnia galeata*, 207
- dark-bellied brent goose (*Branta b. bernicla*), 140
- deer mouse (*Peromyscus maniculatus*), 142
- defecation-dispersed seeds, 16, 19
 determining source, 16
 long-distance dispersal, 5, 7, 8
- Delia* spp., 127
- demographic parameters, variation across species' range, 315, 317, 317
- demographic stochasticity, 354
 range expansion models, 356, 359, 365, 370
- density-dependence, 332
- density-dependent dispersal, 134–48, 438, 439
 buffer effect, 135, 136–8, 139
 condition-dependent dispersal, 146
 differential settlement, 145
 exclusion, 145
 experimental studies, 144
 ideal despotic distribution, 135, 136
 ideal free distribution, 134–5, 135
 mechanisms
 depletion, 140
 interference, 140
 parasitism, 142
 predation, 142
 rank, 141
 territoriality, 141
 observational studies, 142, 144
 between sites, 142, 143
 emigration, 142, 143
 large-scale habitat use, 144
 local buffer effects, 142, 143, 144
 population dynamics, 146–7
 Fisher's model, 173, 174
 processes, 144–6
 seeds, 5
 sources and sinks, 135, 136, 146
 spatially explicit population models (SEPMs), 341
 success-mediated dispersal, 145–6
- density-dependent emigration, 142
Microtus oeconomus dispersal in experimental system, 249, 250, 251, 251
- density-dependent fecundity, 141
 depletion, 140
- Diaeretiella rapae*, 40
- diamond-backed moth (*Plutella xylostella*), 38
- differential settlement, 145
- diffusion models, 64, 354–9
 fragmented habitats, 356–7, 357, 358, 359
 long-distance dispersal, 355–6, 368–9
- Dipsacus sylvestris*, 287
- Discestra trifolii*, 233
- disease outbreaks, 314–15
- dispersal
 causes, 64, 336, 438
 decision mechanisms in animals, 50–1
 definition, 24, 54, 55–6, 73, 328
 ex-natal/extranatal movements, 56
 pre-/postnuptial movements, 56
 detection, 50, 57–9, 58
 animal size influence, 64, 66, 66
 bias, 60, 61, 66
 Ranges V software, 57, 58, 63
 sampling, 59, 59
 volunteer network observations, 59, 60
 local species richness, 196, 197
 recording process, 50
 types, 437–8
- dispersal curves, 3, 431–2
 definition, 4–5
 empirical, 5–8
 empirically fitted models, 8–13, 9, 10
 larger spatial scales, 11, 12, 13–14

- source geometry, 10–11, 11, 13–14
- field observation methods, 15–16
 - determining source, 16
- genetic marker studies, 89–90
- mechanistic models, 8, 14–15
- overlapping
 - methods for determining maternal source, 16
 - molecular study methods, 17
 - statistical methods, 16–17
- seed dispersal
 - local, 18
 - long-distance, 18–21
- shape, 434–7, 440–1
- types of dispersal, 437–8
- dispersal distance
 - aphids, 121, 122
 - fig wasps, 121
 - invasion patterns, 352–3
 - long-distance dispersal, 353–4
 - measurement methods, 353
 - seeds, 3–4, 13
 - spatially explicit population models (SEPMs), 339, 341, 344, 345
- dispersal limitation, 196, 280
- forest, 281
 - gap colonization dynamics, 282
- zooplankton, 207, 209
- dispersal movements, 308–10
- dispersal-related mortality
 - aphids, 124
 - fig wasps, 124
 - spatially explicit population models (SEPMs), 333, 335
- distance-weighted sampling bias, 259, 261
 - finite area adjusted (FAA) model, 261, 262
- disturbed habitats, exotic organisms establishment, 385–6, 387
- DNA markers
 - insect dispersal studies, 38
 - tracking migration response to past climate change, 413, 414, 415
- dominance hierarchies (rank), 141
- downy mildews, 395
 - sexual structures, 407
- zoospore dispersal, 397
- Drepanosiphum platanoidis*, 120
- drift in home range, 56, 57, 58
- earclip tags, 53
- Elisabethiella baijnathi*, 120, 124, 128
- emigration, 238–9
 - butterfly mark–release–recapture studies, 263–4, 265
- empirical dispersal curve, 5–8
- environmental predation, seabird breeding dispersal, 82
- Epilobium hirsutum*, 167
- epizoochory, 18, 19
 - long-distance seed dispersal, 5, 6
- Erica cinerea*, 287, 288
- Erinaceus europaeus* (hedgehog), 413, 414
- Eucalyptus*, 7
- Euphydryas editha* (checkerspot), 267
- Eupristina belagaumensis*, 124
- Evenstad project, 237–8, 240–52
 - experimental manipulations, 240–1, 241, 242
 - experimental study organism, 241
- event–alert (EVAL) system, 57
- EXAMINE, 30
- exclusion, 145
- ex-natal movements, 56
- exotic species *see* human-dispersed organisms
- experimental model systems, 240
- extinction
 - climate change impact, 413, 414
 - determinants of species' range, 305, 314
 - plant metapopulation dynamics, 291, 292
 - role of seed banks, 292
- extranatal movements, 56
- F*-statistics, 91, 92–3
 - insect dispersal, 39
- Fabaceae, seed fat storage, 154
- faecal DNA markers, 51, 89
- faecal isotope tags, 52
- Fagus*, 17
- Fagus sylvatica* (beech), 413, 414
- Fahrenheit's rule, 406
- fairy prion (*Pachyptila turtur*), 81
- Falco tinnunculus* (kestrel), 60
- fat storage in wind-dispersed seeds, 153, 154, 156, 156–7
 - phylogeny, 155
- feather genotyping, 89
- Festuca*, 286
- Ficus burtt-davyi*, 128
- fig (*Ficus* spp.), 116–17, 118
 - Anak Krakatau colonization, 121
 - volatile attractants, 127–8, 129
- fig wasps, 113, 116–18, 117, 129, 437
 - antennal sensilla, 125, 126
 - chemosensory abilities in host finding, 125, 126, 128, 129
 - dispersal distances, 121
 - dispersal event sequence, 118–28, 119
 - directed movement towards hosts, 125–8
 - host-seeking local flights, 128
 - long-distance flight termination, 124
 - sailing with wind, 120–4, 123
 - take-off, 118, 120
 - dispersal-related mortality, 124
 - flight speeds, 114, 117
 - host plant interactions, 117–18
 - ocelli, 124
- fighting, 140
- F_{IS} , 91
- fish, fin clips, 52
- F_{IT} , 91
- forbs, invasion (linear foci) experiments, 284–5, 285, 286
- forest
 - development on oldfields, 296
 - dispersal limitation, 281
 - gap dynamics, 281, 282, 284
 - succession, 295
- fossil record, 413

- founder events, 196
 insect molecular marker studies, 40
 metapopulation dynamics, 205
 seal studies, 79
 zooplankton dispersal, 209
- fox (*Vulpes vulpes*), 56, 336
- Fraxinus americana*, 10, 11
- Fraxinus excelsior*, 9
- freeze-marking, land vertebrates, 52
- freshwater invertebrates, 194–212
 active/passive dispersal methods, 196–7
 bryozoans, 197–8
 dormant propagules, 196–7, 198
 propagule banks, 210–11
 zooplankton, 206–10
- F_{ST}
Cristatella mucedo gene flow study, 200, 201
 derivation, 91–2
 dispersal estimation, 90, 439
 estimation, 92–3
 in natural brown trout population, 98, 100, 101
NUeum (number of migrants) relationship, 94–6, 95, 96
 pairwise estimates
 restricted dispersal detection, 96–8, 99
 testing differences, 101–2
 vicariance differentiation from isolation by distance, 102–4
 temporal variation studies, 106–7
 testing, 93–4
- fulmar (*Fulmar glacialis*), 74
- fungi
 long-distance dispersal by spores, 354
see also biotrophic fungal pathogens
- Gadus morhua* (Atlantic cod), 146–7
- Galapagos cormorant (*Compsahalieu harrisi*), 81
- gap dynamics, 286
 gap attainment ability, 281–2, 284
 gap composition determinants, 282, 283
 plants, 281–4, 294
- gene flow, 89, 272
Cristatella mucedo
 microsatellite studies, 198, 200–2, 202
 mitochondrial DNA studies, 202, 203, 204, 204–5
 insect dispersal studies, 39, 40, 41–2
 patchy populations, 196
 zooplankton, 206, 207, 209–10
see also F -statistics; F_{ST}
- gene-for-gene relationship, biotrophic fungi, 395, 396, 397
 emergence of virulent genotypes, 399
- genetic distance, insect mitochondrial DNA markers, 39
- genetic drift, 196
 zooplankton dispersal, 209
- genetic markers, 51, 89, 436
 direct methods, 90
 indirect methods, 90
 insect dispersal studies, 27, 31, 38–42
 altitudinal movements, 41
 butterflies, 258
 data interpretation, 39–40, 42, 43
 gene flow studies, 39, 40
 geographical barriers to population spread, 41
 large-scale movements, 40–1
 small-scale movements, 40, 40
 overlapping dispersal curves, 17
 seabird/seal dispersal, 74, 78–9
 philopatry, 81
 spread through population, 89
 tracking migration response to past climate change, 413, 414, 415
 zooplankton populations, 206–7
see also allele frequency data
- geographical information systems (GIS), 61, 63
 insect pest invasions forecasting, 30
- Geophilla*, 8
- germinants enumeration, 15–16
 costing study method, 19–21, 20
- global positioning system (GPS)
 land vertebrate tagging, 53–4
 seabird/seals tagging, 77
- Globigerina bulloides*, 436
- Glyceria maxima*, 163, 163
- goldenrod leaf beetle (*Trihabda virgata*), 142
- goshawk (*Accipiter gentilis*), 57, 58
- granulosis virus, 177
- grasshopper (*Chorthippus parallelus*), 413, 414
- grassland
 forb invasion (linear foci) experiments, 284–5, 285, 286
 gap dynamics, 281, 282, 283, 284
 grazing effects, 281, 282
 plant invasion modelling, 285–6
- grazing, 281, 282
- great tit (*Parus major*), 143, 144, 145, 335
- greenhouse gas emissions, 411, 412
- grey plover (*Pluvialis squatarola*), 144
- grey seal (*Halichoerus grypus*), 74, 75, 78, 79, 82, 82
- grey squirrel, 338, 339
- grid cells, spatially explicit population models (SEPMs), 329, 331, 332
- ground beetles, 315, 316
- gulls, 81, 82, 83
- gut contents, insect mark–recapture studies, 29
- gypsy moth (*Lymantria dispar*), 177
 nuclear polyhedrosis virus interaction modelling, 179, 180, 181, 183, 186–9
- habitat, 60
 definition, 328
 modelling problems, 64
 spatially explicit population models (SEPMs), 333, 334, 335, 336
- habitat corridors, 421
- habitat fragmentation, diffusion model, 356–7, 357, 358, 359
- habitat imprinting, 240
- habitat patches
 definition, 328
 dispersal response to climate change, 421, 422
 spatially explicit population model (SEPM)
 representations, 329, 335, 336
- habitat selection, 239, 240, 336
- habitat specialists, 432–3

- habitat tracking, 291, 310, 318, 320, 440
 habitat transience, 438
Haematopus ostralegus (oystercatcher), 141, 142, 145
Haliaeetus albicilla (white tailed sea eagle), 59, 423
Halichoerus grypus (grey seal), 74, 75, 78, 79, 82, 82
 harbour seal (*Phoca vitulina*), 74, 79, 81
 harmonic radar, insect dispersal studies, 27, 30, 31–3, 32, 34, 35
 hawthorn (*Crataegus monogyna*), 423
 heath grasshopper (*Chorthippus vagans*), 422
 heathland succession, 295, 296
 hedgehog (*Erinaceus europaeus*), 413, 414
Helianthemum nummularium, 260
Helicoverpa, 29, 30
Heliothis spp., 30
Heliothis virescens, 28
 Hemiptera, flight polymorphism, 26
Heracleum sphondylium, 164
 herbs
 migration rate, 3, 4
 seed dispersal, 19
 local, 5, 7
 source geometry, 13
 seed-limited abundance, 164, 165
Hesperia comma (silver spotted skipper), 268, 269, 270, 308, 432
 Himalayan thar, 352
 HIV, 314
 home range, 54, 55, 55, 67
 data collection bias, 60
 definition, 56–7, 328
 drift, 56, 57, 58
 prevailing range, 56
 spatially explicit population model (SEPM)
 representations, 329, 331, 332, 333, 335, 336
 home range behaviour, 272
 homing in salmon, 106–7
 homogeneous landscape, 431–2
 homogeneous patches, 433
 Homoptera, flight polymorphism, 26
 honey bees
 dispersal swarms, 26, 29
 harmonic radar studies, 33
 host–parasite coevolution
 Fahrenheit's rule, 406
 powdery mildews with cereals/grasses, 405–7
 host–pathogen models, 178–85
 dispersal, 183–4, 185
 host–pathogen population dynamics, 173–91
 house finch, 352, 367
 human-dispersed organisms, 374–90
 escape from parasitism in new habitats, 388–90, **389**
 habitats vulnerable to invasion, 376, 378–9
 disturbance, 385–6, **387**
 physical models, 379
 island invasions, 375–6, 379–81, **380, 382, 383, 384**, 384–5
 organism reproductive potential, 387–8, **388**
 species traits for successful invasion, 375, 376, 378–9
 proxy data sets, 379
 threats to biodiversity, 374, 375
 translocation projects, 422–3
 transport mechanisms, 374–5, 385, 386
Hura, 7
Hyalopterus pruni, 125
 hybrid zones, 413, 415
 Hymenoptera, dispersal swarms, 26

 ideal despotic distribution, 135, 136, 147
 territoriality, 136, 141
 ideal free distribution, 134–5, 135, 147
 buffer effect concept, 136–8, 139
 interference, 140
 immigration, 239–40
 butterfly mark–release–recapture studies, 263–4, 265
 metapopulations, 293, 294
 Microtus oeconomus in experimental system, 245, 246
 spatially explicit population models (SEPMs), 333
Impatiens, 7
 implanted satellite transmitters, seabird/seal dispersal
 measurement, 76
 inbreeding avoidance, 238, 242, 336, 337, 438
 Microtus oeconomus dispersal in experimental system, 243, 245
 inbreeding depression, 90, 354
 incidence function model, 328
 individual-based single-step models, 433, 434
 infinite alleles mutation model, 104
 infinite island model, 94–5, **95**
 influenza, 314
 initiation phase, 64, 352
 spatially explicit population models (SEPMs), 331–2, 341
 insect dispersal, 24–43
 active/passive features, 25–6
 behavioural observations, 27, 29
 harmonic radar studies, 27, 30, 31–3, 32, 34, 35
 mark–recapture studies, 27–9, **28**
 modelling, 27, 30
 molecular genetic markers, 27, 31, 38–42
 polymorphism/polyphenism, 26
 population monitoring, 25–6
 scanning radar studies, 27
 study aims, 25
 study techniques, 26–42
 direct/indirect, 26–7, 42
 swarms, 26
 tracking problems, 25–6
 traps, 29–30
 triggering factors, 25
 vertical looking radar (VLR), 27, 30–1, 33–8, 36, 37
 see also wind-dispersed insects
 insect pests
 invasions forecasting, 25, 30
 vertical looking radar (VLR) monitoring, 35
 insects
 baculovirus interactions see baculoviruses
 exotic species establishment on islands, **380, 380–1, 382**, 384
 flight behaviour see small insect flight behaviour
 integral kernel-based models, 354
 long-distance dispersal, 362, 364–7, 366, 368, 368–70

- integrodifference model, 364, 365, 366, 369, 369–70
 interference, 140
 intraspecific kleptoparasitism, 140
 introns, insect dispersal studies, 38
 invasion, 350
 biotic resistance, 379
 climate change response, 423–4
 determinants of species' range, 305, 314
 human-dispersed organisms, 375–6
 islands, 375–6, 379–81, **382, 383, 384**, 384–5
 stages ('filters') model, 377, 377–8
 velocity, 305–6, 306
 see also range expansion
 island model, 433
 see also infinite island model
 islands
 biotic resistance to invasion, 379–80
 invasions by human-dispersed organisms, 375–6,
 379–81, **380, 382, 383, 384**, 384–5
 vacant niches, 380
 isolation by distance, 97
 Cristatella mucedo microsatellite studies, 201
 detection, 98, **99**
 in natural brown trout population, 98, 100, 101
 differentiation from vicariance, 102–4, 103
 insect dispersal studies, 40
 isotope tags, 52

Juniperus virginiana, 13
 juvenile dispersal, 55, 56

k-alleles mutation model, 104
 kestrel (*Falco tinnunculus*), 60
 kittiwake (*Rissa tridactyla*), 81, 83
 knot (*Calidris canutus*), 142, 143

Lacerata agilis (sand lizard), 422
Lagopus lagopus (red grouse), 142
 Land Cover Map of Great Britain, 63
 land vertebrates dispersal
 detection, 50, 54–60
 experimental approaches, 64
 maps, 50, 60–3, 62, 66
 markers, 50, 51–4, 52, 66
 animal size effects, 64, 66, 66
 models, 50, 63–4
 study methods, 50
 landscape
 definition, 328
 heterogeneous, 433–4
 homogeneous, 431–2
 types, 431
 landscape-scale studies
 behavioural approaches, 336
 insect mark–recapture studies, 28
 plant dispersal, 279, 280, 290–8, 437
 large blue butterfly (*Maculinea arion*), 423
Larus audouinii (Audouin's gull), 79
Larus occidentalis (western gull), 147
 leaf-colonizing bacteria, 223, 224
 leg rings *see* bands

 Lepidoptera
 baculovirus pest control, 174
 migration behaviour, 29
 see also butterfly dispersal
Leucanthemum vulgare, 284, 285, 285, 286
 life history characteristics
 freshwater bryozoans, 197–8
 gap colonization dynamics, 282, 284
 invasion pattern studies, 352
 light-bellied brent goose (*Branta b. hrota*), 140
 light loggers, seabird/seal dispersal measurement, 77
 light traps, 29, 38
 pest insect invasions forecasting, 30
Limosa limosa (black-tailed godwit), 140, 145, 147
 line sources, seed dispersal empirically fitted models,
 10–11, 11
Liporrhopalum tentacularis, 125, 126, 128
 lizards, 64
 local scale dispersal
 plants, 279, 280–90
 gap dynamics, 281–4, 282, 283
 invasion of new areas, 279, 284–90
 range expansion, 350, 351
 seeds, 5, 7
 dispersal curves, 18
 study methods, 17–18
 locust swarms, 26
Lonchocarpus pentaphyllus, 9
 long-distance dispersal, 350–70, 410
 bacterial spores, 354
 butterflies, 262, 263, 264, 267–71
 diffusion model, 355–6, 368–9
 dispersal curve shape, 434–7
 dispersal distance, 353–4
 during Holocene migration, 416–17
 fungal spores, 354
 biotrophic fungi, 397, 398–401
 integral kernel-based models, 362, 364–7, 366, 368,
 368–9
 range expansion, 350, 351, 352
 seeds, 5–8, 8, 11, 12, 13–14, 18–21, 354, 417
 stratified diffusion model, 359–62, 360, 361, 363, 368–9
 viruses, 354
 lost parasite hypothesis, 389
Lotus comiculatus, 285
Lymantria dispar (gypsy moth), 177
 nuclear polyhedrosis virus interaction modelling, 179,
 180, 181, 183, 186–9

Maculinea arion (large blue butterfly), 423
Mamestra brassicae L. (cabbage moth), 177, 230, 231, 232,
 233
 mammals
 dispersal causality, 64
 seed dispersal
 by defecation, 16, 18
 local, 5
 long-distance, 5, 8, 8, 18
Maniola jurtina (meadow brown), 274
 maps
 land vertebrates dispersal, 50, 60–3, 62, 66

- raster data, 61, 62, 63
vector data, 61, 62
- mark–release–recapture studies, 257
- butterfly dispersal, 258, 259–66, 260, 265
long-distance dispersal events, 262, 263, 266
- density-dependent dispersal, 142, 143
- dispersal distance measurement, 353
- distance-weighted sampling bias, 60, 259, 261, 268
finite area adjusted (FAA) model, 261, 262
- home range concept, 54
- insect dispersal, 27–9, 28
essential criteria, 27
forensic markers, 29
genetic markers, 28–9
landscape-scale studies, 28
recapture problems, 27–8
- mass markings, 59
- modelling approaches, 264
- seabird/seal dispersal, 74, 76–8, 80–1
- volunteer networks, 59
- Markov chains, 310
- matrix areas, 328
spatially explicit population models (SEPMs), 333, 334, 335, 336
- meadow brown (*Maniola jurtina*), 274
- meadow grasshopper, 413
- measles, 314
- measurement of dispersal
land vertebrates, 50, 54–60
seabirds/seals, 73–9
- mechanistic dispersal curve models, 8, 14–15
- Meles meles* (badger), 52
- Melittaea cinxa*, 439
- mer* (mercuric resistance) genes, 221, 222, 222, 223, 224
plasmids, 225, 226
dispersal experiments, 228–9
- metapopulation dynamics, 194–6, 195
Cristatella mucedo, 205, 211–12
plant dispersal, 290–4, 292, 293
differentiation from seed bank emergence, 292, 293
independent dynamics of populations, 291
periodic extinctions, 291, 293
recolonization of unoccupied sites, 291, 292, 293
- seabird/seal dispersal, 80
- synchrony in population fluctuations, 266
- metapopulations, 290, 291, 315, 328, 432–3, 439
butterflies, 258
persistence, 64, 269
- Metrosideros*, 436
- Metrosideros collina*, 436
- Metrosideros polymorpha*, 436
- microsatellites
Cristatella mucedo gene flow, 198, 200–2, 202, 203, 205
 F_{ST} estimation in natural populations
Atlantic salmon, 106–7
brown trout, 98
mutation rates, 104–6
- insect dispersal, 38, 39, 40, 41, 41
- parentage analysis, 89
- pollen dispersal, 89
- seal dispersal, 79
- seed dispersal, 89
- microtransponders (passive integrated transponders; PITs)
butterfly tagging, 259
land vertebrate tagging, 52–3
seabird/seal tagging, 78
- Microtus oeconomus* (root vole), 136, 238, 240
avian predation, 244, 246
dispersal in experimental system, 242–4
Evenstad project study, 241
spatiotemporal population dynamics, 244–5, 245
conceptual model, 245–7, 246
mathematical model, 247–9, 250, 251, 251
transfer movements, 244
- migration
behavioural cues in insects, 29
definition, 24
long-distance seed vectors, 5
rate in terrestrial plants, 3–4
- mildews
distribution of host populations, 398
see also downy mildews; powdery mildews
Milvus milvus (red kite), 59, 344, 423
Mimosa pigra, 286
- minisatellites, insect dispersal studies, 38
- mistletoe (*Phrygilanthus sonoroe*), 292
- mitochondrial cytochrome *b* gene, seabird dispersal studies, 79
- mitochondrial DNA markers
Cristatella mucedo, 202, 203, 204–5
insect dispersal studies, 38, 39
bottlenecks, 40
demographic assessments, 39
founder effects, 40
interpopulation nucleotide divergence data, 39
maximum parsimony methods, 39
seabird dispersal studies, 79
- molecular markers *see* genetic markers
- monk seal, 77
- monkeys, seed dispersal, 8
- moths
harmonic radar studies, 33
pheromones, 126, 126
- Mount St Helens eruption, 295
- moving phase *see* transfer movements
- multilocus probes, 38
- muskrat (*Ondatra zibethicus*), 306, 352
- mymarid wasps, 129
- Myrmica sabuleti*, 422
- Myzus persicae*, 125
- natal dispersal, 54, 55, 55, 56, 437
ex-natal/extranatal movements, 56
proximate factors, 238
ultimate causes, 238–9
- $N_e m$ (number of migrants), 90
Cristatella mucedo gene flow, 200, 201
 F_{ST} relationship, 9, 94–6, 95
- Neogloboquadrina pachyderma*, 436
- northern brown argus (*Aricia artaxerxes*), 432

- northern fur seal (*Callorhinus ursinus*), 81
Nucifraga, 5
nuclear polyhedrosis virus, 175, 176, 177, 179
- oak (*Quercus* spp.), 7, 16, 17, 413, 414
oat (*Avena sativa*), 304
observational studies
 butterfly dispersal, 258
 land vertebrates dispersal, 51
oceanic planktonic foraminifera, 436
olfactory stimuli, 239
 see also volatile attractants
olive baboon (*Papio anubis*), 141
Ondatra zibethicus (muskrat), 306, 352
Orthopteran dispersal swarms, 26
Osmoderma eremita, 27
ostracods, 206
outbreeding depression, 90
oystercatcher (*Haematopus ostralegus*), 141, 142, 145
- Pachyptila turtur* (fairy prion), 81
Pagodroma nivea (snow petrel), 81
Papio anubis (olive baboon), 141
Pararge aegeria (speckled wood), 268
parasitism
 density-dependent dispersal, 142
 exotic species establishment in new habitats, 388–90, 389
 seabird breeding dispersal, 82
parentage analysis, 89
partridge (*Perdix perdix*), 142
Parus caeruleus (blue tit), 143, 144
Parus cristatus (crested tit), 146
Parus major (great tit), 143, 144, 145, 335
passerines, 338–9
passive integrated transponders (PITs) *see* microtransponders
PATCH, 331
patches
 buffer effect, 135, 136–8, 139
 butterfly dispersal
 larval host plants, 258
 mark–release–recapture studies, 259–66
 gene flow, 196
 homogeneous, 433
 ideal free distribution, 134–5, 135
 insect population dynamics, 113
 local species richness, 196, 197
 metapopulation dynamics, 194–6, 195
 Microtus oeconomus dispersal in experimental system, 245, 246, 246
 range distance, 352
 seed dispersal empirically fitted models, 10–11, 11
 sources and sinks, 136
 see also habitat patches
patchy populations, 432–3
 differentiation from metapopulations, 294
 spatially explicit population models (SEPMs), 329, 332, 333
Pedicularis furbishiae, 292–3
Perdix perdix (partridge), 142
Peromyscus maniculatus (deer mouse), 142
Petasites hybridus, 167
pheromone traps for insects, 29
 pest invasion forecasting, 30
philopatry, 147
 British birds, 145
 colonial seabirds, 80, 81, 147
 seals (pinnipeds), 81
Phoca vitulina (harbour seal), 74, 79, 81
Phorodon humuli, 121, 125
photographic identification, seabirds/seals, 78
Phragmites australis, 167
Phrygilanthus sonoroae (mistletoe), 292
phylogenetically independent contrasts (PICs), seed fat storage study, 154, 155, 156–7
Phylogophora meticulosa, 233
Picea engelmannii, 12
Picea glauca, 12
pine sawyer beetle, 367
pine wilt disease, 367
pinnipeds *see* seals
Pinus, 7, 7, 19
Pinus strobus, 9
plague, 352
planktonic foraminifera, 436
plant dispersal, 279–98, 437
 invasion of new areas, 279, 284–90
 forb invasion (linear foci) experiments, 284–5, 285, 286
 modelling studies, 285–7, 288, 289, 289
 landscape scale, 279, 280, 290–8, 437
 established species (metapopulations), 290–4, 292, 293
 pseudocolonization from seed banks, 292
 succession, 294–7
 local scale, 279, 280–90
 gap dynamics, 281–4, 282, 283
 migrational velocity, 3–4
 see also seed dispersal
plant fungal pathogens *see* biotrophic fungal pathogens
plasmids, 220, 221, 222
 dispersal by self-transfer, 226–9
 distribution, 222–3
 vertical persistence, 225–6
platform transmitter terminals (PTTs), 53–4, 76
 size, 76
Platypodium elegans, 9
Plebejus argus (silver studded blue butterfly), 422
Plodia interpunctella, 177
Plutella xylostella (diamond-backed moth), 38
Pluvialis squatarola (grey plover), 144
Poa, 405
Poaceae, seed fat storage, 154
pocket gopher–chewing lice relationship, 406
Polioptila c. californica (California gnatcatcher), 336
pollen diagrams, 314
pollen dispersal, 89–90, 102
 fig wasp behaviour, 121
pollen record, 318, 413, 415, 416, 418
population composition, 318–19
population dynamics, 439

- butterfly dispersal, 266–7
density-dependent dispersal, 146–7
host–pathogen, 173–91
Microtus oeconomus dispersal in experimental system, 244–5, 245
 conceptual model, 245–7, 246
 mathematical model, 247–9, 250, 251, 251
seabird/seal dispersal, 74, 79–80
 see also metapopulation dynamics
population structure, 439
post-fledgling dispersal, 55
postnuptial movements, 56
powdery mildews, 395
 cereals/grasses coevolution, 405–7
 epidemiological study in China, 402, 402–5
 loss of host resistance, 399
 sexual structures (cleistothecia), 407
 wind-dispersed conidia, 397
 long-distance dispersal, 398
predation
 density-dependent dispersal, 142
 seabird breeding dispersal, 82
prenuptial movements, 56
prey depletion, 140
prey disturbance, 140
propagule banks, freshwater invertebrates, 210–11
Prunella vulgaris, 285
Prunus, 17, 127
Pseudomonas, 220, 221
Pseudomonas fluorescens SBW25, 223
 dispersal by phytophagous caterpillars, 229–33, 231, 231, 232
 plant association, 223
 plasmids, 223
 dispersal experiments, 227, 227–9
Psidium aquilinum, 167
Puccinia graminis, 405
Puccinia striiformis, 396, 408
Puccinia striiformis f.sp. *tritici*, 396, 402
 long-distance spore dispersal, 401, 405
 spread of virulence mutations, 399–401, 400, 401
population genetics, 398–401
resistant wheat cultivars
 breakdown of resistance, 399
 production, 398–9
Quaternary climate change, 413
Quaternary ecology, 314
Quercus ilex, 422
Quercus spp. (oak), 7, 16, 17, 413, 414
rabies, 352
radio tags, 55, 56, 60
 butterflies, 259
 insects, 27
 land vertebrates, 53–4
 Microtus oeconomus (root vole), 241
 seabirds/seals, 74
random amplified polymorphic DNA (RAPDs)
 Cristatella mucedo metapopulation dynamics, 205
 insect dispersal studies, 38–9
range, 304–13, 439–40
 biotype accessibility, 305–7
 range centre versus margins, 307
 waiting times for return of favourable conditions, 307
 biotype variation, 315
 community composition changes, 318, 319
 demographic parameter variations, 315, 317, 317
 dynamics, 305–13, 314
 dispersal movements of individuals, 308–10
 simulations, 310–12, 311, 312, 313
 habitat consistency, 315
 habitat tracking, 318
 margins, 307–8, 315
 radial distance, 352
 scattering distance, 352
 shifts due to climate change, 314
 speciation processes, 320
 structure, 304–5
 see also home range
range expansion, 350
 diffusion model, 354–9
 distance versus time curves, 351, 352
 expansion phase, 352
 initial establishment phase, 352
 integral kernel-based models, 362, 364–7, 366
 long-distance dispersal, 350, 351, 352
 satellite colonies, 350, 351
 saturation phase, 352
 short-distance dispersal, 350, 351
 stratified diffusion model, 359–62, 360
Ranges V software, 57, 58, 63
rank (dominance hierarchies), 141
raptors, 51, 58
raster maps, 61, 62, 63
recruitment, 3
 seal population dynamics, 79
 statistical methods, 17
recruitment curve, 5
red deer, 352
red fox, 331
red grouse (*Lagopus lagopus*), 142
red kite (*Milvus milvus*), 59, 344, 423
red squirrel (*Sciurus vulgaris*), 146, 335, 338
remnant populations, 292, 294
remote imaging, raster maps, 63
rescue effect, 315
resource competition, 238, 242, 243
resource patchiness, 113
restriction fragment length polymorphisms (RFLPs)
 dispersal pattern studies, 101, 102
 insect dispersal studies, 38
 mitochondrial DNA, 39
Rhianthus minor, 284, 285, 285, 286, 287, 289, 289, 289–90
Rhopalosiphum padi, 124, 127
rice water weevil, 352
ringed plover (*Charadrius hiaticula*), 141
rings *see* bands
Rissa tridactyla (kittiwake), 81, 83
rodents, local seed dispersal, 5, 7

- root vole *see* *Microtus oeconomus*
 rotifers, 206, 207
 resting egg banks, 210
 rusts, 395
 distribution of host populations, 398
 sexual structures (telia), 407
 wind-dispersed uredospores, 397
 long-distance dispersal, 398

Salmo salar (Atlantic salmon), 106
Salmo trutta (brown trout), isolation by distance, 98, 100, 101
 sampling, dispersal detection, 59, 59
 sand lizard (*Lacerta agilis*), 422
 satellite-based tagging, 53, 76
 butterfly dispersal, 259
 seabird/seal dispersal, 74, 76
 implanted tags, 76
 satellite colonies, 350, 351, 352, 354, 359
Scabiosa, 7
 scanning radar, 27
Sciurus carolinensis (squirrel), 63
Sciurus vulgaris (red squirrel), 146, 335, 338
 sea beet (*Beta vulgaris* spp. *maritima*), 101, 101, 103, 103–4
 sea eagle (*Haliaeetus albicilla*), 59, 423
 seabirds, 72–84
 breeding range changes, 74
 breeding strategy, 72
 colonial breeding, 73, 79
 dispersal mechanisms, 81–3
 philopatry, 80, 81
 dispersal, 72
 genetic analysis, 74, 78–9, 81
 mark–recapture studies, 74, 76–8, 80–1
 measurement, 73–9
 patterns, 80–1
 population dynamics, 74, 79–80
 proximate causes, 81–3
 social behaviour, 73, 83
 foraging range, 72, 76, 83–4, 84
 individual reproductive value, 73, 83
 seals (pinnipeds), 72–84
 breeding range changes, 74
 breeding strategy, 72
 colonial breeding, 73, 79
 dispersal mechanisms, 82, 82
 dispersal, 72
 genetic analysis, 74, 78–9, 81
 mark–recapture studies, 74, 76–8
 measurement, 73–9
 patterns, 80, 81
 philopatry, 81
 population dynamics, 74, 75, 79–80
 proximate causes, 81–3
 social behaviour, 73, 83
 foraging range, 72, 76, 84
 individual reproductive value, 73
 seed banks, 159, 292, 293
 seed dispersal, 3–21, 89, 90, 102
 azimuthal bias, 5
 definition, 4–5
 density dependence, 5
 dispersal curves, 3
 line/small patch sources, 10–11, 11
 distances, 13
 scales, 3–4
 effectiveness measures, 152–3
 empirically fitted models, 8–13, 9, 10
 larger spatial scales, 11, 12, 13–14
 gap colonization dynamics, 281–2, 283, 284
 local, 5, 7, 17–18
 long-distance, 5–8, 8, 11, 12, 13–14, 354, 417
 study methods, 18–21
 microsatellites, 89
 plant migration rate since last glaciation, 415–17
 population expansion models, 286
 trade-offs, 152–68
 vectors, 5–8, 7, 8, 417
 primary/secondary, 5, 6
 wind-dispersed trees, 8–15
 seed examination in litter, 15
 costing of study method, 19–21, 20
 seed-limited abundance, 164–6, 165
 seed persistence in soil, 152, 157–60, 160
 seed traps, 15
 costing of study method, 19–21, 20
 local scale invasion studies, 287, 289, 290
 seedlings examination, 15–16
Senecio vulgaris, 286
Setaria, 7
 shrew, 413, 414
 shrimps, 89
 shrubs
 heathland succession, 295
 seed-limited abundance, 164
Silene dioica, 293
 silver fir (*Abies alba*), 414
 silver spotted skipper (*Hesperia comma*), 308, 432
 silver studded blue butterfly (*Plebejus argus*), 422
Simulium vittatum (black fly), 142
 single locus probes, 38
 single-step individual-based models, 433, 434
 single-stranded conformation polymorphisms (SSCPs),
 insect dispersal studies, 38
 skin slough genotyping, 89
 skipper (*Hesperia comma*), 268, 269, 270
 skunk (*Spinachia spinachia*), 142
 skylark (*Alauda arvensis*), 143, 144
 small insect flight behaviour, 113, 114–16
 dispersal event sequence, 118–28, 119
 directed movement towards hosts, 124–8
 long-distance flight termination, 124
 sailing with wind, 120–4
 take-off, 118, 120
 visual stimuli responses, 125
 volatile attractant responses, 125–8
 snakes, 53
 snow goose (*Anser caerulescens*), 147
 snow petrel (*Pagodroma nivea*), 81
 social behaviour, seabird/seal dispersal, 73, 83
 sooty tern (*Sterna fuscata*), 81

- Sorghastrum nutans*, 166
 source–sink dynamics, 135, 136, 291
 butterfly dispersal, 266, 267
 density-dependent dispersal, 146
 metapopulations, 293, 293
 patchy populations, 294
 plant colonization processes, 281
 spatial autocorrelation, insect molecular marker studies, 39
 spatial contact models, 64
 spatially explicit population models (SEPMs), 64, 327–46, 433–4
 definitions, 328
 dispersal behaviour uncertainty, 342–3
 modelling for management recommendations, 343
 modelling for research, 343–5
 dispersal-related mortality, 333, 335
 dispersal representation, 330, 331, 331–5, 334
 consequences of uncertainty, 338–42, 340, 342
 dispersal distance, 339, 341, 344, 345
 evaluation, 336–7
 initiation phase, 331–2, 341
 moving phase, 333
 parameter values, 337, 338
 stopping phase, 333
 empirical data incorporation, 335
 evaluation of structural assumptions, 335–7
 map data, 60
 population spatial structure representations, 329, 330, 331, 331, 332
 consequences of misrepresentation, 342
 evaluation, 335–6
 primary predictions, 338
 secondary predictions, 337–8
 speciation, 320
 species richness, 196, 197
 speckled wood (*Pararge aegeria*), 268
Spermophilus beldingi (Belding's ground squirrel), 146
Spinachia spinachia (skunk), 142
Spodoptera exempta, 26, 29
Spodoptera spp., 30
 spotted owl, 335, 343
 squirrel (*Sciurus carolinensis*), 63
 squirrel (*Tamiasciurus hudsonicus*), 56
 squirrels, 51
 starling, 352
 statoblasts, 197, 198, 210
 sediment core studies, 205
 stepping stone models, 97, 195–6, 433
 mutation effects, 106
Sterna fuscata (sooty tern), 81
 stratified diffusion model, 354
 long-range dispersal, 359–62, 360, 361, 363, 368–9
 streamer tags, 53
Streptopelia decaocto (collared dove), 308, 309, 352
 stress, seabird breeding dispersal, 83
 success-mediated dispersal, 145–6
 succession, 294–7
 suction traps, 29, 30, 38
 swarms, 26
 tag bias, 60
 tags
 active, 53–4, 59, 64, 66–7
 artificial, 51–2
 butterfly dispersal studies, 258
 dispersal detection, 57
 animal size influence, 64, 66, 66
 genetic *see* genetic markers
 land vertebrates, 50, 51–4, 52, 64, 66–7
 passive, 52–3
 sampling sizes, 59
 seabird/seals, 74, 76
Tamiasciurus hudsonicus (squirrel), 56
Tapesia yallundae, 399
Taraxacum officinale, 4
 terns, 81, 83
 territorial behaviour, 136, 141
 Microtus oeconomus (root vole), 241
 θ (co-ancestry), 92
 thrush (*Turdus philomelos*), 63
Tilia americana, 9
 tracking, 257
 dispersal detection, 57
 insect dispersal studies, 27
 trade-offs, 167–8, 440
 baculovirus genetic modifications for pest control, 178
 colonization–competition, 161–7, 163
 direction–distance for small flying insects, 114–15, 115, 129
 seed dispersal, 152–68
 transfer movements, 239
 spatially explicit population models (SEPMs), 333
 translocation projects, climate change response, 422–3
 transponders *see* microtransponders
 transposons, 224–5
 insect dispersal studies, 38
 traps
 insect dispersal studies, 29–30
 trap networks, 29
 insect pest invasions forecasting, 30
 tree top disease (Wipfelkrankheit), 177
 trees
 heathland succession, 295
 migration rate, 3, 4
 since last glaciation, 415, 416
 seed dispersal, 19
 local, 5, 7
 long-distance, 8, 437, 440
 wind, 8–15, 19–21
 seed-limited abundance, 165
Trichoplusia ni, 189, 190
Trihabda virgata (goldenrod leaf beetle), 142
Triticum ventricosum, 399
Triturus cristatus (crested newt), 413, 414
 tsetse fly, 33
Turborotalita quinqueloba, 436
Turdus merula (blackbird), 63
Turdus philomelos (thrush), 63
Typha latifolia, 167
Typha spp., 167

Cambridge University Press

978-0-521-54931-8 - Dispersal Ecology

Edited by James M. Bullock, Robert E. Kenward and Rosie S. Hails

Index

[More information](#)

458 INDEX

- unweighted pair group method with arithmetic averages (UPGMA), insect molecular marker studies, 39
- uredospores, 397
 - long-distance wind-borne dispersal, 405
 - virulent mutations dispersal, 399, 401
- Uria aalge* (common guillemot), 81
- Uria lomvia* (Brünnich's guillemot), 81
- Ursos arctos* (bear), 413, 414
- vector maps, 61, 62
- vertebrates
 - population dynamics, 237–52
 - spatially explicit population models (SEPMs), 327–46
 - see also* land vertebrates dispersal
- vertical looking radar (VLR)
 - insect dispersal studies, 27, 30–1, 33–8, 36
 - diurnal periodicity, 37, 37
 - target identification, 38
 - insect pest migrations monitoring, 35
- vicariance, isolation by distance differentiation, 102–4, 103
- Viola*, 7
- viruses, long-distance dispersal, 354
- visible markers, land vertebrates, 66
 - artificial, 52
 - natural, 51
 - volunteer sampling networks, 59
- visual stimuli, 239
 - Microtus oeconomus* dispersal in experimental system, 244, 246
 - small insect flight behaviour, 115, 120, 124
 - aphid, 125
 - fig wasp, 125
- volatile attractants
 - aphid chemosensory abilities, 125–6, 127
 - fig tree, 127–8, 129
 - fig wasp chemosensory ability, 125, 126, 129
 - moth pheromones, 126, 126
 - small insect flight behaviour, 115–16, 116, 120, 124, 125–8
- volunteer network observations, 59, 60
- Vulpes vulpes* (fox), 56, 336
- Vulpia fasciculata*, 417, 423
- Wahlund effect
 - Cristatella mucedo*, 201, 212
 - resting propagule banks, 210
- water vole (*Arvicola terrestris*), 414
- western gull (*Larus occidentalis*), 147
- wheat yellow rust disease *see Puccinia striiformis* f.sp. *tritici*
- wheat yellow rust disease-resistance gene introgression, 398–9
- white tailed sea eagle (*Haliaeetus albicilla*), 59, 423
- wind-dispersed fungal spores, 397
 - biotrophic fungi, 398, 402, 407
 - epidemiological study in China, 403, 404
 - long-distance spread of virulent clones, 399–401, 400, 401
- wind-dispersed insects, 25, 113–29
 - small insect flight behaviour, 113, 114–16
 - direction/distance trade-offs, 114–15, 115, 129
 - flight speed/body size relationship, 114, 114
 - plant volatile compound responses, 115–16, 116
 - stimuli for ending flight, 115–16
 - visual stimuli responses, 115
- suction trap monitoring, 29, 30
- vertical looking radar (VLR) monitoring, 37
- wind-dispersed seeds, 296
 - chemical composition, 153–7
 - fat storage, 153, 154–6, 156
 - determining source, 16
 - dispersal distance, 153
 - dispersal effectiveness measures, 153
 - local dispersal, 5, 7
 - long-distance dispersal, 5, 6, 8, 11, 12, 13–14
 - primary dispersal, 19
 - secondary dispersal on snow, 18
 - persistence in soil, 157–60, 160
 - terminal velocity data, 153
- trees
 - data collection methods, 19–21, 20
 - empirically fitted dispersal models, 9, 9–11, 10, 11, 12, 13–14
 - mechanistic dispersal models, 14
- wing tags, 53
- winter dispersal, 55, 56
- Wipfelkrankheit (tree top disease), 177
- wire tags, binary-coded, 53
- zooplankton, 206–10
 - dispersal capabilities, 206, 207
 - ecological studies, 207, 208, 209, 211
 - gene flow, 206, 207, 209–10
 - temporal, 211
 - local adaptation, 211, 212
 - population genetic studies, 206–7
 - sexual reproduction, 212
- zoospores, 397