

C H A P T E R 1

What Is Enumerative Combinatorics?

1.1 How to Count

The basic problem of enumerative combinatorics is that of counting the number of elements of a finite set. Usually we are given an infinite class of finite sets S_i where i ranges over some index set I (such as the nonnegative integers \mathbb{N}), and we wish to count the number $f(i)$ of elements of each S_i “simultaneously.” Immediate philosophical difficulties arise. What does it mean to “count” the number of elements of S_i ? There is no definitive answer to this question. Only through experience does one develop an idea of what is meant by a “determination” of a counting function $f(i)$. The counting function $f(i)$ can be given in several standard ways:

1. The most satisfactory form of $f(i)$ is a completely explicit closed formula involving only well-known functions, and free from summation symbols. Only in rare cases will such a formula exist. As formulas for $f(i)$ become more complicated, our willingness to accept them as “determinations” of $f(i)$ decreases. Consider the following examples.

1.1.1 Example. For each $n \in \mathbb{N}$, let $f(n)$ be the number of subsets of the set $[n] = \{1, 2, \dots, n\}$. Then $f(n) = 2^n$, and no one will quarrel about this being a satisfactory formula for $f(n)$.

1.1.2 Example. Suppose n men give their n hats to a hat-check person. Let $f(n)$ be the number of ways that the hats can be given back to the men, each man receiving one hat, so that no man receives his own hat. For instance, $f(1) = 0$, $f(2) = 1$, $f(3) = 2$. We will see in Chapter 2 that

$$f(n) = n! \sum_{i=0}^n (-1)^i / i!. \quad (1)$$

This formula for $f(n)$ is not as elegant as the formula in Example 1.1.1, but for lack of a simpler answer we are willing to accept (1) as a satisfactory formula. In

2 Chapter 1 What Is Enumerative Combinatorics?

fact, once the derivation of (1) is understood (using the Principle of Inclusion–Exclusion), every term of (1) has an easily understood combinatorial meaning. This enables us to “understand” (1) intuitively, so our willingness to accept it is enhanced. We also remark that it follows easily from (1) that $f(n)$ is the nearest integer to $n!/e$. This is certainly a simple explicit formula, but it has the disadvantage of being “non-combinatorial”; that is, dividing by e and rounding off to the nearest integer has no direct combinatorial significance.

1.1.3 Example. Let $f(n)$ be the number of $n \times n$ matrices \mathbf{M} of zeros and ones such that every row and column of \mathbf{M} has three ones. For example, $f(0) = f(1) = f(2) = 0$, $f(3) = 1$. The most explicit formula known at present for $f(n)$ is

$$f(n) = 6^{-n} \sum \frac{(-1)^\beta n!^2 (\beta + 3\gamma)! 2^\alpha 3^\beta}{\alpha! \beta! \gamma!^2 6^\gamma} \quad (2)$$

where the sum is over all $(n+2)(n+1)/2$ solutions to $\alpha + \beta + \gamma = n$ in non-negative integers. This formula gives very little insight into the behavior of $f(n)$, but it does allow one to compute $f(n)$ much faster than if only the combinatorial definition of $f(n)$ were used. Hence with some reluctance we accept (2) as a “determination” of $f(n)$. Of course if someone were later to prove $f(n) = (n-1)(n-2)/2$ (rather unlikely), then our enthusiasm for (2) would be considerably diminished.

1.1.4 Example. There are actually formulas in the literature (“nameless here for evermore”) for certain counting functions $f(n)$ whose evaluation requires listing all (or almost all) of the $f(n)$ objects being counted! Such a “formula” is completely worthless.

2. A recurrence for $f(i)$ may be given in terms of previously calculated $f(j)$ ’s, thereby giving a simple procedure for calculating $f(i)$ for any desired $i \in I$. For instance, let $f(n)$ be the number of subsets of $[n]$ that do not contain two consecutive integers. For example, for $n = 4$ we have the subsets $\emptyset, \{1\}, \{2\}, \{3\}, \{4\}, \{1, 3\}, \{1, 4\}, \{2, 4\}$, so $f(4) = 8$. It is easily seen that $f(n) = f(n-1) + f(n-2)$ for $n \geq 2$. This makes it trivial, for example, to compute $f(20)$. On the other hand, it can be shown that

$$f(n) = \frac{1}{\sqrt{5}} (\tau^{n+2} - \bar{\tau}^{n+2}),$$

where $\tau = \frac{1}{2}(1 + \sqrt{5})$, $\bar{\tau} = \frac{1}{2}(1 - \sqrt{5})$. This is an explicit answer, but because it involves irrational numbers it is a matter of opinion whether it is a better answer than the recurrence $f(n) = f(n-1) + f(n-2)$.

3. An estimate may be given for $f(i)$. If $I = \mathbb{N}$, this estimate frequently takes the form of an *asymptotic formula* $f(n) \sim g(n)$, where $g(n)$ is a “familiar function.” The notation $f(n) \sim g(n)$ means that $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$. For instance, let $f(n)$ be the function of Example 1.1.3. It can be shown that

$$f(n) \sim e^{-2} 36^{-n} (3n)!.$$

For many purposes this estimate is superior to the “explicit” formula (2).

4. The most useful but most difficult to understand method for evaluating $f(i)$ is to give its *generating function*. We will not develop in this chapter a rigorous abstract theory of generating functions, but will instead content ourselves with an informal discussion and some examples. Informally, a generating function is an “object” that represents a counting function $f(i)$. Usually this object is a *formal power series*. The two most common types of generating functions are *ordinary generating functions* and *exponential generating functions*. If $I = \mathbb{N}$, then the ordinary generating function of $f(n)$ is the formal power series

$$\sum_{n \geq 0} f(n)x^n,$$

while the exponential generating function of $f(n)$ is the formal power series

$$\sum_{n \geq 0} f(n)x^n/n!.$$

(If $I = \mathbb{P}$, the positive integers, then these sums begin at $n = 1$.) These power series are called “formal” because we are not concerned with letting x take on particular values, and we ignore questions of convergence and divergence. The term x^n or $x^n/n!$ merely marks the place where $f(n)$ is written. If $F(x) = \sum_{n \geq 0} a_n x^n$, we call a_n the *coefficient* of x^n in $F(x)$ and write

$$a_n = [x^n] F(x) \quad \text{or} \quad a_n = F(x)|_n.$$

Similarly we can deal with generating functions of several variables, such as

$$\sum_{l \geq 0} \sum_{m \geq 0} \sum_{n \geq 0} f(l, m, n) x^l y^m z^n / n!$$

(which may be considered as “ordinary” in the indices l, m and “exponential” in n), or even of infinitely many variables. In this latter case every term should involve only finitely many of the variables.

Why bother with generating functions if they are merely another way of writing a counting function? The answer is that we can perform various natural operations on generating functions that have a combinatorial significance. For instance, we can add two generating functions (in one variable) by the rule

$$\left(\sum_{n \geq 0} a_n x^n \right) + \left(\sum_{n \geq 0} b_n x^n \right) = \sum_{n \geq 0} (a_n + b_n) x^n$$

or

$$\left(\sum_{n \geq 0} \frac{a_n x^n}{n!} \right) + \left(\sum_{n \geq 0} \frac{b_n x^n}{n!} \right) = \sum_{n \geq 0} \frac{(a_n + b_n) x^n}{n!}.$$

Similarly, we can multiply generating functions according to the rule

$$\left(\sum_{n \geq 0} a_n x^n \right) \left(\sum_{n \geq 0} b_n x^n \right) = \sum_{n \geq 0} c_n x^n,$$

where $c_n = \sum_{i=0}^n a_i b_{n-i}$, or

$$\left(\sum_{n \geq 0} \frac{a_n x^n}{n!} \right) \left(\sum_{n \geq 0} \frac{b_n x^n}{n!} \right) = \sum_{n \geq 0} \frac{d_n x^n}{n!},$$

4 Chapter 1 What Is Enumerative Combinatorics?

where $d_n = \sum_{i=0}^n \binom{n}{i} a_i b_{n-i}$, with $\binom{n}{i} = n!/i!(n-i)!$. Note that these operations are just what we would obtain by treating generating functions as if they obeyed the ordinary laws of algebra, such as $x^i x^j = x^{i+j}$. These operations coincide with the addition and multiplication of functions when the power series converge for appropriate values of x , and they obey such familiar laws of algebra as associativity and commutativity of addition and multiplication, distributivity of multiplication over addition, and cancellation of multiplication (i.e., if $F(x)G(x) = F(x)H(x)$ and $F(x) \neq 0$, then $G(x) = H(x)$). In fact, the set of all formal power series $\sum_{n \geq 0} a_n x^n$ with complex coefficients a_n forms a (commutative) integral domain under the operations just defined. This integral domain is denoted by $\mathbb{C}[[x]]$. (Actually, $\mathbb{C}[[x]]$ is a very special type of integral domain. For readers with some familiarity with algebra, we remark that $\mathbb{C}[[x]]$ is a principal ideal domain and therefore a unique factorization domain. In fact, every ideal of $\mathbb{C}[[x]]$ has the form (x^n) for some $n \geq 0$. From the viewpoint of commutative algebra, $\mathbb{C}[[x]]$ is a one-dimensional complete regular local ring. These general algebraic considerations will not concern us here; rather we will discuss from an elementary viewpoint the properties of $\mathbb{C}[[x]]$ that will be useful to us.) Similarly, the set of formal power series in the m variables x_1, \dots, x_m (where m may be infinite) is denoted $\mathbb{C}[[x_1, \dots, x_m]]$ and forms a unique factorization domain (though not a principal ideal domain for $m \geq 2$).

It is primarily through experience that the combinatorial significance of the algebraic operations of $\mathbb{C}[[x]]$ or $\mathbb{C}[[x_1, \dots, x_m]]$ is understood, as well as the problem of whether to use ordinary or exponential generating functions (or various other kinds discussed in later chapters). In Section 3.15, we will explain to some extent the combinatorial significance of these operations, but even then experience is indispensable.

If $F(x)$ and $G(x)$ are elements of $\mathbb{C}[[x]]$ satisfying $F(x)G(x) = 1$, then we (naturally) write $G(x) = F(x)^{-1}$. (Here 1 is short for $1 + 0x + 0x^2 + \dots$.) It is easy to see that $F(x)^{-1}$ exists (in which case it is unique) if and only if $a_0 \neq 0$, where $F(x) = \sum_{n \geq 0} a_n x^n$. One commonly writes “symbolically” $a_0 = F(0)$, even though $F(x)$ is not considered to be a function of x . If $F(0) \neq 0$ and $F(x)G(x) = H(x)$, then $G(x) = F(x)^{-1}H(x)$. More generally, the operation $^{-1}$ satisfies all the familiar laws of algebra, provided it is only applied to power series $F(x)$ satisfying $F(0) \neq 0$. For instance, $(F(x)G(x))^{-1} = F(x)^{-1}G(x)^{-1}$, $(F(x)^{-1})^{-1} = F(x)$, and so on. Similar results hold for $\mathbb{C}[[x_1, \dots, x_n]]$.

1.1.5 Example. Let $(\sum_{n \geq 0} \alpha^n x^n)(1 - \alpha x) = \sum_{n \geq 0} c_n x^n$, where α is a non-zero complex number. Then by the definition of power series multiplication,

$$c_n = \begin{cases} 1, & n = 0 \\ \alpha^n - \alpha(\alpha^{n-1}) = 0, & n \geq 1. \end{cases}$$

Hence $\sum_{n \geq 0} \alpha^n x^n = (1 - \alpha x)^{-1}$, which can also be written

$$\sum_{n \geq 0} \alpha^n x^n = \frac{1}{1 - \alpha x}.$$

This formula comes as no surprise; it is simply the formula (in a formal setting) for summing a geometric series.

Example 1.1.5 provides a simple illustration of a general principle that, informally speaking, states that if we have an identity involving power series that is valid when the power series are regarded as functions (so that the variables are sufficiently small complex numbers), then this identity continues to remain valid when regarded as an identity among formal power series, *provided* the operations involved in the formulas are well-defined for formal power series. It would be unnecessarily pedantic for us to state a precise form of this principle here, since the reader should have little trouble justifying in any particular case the formal validity of our manipulations with power series. We will give several examples throughout this section to illustrate this contention.

1.1.6 Example. The identity

$$\left(\sum_{n \geq 0} x^n/n! \right) \left(\sum_{n \geq 0} (-1)^n x^n/n! \right) = 1 \quad (3)$$

is valid at the function-theoretic level (it states that $e^x e^{-x} = 1$) and is well-defined as a statement involving formal power series. Hence (3) is a valid formal power series identity. In other words (equating coefficients of $x^n/n!$ on both sides of (3)), we have

$$\sum_{k=0}^n (-1)^k \binom{n}{k} = \delta_{0n}. \quad (4)$$

To justify this identity directly from (3), we may reason as follows. Both sides of (3) converge for all $x \in \mathbb{C}$, so we have

$$\sum_{n \geq 0} \left(\sum_{k=0}^n (-1)^k \binom{n}{k} \right) \frac{x^n}{n!} = 1, \quad \text{for all } x \in \mathbb{C}.$$

But if two power series in x represent the same function $f(x)$ in a neighborhood of 0, then these two power series must agree term-by-term, by a standard elementary result concerning power series. Hence (4) follows.

1.1.7 Example. The identity

$$\sum_{n \geq 0} (x+1)^n/n! = e \sum_{n \geq 0} x^n/n!$$

is valid at the function-theoretic level (it states that $e^{x+1} = e \cdot e^x$), but does not make sense as a statement involving formal power series. There is no *formal* procedure for writing $\sum_{n \geq 0} (x+1)^n/n!$ as a member of $\mathbb{C}[[x]]$.

Although the expression $\sum_{n \geq 0} (x+1)^n/n!$ does not make sense *formally*, there are nevertheless certain infinite processes that can be carried out formally in $\mathbb{C}[[x]]$. (These concepts extend straightforwardly to $\mathbb{C}[[x_1, \dots, x_m]]$, but for simplicity we consider only $\mathbb{C}[[x]]$.) To define these processes, we need to put some additional structure on $\mathbb{C}[[x]]$ —namely, the notion of *convergence*.

6 Chapter 1 What Is Enumerative Combinatorics?

From an algebraic standpoint, the definition of convergence is inherent in the statement that $\mathbb{C}[[x]]$ is *complete* in a certain standard topology that can be put on $\mathbb{C}[[x]]$. However, we will assume no knowledge of topology on the part of the reader and will instead give a self-contained, elementary treatment of convergence.

If $F_1(x), F_2(x), \dots$ is a sequence of formal power series, and if $F(x) = \sum_{n \geq 0} a_n x^n$ is another formal power series, we say by definition that $F_i(x)$ *converges* to $F(x)$ as $i \rightarrow \infty$, written $F_i(x) \rightarrow F(x)$, provided that for all $n \geq 0$ there is a number $\delta(n)$ such that the coefficient of x^n in $F_i(x)$ is a_n whenever $i \geq \delta(n)$. In other words, for every n the sequence

$$\left\{ \text{coeff}_n F_1(x), \text{coeff}_n F_2(x), \dots \right.$$

of complex numbers eventually becomes constant with value a_n . An equivalent definition of convergence is the following. Define the *degree* of a non-zero formal power series $F(x) = \sum_{n \geq 0} a_n x^n$, denoted $\deg F(x)$, to be the least integer n such that $a_n \neq 0$. Note that $\deg F(x)G(x) = \deg F(x) + \deg G(x)$. Then $F_i(x)$ converges if and only if $\lim_{i \rightarrow \infty} \deg(F_{i+1}(x) - F_i(x)) = \infty$.

We now say that an infinite sum $\sum_{j \geq 0} F_j(x)$ has the value $F(x)$ provided that $\sum_{j=0}^i F_j(x) \rightarrow F(x)$. A similar definition is made for the infinite product $\prod_{j \geq 1} F_j(x)$. To avoid unimportant technicalities we assume that in any infinite product $\prod_{j \geq 1} F_j(x)$, each factor $F_j(x)$ satisfies $F_j(0) = 1$. For instance, let $F_j(x) = a_j x^j$. Then for $i \geq n$, the coefficient of x^n in $\sum_{j=0}^i F_j(x)$ is a_n . Hence $\sum_{j \geq 0} F_j(x)$ is just the power series $\sum_{n \geq 0} a_n x^n$. Thus we can think of the formal power series $\sum_{n \geq 0} a_n x^n$ as actually being the “sum” of its individual terms. The proofs of the following two elementary results are left to the reader.

1.1.8 Proposition. The infinite series $\sum_{j \geq 0} F_j(x)$ converges if and only if $\lim_{j \rightarrow \infty} \deg F_j(x) = \infty$. \square

1.1.9 Proposition. The infinite product $\prod_{j \geq 1} (1 + F_j(x))$, where $F_j(0) = 0$, converges if and only if $\lim_{j \rightarrow \infty} \deg F_j(x) = \infty$. \square

It is essential to realize that in evaluating a convergent series $\sum_{j \geq 0} F_j(x)$ (or similarly a product $\prod_{j \geq 1} F_j(x)$), the coefficient of x^n for any given n can be computed using only *finite* processes. For if j is sufficiently large, say $j > \delta(n)$, then $\deg F_j(x) > n$, so that

$$\sum_{n \geq 0} F_j(x) = \sum_{n \geq 0}^{\delta(n)} F_j(x).$$

The latter expression involves only a *finite* sum.

The most important combinatorial application of the notion of convergence is to the idea of power series composition. If $F(x) = \sum_{n \geq 0} a_n x^n$ and $G(x)$ are formal power series with $G(0) = 0$, define the *composition* $F(G(x))$ to be the infinite sum $\sum_{n \geq 0} a_n G(x)^n$. Since $\deg G(x)^n = n \cdot \deg G(x) \geq n$, we see by Proposition 1.1.8 that $F(G(x))$ is well-defined as a *formal* power series. We also see why an expression such as e^{1+x} does not make sense formally; namely, the infinite series

1.1 How to Count

7

$\sum_{n \geq 0} (1+x)^n/n!$ does not converge in accordance with the above definition. On the other hand, an expression like e^{e^x-1} makes good sense formally, since it has the form $F(G(x))$ where $F(x) = \sum_{n \geq 0} x^n/n!$ and $G(x) = \sum_{n \geq 1} x^n/n!$.

1.1.10 Example. If $F(x) \in \mathbb{C}[[x]]$ satisfies $F(0) = 0$, then we can *define* for any $\lambda \in \mathbb{C}$ the formal power series

$$(1 + F(x))^\lambda = \sum_{n \geq 0} \binom{\lambda}{n} F(x)^n, \quad (5)$$

where $\binom{\lambda}{n} = \lambda(\lambda-1)\cdots(\lambda-n+1)/n!$. In fact, we may regard λ as an indeterminate and take (5) as the definition of $(1 + F(x))^\lambda$ as an element of $\mathbb{C}[[x, \lambda]]$ (or of $\mathbb{C}[\lambda][[x]]$; that is, the coefficient of x^n in $(1 + F(x))^\lambda$ is a *polynomial* in λ). All the expected properties of exponentiation are indeed valid, such as $(1 + F(x))^{\lambda+\mu} = (1 + F(x))^\lambda (1 + F(x))^\mu$ (regarded as an identity in the ring $\mathbb{C}[[x, \lambda, \mu]]$, or in the ring $\mathbb{C}[[x]]$ where one takes $\lambda, \mu \in \mathbb{C}$).

If $F(x) = \sum_{n \geq 0} a_n x^n$, define the *formal derivative* $F'(x)$ (also denoted $\frac{dF}{dx}$ or $DF(x)$) to be the formal power series $\sum_{n \geq 0} n a_n x^{n-1} = \sum_{n \geq 0} (n+1) a_{n+1} x^n$. It is easy to check that all the familiar laws of differentiation that are well-defined formally continue to be valid for formal power series. In particular

$$(F + G)' = F' + G'$$

$$(FG)' = F'G + FG'$$

$$F(G(x))' = G'(x)F'(G(x)).$$

We thus have a theory of *formal calculus* for formal power series. The usefulness of this theory will become apparent in subsequent examples. We first give an example of the use of the formal calculus that should shed some additional light on the validity of manipulating formal power series as if they were actual functions of x .

1.1.11 Example. Suppose $F(0) = 1$, and let $G(x)$ be the unique power series satisfying

$$G'(x) = F'(x)/F(x), \quad G(0) = 0. \quad (6)$$

From the function-theoretic viewpoint we can “solve” (6) to obtain $F(x) = \exp G(x)$, where by definition $\exp G(x) = \sum_{n \geq 0} G(x)^n/n!$. Since $G(0) = 0$ everything is well-defined formally, so (6) should remain equivalent to $F(x) = \exp G(x)$ even if the power series for $F(x)$ converges only at $x = 0$. How can this assertion be justified without actually proving a combinatorial identity? Let $F(x) = 1 + \sum_{n \geq 1} a_n x^n$. From (6) we can compute explicitly $G(x) = \sum_{n \geq 1} b_n x^n$, and it is quickly seen that each b_n is a *polynomial* in finitely many of the a_i ’s. It then follows that if $\exp G(x) = 1 + \sum_{n \geq 1} c_n x^n$, then each c_n will also be a polynomial in finitely many of the a_i ’s, say $c_n = p_n(a_1, a_2, \dots, a_m)$, where m depends on n . Now we know that $F(x) = \exp G(x)$ provided $1 + \sum_{n \geq 1} a_n x^n$ converges. If two Taylor series convergent in some neighborhood of the origin represent the same function, then their coefficients coincide. Hence $a_n = p_n(a_1, a_2, \dots, a_m)$ provided $1 + \sum_{n \geq 1} a_n x^n$

8 Chapter 1 What Is Enumerative Combinatorics?

converges. Thus the two polynomials a_n and $p_n(a_1, a_2, \dots, a_m)$ agree in some neighborhood of the origin of \mathbb{C}^m , so they must be equal. (It is well-known that if two complex polynomials in m variables agree in some open set of \mathbb{C}^m , then they are identical.) Since $a_n = p_n(a_1, a_2, \dots, a_m)$ as polynomials, the identity $F(x) = \exp G(x)$ continues to remain valid for *formal* power series.

There is an alternative method for justifying the formal solution $F(x) = \exp G(x)$ to (6), which may appeal to topologically inclined readers. Given $G(x)$ with $G(0) = 0$, define $F(x) = \exp G(x)$ and consider a map $\phi: \mathbb{C}[[x]] \rightarrow \mathbb{C}[[x]]$ defined by $\phi(G(x)) = G'(x) - \frac{F'(x)}{F(x)}$. One easily verifies the following: (a) if G converges in some neighborhood of 0 then $\phi(G(x)) = 0$; (b) the set \mathcal{G} of all power series $G(x) \in \mathbb{C}[[x]]$ that converge in some neighborhood of 0 is dense in $\mathbb{C}[[x]]$, in the topology defined above (in fact, the set $\mathbb{C}[x]$ of polynomials is dense); and (c) the function ϕ is continuous in the topology defined above. From this it follows that $\phi(G(x)) = 0$ for all $G(x) \in \mathbb{C}[[x]]$ with $G(0) = 0$.

We now present various illustrations in the manipulation of generating functions. Throughout we will be making heavy use of the principle that formal power series can be treated as if they were functions.

1.1.12 Example. Find a simple expression for the generating function $F(x) = \sum_{n \geq 0} a_n x^n$, where $a_0 = a_1 = 1$, $a_n = a_{n-1} + a_{n-2}$ if $n \geq 2$. We have

$$\begin{aligned} F(x) &= \sum_{n \geq 0} a_n x^n = 1 + x + \sum_{n \geq 2} a_n x^n \\ &= 1 + x + \sum_{n \geq 2} (a_{n-1} + a_{n-2}) x^n \\ &= 1 + x + x \sum_{n \geq 2} a_{n-1} x^{n-1} + x^2 \sum_{n \geq 2} a_{n-2} x^{n-2} \\ &= 1 + x + x(F(x) - 1) + x^2 F(x). \end{aligned}$$

Solving for $F(x)$ yields $F(x) = 1/(1 - x - x^2)$.

1.1.13 Example. Find a simple expression for the generating function $F(x) = \sum_{n \geq 0} a_n x^n/n!$, where $a_0 = a_1 = 1$, $a_n = a_{n-1} + (n-1)a_{n-2}$ if $n \geq 2$. We have

$$\begin{aligned} F(x) &= \sum_{n \geq 0} a_n x^n/n! \\ &= 1 + x + \sum_{n \geq 2} a_n x^n/n! \\ &= 1 + x + \sum_{n \geq 2} (a_{n-1} + (n-1)a_{n-2}) x^n/n!. \end{aligned} \tag{7}$$

Let $G(x) = \sum_{n \geq 2} a_{n-1} x^n/n!$ and $H(x) = \sum_{n \geq 2} (n-1)a_{n-2} x^n/n!$. Then $G'(x) = \sum_{n \geq 2} a_{n-1} x^{n-1}/(n-1)! = F(x) - 1$, and $H'(x) = \sum_{n \geq 2} a_{n-2} x^{n-1}/(n-2)! = xF(x)$. Hence if we differentiate (7) we obtain

$$F'(x) = 1 + (F(x) - 1) + xF(x) = (1 + x)F(x).$$

The unique solution to this differential equation satisfying $F(0) = 1$ is $F(x) = \exp(x + \frac{1}{2}x^2)$. (As shown in Example 1.1.11, solving this differential equation is a purely formal procedure.)

1.1 How to Count

9

1.1.14 Example. Let $\mu(n)$ be the Möbius function of number theory; that is, $\mu(1) = 1$, $\mu(n) = 0$ if n is divisible by the square of an integer greater than one, and $\mu(n) = (-1)^r$ if n is the product of r distinct primes. Find a simple expression for the power series

$$F(x) = \prod_{n \geq 1} (1 - x^n)^{-\mu(n)/n}. \quad (8)$$

First let us make sure that $F(x)$ is well-defined as a formal power series. We have by Example 1.1.10 that

$$(1 - x^n)^{-\mu(n)/n} = \sum_{i \geq 0} \binom{-\mu(n)/n}{i} (-1)^i x^{in}.$$

Note that $(1 - x^n)^{-\mu(n)/n} = 1 + H(x)$, where $\deg H(x) = n$. Hence by Proposition 1.1.9 the infinite product (8) converges, so $F(x)$ is well-defined. Now apply log to (8). In other words, form $\log F(x)$, where

$$\log(1 + x) = \sum_{n \geq 1} (-1)^{n-1} x^n / n,$$

the power series expansion for the natural logarithm. We obtain

$$\begin{aligned} \log F(x) &= \log \prod_{n \geq 1} (1 - x^n)^{-\mu(n)/n} \\ &= \sum_{n \geq 1} \log(1 - x^n)^{-\mu(n)/n} \\ &= - \sum_{n \geq 1} \frac{\mu(n)}{n} \log(1 - x^n) \\ &= - \sum_{n \geq 1} \frac{\mu(n)}{n} \sum_{i \geq 1} \left(-\frac{x^{in}}{i} \right). \end{aligned}$$

The coefficient of x^m in the above power series is

$$\frac{1}{m} \sum_{d|m} \mu(d),$$

where the sum is over all positive integers d dividing m . It is well-known that

$$\frac{1}{m} \sum_{d|m} \mu(d) = \begin{cases} 1, & m = 1 \\ 0, & \text{otherwise.} \end{cases}$$

Hence $\log F(x) = x$, so $F(x) = e^x$. Note that the derivation of this miraculous formula involved only *formal* manipulations.

1.1.15 Example. Find the unique sequence $a_0 = 1, a_1, a_2, \dots$ of real numbers satisfying

$$\sum_{k=0}^n a_k a_{n-k} = 1 \quad (9)$$

for all $n \in \mathbb{N}$. The trick is to recognize the left-hand side of (9) as the coefficient

10 Chapter 1 What Is Enumerative Combinatorics?

of x^n in $(\sum_{n \geq 0} a_n x^n)^2$. Letting $F(x) = \sum_{n \geq 0} a_n x^n$, we then have

$$F(x)^2 = \sum_{n \geq 0} x^n = 1/(1-x).$$

Hence $F(x) = (1-x)^{-1/2} = \sum_{n \geq 0} \binom{-1/2}{n} (-1)^n x^n$, so

$$\begin{aligned} a_n &= (-1)^n \binom{-\frac{1}{2}}{n} = (-1)^n \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right) \cdots \left(-\frac{2n-1}{2}\right)}{n!} \\ &= \frac{1 \cdot 3 \cdot 5 \cdots (2n-1)}{2^n n!}. \end{aligned}$$

Now that we have discussed the manipulation of formal power series, the question arises as to the advantages of using generating functions to represent a counting function $f(n)$. Why, for instance, should a formula such as

$$\sum_{n \geq 0} f(n) x^n / n! = \exp\left(x + \frac{x^2}{2}\right) \quad (10)$$

be regarded as a “determination” of $f(n)$? Basically, the answer is that there are many standard, routine techniques for extracting information from generating functions. Generating functions are frequently the most concise and efficient way of presenting information about their coefficients. For instance, from (10) an experienced enumerative combinatorialist can tell at a glance the following:

1. A simple recurrence for $f(n)$ can be found by differentiation. Namely, we obtain

$$\sum_{n \geq 0} f(n) x^{n-1} / (n-1)! = (1+x) e^{x+(x^2/2)} = (1+x) \sum_{n \geq 0} f(n) x^n / n!.$$

Equating coefficients of $x^n / n!$ yields

$$f(n+1) = f(n) + n f(n-1), \quad n \geq 1.$$

2. An explicit formula for $f(n)$ can be obtained from $e^{x+(x^2/2)} = e^x e^{x^2/2}$. Namely,

$$\begin{aligned} \sum_{n \geq 0} f(n) x^n / n! &= e^x e^{x^2/2} = \left(\sum_{n \geq 0} \frac{x^n}{n!} \right) \left(\sum_{n \geq 0} \frac{x^{2n}}{2^n n!} \right) \\ &= \left(\sum_{n \geq 0} \frac{x^n}{n!} \right) \left(\sum_{n \geq 0} \frac{(2n)!}{2^n n!} \frac{x^{2n}}{(2n)!} \right), \end{aligned}$$

so that

$$f(n) = \sum_{\substack{i=0 \\ i \text{ even}}}^n \binom{n}{i} \frac{i!}{2^{i/2} (i/2)!}.$$

3. Regarded as a function of a complex variable, $\exp(x + \frac{x^2}{2})$ is a nicely-behaved entire function, so that standard techniques from the theory of asymptotic estimates can be used to estimate $f(n)$. As a first approximation, it is routine