

Cambridge University Press
0521604893 - Hopf Algebras
Eiichi Abe
Frontmatter
[More information](#)

CAMBRIDGE TRACTS IN MATHEMATICS

General editors

H. BASS, H. HALBERSTAM, J. F. C. KINGMAN
J. E. ROSEBLADE, C. T. C. WALL

Hopf algebras

Cambridge University Press
0521604893 - Hopf Algebras
Eiichi Abe
Frontmatter
[More information](#)

EIICHI ABE

Professor of Mathematics, University of Tsukuba

Hopf algebras

translated by Hisae Kinoshita and Hiroko Tanaka

CAMBRIDGE UNIVERSITY PRESS

CAMBRIDGE

LONDON NEW YORK NEW ROCHELLE

MELBOURNE SYDNEY

Cambridge University Press
0521604893 - Hopf Algebras
Eiichi Abe
Frontmatter
[More information](#)

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS
The Edinburgh Building, Cambridge CB2 2RU, UK
40 West 20th Street, New York NY 10011-4211, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
Ruiz de Alarcón 13, 28014 Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

<http://www.cambridge.org>

Copyright © 1977 by Eiichi Abe
Translated by Hiroko Tanaka and Hisae Kinoshita

This English language edition © 1980 Cambridge University Press

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

Originally published in Japanese by Iwanami Shoten, Publishers, Tokyo, 1977
English edition first published by Cambridge University Press, 1980
First paperback edition 2004

A catalogue record for this book is available from the British Library

Library of Congress cataloguing in publication data

Abe, Eiichi, 1927-
Hopf algebras.
(Cambridge tracts in mathematics)
Includes index.
I. Hopf algebras. I. Title. II. Series.
QA613.8.A213 512'.55 79-50912

ISBN 0 521 22240 0 hardback
ISBN 0 521 60489 3 paperback

Contents

	<i>Page</i>
Preface	vii
Notation	xi
1 Modules and algebras	
1. Modules	1
2. Algebras over a commutative ring	16
3. Lie algebras	24
4. Semi-simple algebras	40
5. Finitely generated commutative algebras	49
2 Hopf algebras	
1. Bialgebras and Hopf algebras	53
2. The representative bialgebras of semigroups	65
3. The duality between algebras and coalgebras	78
4. Irreducible bialgebras	90
5. Irreducible cocommutative bialgebras	107
3 Hopf algebras and representations of groups	
1. Comodules and bimodules	122
2. Bimodules and bialgebras	137
3. Integrals for Hopf algebras	144
4. The duality theorem	158
4 Applications to algebraic groups	
1. Affine k -varieties	163
2. Affine k -groups	170
3. Lie algebras of affine algebraic k -groups	186
4. Factor groups	201
5. Unipotent groups and solvable groups	210
6. Completely reducible groups	222

vi	<i>Contents</i>
5 Applications to field theory	
1. K/k -bialgebras	229
2. Jacobson's theorem	243
3. Modular extensions	252
Appendix: Categories and functors	
A.1 Categories	264
A.2 Functors	267
A.3 Adjoint functors	271
A.4 Representable functors	271
A.5 \mathcal{G} -groups and \mathcal{G} -cogroups	274
References	278
Index	281

Preface

Let G be a finite group and k a field. The set $A = \text{Map}(G, k)$ of all functions defined on G with values in k becomes a k -algebra when we define the scalar product and the sum and product of functions by

$$\begin{aligned}(\alpha f)(x) &= \alpha f(x), & (f + g)(x) &= f(x) + g(x), \\ (fg)(x) &= f(x)g(x), & f, g \in A, \alpha \in k, x \in G.\end{aligned}$$

In general, a k -algebra A can be characterized as a k -linear space A together with two k -linear maps

$$\mu: A \otimes_k A \rightarrow A, \quad \eta: k \rightarrow A,$$

which satisfy axioms corresponding to the associative law and the unitary property respectively. If we identify $A \otimes_k A$ with $\text{Map}(G \times G, k)$ where $A = \text{Map}(G, k)$, and if the operations of G are employed in defining the k -linear maps

$$\Delta: A \rightarrow A \otimes_k A, \quad \varepsilon: A \rightarrow k$$

respectively by $\Delta f(x \otimes y) = f(xy)$, $\varepsilon f = f(e)$ for $x, y \in G$ and where e is the identity element of G , then Δ and ε become homomorphisms of k -algebras having properties which are dual to μ and η respectively. In general, a k -linear space A with k -linear maps $\mu, \eta, \Delta, \varepsilon$ defined as above is called a k -bialgebra. Furthermore, we can define a k -linear endomorphism of $A = \text{Map}(G, k)$

$$S: A \rightarrow A, \quad (Sf)(x) = f(x^{-1}), \quad f \in A, \quad x \in G$$

such that the equalities

$$\mu(1 \otimes S)\Delta = \mu(S \otimes 1)\Delta = \eta \circ \varepsilon$$

hold. A k -bialgebra on which we can define a k -linear map S as above is called a k -Hopf algebra. Thus, a k -Hopf algebra is an algebraic system which simultaneously admits structures of a k -algebra as well

as its dual, where these two structures are related by a certain specific law. For a finite group G , its group ring kG over a field k is the dual space of the k -linear space $A = \text{Map}(G, k)$ where its k -algebra structure is given by the dual k -linear maps of Δ and ε . Moreover, kG admits a k -Hopf algebra structure when we take the dual k -linear maps of μ , η , and S . In other words, kG is the dual k -Hopf algebra of $\text{Map}(G, k)$.

If, for instance, we replace the finite group G in the above argument by a topological group and k by the field of real numbers or the field of complex numbers, or if we take G to be an algebraic group over an algebraically closed field k and A is replaced by the k -algebra of all continuous representative functions or of all regular functions over G , then A turns out to be a k -Hopf algebra in exactly the same manner. These algebraic systems play an important role when studying the structure of G . Similarly, a k -Hopf algebra structure can be defined naturally on the universal enveloping algebra of a k -Lie algebra. The universal enveloping algebra of the Lie algebra of a semi-simple algebraic group turns out to be (in a sense) the dual of the Hopf algebra defined above. These constitute some of the most natural examples of Hopf algebras. The general structure of such algebraic systems has recently become a focus of interest in conjunction with its applications to the theory of algebraic groups or the Galois theory of purely inseparable extensions, and a great deal of research is currently being conducted in this area.

It has only been since the late 1960s that Hopf algebras, as algebraic systems, became objects of study from an algebraic standpoint. However, beginning with the research on representation theory through the use of the representative rings of Lie groups by Hochschild–Mostow (*Ann. of Math.* **66** (1957), 495–542, **68** (1958), 295–313) and in subsequent studies (cf. references [4], [7], [8] for Chapter 3), Hopf algebras have been taken up extensively as algebraic systems and also used in applications. On the other hand, in algebraic topology, the concept of graded Hopf algebras was derived at an even earlier date from an axiomatization of the works of H. Hopf relating to topological properties of Lie groups (cf. *Ann. of Math.* **42** (1941), 22–52). Hence the name, ‘Hopf algebra’. For instance, if G is a connected Lie group, the cohomology group $H^*(G)$ or the homology

Preface

ix

group $H_*(G)$ of G with coefficients in a field k has a multiplication or a dual multiplication induced by the diagonal map $d : G \rightarrow G \times G$ ($d(x) = (x, x)$, $x \in G$) and is moreover a commutative k -algebra. In addition, the map $m : G \times G \rightarrow G$ defined via the Lie group multiplication induces the maps

$$\Delta : H^*(G) \rightarrow H^*(G) \otimes H^*(G) \quad \text{or} \quad \Delta : H_*(G) \rightarrow H_*(G) \otimes H_*(G)$$

which respectively make $H^*(G)$ or $H_*(G)$ a k -Hopf algebra. These are graded Hopf algebras and such structures can be defined also for H -spaces, and have been generalized by A. Borel, J. Leary, and others (cf. A. Borel: *Ann. of Math.* **57** (1953), 115–207). For details on the algebraic properties of Hopf algebras, the reader is referred to J. Milnor–J. C. Moore: *Ann of Math.* **81** (1965), 211–64.

This book has as its main objective algebraic applications of (non-graded) Hopf algebras, and an attempt has been made to acquaint the reader with the elementary properties of Hopf algebras with a minimal amount of preliminary knowledge. There is an excellent work on the subject, *Hopf algebras* by Sweedler (Benjamin, 1969), to which this book owes a great deal. But here the central theme will revolve around applications of Hopf algebras to the representative rings of topological groups and to algebraic groups. Some recent developments on the subject have also been incorporated.

This book consists of five chapters and an appendix. Chapter 1 is preparation for the central subject of the book and deals with some basic properties of modules and algebras which become necessary in the sequel. Some simple properties of groups, fields, and topological spaces have been used without proofs. With regard to the solutions to some of the exercises and in the treatment of finitely generated commutative algebras, where I have either omitted or condensed a number of the proofs of well-known theorems, the reader is asked to refer to other texts. In Chapter 2, coalgebras, bialgebras, and Hopf algebras are defined, and their fundamental properties are outlined. Chapter 3 takes up the structure theorem of bimodules and properties of Hopf algebras similar to those of the representative rings of topological groups. The proofs of the existence and uniqueness of the integral of commutative Hopf algebras are due to J. B. Sullivan. In Chapter 4, fundamental properties of affine algebraic groups are

proven through an application of the theory of Hopf algebras. The construction of factor groups, the proof of the decomposition theorem of solvable groups, and the proofs of theorems related to completely reducible groups are respectively due to Mitsuhiro Takeuchi, J. B. Sullivan, and M. E. Sweedler. Although properties pertaining to the representation of affine algebraic groups can be described satisfactorily by such an approach, there are drawbacks as well as merits in pursuing a general theory in this context. For instance, it may distort the overall view of the development of the subject matter. It should be interesting, for instance, to attempt an exposition of the theory of Borel subgroups, which is an important topic in the theory of linear algebraic groups, not touched upon in this book. Chapter 5 contains a brief glance at the Galois theory of purely inseparable extensions. Although there are numerous texts dealing with Galois theory, here I have presented a treatment of the subject by D. Winter. The appendix contains a sketch of the theory of categories, where notions such as categories, functors and, for a category \mathcal{C} , \mathcal{C} -groups and \mathcal{C} -cogroups, which are used throughout this book, are defined.

Since commutative Hopf algebras are precisely the commutative algebras which represent affine group schemes, applications of Hopf algebras also naturally develop along these lines. The reader will do well to refer to other texts on this matter.

The writing of this book was suggested to me by Professor Nagayoshi Iwahori of the Faculty of Science of the University of Tokyo, to whom I would like to express my most sincere gratitude. I am also very grateful to Professors Yukio Doi of Fukui University and Mitsuhiro Takeuchi of the University of Tsukuba for many valuable words of advice which they have given me in preparing the manuscript.

Eiichi Abe
Tokyo, Japan

Notations

Sets

$a \in M$ or $M \ni a$	a is an element of the set M
$M \subset N$ or $N \supset M$	$a \in M$ implies $a \in N$, namely, M is contained in N
$M \cup N; \bigcup_{\lambda \in \Lambda} M_\lambda$	the union of M and N ; the union of M_λ ($\lambda \in \Lambda$)
$M \cap N; \bigcap_{\lambda \in \Lambda} M_\lambda$	the intersection of M and N ; the intersection of M_λ ($\lambda \in \Lambda$)
\emptyset	the empty set
\mathbb{Z}	the set of all integers
\mathbb{N}	the set of all natural numbers

Maps

$f : M \rightarrow N$	a map from the set M to the set N
$x \mapsto y$	the image $f(x)$ of $x \in M$ under the map f is $y \in N$

When $f(M) = N$, f is called a surjection.

If $x, x' \in M$, $x \neq x'$ implies $f(x) \neq f(x')$, then f is called an injection.

For $M' \subseteq M$, we denote the restriction of f to M' by $f|_{M'}$.

If $f : M \rightarrow N$, $g : N \rightarrow P$ are two maps, the composition of f and g is written $g \circ f$, $x \mapsto g(f(x))$.

Logical symbols $A \Rightarrow B$

If proposition A holds, then
proposition B holds

 $A \Leftrightarrow B$ $A \Rightarrow B$ and $B \Rightarrow A$ $\forall x \in M$ for any element x of M