

## Index

- acid phosphatase, in *Erwinia chrysanthemi*, 36–7  
 actinomycetes, 28–9  
 aerial environment, 77–84  
 aerosol dispersal of bacteria, 77–9  
 agglutination of bacteria  
   by plant cell wall components, 140–1  
   in serology, 65  
*Agrobacterium radiobacter*  
   bacteriocin production, 263–4  
   role in biological control, 290–2  
*Agrobacterium rhizogenes*, Ri plasmid, 257–8  
*Agrobacterium tumefaciens*  
   biological control, 290–2  
   crown gall disease, 10, 183–5  
   gene induction, 248, 259–63  
   IAA genes, 245, 261–3  
   motility and chemotaxis, 25, 112–13  
   PHB inclusions, 45  
   pigment, 58  
   rhizosphere colonisation, 85, 112–13  
   serology, 63  
   Ti plasmid, 258–63  
   vascular migration, 123  
 agrocin-84, 290–1, 263–4  
 agropine, 186  
 amino-terminal recognition of secretory proteins,  
   36–7  
 antibiotics  
   bacterial resistance, 264–6, 277  
   biological control of disease, 283–7  
   chemical control of disease, 276  
   genetic control of resistance, 264–6  
 antibodies to plant pathogenic bacteria, 62–6, 142  
 autoradiography, 33, 164–5  
 auxin  
   genetic determination, 244–6, 259–63  
   synthesis, 206–8  
 auxotrophs, analysis of virulence, 233–5  
 avenulamin, 146  
 avirulence genes  
   cloning, 225–6  
   gene-for-gene hypothesis, 223–4  
   gene products, 226–9  
 avirulence mutants, 237–44  
 bacteria–plant associations  
   epiphytic associations, 79–86  
   general aspects, 1  
   hypersensitive reaction, 138–42, 163–4  
 bacterial antagonist, *see* biological control  
 bacterial antigens, 63  
 bacterial attachment  
   during the hypersensitive reaction, 163–4  
   role of fimbriae, 26  
 bacterial classification and nomenclature, 42–9  
 bacterial competition  
   general aspects, 108  
   on phylloplane, 97  
   in rhizosphere, 87–8  
 bacterial entry into plant tissue, 114–20  
 bacterial flagella  
   general aspects, 23–5  
   taxonomic significance, 44–5  
   *see also* motility  
 bacterial gene  
   evolutionary conservation, 247–8  
   identification and investigation, 212–22  
   library, 215–20  
   mapping, 222  
   regulation of expression, 248, 261–2  
   translation product, 222  
 bacterial genes, encoding  
   antibiotic resistance, 264–5  
   avirulence, 223–9  
   bacteriocin production, 263  
   bioluminescence, 266  
   cellulases, 239–41  
   chemotaxis control proteins, 25  
   disease virulence, 233–48  
   extracellular polysaccharides, 243–4  
   hypersensitivity and pathogenicity, 229–33  
   ice nucleation, 91, 249–50  
   pectolytic enzymes, 238–9  
   plant hormones, 244–7  
   proteases, 241–3  
   protein secretion, 35, 37

- siderophore metabolism, 250–2
- toxins, 236–8
- tumour induction, 257–63
- bacterial leaf blight of rice, *see* *Xanthomonas campestris* pv. *oryzae*
- bacterial membrane
  - fatty acids, 70–1
  - ice nucleation complexes, 91
  - lipids, 21
  - pigments, 58–9
  - proteins, 21
  - siderophore uptake, 95, 251
  - see also* recognition, surface receptors
- bacterial migration during infection, 109–14
- bacterial pigments in taxonomy, 58–61
- bacterial plant pathogen
  - comparison to fungal pathogens, 4–6
  - criteria as disease agent, 6–7
  - fine structure, 13–40
- bacterial populations
  - on aerial plant surfaces, 79–84
  - in planta*, after inoculation, 128–30
- bacterial races, 48–9
- bacterial species, 43
- bacterial structure
  - capsule, 22–3
  - cell surface complex, 18
  - changes during infection, 163
  - mesosomes, 18
  - nucleoid, 15–16
  - peripheral ribosomal area, 17–18
  - periplasmic space, 21–2
  - R-bodies, 18
- bacterial taxonomy, 41–76
- bacterial virulence and disease, 172–211
- bactericides, 273–6
- bacteriocin
  - diversity, 287–8
  - experimental uses, 288–9
  - genetic determination, 263–4
  - mode of action, 288
  - production in environment, 289
  - role in biological control, 287–92
- bacteriophage
  - bacterial receptor sites, 27
  - biological control agent, 299–300
  - DNA vector, 217–20
  - lysotype, 61–2
- Bdellovibrio*, use as biological control agent, 300–2
- bean, race/cultivar differentiation, 48
- beta-galactosidase, *in Erwinia chrysanthemi*, 36–7
- biological control
  - antagonist mode of action, 283–302
  - antagonist selection, 279–80
  - general aspects, 278–302
  - in rhizosphere, 87
- bioluminescence, use in bacterial monitoring, 266
- biotypes, 49
- biovars, 49, 57, 184
- black rot of crucifers, *see* *Xanthomonas campestris* pv. *campestris*
- blackleg of potato, *see* *Erwinia carotovora*, *Erwinia chrysanthemi*
- breeding for resistance, 305–8
- capillary assay, of chemotaxis, 109–10
- capsule, 22–3
- carboxyl terminal recognition of secretory proteins, 35
- carboxylic acid bacterial toxins, 197
- carotenoid pigment, bacterial, 58, 60
- cations
  - determination by X-ray microanalysis, 31–2
  - inhibitors and cofactors of pectolytic enzymes, 198–9
  - leakage during the compatible reaction, 154, 157–8
  - leakage during the hypersensitive reaction, 154–7
  - siderophore uptake of ferric ions, 93–7
  - synthesis and activation of bacterial toxins, 195–6
- cell surface membranes, protein partitioning, 35–7
- cell surface complex, 18–27
- cellulase enzymes
  - disease virulence factors, 202
  - secretion in *Erwinia chrysanthemi*, 203
- certification
  - of disease-free plants, 310
  - of disease-free seed, 51, 310–12
- chemical analysis of bacterial cells, 30–3
- chemical control of disease, 273–8
- chemotaxis
  - during *Agrobacterium tumefaciens* infection, 262
  - general aspects, 24–5
  - measurement, 109–10
  - in rhizosphere, 85
  - role in plant infection, 111–14
- classification and nomenclature of bacteria, 42–9
- Clavibacter michiganensis* subsp. *sepedonicum*
  - control in potato, 312
  - ELISA detection, 65
- colonisation of rhizosphere, 85
- compartmentation of cell activities, 33–7
- compatibility
  - evolution, 3–4
  - see also* disease
- compatible reaction
  - changes in plant cell plasmalemma, 154, 157–8
  - general aspects, 126–71
- computer identification, 71–2
- conjugation, DNA uptake by, 219–20
- coronatine, 191–2
- corky root disease of lettuce, 7–8, 304
- cosmid vector, 217–20
- cotton, resistance to *Xanthomonas campestris* pv. *malvacearum*, 306
- crop monoculture, 273
- crown gall disease, *see* *Agrobacterium tumefaciens*

cultivar resistance, screening and evaluation, 308–9

cultural practices  
 aerosol dispersal of bacteria, 79  
 disease control, 312–13

cytokinin synthesis  
 genetic control, 245–7  
 virulence factor, 206–8

differential growth media, 53

disease  
 bacterial pathogenicity factors, 173  
 bacterial virulence factors, 173  
 biological control, 278–302  
 causal agents, 4–7  
 chemical control, 273–8  
 diagnosis, 50–1  
 environmental effects, 172  
 genetical control, 303–9  
 induction, 172–4  
 insect transmission, 97–103  
 major types, 174–86  
 sanitary control, 309–13  
 specificity of induction by toxins, 189–90

disease-specific genes, *see* virulence genes

DNA  
 preparation of fraction, 214–16  
 probe, 67–9  
 restriction fragment profile analysis, 66–7  
 sequence analysis, 222  
 vector molecules, 215–18

electron probe X-ray microanalysis, 31–2

electrolyte leakage from plant cell, 155–6

elicitors  
 avirulence gene products, 227–9  
 induction of the hypersensitive reaction, 144–5

enzyme-linked immunosorbent assay (ELISA), 63, 65

epiphytic bacteria  
 bacteria–plant associations, 1, 80–1  
 induction of frost damage, 92–3  
 plant infection, 107–14  
 populations, 79–80, 93  
 seasonal variation, 81–4  
 siderophore production, 96–7  
 sites of multiplication, 81–4

*Erwinia amylovora*  
 annual growth cycle, 81–2  
 biological control, 280–3, 297–8  
 chemotaxis, 114  
 economic importance, 10–11  
 extracellular polysaccharide production, 204–5  
 floral colonisation, 80, 82–3  
 floral infection, 118–20  
 growth media, 53–4  
 host resistance, 304  
 immunogold labelling, 34  
 nomenclature, 42  
 streptomycin resistance, 264  
 vector transmission, 98–100

*Erwinia carotovora*  
 bacteriocin production, 290  
 biological control by predation, 302  
 DNA probe, 68  
 economic importance, 10–11  
 in irrigation water, 87  
 lenticel entry into potato, 118  
 soft rot disease, 181–2  
 suppression in rhizosphere, 295

*Erwinia chrysanthemi*  
 cellulase production, 203  
 pectate lyase genes, 239  
 pectolytic enzyme production, 200–1  
 protease production, 203  
 protease genes, 241–2  
 protein secretion, 35–7  
 siderophore production, 252–2  
 soft rot disease, 180–2

*Erwinia herbicola*  
 antibiotic production, 285–7  
 auxin synthesis, 246  
 biocontrol agent, 280–3, 297–8  
 chemotaxis, 114  
 ice-nucleation activity, 89–90, 97  
 ice-nucleation genes, 250

*Erwinia stewartii*  
 extracellular polysaccharide genes, 243–4  
 extracellular polysaccharide production, 23, 205–6  
 insect transmission, 101

*Escherichia coli*, siderophore production, 94–5

evolution  
 bacterial pathogenicity, 2–3  
 conservation of *hrp* genes, 231–2  
 conservation of pathogenicity and virulence genes, 247  
 plant resistance, 3–4

extracellular enzymes  
 genetical determination, 238–43  
 production by bacteria, 197–204

extracellular polysaccharide  
 bacterial capsule, 22–3  
 genetical determination, 243–4  
 toxins, 197  
 virulence factors, 202–6

fastidious prokaryotes  
 insect transmission, 102–3  
 selective media, 56  
 vascular migration, 123  
 vascular transport, 122–3  
 vascular wilt disease, 178–80

fatty acid composition, 70–1

fimbriae, 25–7

fireblight disease  
 annual cycle, 82  
 biological control, 280–3  
 economic importance, 11  
 genetical resistance, 304  
*see also Erwinia amylovora*

flagella, 23–5

- floral infection, 118–20  
 fluorescent pseudomonads  
   LOPAT classification, 56  
   PHB inclusions, 45  
   pigments, 58–9  
   rhizosphere colonisation, 85–6  
   specific growth media, 53–4  
   suppression of soil pathogens, 292–5  
   *see also* ice nucleation, siderophore production,  
   *Pseudomonas syringae*, *Pseudomonas fluorescens*  
 freezing spectra of plant pathogenic bacteria, 89  
 frost damage by bacteria  
   general aspects, 92–3  
   leading to bacterial infection, 115–17  
   use in competition assessment, 97  
   *see also* ice nucleation
- gel electrophoresis, 69–70  
 gel immunodiffusion, 64–5  
 gene-for-gene hypothesis, 223–4  
 genes, *see* bacterial genes, plant genes  
 genetic analysis of plant pathogenic bacteria, 212–72  
 genetic control of plant disease, 303–9  
 genomic fingerprinting, 67  
 Gram-negative bacteria  
   cell envelope, 19–22  
   general fine structure, 15  
   protein secretion, 35–7  
   siderophore production, 94–5  
   taxonomy, 43–4  
 Gram-positive bacteria  
   cell envelope, 19–20  
   taxonomy, 43–4  
 growth media in diagnosis, 53–6
- halo blight disease, 175–6, 311  
 hawthorn, floral infection by *Erwinia amylovora*, 118–20  
 histochemical stains, 33  
 host/pathogen specificity, 129, 174  
 hydathode entry of bacteria, 115, 118  
 hypersensitive reaction  
   activation of plant genes, 158–9  
   bacteria–plant recognition, 139–42  
   bacterial changes, 161–5  
   biochemical interactions, 142–52  
   general aspects, 126–71  
   genetic determination, 223–33  
   *hrp* genes, 229–33  
   factors affecting induction and expression, 135–7  
   induction, 133–4  
   plant cell changes, 138, 152–61, 166  
   resistance mechanism, 165–6  
   test for pathogenicity, 131–2  
   time course, 134–5  
 HR, *see* hypersensitive reaction
- ice nucleation  
   genes, 91, 249–50  
   phenotypic aspects, 88–93  
 identification of bacteria  
   epiphytes, 51  
   from infected tissue, 50  
   in physical environment, 52  
 immobilisation of bacteria, during the HR, 162–4  
 immunocytochemistry, 33–4, 65–6  
 incompatibility  
   bacteria–plant interaction, 126–71  
   evolution, 3–4  
 indoleacetic acid, *see* auxin  
 infection of plant tissue, 107–25  
 insect vectors, 97–103  
 isoflavonoid phytoalexins, 149–50  
 invertebrate transmission of disease, 97–103  
*in vitro* systems, experimental uses, 132–3
- Koch's postulates, 6–8
- lectin agglutination of bacteria, 140–1  
 lenticel entry of bacteria, 115–18  
 light, effects on phytoalexin production, 150–1, 166
- lipids  
   bacterial membrane, 21  
   fatty acid composition in bacteria, 70–1  
   ice nucleation sites, 91  
   peroxidation in plant cell plasmalemma, 153–5
- lipopolysaccharide  
   insertion in outer membrane, 19–20  
   surface vesicles, 27
- log-normal distribution of bacteria  
   on phylloplane, 79  
   in rhizosphere, 86
- LOPAT grouping of fluorescent pseudomonads  
   HR test, 131  
   taxonomy, 56  
 lysotype, 49, 61–2
- mesosomes, 18  
 membrane, *see* bacterial membrane, plant cell  
   plasmalemma
- meristem culture of potato, 312  
 microelectrode measurement of membrane potential, 155, 157  
 micropropagated plants, 132–3  
 MLO, *see* mycoplasma-like organisms
- mollicutes  
   taxonomic status, 91  
   *see also* mycoplasma-like organisms,  
   *Spiroplasma*
- momilactone, 146  
 motility of plant pathogenic bacteria  
   during the hypersensitive reaction, 165  
   flagellar activity, 24–5  
   measurement, 109–10  
   migration within plant tissue, 121–2  
   role in plant infection, 110–11  
   role in root colonisation, 85
- monoclonal antibodies, 64  
 multiple gene action, 307–9

mycoplasma-like organisms  
 fine structure, 29–31  
 insect transmission, 103  
 motility and chemotaxis, 24–5  
 nutritional requirements, 46  
 taxonomy, 45–6, 68  
*see also* mollicutes

necrotic diseases, 174–80

nopaline, 186, 259–63

nucleoid, of bacterial cell, 15–17

numerical taxonomy, 71–2

octopine, 186, 259–63

oncogenic bacteria, 174–5, 183–6

opines  
 genetic determination, 257–63  
 synthesis by transformed plant cells, 185

opportunistic pathogens  
 HR test for pathogenicity, 131  
 soft-rot disease, 181, 183

oxygen  
 effect on phytoalexin production, 151–2  
 induction of plant cell damage, 153, 159

PAGE, *see* polyacrylamide gel electrophoresis

parasitic bacteria, 1, 3

pathogenicity  
 criteria, 6–7  
 definition, 173  
 evolution, 2  
 inoculation responses, 126–8, 131–2  
 testing, 52, 80, 127

pathovars, 46–8

pectate lyase  
 secretion in *Erwinia chrysanthemi*, 35–7  
*see also* pectolytic enzymes

pectolytic enzymes  
 genetic determination, 238–9  
 virulence factors, 198–201

PEL, *see* pectate lyase

pepper, race/cultivar differentiation, 48, 306–7

peripheral ribosomal area of bacterial cell, 17–18

phage, *see* bacteriophage

PHB, *see* polyhydroxybutyrate

phaseollin, 146

phaseolotoxin, 192–3

phloem-limited bacteria  
 insect transmission, 100, 103  
 vascular disease, 179

phylloplane  
 epiphytic bacteria, 79–81  
 siderophore production, 96–7

phyllosphere, 79

phytoalexins  
 detection and quantitation, 147  
 diversity, 145  
 elicitor monitoring, 144–5  
 environmental effects, 150–2  
 in plant resistance, 165–6  
 synthesis, 149–50

phytopathogen  
 definition, 4  
 inoculation test, 126–7

pigments  
 in bacterial taxonomy, 58–61  
 siderophores, 93–7

pili, *see* fimbriae

plant cell death  
 during disease, 128–31, 174–5  
 in the hypersensitive reaction, 128–31, 138,  
 152–3, 160–1, 166

plant cell fine structure  
 mesophyll cell changes during halo blight  
 disease, 176–7  
 pathological changes during the hypersensitive  
 reaction, 160–1  
 potato tuber cell changes during soft rot  
 disease, 181–3

plant cell plasmalemma  
 effect of bacterial toxins, 194  
 effect of plant pathogenic bacteria, 153–8  
 electrolyte leakage, 155–6  
 lipid peroxidation, 153–5  
 membrane potential, 155  
 water efflux and plasmolysis, 156

plant cell wall components  
 elicitors, 144–5, 149  
 recognition factors, 140–1

plant disease, *see* disease

plant genes, encoding  
 bacterial attachment sites, 262–3  
 resistance response, 158–9

plant growth-promoting rhizobacteria  
 role in biological control, 294–5  
 soil competition, 87–8

plant hormones  
 role in tumour diseases, 206–8  
 bacterial genes, 244–7, 259  
*see also* auxin, cytokinin

plant pathogenic bacteria  
 economically important species, 9–10  
 relation to crop monoculture, 7–8, 273  
*see also* parasitic bacteria, phytopathogen

plant resistance  
 evolution, 3  
 response to bacteria, 126–71  
*see also* resistance response genes, resistance  
 genes

plant signal molecules, 248

plasmid  
 electron microscope observation, 254–5  
 isolation and characterisation, 252–4  
 in plant pathogenic bacteria, 252–66  
 vector molecules, 215–18  
*see also* cosmid, plasmid genes, Ri plasmid, Ti  
 plasmid

plasmid genes, encoding  
 antibiotic resistance, 264–6  
 bacteriocin production, 263  
 pathogenicity and virulence factors, 255–63

polyacrylamide gel electrophoresis, 69–70

- polyclonal antibodies, 63–7  
 polygalacturonase,  
   mode of action, 198–201  
   secretion of *Erwinia chrysanthemi*, 35–7  
 polygenes, in disease resistance, 307–9  
 polyhydroxybutyrate inclusions, 17–18, 45  
 polypeptide analysis, 69–70  
 polysaccharide, *see* extracellular polysaccharide  
 potato  
   biological control of *Erwinia carotovora*, 295, 302  
   lenticel entry of bacteria, 118  
   meristem culture, 312  
   phytoalexins, 146, 151–2  
   resistance to soft rot, 304, 307–8  
   soft rot diseases, 11, 180–2  
 predatory antagonists, 87–8, 299–302  
 protease  
   genes, 241–3  
   secretion in *Erwinia chrysanthemi*, 35–7, 203  
   virulence factor, 202  
 protein partitioning by bacterial surface  
   membranes, 35–7  
 proton influx during the hypersensitive reaction, 155–7  
 proton pump, bacterial effect on, 154–5, 158  
 prototroph, analysis of virulence, 233–5  
*Pseudomonas andropogonis*, toxin production, 196–7  
*Pseudomonas fluorescens*  
   colonisation of rhizosphere, 85  
   ice nucleation activity, 90–1  
   ice nucleation genes, 249–50  
   non-pathogenic response after leaf infiltration, 126–7  
   siderophore production, 96  
   use as biological control agent in rhizosphere, 294–5  
   *see also* fluorescent pseudomonads  
*Pseudomonas melophthora*, insect transmission, 101–2  
*Pseudomonas putida*  
   colonisation of rhizosphere, 85  
   use as biological control agent in rhizosphere, 294–5  
   *see also* fluorescent pseudomonads  
*Pseudomonas savastanoi* subsp. *savastanoi*  
   cytokinin genes, 245, 247  
   indoleacetic acid genes, 244–6  
   insect transmission, 102  
   secretion of plant hormones, 207–8  
   tumour disease, 185–6  
*Pseudomonas solanacearum*  
   attachment and incompatibility, 138–9  
   avirulent mutants in biological control, 298–300  
   biovars, 57  
   cellulase genes, 241  
   cellulase secretion, 202  
   extracellular polysaccharide, 205  
   fimbriae, 25  
   major plant pathogen, 10  
   races, 49  
   surface membrane proteins, 69  
   virulence mutants, 235  
*Pseudomonas syringae*  
   electron microscopy, 16  
   epiphytic populations, 67, 81  
   ice nucleation activity, 90–3, 97  
   ice nucleation genes, 249–50  
   site competition between strains, 298  
   toxin production, 191–6  
   use as biological control agent, 280–3  
   *see also* fluorescent pseudomonads  
*Pseudomonas syringae* pv. *glycinea*  
   cloning and analysis of *avrA* gene, 225–8  
   motility and plant infection, 110–12, 121–2  
   race-cultivar interactions, 223–4  
*Pseudomonas syringae* pv. *mors-prunorum*,  
   epiphyte, 107  
   numerical taxonomy, 72  
   pathovar status, 47  
*Pseudomonas syringae* pv. *papulans*, antibiotic  
   resistance, 264  
*Pseudomonas syringae* pv. *phaseolicola*  
   attachment to surfaces, 26  
   bacteriophage receptor sites, 27  
   biochemical differentiation, 57  
   certification of bean seed, 311  
   DNA probe, 68–9  
   halo blight disease, 175–6  
   *hrp* genes, 231  
   phylloplane population, 82–4  
   phytoalexin induction, 149  
   race-cultivar interactions, 48  
   surface vesicles, 27  
   toxin production, 190–3  
*Pseudomonas syringae* pv. *syringae*  
   biochemical differentiation, 57  
   certification of seed, 311  
   certification of tomato transplants, 310  
   induction of frost damage, 92–3  
   minimum threshold level, 108  
   numerical taxonomy, 72  
   pathovar status, 47  
   phylloplane populations, 82–4  
   siderophore production on phylloplane, 96–7  
   surface pathogenicity factor, 141–2  
   toxin production, 193–5  
   *see also* fluorescent pseudomonads  
*Pseudomonas syringae* pv. *tabaci*  
   elemental composition, 31–2  
   host resistance, 305–6  
   mutagenesis of toxin genes, 237–8  
   occurrence of flagella during disease  
     development, 121–2  
   toxin production, 191, 195–6  
   uptake of Ni<sup>63</sup>, 34  
*Pseudomonas syringae* pv. *tomato*  
   aerosol/rain dispersal, 78  
   certification of tomato transplants, 310  
   DNA probe, 68  
   epiphytic survival, 60

pyoverdines  
 bacterial pigments, 58–9  
 genetic determinat, 251  
 pyridine-derivatives, bacterial pigments, 58

races, of plant pathogenic bacteria, 48–9  
 race-specific resistance, 128  
 R-bodies, 18  
 rain dispersal of bacteria, 77–9  
 receptors, *see* surface receptors  
 recognition  
 activation of the hypersensitive reaction, 139–42  
 of amino and carboxyl terminals during protein secretion, 35–6  
 of avirulence gene products, 226–7  
 prevention by extracellular polysaccharide, 203

resistance genes  
 determining incompatibility, 223–4  
 in biological control, 303–9  
*see also* cultivar resistance

resistance response genes, 158–9  
 rhizobitoxine, 196–7  
 rhizosphere  
 bacterial competition and predation, 87–8  
 colonisation, 84–5  
 pathogen populations and disease, 86  
 siderophore production, 96, 294–5  
 specificity of bacteria-plant associations, 85–6

R1 plasmid, 258

rice  
 assessment of cultivar resistance, 309  
 bacterial leaf blight, 9, 10  
 bacterial entry via hydathodes, 118  
*see also* *Xanthomonas campestris* pv. *oryzae*

rishitin, 146, 151–2  
 RNA-DNA probe, 69  
 root, *see* rhizosphere

sanitary procedures for disease control, 309–13

saprophyte  
 chemotactic response, 114  
 leaf inoculation response, 126–8

seasonal variation in epiphytic populations, 81–4

seedborne pathogens  
 certification of seed, 310–12  
 detection and identification, 51–2

selective growth media, 53–6  
 serotype, 49, 62–6  
 siderophore  
 general features, 93–6  
 genetic control, 250–2  
*in vitro* production, 293–4  
 phylloplane production, 96–7  
 rhizosphere production, 96, 294  
 role in biological control, 292–5

signal peptide sequences, 36  
 site competition, in biological control, 296–9  
 soft rot diseases, 180–3  
 soil/water environment, 84–8  
 somaclonal variation, 305

somatic hybridisation, in genetical control of disease, 307–8

soybean, race/cultivar differentiation, 48, 223–4

spheroplast formation, 30

*Spiroplasma*  
 fine structure, 29  
 insect transmission, 103  
 selective growth medium, 55–6  
 taxonomy, 45–6  
*see also* mollicutes

sterilisation of equipment, 313

stomatal entry of bacteria, 115, 117–18

*Streptomyces*, 28–9

surface receptors  
 chemotaxis, 24, 261–2  
 membrane proteins, 21  
 recognition during the hypersensitive reaction, 140–2  
 siderophore uptake, 95  
 sites for attachment to plant surfaces, 26  
 sites for viral attachment, 27, 251  
*see also* bacterial membrane, recognition

syringomycin/syringotoxin, 193–5

tabtoxin, 195–6, 237–8

tagetitoxin, 192

Ti plasmid, 185, 245, 257–63

tobacco  
 leaf inoculation, 126–8  
 pathogen races, 48  
 resistance to wildfire disease, 305–6  
 tolaasin, 195

toxaasin, 195

toxins, produced by plant pathogenic bacteria  
 bacterial resistance to toxins, 190–1  
 characterisation of individual toxins, 191–7  
 criteria as virulence factors, 187  
 genetic determination, 236–8  
 specificity and function, 189–90  
 structure and biosynthesis, 188–9

transduction, 218–20

transduction in bacteria, 219

transfection of bacterial cells, 218–19

transformation of bacterial cells, 218–19

transformation of plant cells by *Agrobacterium tumefaciens*, 184–5, 257–63

transposon mutagenesis, 37, 221

triparental mating, 220, 228

tumour diseases  
 general characteristics, 183–6  
 genetic determination of pathogenicity, 244–7, 257–63

vascular migration of bacteria, 122–3

vascular wilt and yellows diseases, 178–80

vector molecules, 215–20

virulence  
 definition, 173  
 factors causing disease symptoms, 173–208

virulence genes, 229, 233–48

- wild fire disease, *see Pseudomonas syringae* pv. *tabaci*
- wilt disease, 178–9
- wound entry of bacteria, 115–16
- Xanthomonas campestris*
- epiphytic survival, 81
  - ice nucleation activity, 92–3
  - plasmid DNA restriction fragments, 67
  - polypeptide analysis, 69
  - seed contamination, 312
- Xanthomonas campestris* pv. *campestris*
- black rot of crucifers, 180–1
  - seed transmission, 311–12
- Xanthomonas campestris* pv. *citri*
- biotypes, 49, 67
  - DNA restriction fragment profile analysis, 67
  - isozymes, 70
  - major pathogen, 10
  - rain dispersal, 77–8
- Xanthomonas campestris* pv. *malvacearum*
- elicitation of phytoalexin production, 147–8
  - induction of the hypersensitive reaction, 150–1
- Xanthomonas campestris* pv. *oryzae*
- assessment of rice cultivar resistance, 309
  - geographic spread, 9
  - lysotypes, 61–2
  - major pathogen, 10
- Xanthomonas campestris* pv. *vesicatoria*
- entry via stomata, 117–18
  - rain dispersal, 78
  - resistance in pepper, 306–7
- xanthomonadin pigment, 58
- xylem-limited bacteria
- insect transmission, 100, 102
  - selective medium, 55–6
  - vascular wilt disease, 178–9
- Xylella fastidiosa*, *see* xylem-limited bacteria
- yellowing disease, 179–80