
Author index

Note: with references having more than three authors, the names of all authors appear in this index, but only the first authors appear in the in-text citations.

- Aaronson, D., 231
 Abayomi, K., 542
 Achen, C., 194, 484
 Afshartous, D., 485
 Agodini, R., 229
 Agresti, A., 17, 26, 105, 131
 Ainsley, A., 194
 Aitkin, M., 484
 Akaike, H., 105, 527
 Albert, A., 105
 Albert, J., 385
 Aldy, J., 198
 Almond, D., 231
 Amemiya, T., 131
 Anderson, D., 131
 Anderson, J., 105
 Angrist, J., 229, 230
 Ansolabehere, S., 197, 321
 Ashenfelter, O., 231
 Asimov, D., 563
 Atkinson, A., 74
- Bafumi, J., 74, 105, 321
 Baker, S., 321
 Balke, A., 231
 Ball, S., 194, 231
 Bannerjee, S., 298
 Barnard, J., 230, 297, 543
 Barry, S., 342
 Bates, D., 277, 297, 574
 Beal, S., 558
 Beck, N., 248
 Becker, R., 574
 Belin, T., 542
 Benane, S., 485
 Bentler, P., 485
 Berger, J., 276
 Berk, R., 49
 Berliner, L., 298
 Berman, B., 297
 Bernard, H., 342
 Bernardo, J., 277, 413
 Bertin, J., 562
- Besag, J., 194, 298
 Best, N., 105, 298, 527, 574
 Bird, C., 294, 298
 Blackman, C., 485
 Blalock, H., 74
 Blanchard, S., 527
 Blattner, F., 132
 Bloom, H., 229
 Bock, D., 26, 49
 Bock, R., 276
 Bogart, W., 231
 Bogatz, G., 194, 231
 Bolger, N., 194
 Boscardin, W., 297, 434
 Boshuizen, H., 542
 Bosker, R., 245, 247, 276, 454
 Box, G., 74, 276, 413, 454, 526
 Bradley, R., 132
 Bradlow, E., 231
 Brainard, J., 26
 Brewer, D., 485
 Bring, J., 74
 Brookmeyer, R., 194
 Brooks, S., 342, 373
 Brooks-Gunn, J., 49, 229
 Brown, C., 248
 Browne, W., 297, 485
 Browner, W., 26
 Bryk, A., 247, 276, 297, 454
 Buenconsejo, J., 229
 Buja, A., 563
 Burmaster, D., 26
 Bush, R., 165, 526
 Butler, M., 230
- Campbell, D., 194, 230
 Campbell, J., 233, 298
 Campbell, S., 194
 Card, D., 231
 Carlin, B., 105, 276, 298, 373, 413, 434
 Carlin, J., 26, 49, 105, 151, 165, 248, 276, 298, 346, 385, 386, 413, 434, 526, 574
 Carroll, R., 74, 230
 Casella, A., 131, 342
 Casella, G., 245, 484, 501
 Catchpole, E., 342

- Chambers, J., 49, 562, 574
 Chang, H., 526
 Chapman, R., 131
 Chay, K., 231
 Chen, M., 385
 Chib, S., 385
 Chipman, H., 298
 Clarke, B., 485
 Clayton, D., 342, 542
 Cleveland, W., 49, 97, 562
 Clinton, J., 133
 Clogg, C., 542
 Cochran, W., 26, 229, 292, 297, 454
 Cole, N., 276
 Collins, R., 386
 Congdon, P., 373, 574
 Cook, D., 563
 Cook, R., 8, 105, 485, 526
 Cook, S., 165, 434
 Cook, T., 194
 Costa, D., 198
 Coull, B., 17, 26
 Cox, D., 74, 194
 Cox, G., 197
 Crainiceanu, C., 230, 298
 Cramer, C., 105
 Criqui, M., 74
 Cromwell, B., 231
 Cui, Y., 298
- D'Agostino, R., 229
 Daniels, M., 297
 Datta, G., 298
 David, M., 542
 Davidian, M., 247
 Dawid, A., 194
 De Leeuw, J., 8, 245, 247, 248, 276, 297, 485
 De Veaux, R., 26, 49
 Dehejia, R., 229, 231
 Dellaportas, P., 385
 Dempster, A., 297, 413
 Dey, D., 526
 Diamond, A., 229
 Diffendal, G., 542
 Diggle, P., 231
 Dismukes, R., 297
 Dobson, A., 131
 Dodhia, R., 563
 Donohue, J., 321
 Dorman, P., 198
 Drake, C., 229
 Dreze, J., 512
 Du, J., 230
 DuMouchel, W., 277
 Duncan, D., 131
- Dunn, G., 542
 Dunne, A., 558
 Dunson, D., 298
 Dyke, G., 194
 Dynarski, M., 229
- Earls, F., 342
 Efron, B., 277, 413
 Ehrenberg, A., 563
 Ehrenberg, R., 485
 Elliott, D., 485
 Endicott, J., 229
 Erdos, P., 334
 Erikson, R., 133, 197, 298
 Evans, W., 230
 Everitt, B., 574
- Fagan, J., 131, 342
 Fair, R., 298
 Fay, R., 298, 542
 Felton, J., 51
 Fienberg, S., 131
 Finkelstein, M., 543
 Firth, D., 105
 Fisman, R., 322
 Forbes, A., 49
 Fournier, D., 574
 Fox, J., 49, 105, 484, 562, 574
 Frangakis, C., 194, 230
 Friedlander, D., 229
 Friedman, J., 132, 298
 Friedman, M., 132
 Friendly, M., 563
 Frolich, M., 229
- Garfinkel, I., 248, 543
 Gatsonis, C., 277
 Gawron, V., 297
 Gelfand, A., 298, 385, 413, 434, 526
 Gelman, A., 26, 49, 74, 105, 131, 151, 165, 194, 197, 230, 248, 276, 297–299, 321, 342, 346, 373, 385, 413, 434, 484, 501, 526, 542, 543, 563, 574
 Geoghebur, Y., 527
 George, E., 298
 Gerber, A., 233
 Geweke, J., 165, 434
 Gilks, W., 413, 527, 574
 Gilovich, T., 26
 Giltinan, D., 247
 Girosi, F., 298
 Glickman, M., 230, 277
 Goegebeur, Y., 105
 Goldstein, H., 247, 276, 454, 485
 Graddy, K., 230
 Graubard, B., 485
 Graves, T., 574

AUTHOR INDEX

603

- Green, B., 245, 501
 Greenland, S., 74, 231, 294, 298
 Greenstone, M., 231
 Guldemond, H., 454
 Gustafson, P., 298, 485

 Haavelmo, T., 230
 Hagstrom, P., 198
 Hahn, J., 230
 Hahn, T., 194
 Hale, R., 294, 298
 Hambleton, R., 321
 Hamermesh, D., 51
 Han, W., 49
 Handcock, M., 342
 Hansen, B., 229, 230
 Hanushek, E., 485
 Harrell, F., 74
 Harris, J., 277
 Hasbun, R., 229
 Hastie, T., 132, 298
 Hauer, E., 131
 Heckman, J., 194, 229, 231
 Heinze, G., 105
 Heisey, D., 454
 Heitjan, D., 153, 542
 Henderson, C., 276
 Herring, A., 298
 Herriot, R., 298
 Higdon, D., 194, 298
 Hill, B., 413
 Hill, C., 229
 Hill, J., 49, 194, 229, 230
 Hills, S., 385
 Hirano, K., 194, 229, 230
 Hodges, J., 298, 385, 434, 527
 Hoenig, J., 454
 Hoeting, J., 298
 Hoff, P., 342
 Honaker, J., 542
 Hoogerheide, L., 512
 Hoppin, J., 298
 House, D., 485
 Hox, J., 276
 Huang, Z., 197, 434
 Hunter, J., 454
 Hunter, W., 454
 Hurley, C., 563

 Ibrahim, J., 385
 Ichimura, H., 194, 229
 Imai, K., 131, 229
 Imbens, G., 194, 229, 230
 Iyengar, S., 322

 Jackman, S., 133, 321
 Jackson, C., 298
 Jackson, P., 276, 342
 Jacob, B., 230
 Jakulin, A., 105
 James, W., 277
 Jaynes, E., 413
 Jeffreys, H., 413
 Jenkyn, J., 194
 Joffe, M., 229
 Johnsen, E., 342
 Johnson, V., 342
 Jones, K., 485
 Joseph, A., 542

 Kahn, M., 198
 Kaldor, J., 342
 Kamenica, E., 322
 Kane, T., 458, 485
 Karagas, M., 230
 Karim, M., 342, 385
 Kashy, D., 194
 Kass, R., 297, 373, 413
 Katz, J., 197, 230, 248
 Kazarow, K., 194
 Kedar, O., 248
 Keller, M., 229
 Kelling, G., 342
 Kempthorne, O., 276
 Kendziorowski, C., 132
 Kenny, D., 194
 Kenward, M., 231
 Kerman, J., 151, 413, 574
 Killworth, P., 342
 King, G., 49, 151, 197, 229, 298, 485, 542
 Kirk, R., 501
 Kiss, A., 131, 321, 342, 484, 543
 Klebanov, P., 229
 Kleibergen, F., 512
 Kleiner, B., 49, 562
 Knook, D., 542
 Korn, E., 485
 Kramer, B., 321
 Krantz, D., 26, 198
 Kreft, I., 8, 245, 247, 248, 276, 297, 485
 Krueger, A., 230, 231
 Kutner, M., 49
 Kwan, E., 563

 Lahiri, P., 298
 LaLonde, R., 229, 231
 LaMotte, L., 245
 Lancaster, T., 230, 373
 Landwehr, J., 105
 Lane, P., 501
 Lange, K., 132
 Lavori, P., 229
 Leamer, E., 194
 Lechner, M., 229

- Lee, D., 230, 231
 Lefgren, L., 230
 Lenth, R., 454
 Lessafre, E., 105
 Levin, B., 543
 Levine, D., 231
 Levitt, S., 197
 Levy, M., 542
 Lewis, C., 276, 342
 Lewis, J., 386
 Leyland, A., 276
 Liang, K., 131
 Liaw, F., 229
 Liebman, J., 131, 321, 543
 Ligges, U., 574
 Lin, C., 198
 Linden, L., 69, 230
 Lindley, D., 277, 385, 413
 Linton, O., 229
 Little, R., 132, 542
 Little, T., 321
 Liu, C., 132, 385, 434, 542
 Lohr, S., 454
 Loken, E., 321
 Longford, N., 276, 484
 Lord, F., 321
 Louis, T., 276, 277, 413
 Lovell, J., 131
 Lu, H., 105
 Lu, K., 298
 Lunn, D., 527, 574

 Ma, G., 542
 Mack, S., 542
 MacLehose, R., 298
 Maddala, G., 131, 512
 Madigan, D., 298
 Mahboobeh, S., 194
 Maiti, T., 298
 Mallows, C., 105, 527
 Manski, C., 231
 Martin, A., 321, 574
 McCabe, G., 49
 McCarty, C., 342
 McCarty, N., 321
 McCullagh, P., 105, 131, 501
 McCulloch, C., 245, 484, 501
 McCulloch, R., 230, 297, 298
 McDonald, G., 76
 McDonald, J., 563
 McFadden, D., 132
 McKeague, I., 543
 Meng, X., 165, 297, 413, 433, 434, 526, 542
 Metropolis, N., 385
 Meyers, M., 543
 Michalopoulos, C., 229

 Milliff, R., 298
 Mitchell, J., 51
 Molenberghs, G., 247, 276
 Mollie, A., 298
 Montgomery, D., 454
 Moore, D., 49
 Moretti, E., 230
 Morgan, B., 342
 Morris, C., 277, 413
 Moskowitz, A., 153
 Mosteller, F., 74, 132, 165, 526
 Murphy, R., 231
 Murrell, P., 49, 562, 574

 Nachtsheim, C., 49
 Neal, R., 373
 Nelder, J., 105, 131, 501
 Nepomnyaschy, L., 248
 Nero, A., 277
 Neter, J., 49
 Newman, M., 342
 Newman, T., 26
 Newton, M., 132
 Neyman, J., 194
 Ng, J., 131
 Nolan, D., 49, 74, 299
 Normand, S., 230, 277
 Novick, M., 276, 321, 342
 Nychka, D., 298

 O'Hara, B., 574
 O'Keefe, S., 229
 O'Malley, A., 298
 Oakes, J., 512
 Oudshoorn, C., 542

 Palfrey, T., 131, 342
 Palmer, S., 231
 Pardoe, I., 49, 105, 133, 485, 526
 Park, D., 105, 321
 Parker, A., 51
 Pasarica, C., 563
 Pauler, D., 413
 Pearl, D., 194
 Pearl, J., 231
 Peer, J., 297
 Persico, N., 74
 Peto, R., 386
 Pickles, A., 542
 Pinheiro, J., 277, 297, 574
 Pittau, M. G., 105
 Plummer, M., 574
 Pollock, M., 485
 Poole, K., 230, 321
 Postlewaite, A., 74
 Pourahmadi, M., 297
 Pratt, J., 485

AUTHOR INDEX

605

- Pregibon, D., 105
 Price, P., 198, 277, 485

 Quagliarello, V., 229
 Quinn, K., 321, 574

 Rabe-Hesketh, S., 574
 Racine-Poon, A., 385
 Raftery, A., 298, 342
 Raghunathan, T., 542
 Ramsey, F., 8, 26, 49, 484
 Rasbash, J., 297, 485
 Raudenbush, S., 194, 247, 276, 297, 342, 454
 Raykov, T., 485
 Reiter, J., 230
 Renyi, A., 334
 Richardson, C., 298
 Richardson, S., 413
 Richmond, C., 132
 Ridder, G., 194, 229
 Ripley, B., 165, 298, 574
 Ritov, Y., 229
 Rivers, D., 133
 Roberts, G., 434
 Robins, J., 194, 229, 231, 542
 Robins, P., 229
 Robinson, G., 245, 248, 276, 485
 Rockoff, J., 458, 485
 Rodden, J., 321
 Romero, D., 133
 Rosenbaum, P., 194, 229, 231
 Rosenbluth, A., 385
 Rosenbluth, M., 385
 Rosenstone, S., 298
 Rosenthal, H., 230, 321
 Rosenthal, R., 485
 Rosnow, R., 485
 Ross, C., 75
 Rotnitzky, A., 229
 Rubin, D., 26, 105, 151, 165, 170, 194, 229, 230, 276, 297, 298, 346, 373, 385, 413, 434, 485, 526, 542, 574
 Ruppert, D., 74, 230, 298

 Sacerdote, B., 194
 Sahu, S., 434
 Salganik, M., 342, 413
 Sampson, R., 194, 342
 Samuhel, M., 542
 Sargent, D., 298, 434, 527
 Schafer, D., 26, 49, 484
 Schafer, J., 8, 542
 Schemper, M., 105
 Schenker, N., 542
 Scheve, K., 542
 Schlessman, J., 74

 Schultz, B., 542
 Schwing, R., 76
 Scott, A., 454
 Searle, S., 245, 276, 484, 501
 Sekhon, J., 229
 Shadish, W., 194
 Shao, Q., 385
 Sheiner, L., 558
 Shelley, G., 342
 Shen, W., 277
 Shinharay, S., 526
 Shively, W., 248
 Shoemaker, A., 105
 Shor, B., 105, 321
 Silver, R., 454
 Silverman, D., 74
 Simonson, I., 322
 Simonton, D., 133
 Skaug, D., 574
 Skogan, W., 342
 Skrondal, A., 574
 Sleight, P., 386
 Smith, A., 277, 385, 413
 Smith, J., 229
 Smith, T., 454
 Snedecor, G., 26, 292, 297
 Snijders, T., 245, 247, 276, 454
 Snyder, J., 197, 321
 Sobel, M., 194, 231, 512
 Solenberger, P., 542
 Spiegelhalter, D., 105, 413, 527, 542, 574
 Spitzer, E., 131
 Staiger, D., 458, 485
 Stanley, J., 194
 Stein, C., 277
 Stern, H., 26, 105, 151, 165, 276, 298, 346, 385, 413, 434, 526, 542, 574
 Sternberg, S., 527
 Stewart, C., 197
 Stigler, S., 49, 74, 132, 159, 277
 Stinson, M., 51
 Stone, M., 526
 Strathdee, S., 194
 Stuart, E., 229
 Sturtz, S., 574
 Su, Y. S., 105
 Subramanian, S., 485
 Summers, A., 485
 Swayne, D., 563

 Tan, W., 413
 Tanner, M., 385
 Taylor, J., 132
 Teller, A., 385
 Teller, E., 385
 Terry, M., 132

- Thayer, D., 276
 Thistlethwaite, D., 230
 Thomas, A., 527, 574
 Thomas, N., 229
 Thurstone, L., 132
 Tiao, G., 276, 413
 Tibshirani, R., 132, 298
 Tinbergen, J., 230
 Tobin, J., 132
 Todd, P., 194, 229, 230
 Tomz, M., 485
 Tosteson, T., 230
 Trevisani, M., 105
 Triest, R., 542
 Trochim, W., 454
 Troxel, A., 542
 Tsai, W., 543
 Tsui, K., 132
 Tsutakawa, R., 297
 Tu, W., 230
 Tuerlinckx, F., 105, 485, 527
 Tufte, E., 562
 Tukey, J., 74, 132, 165, 245, 501, 563
 Tukey, P., 49, 562
 Tustakawa, R., 413
 Tversky, A., 26

 Vaida, F., 527
 Vallone, R., 26
 Van Buuren, S., 542
 van der Klaauw, W., 230
 van der Linde, A., 105
 van der Linden, W., 321
 van Dijk, H., 512
 van Dyk, D., 131, 229, 434
 van Geen, A., 105
 Van Hoewyk, J., 542
 Van Mechelen, I., 105, 527
 Varadhan, R., 194
 Vehovar, V., 542
 Velleman, P., 26, 49
 Venables, W., 298, 574
 Verbeke, G., 247, 276
 Versteeg, R., 105
 Vikram, H., 229
 Viscusi, W., 198
 Vlahov, D., 194
 Volinsky, C., 298
 Von Krosigk, C., 276
 von Mises, R., 27

 Wahba, S., 229, 231
 Wainer, H., 231, 321, 485, 562
 Wakefield, J., 413
 Waldfogel, J., 49, 229
 Walker, S., 131
 Waller, L., 298

 Wallis, W., 132
 Wand, M., 298, 563
 Wang, N., 543
 Wang, X., 485
 Warnes, G., 574
 Wasserman, L., 413
 Wasserman, W., 49
 Wedderburn, R., 131
 Weidner, R., 485
 Weisberg, S., 8, 49, 526
 West, M., 298
 West, V., 131, 321, 543
 Whang, W., 153
 Wherry, R., 49
 Wideman, L., 542
 Wiens, B., 132
 Wikle, C., 298
 Wilkinson, L., 563
 Wilks, A., 574
 Wilson, J., 342
 Witte, J., 294, 298
 Wittenberg, J., 485
 Wlezien, C., 298
 Wolfe, B., 485
 Wolfe, R., 248
 Wolfers, J., 321
 Wolfram, C., 197
 Wong, W., 385
 Woolridge, J., 49, 132, 230
 Wright, G., 321
 Wu, Y., 434

 Xia, H., 298
 Xiaofeng, L., 454
 Xu, R., 485

 Yates, F., 501
 York, J., 298
 Yusuf, S., 386

 Zanutto, E., 230
 Zaslavsky, A., 277, 298, 413, 433, 542
 Zeger, S., 342, 385
 Zellner, A., 131
 Zeng, L., 229
 Zhao, L., 229
 Zheng, T., 342, 413
 Zheng, Y., 105
 Zhou, A., 230
 Zhou, X., 230
 Zimmerman, P., 231
 Zivot, E., 512
 Zorn, C., 105

Subject index

- 1.6, factor for converting from logit to
probit, 86, 118, 131
- 100,000 iterations, why we do not
recommend, 369
- 2 standard deviations, scaling by, 56–57
- 2.8 standard errors for power calculation,
441
- 5 ways to write the same model, 262–265
- 55,000 residents desperately need your
help, 23–26
- 6 quick tips, 547–549
- 8 schools, 430–434
- ability parameter, in item-response model,
315–320
- abline(), 353, 520
- Academy awards, 133
- accessing the simulations from a Bugs fit,
358
- AD Model Builder, multilevel modeling in,
572–573
- adding a predictor can increase the
residual variance, 480–481
- additivity, 46
- adjusted parameters, 382, 420, 422
compared to raw parameters, 423
- adjusted R, 475
- Advanced Placement exams, 463–464
- age guessing, 299
- age, as regression input, 66, 67
- Akaike information criterion (AIC), 105,
525–527
- aliasing, a form of nonidentifiability, 318,
321
- American Indians, 340
- analysis of covariance, 496–498
- analysis of variance (ANOVA), 487–502
additive data decomposition, 488
balance, 489
classical, 487–490
contrasts, 462–466, 496–498
degrees of freedom, 488, 499
F-test, 489
generalized linear models, 491
graphical display, 492–495, 497, 499,
500
mean square, 489
method of summarizing a multilevel
model, 492–494
model comparison and, 489
multilevel models and, 490–502
nested designs, 496
sum of squares, 488
unbalanced designs, 496
- analytic calculations and simulation, 148
- Annenberg Election Survey, 311
- apply(), 44, 353, 477
- arm package, 566
- array(), 308
- arsenic in drinking water in Bangladesh,
see wells
- as.bugs.array(), 400, 413
- as.vector(), 348
- attach.bugs(), 305, 352, 358
- augmented data
multilevel model, 397
regression with prior information, 393
- autoregression, 163–165
- available-case analysis, 532
- average causal effect, definition, 173
- average predictive comparison, *see*
predictive comparison
- axes of graphs, 553
- B-K plot, 312, 313, 321
- balance, *see* causal inference
- Bangladesh, *see* wells
- basement, 254
- bayesglm(), 104, 110
- Bayesian inference, 142, 276, 345–348,
392–414
calibration, 433
censored data, 405–408
classical regression, 392
debugging, 434
exchangeability, 347
Gibbs sampler, 397–402
informal, 143
informative prior distribution,
392–393, 427–434
Metropolis algorithm, 408–409
multilevel model, 393–402, 413
picture, 395, 396
prior distribution, 427–434

- variance parameters, 432–434, 499–501
- separation in logistic regression, 104
- social networks model, 409–413
- bayespplr(), 110
- beauty and teaching evaluations, 51
- before-after graph, 555
- behavioral learning experiment, 515–524
- beta-binomial model, 321
- binary data and count data, 117
- binary outcomes, 79
- binned residual plot, 97–101, 105, 559
- binomial distribution, 16
 - bounding trick in Bugs, 382
 - compared to Poisson, 112
- binomial-normal model, 321
- births, 27, 137–139
- black box, 190
- bounding parameters in Bugs, 383
- bounding trick for binomial model in Bugs, 382
- brlr package, 104
- broken windows, 325
- Bugs, 11, 305, 345–386, 567, 574
 - bounding parameters, 383
 - bounding trick for binomial model, 382
 - brute force, why we do not recommend, 369
 - calling from R, 350–352
 - censored data, 405–406
 - classical regression, 360
 - compared to lmer(), 386
 - compared to R, 399–402
 - compiling, 416
 - complete-pooling model, 360
 - confidence-building, 415–417
 - current problems with, 417
 - data, 350, 356, 416
 - data, parameters, and derived quantities, 366–369
 - dbin(), 381
 - dcat(), 383
 - debugging, 415–417
 - derived quantities, 367
 - detailed example, 348–359
 - dgamma(), 430
 - dlnorm(), 383
 - dlogis(), 384
 - dmnorm(), 376
 - dnorm(), 353, 354
 - dt(), 372, 384
 - dunif(), 355
 - dwish(), 377
 - each object can be defined at most once, 368
 - equals(), 384, 405
 - example of graphical output, 351
 - example of slow convergence, 421
 - example of tabular output, 351
 - examples, 353
 - fake-data simulation, 363–365
 - finding errors in a model, 373
 - general principles, 366–369
 - group-level model, 354
 - help, 353, 373, 565
 - I(), 383, 384, 405
 - individual-level model, 353
 - initial values, 350, 356, 369–370, 416
 - inprod(), 361, 378, 379
 - interpreting inferences from fitted model, 352
 - inverse(), 377
 - inverse-variance parameterization, 354, 368
 - knowing when to give up, 419
 - latent data for generalized linear models, 384
 - linear transformation to speed convergence, 419–427
 - logistic regression, 381–382
 - latent-data formulation, 384
 - logit(), 381
 - looping indexes, 353, 367
 - max(), 381
 - mean(), 382
 - min(), 381
 - missing data, 367, 529
 - model as specifications, not instructions, 368
 - model too big for, 336
 - modeled data and parameters, 366, 367
 - multilevel model, 375–386
 - logistic, 381–382
 - non-nested, 380–382
 - Poisson, 382
 - varying intercepts and slopes, 375–379
 - multiple group-level predictors, 379–380
 - multiple varying coefficients, 378
 - n.thin, 518
 - nested subscripts, 372
 - no overwriting in model, 368
 - no-pooling model, 360
 - non-nested model, 380–381
 - noninformative prior distribution, 355
 - nonlinear and nonadditive models, 371
 - number of iterations and sequences, 356–358, 369

AUTHOR INDEX

609

- open-ended modeling, 370–372
- ordered categorical regression, 383
- overdispersed Poisson regression, 382
- parameters, 350, 356
- postprocessing inferences in R, 359
- pow(), 355
- practical advice, 357, 369–370, 415–434, 547–549
- prediction, 361–363
- prior distribution, 354
- \hat{R} , 358, 369
- raw (unscaled) parameters, 377
- redundant parameters, 419–427
- replicated data for predictive checking, 518
- robit regression, 384
- scaled inverse-Wishart model, 376–380
- sd(), 460
- simple multilevel model, 350
- slow with large datasets, 418
- stochastic learning model
 - logarithmic link, 521
 - logistic link, 517
 - multilevel, 522–524
- thinning, 418
- too slow for looping in power calculation, 451
- unequal variances, 371–372
- unmodeled data and parameters, 366, 367, 378
- varying-intercept model, 350, 361
- varying-intercept, varying-slope model, 375–379
 - group-level predictors, 379–380
 - modeling the correlation, 376
 - no correlation, 375
 - scaled inverse-Wishart, 376
 - why you should learn Bugs, 345
 - Wishart model, 376–380
- bugs(), 350, 565, 567
- building generalized linear models, 125–127
- building regression models, 68–73
- c(), 350
- calibration
 - Bayesian inference, 433
 - plot, 309, 558
- calling Bugs from R, 350–352
- Caltech, 332
- cancer and nutrients, 294
- categorical variables as regression inputs, 66
- Cauchy distribution, 428, 430
- causal inference, 95, 167–198, 549
- average causal effect, 173, 180, 205
- balance, 181, 184–186, 199–204, 208, 209, 212
 - plot, 202, 229
- causes of an effect, 187
- confounding covariate, 169, 176, 181, 184, 196, 200, 202–203, 207, 212–213, 215
- constructed observational study, 210, 230, 231
- continuous treatment, 177
- controlling for post-treatment variable, don't do it, 177, 187–194, 227, 229
- counterfactual, 170, 173, 184, 185, 188, 201, 206, 218
- definition, 170
- difference-in-differences, 228, 231
- direct and indirect effects, 191–194
- effect of causes, 187
- effect of the treatment on the controls and treated, 205, 206, 211
- exclusion restriction, 216–218
- experiment, *see* randomized experiment
- external validity, 174
- fixed effects, 231
- fundamental problem, 171–172, 191
- hypothetical intervention, 187
- ignorability, 182–184, 186, 207, 212, 216–217, 225, 228, 231, 232
- infinite regress of causation, 187
- instrumental variables, 215–226, 230–231, 233
 - assumptions, 216–218, 220–221, 225–226, 233
 - derivation, 218–221
 - identifiability, 220–221, 224
 - multilevel model, 509–511
 - standard error, 222
 - two-stage least squares, 221
 - Wald estimate, 219, 221
 - weak instrument, 225
- intent-to-treat (ITT) effect, 216, 219, 220, 222
- internal validity, 174
- local average treatment effect (LATE), 219–220, 229, 233
- mapping your study to an ideal randomized experiment, 187–188
- matching, 206–212
 - model checking, 208
 - nearest-neighbor, 207
 - propensity score, 230
 - R packages for, 230

- model extrapolation, 169, 185, 201, 209, 213
- monotonicity, 216–217
- multilevel model, 178, 210, 212, 227–228, 503–512
- multiple treatments, 177, 186, 187
- natural experiment, 187, 225
- nonignorable assignment mechanism, 183
- observational study, 172, 181–188
 - design of, 437
- overlap, 184–186, 199–204, 209, 212–215
- poststratification, 178, 181, 206
- potential outcome, 168, 170, 171, 183, 186, 189, 191, 219
 - close substitutes, 171–172
- predictive comparison, 167, 168
- principal stratification, 192
- propensity score matching, 207–210, 230, 232
 - assumptions, 207
 - model checking, 208
 - R packages for, 230
 - standard errors, 212
- propensity score weighting, 211, 229
- randomized encouragement, 216
- randomized experiment, 171–181, 183
 - completely randomized, 175, 182, 183
 - design of, 437
 - paired comparisons, 175
 - randomized blocks, 175
- regression, 167–198
- regression discontinuity, 212–215, 232
- selection bias, 168
- selection model, 231
- selection on observables, 183
- sensitivity analysis, 231
- simultaneous causality, a meaningless concept?, 188
- structural equation model, 226
- subclassification, 204–207, 210–211, 229
- support of the data, 169
- SUTVA, stable unit treatment value assumption, 178
- treatment interaction, 178–180, 189, 205, 214–215
- treatment variable, definition, 186
- weighting, 205
- cbind(), 43, 146, 157, 361, 529
- CD4, 249, 277, 373
- censored data, 402–408
 - Bayesian inference, 405–408
 - Bugs, 405–406
 - Gibbs sampler, 406–408
 - imputation for, 540–541
 - likelihood, 404
 - maximum likelihood, 404–405
 - naive regression estimates, 403–404
 - picture, 541
 - R, 404–408
- Census, 277, 301, 308
- centering, 55–57, 419, 464–466
 - changes the correlation of group-level intercepts and slopes, 288–289
 - interactions and, 93
 - non-nested model, 292
- Central Limit Theorem, 13–14, 27
 - regression coefficients, 14
- chick brains, magnetic fields and, 481–484
- child care, 189–193, 201, 224
- child support enforcement in cities, 237–241
- children with HIV, 249, 443–447
- children's test scores, 31–51, 55–57
- classical models for regression coefficients, 293
- classical regression, as a special case of multilevel modeling, 258
- cluster sampling, 437, 447–449
- clustered data, 237–241
- cockroaches, 126–127, 161–163
- coef(), 43, 156, 260, 267, 280, 352
- coin flipping, patterns in, 554
- collinearity, 68, 255, 393, *see also* identifiability
- colMeans(), 520
- colochos, canasta de, vii, xxii
- colors(), 43
- colSums(), 411
- combining inferences from multiple imputations, 542
- combining separate local regressions, 263
- comparing two surveys, 442
- comparisons, graphs as, 552–553
- compiling Bugs, 416
- complete pooling, 247, 252–259, 270, 348
 - Bugs model, 360
 - problems, 256
 - special case of multilevel modeling, 258
- complete-case analysis, 531
- compound models
 - imputing a variable that can be positive or zero, 537–538
 - simulation of, 150–151, 537–538
- computer display, 565, 566
- computing, 9–11, 547, 565–574
 - debugging, 415–417
 - instead of matrix algebra, 8

AUTHOR INDEX

611

- multilevel models, 345–434
 - practical advice, 415–434
- confidence intervals, 18–20
- continuous data, 18
- discrete data, 18
- proportions, 18
- confidence-building, 415–417
- confounding covariate, 169, 176, 181, 184, 196, 200, 202–203, 207, 212–213, 215
- congressional elections, 76, 144–148, 197, 213, 233
- conjunctive item-response or ideal-point model, 319
- connect times on the web, 492–493
- Connecticut, what’s the matter with, 310–314
- constant term, 251, 349
- constraining a batch of coefficients to sum to zero, 326
- constructed observational study, 210
- constructive choice models, 127–131
- contextual effect, 481
- continuous and discrete predictors, 66
- continuous probability simulation, 152
- contrasts, 462–466
 - analysis of variance, 496–498
 - computing in R, 464
- convergence of iterative simulations
 - monitoring, 352
 - pictures, 357
- correlation, 57–59, 265, 389
 - coefficient estimates, 40
 - graph, 340
 - group-level intercepts and slopes, 279, 287–289
 - individual-level variables and group-level errors, 481
 - modeling in Bugs, 376–380
 - scaled inverse-Wishart model, 287
- cost-benefit analysis, 128, 153, 454–455
- costs and benefits of multilevel modeling, 246–247
- count data and binary data, 117
- counterfactual, 170, 173, 184, 185, 188, 201, 206, *see also* potential outcome
- counterfactual and predictive
 - interpretations of regression, 34
- covariance, *see* correlation
- coverage of confidence intervals, 156
- cows, 196
- C_p , 527
- curve(), 43, 353
- cutpoints, for ordered logit or probit, 119–120, 332
 - estimating in Bugs, 383
- data cleaning, for social networks survey, 333
- data for examples, 11
- data matrix, 37, 238, 239
 - group-level, 239, 240, 243
 - imputation, 541
 - individual-level, 239, 243
 - non-nested data, 243
- data model, 347
- data reduction, 209
- data sent to Bugs, 350, 356, 416
- data subsetting, 326, 357, 547
 - speeding computation, 418
- data.frame(), 48, 140, 452, 535
- dbin(), 381
- dcat(), 383
- death penalty, 19, 116, 243–244, 320–321, 540–541
- debugging, 415–417, 434
 - diagram of general strategy, 416
- decision analysis, well-switching example, 127–131
- default line in graph, 556, 560
- defaults in R functions, 452
- degree distribution, estimated for men and women, 337
- degrees of freedom, 41, 372, 488
 - inverse-Wishart distribution, 286
- Democrats and Republicans, 310–314
- dependent variable, *see* outcome
- derived quantities, 366, 367
- design of sampling and experiments, 437–455
- details in graphs, 560
- deterministic or random imputation, 534, 537
- deterministic part of a regression model, 31
- deviance, 100, 105, 113, 524–526
- deviance information criterion (DIC), 105, 525–527
 - instability in computation of, 526
- dgamma(), 430
- diagnostics
 - external validation, 48
 - residuals, 40, 47, 97–101
 - simulation-based, 155–166, 513–524
- diarrhea, zinc, and HIV, 443–447
- difference-in-differences estimation, 228, 231
- difficulty parameter, in item-response model, 315–320
- dim(), 147
- dimnames(), 400
- discrepancy variable, 513

- discrete and continuous predictors, 66, 107
discrete probability simulation, 152
discrimination parameter, in item-response model, 316
disjunctive item, 319
display(), 38–39, 565
distance to the nearest safe well, 88
distribution, 13–16
 binomial, 16
 bounding trick in Bugs, 382
 Cauchy, 428, 430
 folded noncentral t, 428
 gamma, 335
 log=TRUE option in R, 405
 logistic, 85
 Bugs, 384
 lognormal, 15, 383
 multivariate normal, 15
 negative binomial, 115, 336
 normal, 13–15, 263
 computing in R, 405
 truncated, 407
 Poisson, 16, 110–116, 335
 scaled inverse-Wishart, 284–287, 376–380
 t, 124, 372, 428
 truncated normal, 407
 Wishart, 284–287, 298, 376–380
divide by 4 rule, for interpreting logistic regression, 82
dividing by two standard deviations, 56–57
dlnorm(), 383
dlogis(), 384
dmnorm(), 376
dnorm(), 353, 354, 404
dog experiment, 515–524
 model comparison, 526
 observed and replicated data, 516, 523
Douglas, William, 318
dt(), 372, 384
dummy variable, *see* indicator
dunif(), 355
dwish(), 377
dynamic graphics, 563
earnings
 height and, 50, 53–54, 75, 126, 287, 290–292
 logarithmic models, 59–65
 mixed discrete/continuous data, 126
econometrics and biostatistics, 231
education as categorical input variable, 95
educational children's television programs,
 see Electric Company *and*
 Sesame Street
educational testing, 317, 430–434
effect size, why more important than
 sample size, 439
effective sample size, 352
elasticity, 64, 76
election forecasting, graphs of, 557, 562
election fraud, 23–26
Electric Company experiment, 174–186
 graph of data, 552
 multilevel model, 503–505
Emacs, 565
equal variance of errors, 46
equals(), 384, 405
Erdos-Renyi model, 334
error rate, 99
errors, distinguished from residuals, 387
exam scores, actual vs. guessed, 558, 559
examples, data for, 11
exchangeability and prior distribution, 347
exclusion restriction, *see* causal inference,
 instrumental variables,
 assumptions
expected(), 123
experiment, *see* causal inference,
 randomized experiment
experimental design, 437–455
experimental economics, 331
explained variance, *see* R^2
exploratory data analysis, 551
exposure, in Poisson model, 111–113
expression(), 148, 353
external validation, 48
 election polls, 309
external validity, 174
F-test, 489
factor, 68
factor analysis, 296
factor(), 255, 349
fake-data simulation, 50, 155–158
 Bayesian inference and, 434
 checking coverage of intervals, 365
 multilevel model, 363–365
 residual plots, 157–158
 sample size and power calculations,
 449–454
 using Bugs, 363–365
feeling thermometer, 86
figure skaters, 248
finite-population variance, 459–462, 499
 analysis of variance (ANOVA), 491
 Bugs, 460
 graph, 461
 unmodeled coefficients, 462
fit many models, 547
fitted(), 158
fitting multilevel models, 259–262, 345–434

AUTHOR INDEX

613

- fitting the wrong model, 165
- fixed effects, 231, 244–246, 259, 261
 finite-population inferences and, 461
 many definitions, 245, 248
 R, 260
 why we avoid the term, 2, 226–228, 245
- fixef(), 260, 280
- flight-simulator experiment, 289–290, 297, 464–466
 analysis of variance, 488
 Bugs model, 380
 superpopulation and finite-population variance, 459–460
- folded noncentral t distribution, 428
- forecasting elections, 294–296
- Fragile Families study, 238
- function(), 350, 401
- functions in R, 139, 147, 151, 404, 534, 535
- fundamental problem of causal inference, 171–172, 191
- gain scores, 177, 195
- Gallup Report, 560
- GAMM package, 567
- gamma distribution, 335
- generalized additive model, 298, 567
- generalized estimating equations, 248
- generalized linear model, 109–133
 analysis of variance (ANOVA), 491, 493–494
 binomial, 116–117
 building, 125–127
 cutpoints, 119–120
 deviance, 100, 105, 113
 latent-data formulation, 384
 logistic, 79–108
 logistic-binomial, 109, 116–117
 multilevel, 325–342
 multinomial logit and probit, 110, 119–124
 negative binomial, 115
 ordered logit, 119–124
 Bugs, 383
 multilevel, 331–332
 others, 127
 Poisson, 109–116
 Bugs, 382
 compared to binomial, 112
 exposure, 111–113
 overdispersion, 382
 prediction, 272
 predictive comparisons, 466–473
 probit, 109, 118–119
 probit or logit, 118
 robit as robust alternative to logit and probit, 124–125
 robust, 110
 simulation, 148–151
 thresholds, 119–120
- Gibbs sampler, 385, 397–402
 censored data, 406–408
 linear transformation to speed convergence, 419–427
 model building and, 402
 multilevel model, 398–402
 picture, 398
 programming in R, 399–402, 412
 redundant parameters, 419–427
 slow convergence, 424
 social networks model, 410
 updating functions in R, 399–402
- glm(), 79, 110, 565
- global variables in R, 400, 412
- goodness-of-fit, *see* model checking
- grades, predicting, 157
- graphics, 42–45, 548, 551–563
 comparisons, 552–553
 general advice, 562
 instead of tables, 328, 337
 jittering, 32
 no single graph does it all, 552–553
 plotting regression coefficients, 337, 341
 plotting symbols, 555
 R, 562
 scatterplot with regression lines superimposed, 35, 42–45
 shape of the plotting region, 556
 showing fitted models, 551, 553
 symbols and auxiliary lines, 554
 theory of, 562
 why, 551
- group, 261
- group indicators, 264
- group- and individual-level data matrices, 239, 243
 imputation, 541
- group-level predictors, 265–269, 271
 along with group indicators, 269, 293, 463–464, 498
 Bugs model, 361
 varying-intercept, varying-slope models, 280
- group-level standard deviation, 270
- guessing ages, 299
- guessing, in item-response model, 319
- handedness, 66
- hard constraint, 257
- hazard regression, 298

614

height, 139
 earnings and, 50, 53–54, 75, 126, 287, 290–292
 logarithmic models, 59–65
 mixture model for, 14
 parents and children, 58
 weight and, 41, 74, 402–408

help in Bugs, 565
 help in R, 405, 565, 567

heteroscedasticity (unequal variances), 297

hett package for robust regression, 110, 124, 133, 567

hist(), 137, 536, 562

histograms, 561

HLM, multilevel modeling in, 573

holding other predictors constant, difficulty of, 34

homeless people, 333

hot-deck imputation, 538

how many groups, 275–276

how many observations per group, 275–276

how many x's survey, 332–342

hyperparameter, 1, 258

hypothesis testing, 20–26

I(), 215, 383, 384, 405, 538

ideal-point model, 314–321
 multilevel, 316
 picture, 315
 redundant parameters in Bugs, 426
 two-dimensional, 319

identifiability, 419, 420
 Bayesian regression and, 393
 categorized predictors and, 68
 causal inference, 170
 constant term in non-nested models, 381
 ideal-point model, 318
 instrumental variables, 220–221, 224
 item-response model, 315
 likelihood and, 392
 linear regression, 68
 logistic regression, 86, 104, 107
 social networks model, 336

ifelse(), 126, 150, 384, 403, 534

ignorability, 182–184, 186, 231, 530, 542,
see also causal inference

imbalance, *see* causal inference, balance

imputation, *see* missing data

impute(), 535

income and voting, 79–84, 105, 107, 310–314

incremental cost-effectiveness ratio, 153

incumbency, 197, 233

independence of errors, 46

independent variable, *see* input variable

AUTHOR INDEX

index number, don't graph by, 553

index variable, 67, 238, 252
 creating in R, 348
 non-nested models, 289

indicator variable, 67, 238, 244–246, 255
 default, reference, or baseline condition, 68

individual- and group-level data matrices, 239, 243
 imputation, 541

Infant Health and Development Program,
see child care

inference, *see* statistical inference

informative prior distribution, 392–393

initial values sent to Bugs, 350, 356, 369–370, 416
 restricted parameter space, 384

inprod(), 361, 378, 379

input variables, as distinguished from predictors, 37, 466

install.packages(), 567

instrumental variables, *see* causal inference

interactions, 34–36, 242, 453
 centering the input variables, 93
 graphing, 36, 94, 313
 logistic regression, 92–96
 predictive comparisons and, 469
 sample size and, 438
 treatment effects, 178–180, 189, 205
 varying slopes as example of, 282–283
 when to look for, 36

intercept, 33, 35, *see* varying intercepts and slopes

intermediate outcome, *see* causal inference, controlling for post-treatment variable

internal validity, 174

interpreting regression coefficients, *see* linear regression, logistic regression, generalized linear model, *and* multilevel model

interquartile range (IQR), 70

intraclass correlation (ICC), 258, 448

inverse variance, used in dnorm() in Bugs, 354–355

inverse(), 377

inverse-gamma distribution, why we do not use as prior distribution for variance parameters, 430–434

inverse-logit function, 80

invlogit(), 149

item-response model, 314–321
 guessing, 319
 multilevel, 316
 picture, 315, 317
 redundant parameters in Bugs, 426

AUTHOR INDEX

615

- two-dimensional, 319
- iterative regression imputation, 539
- iterative simulation, 408–409
- Jacobian for nonlinear transformations, 409, 430
- Jaycees, 334, 335, 340
- `jitter.binary()`, 89
- jittering, 32, 554
- knowing when to give up, 419
- Lac Qui Parle County, 253
- large regression with correlated errors, 265
- latent-data formulation for logistic regression, 85–86, 120
- latin square, 292, 297, 497–501
- Latinos, hypothetical survey of, 454
- least squares, 39, 387–390
 - augmented data and multilevel model, 396–397
 - weighted, 389
- legislative redistricting, 555
- `length()`, 157
- level, 68
- `lgamma()`, 411
- `library()`, 567
- likelihood, 347, 387–414
 - censored data, 404
 - generalized linear model, 389–390
 - inferential uncertainty and simulation, 392
 - logistic regression, 389
 - picture, 390, 391, 395, 396
 - Poisson regression, 390
 - social networks model, 409–413
 - surface, 390–392
- linear predictor, 79, 305
- linear regression, 31–77, 387–390
 - assumptions, 45–47
 - Bayesian inference, 346
 - binary predictor, 31
 - Bugs model, 360
 - compared to principal component line, 57–58
 - continuous predictor, 32
 - correlation and, 57–59
 - counterfactual interpretation, 34
 - diagnostics, 45–47
 - displaying several, 73–74
 - fitting in R, 38–39
 - general principles for building models, 69
 - inferential uncertainty, 40
 - interactions, 34–36
 - interpreting coefficients, 33–34
 - interactions, 35–36
 - least squares, 39
 - matrix notation, 388
 - missing-data imputation, 533–538
 - multiple predictors, 32–34
 - notation, 37–38
 - one predictor, 31–32
 - picture of matrix, 37
 - prediction, 47–49, 70–73, 272
 - predictive interpretation, 34
 - sample size and power calculations, 446, 451
 - simulation, 140–148
 - standard error, 40
 - statistical inference, 37–42
 - transformation, 53–77
 - validation, 47–49
- linear transformation, 14, 53–54, 88, 95, 122, 294
 - centering a batch of multilevel coefficients, 464–466
 - prior distribution and, 355
 - speeding convergence of Gibbs sampler, 419–427
- linearity, 46
- link function, 109
- `list()`, 143, 350
- `lm()`, 38–39, 402, 565
- `lme4` package, 566
- `lmer()`, 259–262, 266, 267, 277, 566, 574
 - compared to Bugs, 386
 - limitations, 262, 304, 345
 - logistic regression, 302
 - non-nested model, 289
 - six quick examples, 568–569
 - varying intercepts and slopes, 279–289
- local average treatment effect (LATE), 219–220, 229, 233
- `log()`, 59
- log-log model, 64
- `log10()`, 61
- logarithmic transformation, 59–65, 98, 252
 - even when not necessary, 65
 - interpreting regression coefficients, 60
 - interpreting variance parameters, 327
 - picture, 60
 - why we usually don't use base 10, 60–61
- logistic distribution, 85
 - Bugs, 384
 - close to normal with standard deviation 1.6, 86, 118, 131
- logistic regression, 79–108
 - binned residual plot, 105
 - Bugs, 381–382
 - latent-data formulation, 384

- choice models in one and multiple dimensions, 128–131
- compared to probit, 118, 129
- computing using `lmer()`, 302
- deviance, 100, 105
- divide by 4 rule for interpreting coefficients, 82
- graph of coefficient estimates, 306
- graphs of data and fitted curves, 307
- ideal-point model, 314–320
- identifiability, 86, 104, 107
- inference, 83
- interactions, 92–96
- interpreting coefficients, 81–84, 89
- item-response (Rasch) model, 314–320
- latent-data formulation, 85–86, 384
- logit and logit^{-1} functions, 80
- missing-data imputation, 533–538
- multilevel, 301–323
 - Bugs, 381–382
 - formula, 302, 303
 - graphing, 304–310
 - interpreting coefficients, 304
 - non-nested, 302–304, 320–321
 - overdispersion, 320–321
- odds ratios, 82–83
- pictures, 80
- plotting data and fitted curve, 80, 84
- predictive comparisons, 81
- prediction, 272
- predictive comparisons, 101–104, 466–473
- propensity score, 207–208
- separation, 104, 107
- simulation, 148
- standard error, 83
- two predictors, 90–92
- varying-intercept, varying-slope model, 310–314
- wells in Bangladesh, 86–92
- logistic-binomial model, 109, 116–117
 - overdispersion, 116
- logit, *see* logistic regression
- `logit()`, 381
- lognormal distribution, 15, 383
- looping indexes in Bugs, 353, 366, 367
- looping, for power calculation, 452
- lowess, 298
- lurking variable, 169

- magnetic fields and brain functioning, 481–484
- Mahalanobis distance, 207
- many predictors, multilevel models for, 293–296

- maps, 556–557
- Markov chain Monte Carlo (MCMC), 408–409
- MASS package, 122, 567, 574
- matching, 206–212
 - missing-data imputation, 538
 - propensity score, 207–210, 232
 - R packages for, 230
- `matching()`, 208
- maternal IQ, 32
- matrix notation, 37–38, 284–287
- matrix of predictors
 - group-level, 252
 - individual-level, 251
- matrix of simulations, 146, 149, 353, 358
- Matrix package, 259
- `max()`, 381
- maximum likelihood, 388–390
 - censored data, 404–405
 - generalized linear model, 389–390
 - logistic regression, 389
- MCMCpack, 567
- `mean()`, 56, 359, 382, 477
- mediator, *see* causal inference, controlling for post-treatment variable
- men and women, 337
- mesquite bushes, 68–73
- meta-analysis, 386, 438
- Metropolis algorithm, 385, 408–409
 - picture, 408
 - social networks model, 410
- midterm and final exams, 157
- millet crop, 292
- millimeters, inches, and miles, 53
- `min()`, 381
- missing at random (MAR), 530, 542
 - impossibility of verifying, 531
- missing completely at random (MCAR), 530
- missing data in R and Bugs, 529
- missing not at random (MNAR), 530
- missing values, not allowed in unmodeled data, 416, 529
- missing-data imputation, 333, 529–543
 - available-case analysis, 532
 - Bugs, 367
 - complete-case analysis, 531
 - congressional elections, 145
 - deterministic or random, 534, 537
 - iterative, 539
 - many variables, 539–540
 - matching, 538
 - model-based, 540–541
 - models, 530–531
 - multilevel data structures, 541
 - one variable, 533–538

AUTHOR INDEX

617

- simple methods, 531–533
 - topcoding, 534
- Mississippi, as poor state, 313
- mixed discrete/continuous data, 126, 537
- mixed effects, *see* random effects
- MLWin, multilevel modeling in, 573
- mnp package, 110
- model checking, *see* posterior predictive checks *and* residuals
 - using simulation, 158–165, 513–524
- model comparison, 524–526
- model extrapolation, 169, 185, 201, 209, 213
- model-based imputation, 540–541
- modeled data and parameters, 367
- modeling the coefficients of a large regression model, 264
- monotonicity, *see* causal inference,
 - instrumental variables,
 - assumptions
- month of arrest, 21, 331
- more than two varying coefficients, 285
- mothers and children, 31–51, 55–57
- motivations for multilevel modeling, 6–8, 246–247
- mtext(), 520
- multilevel model, 1, 237–342, 463
 - alternative to selecting regression predictors, 294
 - analysis of variance (ANOVA), 490–502
 - assumptions, 247
 - Bayesian inference, 393
 - Bayesian perspective, 346
 - building, 293–296
 - Gibbs sampler, 402
 - building from classical regression, 270
 - causal inference, 503–512
 - combining regression inputs, 293–296
 - compared to classical regression, 463
 - comparison to simpler methods, 310
 - complexity, 246
 - computing, 345–434
 - equivalent sample size, 258, 268
 - factor analysis and, 296
 - fake-data simulation, 363–365
 - fitting in Bugs, 345–386
 - fitting in R, Stata, SAS, and other software, 568
 - fitting using lmer(), 259–262
 - five ways to write, 262–265
 - generalized linear model, 325–342
 - Gibbs sampler, 398–402
 - programming in R, 399–402
 - graphing, 304–310
 - group-level predictors, 265–269, 568
 - group-level variance, superpopulation
 - and finite-population, 459–462
 - how many groups needed, 275–276
 - how many observations per group needed, 275–276
 - imputation at different levels, 541
 - inference for groups with no data, 306
 - instead of comparing significance levels, 482
 - instrumental variables, 509–511
 - interpreting coefficients, 268
 - least squares with augmented data, 396–397
 - logistic regression, 301–323, 568
 - Bugs, 425
 - computing using lmer(), 302
 - non-nested, 302–304, 320–321
 - overdispersion, 320–321
 - matrix notation, 284
 - negative binomial, 332–342
 - non-nested, 289–293, 569
 - Bugs, 380–382, 424
 - identifiability of constant term, 381
 - negative binomial, 332–342
 - redundant parameters, 421–423
 - notation, 251–252
 - ordered logistic regression, 331–332
 - plot of group-level estimates and fitted regression line, 266, 307
 - Poisson (overdispersed), 332–342, 382, 568
 - pooling, 252–259
 - prediction, 272–275, 361–363
 - predictive comparisons, 470
 - prior distribution for variance parameters, 427–434, 499–501
 - R^2 , 473–477
 - redundant parameters, 420
 - sample size and power calculations, 447–454
 - six quick examples, 568–569
 - small number of groups, 431–432, 461
 - statistical significance, 271
 - summarizing and displaying, 261
 - understanding and summarizing, 457–486
 - variance parameters, 480–481
 - varying intercepts and slopes, 279–289, 568
 - varying slopes without varying intercepts, 283–284
 - varying the number of groups, 330
- multilevel modeling
 - costs and benefits, 9, 246–247
 - motivations, 6–8, 246
 - when most effective, 270

618

multilevel structures, 237–249
 data matrix, 238–240, 243
 imputation, 541
 multinomial logit and probit models, 110,
 119–124
 storable votes, 120–124
 multiple chains, necessary to monitor
 mixing, 356–358
 multiple comparisons, why we do not
 worry about, 22, 484–485
 multiple imputation, combining inferences,
 542
 multiplicative model, 59
 multivariate imputation, 539–540
 multivariate normal distribution, 15
 mvnorm(), 143

 n.chains, number of chains when running
 Bugs, 356–358, 369
 n.eff, effective sample size of Bugs fit, 352,
 358
 n.iter, number of iterations when running
 Bugs, 356–358, 369
 n.sims, 143
 n.thin option in Bugs, 518
 NA, missing value in R and Bugs, 50, 362,
 529
 naming inputs, 62
 National Election Study, 73, 311, 342, 385
 National Longitudinal Survey of Youth,
 210
 National Supported Work, 231
 natural log, 60–61
 ncol(), 361, 519
 negative binomial distribution, 115, 336
 multilevel model, 332–342
 neighborhoods and crime, 325, 342
 nested subscripts in Bugs, 372
 networks, 297, 333
 New York City schools, 458–459
 Newcomb's speed of light data, 159
 Nicoles, 333, 335, 340
 nmatch package, 230
 no data, multilevel inference for groups
 with, 306
 no pooling, 247, 252–259, 270, 349
 Bugs model, 360
 overestimates between-group
 variation, 253
 picture, 253
 problems, 256
 special case of multilevel modeling,
 258
 non-nested models, 241–244, 248–249,
 289–293
 Bugs, 380–381

AUTHOR INDEX

varying intercepts and slopes, 291
 where to put the intercept or
 constant term, 381
 nonidentifiability, *see* identifiability
 noninformative prior distribution, 347, 355
 Bugs, 354, 355
 nonlinear prediction, 147
 nonparametric regression, 297
 normal distribution, 13–15, 263
 computing in R, 405
 estimated regression coefficients, 15,
 40, 83
 inverse-variance parameterization in
 Bugs, 354–355
 regression errors, 46
 truncated, 407
 notation, 263
 capital letters for matrices, lowercase
 for vectors and scalars, 167, 252,
 376, 383
 cluster sampling, 447
 linear regression, 37–38
 multilevel model, 251–252
 parameters and probability
 distributions, 13
 varying intercepts and slopes, 284
 nrow(), 519
 number of iterations when running Bugs,
 356–358, 369
 number of observations and groups
 needed, 275–276, 278
 number of sequences when running Bugs,
 356–358, 369
 numerical optimization in R, 405
 nutrients and cancer, 294
 NYPD stops, *see* police stops

 O'Connor, Sandra Day, 318
 observational study, *see* causal inference
 odds ratios and logistic regression, 82–83
 offset, in Poisson regression, 112, 326, 382
 Ohio, as intermediate state, 313
 Olympics, 248, 485
 omitted variable, 169
 omniscience, 195
 one-way analysis of variance, 494
 open-ended modeling in Bugs, 370–372
 OpenBugs, 11, 565, 574, *see also* Bugs
 optim(), 405, 413
 optimal design, 455
 options(), 561
 order(), 519
 ordered and unordered categorical
 outcomes, 119, 123
 ordered logistic model, 119–124
 fitting in Bugs, 383

AUTHOR INDEX

619

- multilevel, 331–332
 - storable votes, 120–124
- outcome, 37, 251
- outer(), 411
- overdispersion, 21, 114–116, 320
 - adjusting standard errors, 115, 117
 - groups in the social network, 338
 - multilevel Poisson model, 335–336, 382, 409–413
 - simulation, 150
 - variance components and, 325
- overlap, *see* causal inference
- pain scores, observed vs. expected, 558
- panel-corrected standard errors, 248
- par(), 305
- parameter expansion, *see* redundant parameters
- parameters saved from Bugs, 350, 356
- partial pooling, 252–259, 394, 477–480
 - Bayesian, 394
 - formula, 253, 258, 269, 477
 - graph, 479
 - group-level predictors and, 269
 - picture, 253
 - plotting data and fitted lines, 257, 266
 - set of regression predictors, 295
 - summarizing a fitted multilevel model, 477–480
- partisan bias, 555
- paste(), 353
- pch, 43
- p_D , effective number of parameters in a Bayesian inference, 525
- phase diagram for decision analysis, 130
- plot(), 43, 350
- plots of replicated datasets, 160, 163
- pmin(), 478
- pnorm(), 404
- points(), 43
- Poisson model, 16, 109–116, 335
 - checking using simulation, 161–163
 - compared to binomial, 112
 - exposure, 111–113
 - interpreting coefficients, 111
 - multilevel, 325–331
 - offset, 326
 - overdispersion, 114–116
 - police stops, 112–116
 - zero-inflated, 126–127
- police stops, 5–6, 21, 112–116, 325–331, 342
 - Bugs model, 382
 - graph, 328
- political ideology, 73
- political party identification, 73–74
- pollution, 76
- polr(), 110, 122
- pooling, *see* complete pooling, no pooling, partial pooling
- pooling factor, 478–480
- posterior distribution
 - picture, 395, 396
 - programming in R, 411
 - social networks model, 409
- posterior predictive checks, 158–165, 513–515
 - data display for dog example, 516, 523
 - numerical summary, 521
 - time plot, 519, 520, 522
 - using Bugs, 518
 - using R, 518
- posterior uncertainty, 149
- postprocessing Bugs output, 359
- poststratification, 178, 181, 206, 301–310
 - formula, 301, 308
 - R code, 308
- potential outcome, 168, 171, 183, 186, 189, 191, 219
 - close substitutes, 171–172
 - interpreting regression coefficients, 34
- pow(), 355
- power calculation, 437–455
 - 2.8 standard errors, 441
 - classical, 439–447
 - general concerns, 439
 - inference for continuous outcomes, 443–447
 - inference for linear regression, 451
 - inference for proportions, 439–443
 - inference for regression coefficients, 446–447
 - inherently speculative, 445, 447
 - multilevel models, 447–454
 - pictures, 440, 441
 - unequal sample sizes, 443
- pre-election polls, 560
- precinct, 325
- predict(), 48, 115, 208, 535
- prediction, 47–49, 68–73
 - Bugs, 361–363
 - interpreting regression coefficients, 34
 - model checking and, 513–515
 - multilevel model, 272–275, 361–363
 - new observations and new groups, 272–275, 361–363
 - nonlinear, 274
- predictive checks, *see* posterior predictive checks
- predictive comparison, 81, 101–104, 167, 168, 466–473, 485

- comparing models, 472–473
- formula, 466
- general approach, 468
- general notation, 103
- graph, 467, 468
- interactions and, 103
- model summary, 471
- predictive simulation, 140, 147–151
 - binomial distribution, 149
 - generalized linear models, 148–151
 - latent logistic distribution, 149
 - linear regression, 140–148, 152
 - model checking and, 158–165, 513–524
- predictive standard deviation, 274
- predictive uncertainty, simulation of, 140
- predictors, as distinguished from input variables, 37, 466
- presidential elections, 3–4, 79–84, 294, 301–314, 493, 557, 560
- principal component line, 57
- principles of modeling in Bugs, 366–369
- print(), 350
- prior distribution, 143, 345–348, 413, 427–434
 - Bugs, 354
 - effect on posterior distribution, picture, 430, 432
 - informative, 392–393
 - scale, 430
 - inverse-gamma, why we do not use, 430–434
 - noninformative, 347, 354, 355
 - picture, 395, 396
 - provisional nature of, 347
 - scale, 355
 - uniform, 428–429
 - variance parameters, 432–434, 499–501
 - weakly informative, 431–432
 - Wishart model, 377
- prison sentences, example for predictive comparisons, 470
- probability, *see* distribution
- probability models, simulation of, 137–140
- probability of a tied election, 148
- probit model, 109, 118–119
 - compared to logit, 118, 129
- programming in R, 567
- Progres, 508–509
- propagating uncertainty, 142, 152
- propensity score matching, *see* causal inference
- provisional nature of prior distributions, 347
- psychological experiment of pilots on flight simulators, 289–290
- quantile(), 141, 359
- quasibinomial family, 117
- quasipoisson family, 115
- quick tips, 547–549
- R, 10–11, 298, 565, 574
 - abline(), 353, 520
 - apply(), 44, 353, 477
 - arm package, 566
 - array(), 308
 - as.bugs(), 413
 - as.bugs.array(), 400
 - as.vector(), 348
 - attach.bugs(), 305, 352, 358
 - bayesglm(), 110
 - Bayesian inference for social networks model, 409–413
 - bayespolr(), 110
 - brlr package, 104
 - bugs(), 350, 567
 - byesglm(), 104, 566
 - c(), 350
 - calling Bugs from, 350–352
 - cbind(), 43, 146, 157, 361, 529
 - censored data, 404–408
 - coef(), 43, 156, 260, 267, 280, 352
 - colMeans(), 520
 - colors(), 43
 - colSums(), 411
 - console, 565
 - curve(), 43, 353
 - data.frame(), 48, 140, 452, 535
 - default values in functions, 452
 - digits, 561
 - dim(), 147
 - dimnames(), 400
 - display(), 38–39, 565
 - dnorm(), 404
 - expected(), 123
 - expression(), 148, 353
 - factor(), 255, 349
 - fitted(), 158
 - fixef(), 260, 280
 - function(), 350, 401
 - GAMM package, 567
 - Gibbs sampler, 399–402
 - censored data, 406–408
 - glm(), 79, 110, 565
 - global variables, 400, 412
 - graphics, 562
 - graphics window, 565
 - graphing models fit in Bugs, 352
 - help, 405, 565, 567

AUTHOR INDEX

621

- hett package for robust regression, 110, 124, 133, 567
- hist(), 137, 536, 562
- I(), 215, 538
- ifelse(), 126, 150, 384, 403, 534
- impute(), 535
- install.packages(), 567
- invlogit(), 149
- jitter.binary(), 89
- length(), 157
- lgamma(), 411
- library(), 411, 567
- list(), 143, 350
- lm(), 38–39, 402, 565
- lme4 package, 566
- lmer(), 259–262, 266, 267, 277, 566, 574
 - limitations, 262, 304, 345
 - logistic regression, 302
 - non-nested model, 289
 - six quick examples, 568–569
 - varying intercepts and slopes, 279–289
- log(), 59
- log10(), 61
- log=TRUE option, 405
- MASS package, 122, 567, 574
- matching(), 208
- Matrix package, 259
- MCMCpack package, 567
- mean(), 56, 359, 477
- mnp package, 110
- mtext(), 520
- mvrnorm(), 143
- NA, 50
- ncol(), 361, 519
- nmatch package, 230
- nrow(), 519
- optim(), 405, 413
- options(), 561
- order(), 519
- outer(), 411
- par(), 305
- paste(), 353
- pch, 43
- plot(), 43, 350
- pmin(), 478
- pnorm(), 404
- points(), 43
- polr(), 110, 122, 566
- predict(), 48, 115, 208, 535
- print(), 350
- probit family, 118
- programming, 567
- quantile(), 141, 359
- quasibinomial family, 117
- quasipoisson family, 115
- R2WinBUGS, 565
- ranef(), 260, 280
- range(), 352, 520
- rbinom(), 137, 149
- read.dta(), 411
- read.table(), 49, 348
- rep(), 452
- replicate(), 139, 147
- replicated data for predictive checking, 518
- return(), 401
- rnegbin(), 150
- rnorm(), 106, 141, 155, 356, 401, 407
- rnorm.trunc(), 407
- rowMeans(), 180
- rowSums(), 147, 411
- rpois(), 150
- runif(), 150, 353, 356
- rwish(), 377
- sample(), 138, 278, 418, 452, 534
- sapply(), 151
- save(), 362
- sd(), 56, 462
- se.coef(), 156, 565
- se.fixef(), 261
- se.ranef(), 261
- sem package, 223
- sigma.hat(), 273, 565
- sim(), 43, 142, 143, 392, 565
- sorting, 519
- subset option in lm() and glm(), 107, 126, 538
- sum(), 147
- summary(), why we don't use, 39
- table(), 353
- tlm(), 124, 133, 567
- tsls(), 223
- unique(), 348
- updating functions for Gibbs sampler, 399–402
- var(), 477
- R functions, 44, 139, 147, 151, 404, 534, 535
- R^2 , 41, 49, 62, 485
 - adjusted, 475
 - Bayesian definition, 475
 - classical definition, 474
 - computation, 476
 - each level of a model, 474
 - interpretation for model with no constant term, 349
 - multilevel, 473–477
 - pictures, 42
 - why we do not define in terms of model comparison, 474

622

R2WinBUGS package, 565
 \hat{R} for summarizing convergence of Bugs fit, 352, 358, 369
 radon, 3, 36, 252–283, 348–369, 480–481
 random effects, 244–246, 259
 many definitions, 245, 248
 R, 260
 superpopulation inferences and, 461
 why we avoid the term, 2, 245
 random imputation, 534, 537
 randomized experiment, 171–181, 183, *see also* causal inference
 ranef(), 260, 280
 range(), 352, 520
 Rasch model, 315–320
 ratings, 298
 ratio of parameters, 152
 raw (unscaled) parameters, 377
 compared to adjusted parameters, 423
 rbinom(), 137, 149
 read.dta(), 411
 read.table(), 49, 348
 recall, in social networks survey, 339
 red states and blue states, 310–314
 redundant parameters, 419–427
 additive, 316, 326, 336, 382, 412, 419–423, 464–466
 future implementations, 427
 item-response and ideal-point models, 316
 multiplicative, 424–427
 Bugs, 424, 425
 social network model, 336
 reference condition, in classical no-pooling regression, 349
 reference model, 347
 regression, *see* linear regression, logistic regression, generalized linear models
 regression coefficients, graph, 337, 341
 regression discontinuity, 212–215
 regression to the mean, 57–59
 rejection, not the goal of model checking, 524
 rep(), 452
 repeated measurements, 241–243
 graph, 450
 replicate(), 139, 147
 replicated data, for predictive checking, 514
 replicated datasets, plotted, 160, 163
 residuals, 40, 97–101
 binned, 97–101, 559
 distinguished from errors, 387
 plot, 47, 48, 97, 114, 558

AUTHOR INDEX

plot vs. predicted values, not vs. observed values, 157, 158
 social network model, 341–342
 square root, for Poisson model, 341–342
 standard deviation of, 41
 return(), 401
 rnegbin(), 150
 rnorm(), 106, 141, 155, 356, 401, 407
 rnorm.trunc(), 407
 robit regression, 124–125, 133, 320
 Bugs, 384
 generalization of logit and probit, 125
 latent-data formulation, 384
 picture, 124
 robust regression, 110, 131
 rodents, 106, 248, 322
 rowMeans(), 180
 rowSums(), 147, 411
 rpois(), 150
 runif(), 150, 353, 356
 rwish(), 377
 S and S-Plus, *see* R
 sample size and interactions, 438
 sample size calculation, 437–455
 2.8 standard errors, 441
 classical, 439–447
 general concerns, 439
 inference for continuous outcomes, 443–447
 inference for linear regression, 451
 inference for proportions, 439–443
 inference for regression coefficients, 446–447
 inherently speculative, 445, 447
 multilevel models, 447–454
 pictures, 440, 441
 unequal sample sizes, 443
 sample(), 138, 278, 418, 452, 534
 sampling, design for, 437–455
 sapply(), 151
 SAS
 code for matching, 230
 multilevel modeling in, 570–571
 save(), 362
 scale of prior distribution, 355
 scale-up model for estimation in a social network, 333
 scaled inverse-Wishart distribution, 284–287, 298, 376–380
 Scalia, Antonin, 318
 scaling of predictors, 53
 scatterplot
 advice, 553–559

AUTHOR INDEX

623

- data and regression lines
 - superimposed, 35
 - sd(), 56, 460, 462
 - se.coef(), 156, 565
 - se.fixef(), 261
 - secret weapon, 73–74
 - income and voting, 311
 - pictures, 19, 74, 84
 - selection bias, 168, 231, *see also* causal inference
 - selection on observables, *see* ignorability
 - sem package, 223
 - separation in logistic regression, 104, 107, 122
 - Sesame Street, 196, 216–220, 231, 509–511
 - sex ratio of births, 27, 137–139
 - shrinkage and partial pooling, 477
 - sigma.hat(), 273, 565
 - significance, *see* statistical significance
 - significant digits and uncertainty, 561
 - sim(), 43, 142, 143, 392, 565
 - simple and complex models, 416
 - simulation, 19–20, 137–166
 - combined with analytic calculations, 148
 - comparing simulated to actual data, 158–165
 - compound models, 150–151, 537–538
 - coverage of confidence intervals, 156
 - displaying uncertainty in a fitted model, 149
 - fake data, 50, 155–158
 - generalized linear models, 148–151
 - how many draws are needed, 153
 - logistic regression, 148
 - matrix of simulated parameters and predictions, 146, 149
 - nonlinear predictions, 144–148
 - overdispersed models, 150
 - posterior predictive checks, 158–165, 513–524
 - predictive, 148–151
 - probability models, 137–140
 - regression inferences, 140–148
 - replicated datasets, plotted, 160, 163
 - saved as vectors and matrices, 353, 358
 - why necessary, 141
 - slope, *see* varying intercepts and slopes
 - small multiples plot, 255, 257, 266, 291, 560
 - logistic regression, 307
 - small-area estimation, 301–310
 - smoking, 36, 241–243
 - data matrix, 242, 243
 - Social Indicators Survey, 529–543
 - social networks, 332–342
 - Bayesian inference, 409–413
 - graph of data, 335
 - group sizes, 339
 - predicted from demographics, 337
 - residuals, 341–342
 - soft constraint, 257
 - software, 565–574, *see also* R, Bugs, Stata, SAS, SPSS, AD Model Builder, HLM, MLWin
 - data and code for examples, 11
 - getting started, 565
 - multilevel modeling, 573
 - S and S-Plus, *see* R
 - WinBugs and OpenBugs, *see* Bugs
 - speed dating experiment, 322, 323
 - speed of light, 159
 - splines, 298
 - split-plot latin square, 498–501, 509
 - SPSS, multilevel modeling in, 572
 - square root transformation, 249, 535
 - standard deviation, *see* variance
 - standard error, 17, 40
 - picture, 40
 - proportions, 17
 - standardizing predictors, 54–57, 96
 - Stata
 - multilevel modeling in, 569–570
 - reading in data from, 50, 411
 - state-level opinions from national polls, 4–5, 301–310, 493–494
 - Bugs model, 381
 - statistical inference, 16–17, 37–42
 - graph of uncertainty, 40, 83
 - measurement error model, 16
 - sampling model, 16
 - standard error, 40, 83
 - statistical significance, 42, 69, 83, 94
 - limitations of, 481–484
 - multilevel model, 271
 - problems with, 22–23
 - sample size and power, 440
 - stochastic learning in dogs, 515–524
 - Bugs model, 517, 521–524
 - model comparison, 526
 - stop and frisk, *see* police stops
 - storable votes, 120–124, 331–332, 386
 - Bugs model, 383
 - data and fitted curves, 121
 - strategy of debugging, 416
 - structural equation modeling, 231
 - subclassification, 206–207, 229
 - subset option in lm() and glm(), 107, 126, 538
 - subsetting data, 326, 357, 547
 - speeding computation, 418

624

sum of squares
 analysis of variance, 488
 least squares estimation, 387

sum(), 147

superpopulation, 167, 173
 analysis of variance, 491, 500
 variance, 459–462
 computing in Bugs, 460
 graph, 461

Supreme Court voting, ideal-point model
 for, 317

survey design, 437

survey weighting, 301–310

switches, 165

t distribution, 124, 131, 372, 428

table(), 353

tables, 563

teachers, effect of, 459

teaching evaluations and beauty, 51

test summary, 513–515
 graphical, 160, 163, 516, 519, 522, 523
 numerical, 23, 159, 161, 521

text editor, 565

thinning in Bugs, 418, 518

thresholds, for ordered logit or probit,
 119–120

tied election, probability of, 148

time series, 297
 checking a fitted model, 163–165
 cross-sections, 243–244, 248

tlm(), 124, 133, 567

tobit model, 126, 132

topcoding, for missing-data imputation,
 534

traffic accidents, 110–111

transformation, 53–77, 548
 idiosyncratic, 65
 linear, 53–54
 logarithmic, 59–65, 98, 252
 interpreting variance parameters,
 327
 square root, 65, 249, 535

treatment effect, *see* causal inference

true values in fake-data simulation, 155,
 363

truncated normal distribution, 407

tsls(), 223

twins, 138

two-factor experiment, 289–290

two-level classical regression, 240, 248, 270

two-stage least squares, *see* causal
 inference, instrumental variables

two-stage model for mixed discrete/
 continuous data, 126, 537

two-way analysis of variance, 495, 496

AUTHOR INDEX

U.S. Census, 277, 301, 308

UCLA, 332

Ulysses, 339

Umacs, 337, 410–413, 567

uncertainty, as distinguished from
 variability, 457–459

uncontested elections, as missing data, 145

underdispersion, 21, 22

unemployment series, graph of data and
 replications, 163–165

unexpected patterns, discovering through
 graphs, 551

unexplained variance, *see* R^2

unique(), 348

units, 37, 251, 553

unmodeled data and parameters, 364, 367,
 378

unordered categorical regression, 124

updating functions in R, 399–402

uranium, as county-level predictor in
 radon model, 266

utility theory, 128

validation, 47–49

validity, 45, 174

value added by schools, 458–459, 485

value function, 128

value of a statistical life, 197

var(), 477

variability, as distinguished from
 uncertainty, 457–459

variance
 explained and unexplained, 41,
 473–477
 group level, 270
 models for, 297
 multiple error terms, 264
 non-nested models, 290, 291
 predictive, 274
 ratio of between to within, 258
 residual, 41
 superpopulation and
 finite-population, 459–462

varying intercepts, *see* multilevel model

varying intercepts and slopes, 1, 237,
 279–289, 549
 Bayesian perspective, 346
 Bugs, 375–379
 computing using lmer(), 282
 graph, 450
 group-level predictors, 280, 379–380
 interactions, 282–283
 logistic regression, 310–314
 non-nested model, 291
 notation, 284
 pictures, 238

AUTHOR INDEX

625

- varying slopes without varying intercepts,
283–284
- vector of simulations, 353, 358
- vector-matrix notation, 37–38, 284–287
 - Bugs, 361
- Vietnam War draft lottery, 225–226, 230
- visual and numerical comparisons of
 - replicated to actual data, 164
- voting and income, 79–84, 105, 107,
310–314

- Wald estimate for instrumental variables,
219, 221
- wavelets, 298
- weight
 - age and, 75
 - example of a lognormal distribution,
15
 - height and, 74, 402–408
- weighted average, 19
- weighted least squares, 389
- wells in Bangladesh, 86–92, 105, 193
 - arsenic levels, 90
 - choice models, 127–131
 - map, 87
- when does multilevel modeling make a
difference, 247
- WinBugs, *see* Bugs
- WinEdt, 565
- Wishart distribution, 284–287, 298,
376–380
- world wide web connect times, 492–493

- χ^2 test, 25–26, 114

- z-score, 50, 54
- zero-inflated Poisson model, 126–127
- zinc and HIV, 443–447