

Index

- abdominal aortic aneurysm, 86–8
- abdominal CTs, 63–4
- abdominal paracentesis, 273–4
- abdominal trauma
blunt renal trauma, 117
Chance fracture of L4, 114–15
diaphragm rupture, 104–5
focused abdominal sonogram for trauma (FAST), 106
liver trauma, 110–12
splenic laceration, 106–9
- abdominal X-rays, 60, 62
bowel gas pattern, 48, 60
calcification, 61
solid organs, 60
- acalculous cholecystitis, 276–7
- achalasia, 43, 78–9
- acoustic neuroma, 223–4
- air space shadowing, 15
- airway management, 155
- anaesthesia
challenges, 210
diagnostic radiology, 211
hazards to anaesthetists, 210
monitoring, 210
safe management of patients, 210
sedation versus anaesthesia, 211–12
- angiography, *see* CT angiography *and* MR angiography
- angiotensin II, 85
- anisotropy, 289
- ankylosing spondylitis, 150–2
- anterior mediastinal masses, 41–2
- aortic aneurysm rupture, 86
- aortic arch, 57
- aortic dissection, 38
- aortic injury, 97–9
- appendicitis, 74
- appendololiths, 74
- ARDS, 49–50
- aspiration of foreign body, 23–4
- atrial enlargement, 46
- atrial septal defect, 47–8
- biliary stent, 235–6
- bladder injuries, 118–19
- blunt abdominal trauma, *see* abdominal trauma
- blunt chest trauma, 100–2
- Boerhaave's syndrome, 33
- bowel gas pattern, 60
extraluminal gas, 72–3
small bowel infarction, 92
small bowel obstruction, 65–7
- bowel trauma, 114–15
- brain glioma, 203–4
- brain injury, 205–8
- caecal volvulus, 70
- calcification, 61, 74
- Caplan's syndrome, 147
- cardiac enlargement, 34–5, 44–8
- cardiac injury, 103
- cavitating malignancy, 55
- central line insertion, 271–2
- central pontine myelinolysis (CPM), 227
- cerebral contusions, 206
- cerebral infarction, 195–9
- cerebral metastasis, 200–2
- cervical spine
airway management with fractures, 155
ankylosing spondylitis, 150–2
clay shoveler's fracture, 169–70
craniocervical junction, 144–5
diffuse idiopathic skeletal hyperostosis (DISH), 154
extension teardrop fracture, 163
flexion teardrop injury, 164–5
hangman's fracture, 161–2
imaging, 130–3
importance in intubation, 130
injury, 139–41
Jefferson fracture, 157–8
lateral view, 138–9
locked facet injuries, 166–8
occipito-atlantal dissociation, 156
odontoid fracture, 159
open mouth AP radiograph, 136–7
pre-cervical haematoma, 136
rheumatoid arthritis, 145–9
segments, 130
soft tissue contour, 134
unconscious/obtunded patients, 140
- cervical spondylosis, 143, 225–6
- Chance fracture of L4, 114–15
- chest trauma
aortic injury, 97–9
blunt chest trauma, 100–2
cardiac injury, 103, 285
pulmonary contusion, 102–3
steering wheel injuries, 100

Index

- chest X-rays
 air space shadowing, 15
 bones, 6
 diaphragm, 5
 heart and mediastinum, 4–5
 interpretation, 6–8
 interstitial pattern
 shadowing, 26
 labels, 2
 large airways, lungs and
 pleura, 3, 4
 opacification, 7, 8, 14, 17
 projection and patient
 position, 2–3
 quality of film, 3
 review areas, 6, 57
 side marker, 3
 silhouette sign, 7
 soft tissues, 6
- children, 211, 220
- cholangiogram, 235–6
- cholecystitis, 276–7
- cirrhosis, 238–9, 273
- claustrophobia, 211
- clay shoveler's fracture, 169–70
- Clostridium difficile*, 68
- colitis
 necrotizing enterocolitis, 76
 pan colitis, 72
 pseudomembranous, 68–9
- collapsed lung, 29
- colonic diverticulitis, 80
- colo-rectal stents, 241
- compartment syndrome, 123–5
- competency-based training
 and assessment, xvi
- contrast medium, 175, 215
- cranio-cervical junction, 144–5
- CT angiography, 85, 247–8
 complications, 245
- CT-guided lung biopsy, 233–4
- CT pulmonary angiography,
 36–7
- CT scans
 acute subdural haematoma,
 187–9
 analysis, 176, 181
 attenuation, 173
- brain injury, 205–8
 cerebral infarction, 195–9
 cerebral metastasis, 200–2
 extradural haematoma,
 190–1
 Hounsfield units, 173
 image formation, 173
 intracerebral haematoma,
 192–4
 intravenous contrast
 medium, 175
 malignant brain glioma,
 203–4
 MRI versus CT, 216
 multi-slice CT, 174
 protocol variables, 175–6
 spiral CT, 174
 subarachnoid haemorrhage,
 182–3
 tissue density, 173
 windowing, 173–4
- degenerative cervical
 spondylosis, 225–6
- dens fracture, *see* odontoid
 fracture
- dextrocardia, 3
- diagnostic radiology, 211
- diaphragm rupture, 104–5
- diffuse idiopathic skeletal
 hyperostosis (DISH), 154
- discitis, 230–1
- diverticular abscess, 80–1
- Doppler ultrasound, 254
- duplex ultrasound, 246, 254
- echocardiography, 279
 focused echocardiography,
 283
 left ventricular function,
 281–2
 pre-operative, 282
 stress echocardiography, 282
 training, 283
 transoesophageal (TOE),
 282–3
 transthoracic (TTE), 279–81
- Eisenmenger syndrome, 47
- emergency patients, 211
- endotracheal tubes, 40
- endovascular aneurysm repair
 (EVAR), 87–8
- epidural abscess, 230–1
- extension teardrop fracture,
 163
- faecoliths, 74
- fasciotomy, 123
- fat embolism, 123
- femoral fractures, 122–5
 intertrochanteric fracture of
 the hip, 126–7
- femoral nerve block, 301–2
- flexion teardrop injury, 164–5
- focused abdominal sonogram
 for trauma (FAST), 106
- focused echocardiography,
 283
- FRCA examination, xv–xvi
- gadolinium, 215
- gallstones, 60
- Glasgow Coma Score (GCS),
 183–4
- glioma, 203–4
- haemothorax, 103
- Hampton's hump, 36
- hangman's fracture, 161–2
- head injuries, 190, 205–7
- hiatal hernia, 43
- hip fractures, 126–7
- Hounsfield units, 173
- HU values, *see* Hounsfield
 units
- ileocaecal valve, 71
- inferior vena cava (IVC) filters,
 242–3, 245
- intensive care patients, 220
- intensive care units
 echocardiography, *see*
 echocardiography
 ultrasound, 255

- interstitial pattern shadowing, 26
- intertrochanteric fracture of the hip, 126–7
- interventional radiology, 232
- intracerebral haematoma, 192–4
- intracranial sinus thrombosis, 229
- intravenous contrast medium, 175
- intussusception, 77
- Jefferson fracture, 157–8
- juvenile rheumatoid arthritis, 148–9
- kidneys, 60
- kidney stones, 60, 61
- kidney trauma, 117
- kyphoscoliosis, 56
- lap belt injury, 114
- large bowel obstruction, 70–1
- laryngoscopy, 130
- left ventricular function, 281–2
- leg ischaemia, 246
- liver, 60
cirrhosis, 238–9, 273
portal hypertension, 238–9
- liver trauma, 110–12
- lobar collapse, 19–21
- locked facet injuries, 166–8
- lung biopsy, 233–4
contraindications, 233
- lung cavitation, 54–5
- lung collapse, *see* lobar collapse *and* collapsed lung
- malignant brain glioma, 203–4
- mediastinal masses, 41–2
- micro-shock, 220
- miliary nodules, 27–8
- mitral valve disease, 46
- MR angiography, 85, 215, 246–9
complications, 248–9
- MRI
anaesthesia, 219–20
anaesthetic monitoring, 216
bioeffects of electromagnetic radiation, 218
contraindications, 218
contrast agents, 215
CT versus MRI, 216
hazards, 218, 221
image formation, 212–13
intensive care, 220
magnetic field strength and ferromagnetic attraction, 216–17
main uses, 212
micro-shock, 220
monitoring patients, 219
tissue contrast, 213–15
- multi-slice CT, 174
- nasogastric tubes, 39
- National Radiological Protection Board (NRPB), 218
- necrotizing enterocolitis, 76
- needle gauge, 265
- needle visualization, 259–69
problems and tips, 268–9
- needling technique, 292–3
- neuronal tissue, 293–4
- obesity, 293, 294
- occipito-atlantal dissociation, 156
- odontoid fracture, 159
- oesophageal stents, 240–1
- oesophagectomy, 43
- oesophagus, 78
- opaque hemithorax, 28–9
- open book pelvic fractures, 120
- orotracheal intubation, 130
- pan colitis, 72
- pancreatitis, 60, 89–91
- paralytic ileus, 71
- patient archiving and communication systems (PACS), xiii
- pelvic trauma
bladder injuries, 118–19
Chance fracture of L4, 114–15
complications of fractures, 120–1
femoral fractures, 122–5
intertrochanteric fracture of the hip, 126–7
- percutaneous tracheostomy, 275
- percutaneous transhepatic cholangiogram (PTC), 235–6
- pericardial effusion, 44, 285
- pericardial tamponade, 103, 285
- peripheral nerve stimulation, 287, 293–5
- Pneumocystis* pneumonia, 17–18
- pneumomediastinum, 33
- pneumonia, 14–16
- pneumoperitoneum, 77
- pneumothorax, 3, 31–2, 103
lung biopsy, 233–4
- portal hypertension, 238–9
- porto-systemic shunts, 233, 239
- pre-operative assessment, xix–xx
- pre-operative echocardiography, 282
- pseudomembranous colitis, 68–9
- pulmonary contusion, 102–3
- pulmonary embolism, 36–7, 286
inferior vena cava (IVC) filters, 242–5
- pulmonary fibrosis, 25

Index

- pulmonary hypertension, 46–8
- pulmonary nodules, 27–8, 52–5
- pulmonary oedema, 33–5
- pulmonary venous hypertension, 34
- radio-isotope ventilation/perfusion scanning, 37
- Ranson criteria, 90
- regional anaesthesia, *see* ultrasound-guided regional anaesthesia
- Reid's lobe, 60
- renal artery stenosis, 84–5
- renal tract stones, *see* kidney stones
- renal trauma, 117
- rheumatoid arthritis, 145–9
- ring enhancing lesions, 203–4
- scoliosis, 56
- sedation, 211–12
- sigmoid volvulus, 70
- silhouette sign, 7
- sinus thrombosis, 229
- small bowel infarction, 92–3
- small bowel obstruction, 65–7
- spinal stenosis, 225–6
- spiral CT, 174
- spleen, 60
- splenic laceration, 106–9
- spondylosis, 143, 225
- stress echocardiography, 282
- stroke, *see* cerebral infarction
- subarachnoid haemorrhage, 182–3
anaesthetic management, 183–5
common causes, 185
complications, 186
further investigations, 185–6
- subdural haematoma, 187–9
- superior sagittal sinus thrombosis, 229
- tamponade, 285
- tension pneumothorax, 31–2
- thrombo-embolic disease inferior vena cava (IVC) filters, 242–3, 245
- tracheostomy, 275
- transjugular intrahepatic porto-systemic shunt (TIPS), 237–9
- transoesophageal echocardiography (TOE), 282–3
- transthoracic echocardiography (TTE), 279–81
- transversus abdominus plane (TAP) block, 303–4
- trauma
abdominal, *see* abdominal trauma
chest, *see* chest trauma
pelvic, *see* pelvic trauma
- ultrasound
abdomen, 256–7
abdominal paracentesis, 273–4
acalculous cholecystitis, 276–7
advantages and disadvantages, 259
central line insertion, 271–2
coarse gain, 253
colour Doppler and power Doppler, 254
depth control, 254
Doppler ultrasound, 254–5
duplex Doppler, 254
focused abdominal sonogram for trauma (FAST), 106
image display, 253–4
image formation, 251–2
intensive care units, 255
musculoskeletal, 259
optimizing the image, 255
percutaneous tracheostomy, 275
probe frequency, 253
- thoracic, 255–6
time gain compensator (TGC), 253
use in anaesthesia, 251
vascular, 257–9
- ultrasound-guided procedures, 259–60
angles of approach of the needle, 267
background medium, 266
checks prior to passing the interventional needle, 268
depth, 263
focus, 262
hand position, 264–5
initial scan, 263–4
longitudinal or transverse approach, 266–7
needle gauge, 265
needle visualization, 265, 266
objectives, 260
probe covers and patient preparation, 264
probe frequency, 262
probes, 261–2
problems and tips, 268–9
time gain compensator (TGC), 263
- ultrasound-guided regional anaesthesia, 287–8
anatomy, 295
anisotropy, 289
appearance of nerves, 293–4
application, 294–5
femoral nerve block, 301–2
frequency, resolution and depth, 288–9
interscalene approach to the brachial plexus, 299
machines, 288
needles and needling technique, 292–3
patients, 293
techniques, 295–8
transversus abdominus plane (TAP) block, 303–4
- unconscious patients, 140
- ureteric obstruction, 82–3
- Valsalva manoeuvre, 275
- Volkman contracture, 123