

Cambridge University Press

978-0-521-76058-4 - Next Generation Antidepressants: Moving Beyond Monoamines to Discover Novel Treatment Strategies for Mood Disorders

Edited by Chad E. Beyer and Stephen M. Stahl

Index

[More information](#)

Index

- 5-HT. *See* serotonergic system; serotonin
- 5-HTT. *See* serotonin transporter
- 5-hydroxy-indole acetic acid (5-HIAA)
levels in CSF of depressed patients, 13
- acetylcholine (ACh) system
target for treatment strategies, 20–1
- addictive behavior, 18
- adrenalin 2A receptor (ADRA2A) gene, 125
- agomelatine, 5, 19, 29
- α_2 -adrenoceptors
antidepressant targets, 19
- amitriptyline, 14, 102
- AMPA receptors
antidepressant targets, 21
- anhedonia, 30
animal models, 31
definition, 71
objective laboratory measurement, 82–3
- anhedonia as a depressive endophenotype, 71, 73–83
biological plausibility, 75–6
clinical plausibility, 73–5
cosegregation, 82
familial association, 80
future research directions, 84–5
heritability, 81–2
objective laboratory measurement, 82–3
role of anhedonia in depression, 83–4
specificity, 76–8
state-independence, 78–9
- animal models of depression
affective and cognitive disturbances, 52
anhedonia, 31
barriers to development of novel treatments, 39
brain abnormalities in depression, 35–7
challenges, 30
- chronic antidepressant treatment, 34
- chronic mild stress
model, 33
- current models, 31–3
- forced swim test, 31–3
- forced swim test
modifications, 34
- gender differences, 35
- genetically modified mouse models, 33
- in-vivo brain monitoring, 37–9
- limitations, 30, 39
- modeling depressive symptoms, 30–1
- Morris water maze test, 34
- need for novel models, 29–30
- novel animal models, 33–4
- novelty-induced hypophagia
model, 34
- open space swim test, 34
- preclinical models in translational research, 46–7
- range of approaches, 31
- requirements and aims, 30–1
- stress response, 34–5
- tail suspension test, 31–3
- translational neuroimaging, 49–50
- withdrawal of drugs of abuse, 33
- anticipation (genetic), 96
- antidepressant metabolism
effects of individual genetic variations, 122–3
- antidepressant therapeutic targets
effects of genetic variations in, 123–5
- antidepressant treatment
approaches
combination and augmentation strategies, 3–4
looking beyond monoamines, 7
- new formulations of old medications, 4–5
- staging of major depression, 4
- successive monotherapies, 3
- unmet therapeutic needs, 3
- antidepressant treatment
strategies, 15–23
- 5-HT1A/1B receptor targets, 18
- 5-HT2A/2C receptor targets, 18–19
- 5-HT2C receptor blockers, 5
- 5-HT3/5A/7 antagonism, 19
- acetylcholine (ACh) system, 20–1
- α_2 -adrenoceptor targets, 19
- AMPA receptor
modulators, 21
- atypical antipsychotics, 5
- DA/NE release stimulator (bupropion), 18
- dual 5-HT/NE reuptake inhibitors, 17
- GABA-related
approaches, 21
- glutamate system, 21
- histamine H3 antagonist and SSRI, 19
- HPA axis-related
treatment, 20
- neurokinin receptor
antagonists, 20
- NMDA receptor
antagonists, 21
- non-monoaminergic
approaches, 23
- norepinephrine dopamine disinhibitors, 5
- novel agents with new targets, 7
- selective NE reuptake inhibitors, 16–17
- substance P as target, 20
- triple monoamine
reuptake inhibitors, 17–18
- ultraselective serotonin reuptake inhibitor (escitalopram), 15–16

Cambridge University Press

978-0-521-76058-4 - Next Generation Antidepressants: Moving Beyond Monoamines to Discover Novel Treatment Strategies for Mood Disorders

Edited by Chad E. Beyer and Stephen M. Stahl

Index

[More information](#)

Index

- antidepressants
 animal models of chronic treatment, 34
 discovery of tricyclic antidepressants (TCAs), 2
 early discoveries, 1–2
 effectiveness in major depression, 13
 efficacy of interventions, 28
 history of development, 28–9
 limitations of current medications, 121
 novel drug targets, 29
 proven efficacy in trials, 121
 time lag to take effect, 2, 29
- apolipoprotein E epsilon4 gene, 125
- aprepitant, 20, 29
- atypical antipsychotics
 treatment of depression, 5
- BDNF (brain-derived neurotrophic factor), 3
 functions, 58
 gene, 96, 125
 gene polymorphism, 58
- Beck Depression Inventory (BDI), 77
- benzodiazepines
 co-administration with SSRIs, 21
 for patients with depression, 21
- beta 3 agonists, 23
- beta adrenalin receptor (betaARs) gene, 125
- biomarkers
 applications for neuroimaging techniques, 50
 cognitive deficits in mood disorders, 51–2
 cognitive impairment, 52
 disease biomarkers, 50, 56
 HPA axis dysfunction under stress, 57–8
 negative bias, 52
 neuroimaging of negative bias, 50–1
 patient selection and stratification, 55–6
 pharmacodynamic, 54–5
 pharmacogenomics, 56
 pharmacokinetic, 54–5
 placebo effect, 60
 preclinical animal models, 52
- presymptomatic diagnosis in surrogate populations, 56–7
 role of inflammation in depression, 58–9
 role of neurotrophins, 58
 surrogate biomarkers, 46
 target validation biomarkers, 52–3
 target–compound interaction, 54
 vascular depression, 59–60
- bipolar disorder
 distinction from unipolar depression, 7
 DSM-IV-TR criteria, 45
 functional disorders, 45–6
- bipolar II disorder, 7
- blood–brain barrier (BBB), 55
- BOLD (blood oxygen level-dependent) imaging, 49
- brain imaging
 abnormalities in depression, 35–7
- brain monitoring
 animal models of depression, 37–9
- brain-derived neurotrophic factor. *See* BDNF
- bupropion, 15, 17, 21
 catecholamine (DA/NE) release stimulator, 18
 hydrobromide salt formulation, 4
- California Psychological Inventory, 81
- Cambridge Neuropsychological Test Automated Battery (CANTAB), 52
- candidate gene studies
 MDD, 93–4
- catecholamine (DA/NE) release stimulator
 bupropion, 18
- catechol-*O*-methyltransferase (COMT) gene, 125
- cerebrovascular disease
 vascular depression, 59–60
- chronic mild stress model (animals), 33
- citalopram, 3, 14, 28, 58, 102
 comparison with escitalopram, 16
 pharmacology, 15
- classification of mood disorders, 71
 endophenotypic approach, 71
 limitations of diagnostic criteria, 71
See also DSM-IV
- clomipramine, 35
- cognitive deficits in mood disorders
 neuroimaging biomarkers, 51–2
- cognitive impairment
 biomarkers, 52
 preclinical animal models, 52
- Composite International Diagnostic Interview (CIDI), 120
- copy-number variation (CNV)
 MDD risk studies, 95
- corticotropin-releasing factor (CRF), 20
- cortisol
 glucocorticoid receptor antagonists, 112–14
- COX-2 inhibitors, 23
- cytochrome P450 enzymes (CYPs)
 effects of individual genetic variations, 122–3
- depression, 4
 clinical effectiveness of antidepressants, 13
 clinical symptoms, 30
 comorbidity, 29
 etiology, 29
 health consequences, 120
 hypotheses of, 30
 monoamine hypothesis, 2, 13–14, 30
 morbidity and mortality, 12
 neurotransmitter receptor sensitivity hypothesis, 2
 prevalence, 1, 12, 28, 120
 psychiatric comorbidity, 13
 staging, 4
 subtypes, 12–13
See also bipolar disorder; major depressive disorder (MDD); unipolar depression.
- depression endophenotypes.
See anhedonia endophenotypes

Index

- depressive disorders
 as a spectrum, 7
 diagnostic challenges, 7–11
 desimipramine, 17
 desipramine, 102, 123
 desvenlafaxine, 5, 15, 103
 dexamethasone suppression
 test (DST), 57
 diagnosis
 difficulties with mood
 disorders, 7–11
 distinction between unipolar
 and bipolar depression, 7
 mood disorders as a
 spectrum, 7
 DISCI (Disrupted in
 Schizophrenia-1) gene, 96
 disease biomarkers, 50, 56
 DNA structural variation
 MDD risk studies, 95–6
 dopamine (DA)
 DA/NE release stimulator
 (bupropion), 18
 monoamine hypothesis of
 depression, 2
 dopamine D2, D3, D4 receptor
 (DRD2, DRD3, DRD4)
 genes, 125
 dopamine transporter (DAT)
 blockade, 2
 dopamine transporter (DAT1)
 gene, 125
 dorsal raphe nucleus, 14
 drug discovery and development.
See translational
 neuroimaging; translational
 research
 DSM-IV
 heterogeneity of clinical
 phenotypes, 71
 DSM-IV-TR
 criteria for mood
 disorders, 45
 dual 5-HT/NE reuptake
 inhibitors, 17
 duloxetine, 15, 103
 dysphoria, 30
 endophenotypes
 anhedonia as a depressive
 endophenotype, 73–84
 biological plausibility, 72,
 75–6
 clinical plausibility, 72, 73–5
 cosegregation, 72, 82
 criteria, 72–3
 definition, 71–2
 familial association, 72, 80
 future research directions,
 84–5
 heritability, 72, 81–2
 in MDD, 97–8
 limitations of current
 classification systems, 71
 measurement, 72–3, 82–3
 neuroticism, 98
 specificity, 72, 76–8
 state-independence, 72, 78–9
 validation, 72–3
 value in psychiatric research,
 71–2
See also biomarkers.
 endophenotypic approach, 71–2
 environmental stress
 effects of, 57–8
 epigenetics, 96–7
 epigenomics, 96–7
 escitalopram, 15–16
 clinical trials in depressed
 patients, 16
 comparison with
 citalopram, 16
 comparison with
 venlafaxine, 16
 pharmacology, 15
 safety, 16
 tolerability, 16
 eszopiclone, 21
 etiology of depression, 29
 familiarity studies
 MDD, 90–1
 fatty acid amide hydrolase
 inhibitors, 23
 fluoxetine, 2, 14, 17, 28, 34, 55,
 58, 59, 60, 102, 103
 fluvoxamine, 14, 28, 55, 58, 102
 fMRI (functional MRI), 49
 characterization of mood
 disorders, 49
 use in drug discovery and
 development, 49
 use in preclinical animal
 models, 49–50
 forced swim test
 animal model of depression,
 31–3, 34
 galanthamine, 21
 gamma-aminobutyric acid
 (GABA)
 antidepressant target, 21
 gender differences
 animal models of
 depression, 35
 genetic association studies
 MDD, 93–5
 genetic linkage studies
 MDD, 93
 genetic variations
 enzymes which metabolize
 ADs, 122–3
 therapeutic targets of ADs,
 123–5
 genetically modified mice
 models of depression, 33
 transgenic models of mood
 disorders, 53–4
 genomewide association studies
 (GWAS), 94–5
 glucocorticoid receptor
 antagonists, 112–14
 glutamate system
 antidepressant targets, 21
 glutamatergic targets, 108–11
 glutathione-S-transferase
 (GST) gene, 123
 G-protein beta3-subunit
 (Gbeta3) gene, 125
 Hamilton depression rating
 scale (HAM-D scale),
 12–13
 heritability studies
 MDD, 91–3
 heterogeneity of clinical
 phenotypes, 71
 heterogeneity within
 classification criteria,
 71
 histamine H3 antagonist and
 SSRI, 19
 Huntington's disease (HD)
 association with MDD, 96
 hydroxyl bupropion, 15
 hypothalamic–pituitary–
 adrenal (HPA) axis, 7
 candidate genes for MDD
 risk, 94
 glucocorticoid receptor
 antagonists, 112–14
 melanocortin receptor
 antagonists, 115–16
 role in depression, 30
 role in mood disorders,
 57–8
 role in the stress
 response, 112

Cambridge University Press

978-0-521-76058-4 - Next Generation Antidepressants: Moving Beyond Monoamines to Discover Novel Treatment Strategies for Mood Disorders

Edited by Chad E. Beyer and Stephen M. Stahl

Index

[More information](#)

Index

- targets for drug development, 112–16
 treatment target, 20
 vasopressin receptor antagonists, 114–15
- idazoxane, 19
 imipramine, 17, 28, 35, 46, 102
 Implicit Association Test (IAT), 79
 improving treatments. *See* antidepressant treatment approaches
 individualized medicine. *See* pharmacogenomics
 inflammation
 role in depression, 58–9
 Investigational New Drug requirements, 54
 iproniazid, 2, 46
 isoniazid, 2
- ketamine, 21
- L-5-methyl-tetrahydrofolate (MTHF), 7
 lithium, 55, 58
 LSD, 13–14
 Lu 21004, 18, 19
 Lu AA 24530, 19
- major depression, 12–13
 clinical effectiveness of antidepressants, 13
 depression subtypes, 12–13
 morbidity and mortality, 12
 prevalence, 12
 psychiatric comorbidity, 13
 staging, 4
 major depressive disorder (MDD)
 adoption studies, 92–3
 association with rare genetic variations, 96
 candidate gene studies, 93–4
 common gene hypothesis, 96
 copy-number variation (CNV) studies, 95
 disease burden, 70
 DSM-IV-TR criteria, 45
 endophenotypes, 97–8
 familiarity studies, 90–1
 family studies, 97
 functional disorder, 45–6
 genetic association studies, 93–5
- genetic linkage studies, 93
 genomewide association studies (GWAS), 94–5
 heritability studies, 91–3
 heterogeneity within classification criteria, 71
 identifying depressive subtypes, 97–8
 lack of response to drug treatment, 46
 multiple phenotypes hypothesis, 97
 prevalence, 70
 rare genetic variants approach, 96
 role of trinucleotide repeats, 96
 studies of inheritance patterns, 91
 twin studies, 91–2
 median raphe nucleus, 14
 medicinal chemistry
 glucocorticoid receptor antagonists, 112–14
 glutamatergic targets, 108–11
 melanocortin receptor antagonists, 115–16
 non-monoamine based targets, 108–16
 orphanin FQ/nociceptin receptor agonists, 111–12
 potential multitarget approaches, 104–8
 SNRIs, 103
 SSRIs, 102–3
 SSRIs combined with 5-HT1A receptor antagonism, 104–6
 targets in the HPA axis, 112–16
 tricyclic antidepressants (TCAs), 102
 triple reuptake inhibitors, 106–8
 vasopressin receptor antagonists, 114–15
 melanocortin receptor antagonists, 115–16
 milnacipran, 15, 58, 103
 Minnesota Multiphasic Personality Inventory, 81
 mirtazapine, 19
- monoamine hypothesis of depression, 2, 13–14, 30
 5-HIAA in CSF of depressed patients, 13
 evidence from post-mortem brain tissue, 13
 evidence from therapeutic interventions, 13–14
 role of serotonin in depression, 13–14
 monoamine oxidase (MAO) gene, 125
 monoamine oxidase (MAO) inhibitors, 28
 mood disorders
 as a spectrum, 7
 diagnostic challenges, 7–11
 DSM-IV-TR criteria, 45
 pharmacogenomics, 53
 possible progressive nature, 7–11
 prevalence, 45
 Morris water maze test, 34
 mouse
 genetic models of depression, 33
 transgenic models of mood disorders, 53–4
 MRI (magnetic resonance imaging), 49
 MRS (magnetic resonance spectroscopy)
 PK/PD studies, 55
 multidrug-resistance gene (MDR1), 123
- National Comorbid Study (NCS), 120
n-back task, 51
 nefazodone, 19
 negative bias
 biomarkers, 52
 in depressed patients, 50–1
 neuroimaging biomarkers, 50–1
 preclinical animal models, 52
 neuroimaging
 biomarker research applications, 50
 brain abnormalities in depression, 35–7
 role in translational research, 46
 See also translational neuroimaging

Cambridge University Press

978-0-521-76058-4 - Next Generation Antidepressants: Moving Beyond Monoamines to Discover Novel Treatment Strategies for Mood Disorders

Edited by Chad E. Beyer and Stephen M. Stahl

Index

[More information](#)

Index

- neurokinin receptor
antagonists, 7, 20
- neuropsychological deficits in
mood disorders
neuroimaging biomarkers,
51–2
- neuroticism endophenotype, 98
- neurotransmitter receptor
sensitivity hypothesis of
depression, 2
- neurotrophic factors, 3
- neurotrophins
role in depression, 58
- NMDA receptors
antidepressant targets, 21
- nomifensine, 17
- non-monoaminergic treatment
approaches, 23
- noradrenergic system
and reboxetine, 16
- norepinephrine (NE)
DA/NE release stimulator
(bupropion), 18
monoamine hypothesis of
depression, 2
role in depression, 16
selective NE reuptake
inhibitors, 16–17
- norepinephrine dopamine
disinhibitors, 5
- norepinephrine transporter
(NET) blockade, 2
- norepinephrine transporter
(NET) gene, 125
- nortriptyline, 102
- novelty-induced hypophagia
animal model, 34
- open space swim test
animal model of depression,
34
- orosomucoid 1 and 2 (ORM1
and ORM2) genes, 123
- orphanin FQ/nociceptin
receptor agonists,
111–12
- P38 kinase inhibitors, 23
- Parkinson's disease, 96
- paroxetine, 14, 20, 34, 55, 102
- patient selection and
stratification biomarkers,
55–6
disease biomarkers, 56
pharmacogenomics, 56
PDE inhibitors, 23
- personalized medicine. *See*
pharmacogenomics
- PET (positron emission
tomography), 48
in vivo PK/PD studies,
54–5
role in drug discovery and
development, 48
target–compound
interaction biomarkers, 54
use in preclinical animal
models, 49–50
- pharmacodynamic biomarkers,
54–5
- pharmacodynamics, 50
genetic basis for individual
variations, 123–5
- pharmacogenetics
genetic basis for individual
responses, 121–5
- pharmacogenomic panels, 125
- pharmacogenomics, 53
barriers to clinical
application, 125
clinical decision-making
process, 125–6
clinical testing process, 125
definition, 53
genetic basis for individual
responses, 121–5
mood disorders, 53
pilot studies of individualized
medicine, 126
potential for individualized
medicine, 125
tools for interpreting results,
125–6
translational research
applications, 53
use in patient stratification, 56
- pharmacokinetic biomarkers,
54–5
- pharmacokinetics
genetic basis for individual
variations, 122–3
- phMRI (pharmacological
MRI), 49
- pindolol, 18, 29
- placebo effect
neuroimaging, 60
- preclinical animal models.
See animal models
- presymptomatic diagnosis
neuroimaging in
surrogate populations,
56–7
- progression of mood disorders,
7–11
- proteomics, 53
- proton [¹H] spectroscopy, 55
- psychosocial stressors
effects of, 57–8
- quetiapine, 5
- radiolabeling of drugs, 54–5
- reboxetine, 15, 29, 59
and the 5-HT system, 16
and the NE system, 16
in combination with
SSRIs, 17
long-term efficacy, 17
selective NE reuptake
inhibitor, 16–17
tolerability, 17
versus SSRIs, 17
versus TCAs, 16–17
- reserpine, 14, 102
- risperidone, 19
- selective NE reuptake
inhibitors, 16–17
- sense reversal, 52
- Sequenced Treatment
Alternatives to Relieve
Depression study. *See*
STAR*D study
- serotonergic system
and reboxetine, 16
and SSRIs, 14
- serotonin
evidence for role in
depression, 13–14
levels in post-mortem brain
tissue, 13
monoamine hypothesis of
depression, 2
- serotonin receptors
1A/1B receptor targets, 18
2A receptor gene, 125
2A/2C receptor targets,
18–19
2C receptor blockers, 5
3/5A/7 receptor antagonism,
19
- serotonin transporter
blockade, 2
effects of genetic
polymorphisms, 123–4
gene polymorphism, 53
MDD candidate gene
studies, 94

- SERT. *See* serotonin transporter
- sertraline, 102
- Sigma 1 agonists, 23
- smoking cessation and depression, 21
- Snaith–Hamilton Pleasure scale, 76
- SNRIs (selective norepinephrine reuptake inhibitors), 28–9
- SNRIs (serotonin/norepinephrine reuptake inhibitors), 103
- SSR/NRI combined approach, 28–9
- SSRIs (selective serotonin reuptake inhibitors), 102–3
- and the serotonergic system, 14
- combined with 5-HT1A receptor antagonism, 104–6
- combined with histamine H₃ antagonist, 19
- combined with reboxetine, 17
- comparative efficacy in major depression, 14
- delayed onset of action, 14
- efficacy compared with TCAs, 14
- fluoxetine, 2
- introduction of, 28–9
- long-term efficacy, 14–15
- relapse features, 14–15
- tolerability, 14
- versus reboxetine, 17
- St. John's wort, 123
- staging
- major depression, 4
- unipolar depression, 4
- STAR*D study, 3, 70
- stress
- role in depression, 30
- stress response
- animal models, 34–5
- substance P
- antidepressant targeting, 20
- suicide and depression, 29, 30, 120
- tail suspension test
- animal model of depression, 31–3
- target validation biomarkers, 52–3
- target–compound interaction
- biomarkers, 54
- TCAs (tricyclic antidepressants), 28, 102
- discovery of, 2
- efficacy compared with SSRIs, 14
- versus reboxetine, 16–17
- Temperament and Character Inventory (TCI), 80
- therapeutic dose range, 55
- therapeutic interventions
- evidence for the monoamine hypothesis of depression, 13–14
- tianeptine, 29
- transgenic animal models, 53–4
- translational neuroimaging, 47–50
- biomarker research
- applications, 50
- biomarkers of cognitive deficits, 51–2
- biomarkers of negative bias, 50–1, 52
- disease biomarkers, 50, 56
- HPA axis dysfunction under stress, 57–8
- MRS studies, 55
- patient selection and stratification biomarkers, 55–6
- pharmacodynamic biomarkers, 54–5
- pharmacogenomics
- applications, 53
- pharmacokinetic biomarkers, 54–5
- placebo effect, 60
- preclinical animal models, 52
- presymptomatic diagnosis in surrogate populations, 56–7
- radiolabeling of drugs, 54–5
- role of inflammation in depression, 58–9
- role of neurotrophins, 58
- target validation biomarkers, 52–3
- target–compound interaction biomarkers, 54
- therapeutic dose range, 55
- transgenic mouse models, 53–4
- vascular depression, 59–60
- translational research
- aims, 46
- drug discovery and development, 46
- lack of response to drug treatment, 46
- pharmacogenomics
- applications, 56
- preclinical animal models, 46–7
- role of neuroimaging techniques, 46
- surrogate biomarkers, 46
- trazodone
- new formulation, 4
- tricyclic antidepressants. *See* TCAs
- triiodothyronine, 18
- trinucleotide repeats
- role in MDD, 96
- triple monoamine reuptake inhibitors, 5, 15, 17–18, 106–8
- tryptophan hydroxylase (TPH) gene, 125
- TSNAX* (Translin-associated factor X) gene, 96
- twin studies
- inheritance of MDD risk, 91–2
- UDP-glucuronosyltransferases (UGTs) genes, 123
- unipolar depression
- distinction from bipolar disorder, 7
- staging, 4
- uridine, 23
- vascular depression, 59–60
- vasopressin, 19
- vasopressin receptor
- antagonists, 114–15
- venlafaxine, 5, 15, 16, 29, 103, 123
- vilazodone, 18
- Wisconsin Card Sorting Test (WCST), 51
- Wisconsin General Testing Apparatus (WGTA), 52
- withdrawal of drugs of abuse
- animal models, 33
- depressive symptoms, 30