

Index

Boldface page numbers indicate definitions of terms or statements of results

- abelian group
 - complete, **14**
 - separated, **14**
- absolute Frobenius lift, **265**
- absolute value, *see also* norm
- additive value group, **16**
- adjoint form (of a twisted polynomial), **86, 246**
- analytic elements, **144, 156**
- approximation lemmas, 20, 30, 147, 206
- Artin–Hasse exponential series, **162, 293**
- Artin–Schreier extension, **51, 317, 334, 341**
- Artin–Schreier theory
 - nonabelian, 259, 315, 357
- average norm (of a difference module), **281**

- B*-pair, 363
- Bézout domain, **276**
- Baer sum, **82**
- Banach contraction mapping theorem, 41, 269
- Banach space, **23**
- Barsotti–Tate group, *see also* *p*-divisible group
- base change
 - of a difference module, **244**
 - of a differential module, **81, 98, 107, 178**
- Berkovich spaces, 8
- Bessel differential equation, **120, 328**
- binomial series, 50
- Birkhoff factorization, 149
- bounded homomorphism, **14, 94**
- bounded Robba ring, 217, **263, 359**

- Calabi–Yau variety, 355
- Cauchy sequence, **14**
- Cauchy theorem, **120**
 - p*-adic, **154, 302**
- center (of a Frobenius lift), **265**
- change-of-basis matrix, **81, 244**
- clean differential module, **214**
- Cohen structure theorem, 333
- coherent locally free module, **144**
- compactness (of a closed interval), **136, 203, 208**
- companion matrix, **64**
- compatible seminorm on a module, **17**
- complete abelian group, **14**
- completion (of an abelian group), **15**
- connection, 81, 91
- constant subring (of a differential ring), **78**
- constituents (of a differential module), **80**
- contraction mapping theorem, 41, 269
- convexity, 190
- Cramer’s rule, 66
- Crew’s conjecture, 335
- crystalline cohomology, 353
- cyclic vector
 - of a difference module, **246**
 - of a differential module, **84**
 - theorem, **84, 89, 246, 260, 311**
- cyclotomic character, **362**

- de Rham cohomology, 347
- de Rham representation, 336, **362**
- decomposition theorem
 - for difference modules, **252**
 - refined strong, **179, 214**
 - refined visible, **113**
 - strong, **177**
 - visible, **107**
- derivation of rational type, **164**
- Dieudonné–Manin classification theorem, **258, 283, 316, 332**

- difference
 - field, **244**
 - difference-closed, **247**
 - isometric, **249**
- Galois theory, 258
- ideal, **244**
- module, **244**
- ring, **244**
- differential
 - Galois theory, 91
 - ideal, **78**
 - module, **78**
 - normed, **96**
 - on an open annulus, **152**
 - trivial, **78**
 - unipotent, **78, 153**
 - ring or field, **77**
 - slopes, **211**
- dimensioned constants, 27
- discrepancy
 - of a differential module, **203**
 - of a function, **202**
- discretely valued field, **16**
- dual
 - of a difference module, **245**
 - of a differential module, **82**
- duality for convex functions, 43
- dualizable difference module, **245**
- Dwork
 - exponential series, 181, **292**
 - trick, 273, 275, 280, 281, **295**,
329, 336
- eigenvalues
 - prepared, **123**
 - weakly prepared, **123**
- elementary divisors, **63**
- elementary matrix, **55**
- elementary row or column
 - operation, **56**
- elliptic curve, 3
- elliptic integral, 5
- equivalence (of *p*-adic exponents), **224**
- étale cohomology, 5, 354
- étale difference module, *see also* unit-root
difference module
- étale fundamental group, 323
- exponents
 - of a differential module, **122**
 - of a *p*-adic differential module, **226**
- extended Robba ring, **282**
- exterior power, **57, 82**
- exterior product \wedge , 57
- extrinsic generic radius of convergence, 157

- Fekete's lemma, 94
- field of norms, 357

- filtration
 - lower numbering, **52**
 - upper numbering, **52, 321**
- finite local monodromy (*p*-adic), **318**
- formal adjoint (of a twisted polynomial), **86**,
246
- Fréchet completeness, 139
- Frobenius
 - antecedent, **173, 181, 213, 225, 303, 341**,
342
 - descendant, **171, 181, 193, 196, 206**
 - lift, **265**
 - absolute, **265**
 - standard, **297**
 - zero-centered, **266**
 - off-center, 181, 196
 - structure (on a differential module), 160,
291, 327
- Fuchs theorem, **125**
 - p*-adic, **221, 231, 302**
- Fuchsian singularities, 130
- full spectrum, **100**
- fundamental solution matrix, **123, 221, 234**,
295, 302, 308

- Galois cohomology, 363
- Galois descent, 129, 173
- Gauss
 - (semi)norm, **26, 138, 282**
 - for twisted polynomials, **104**
 - (semi)valuation, **26**
 - for twisted polynomials, **104**
- Gauss–Jordan elimination, 63, 141
- Gauss–Manin connection, 351
- generalized power series, 31, 282
- generic Newton polygon, **266, 280**
- generic point, **160**
- generic radius of convergence, **156**
 - extrinsic, **157**
 - intrinsic, **157**
- Gevrey functions, 130
- gluing lemma, **142, 208**

- Hadamard three-circle theorem, 139
- Hahn–Banach theorem, 32
- Hasse–Arf theorem, 322
- Hasse's theorem, 3
- henselian field, **46, 263**
- Hensel's lemma, 41
- Hermitian transpose, **56**
- Herr complex, **361**
- Hilbert–Noether theorem, 173, 314, 359
- Hodge polygon
 - of a difference module, **254**
 - of a matrix, **63**
- Hodge–Newton decomposition, **65**
 - for difference modules, **254**

- homomorphism
 - bounded, **14**
 - isometric, **14**
 - horizontal
 - element (of a differential module), **78**
 - morphism (of differential modules), **81**
 - Horn inequalities, 71
 - hypergeometric
 - differential equation, **5**, 120, 159, 222, 348
 - series, **5**, 159
 - Igusa polynomial, **6**
 - index
 - of a differential module, **126**
 - of nilpotency, **123**
 - inertia group, **49**, 323
 - intrinsic
 - generic radius of convergence, **157**
 - radius, **157**
 - subsidiary generic radii of convergence, **162**
 - invariant factors, **63**
 - inverse difference ring, **245**
 - irregularity, **119**, 162
 - isoclinic difference module, **250**
 - isometric
 - difference field, **249**
 - homomorphism, 14
 - Jordan–Hölder constituents, **80**
 - Kiehl–Tate theorems, 149
 - Kummer theory, 50
 - Lang torsor, 259
 - Lefschetz
 - principle, 119
 - trace formula, 5
 - limit (of a sequence), **14**
 - Liouville number, **219**
 - Littlewood–Richardson
 - numbers, 71
 - local horizontal section, **154**
 - locally free module, **84**
 - lower convex hull, **36**, 105
 - lower numbering filtration, **52**
 - Mal’cev–Neumann series, 31, 282
 - master factorization theorem, **39**, 104, 140, 142, 246
 - matrix of action, **80**, **244**
 - maximally complete field, **28**
 - metrically equivalent seminorms, **14**
 - minor (of a matrix), **61**
 - mirror symmetry, 355
 - monodromy transformation, **122**
 - multiplicative
 - seminorm, **15**
 - value group, **16**
 - mysterious functor, 363
 - Newton above Hodge theorem, **64**
 - for difference modules, **254**
 - Newton polygon
 - generic, **266**, **280**
 - of a difference module, **254**, **279**
 - of a matrix, **64**
 - of a polynomial, **36**
 - of a power series, **138**
 - of a twisted polynomial, **104**, 246
 - semicontinuity theorem, **266**, 280, 311, 329
 - special, **266**, **280**
- Newton’s method, 264
 - nilpotency index, **123**
 - nonarchimedean
 - group, ring, field, **16**
 - seminorm, **16**
 - norm, **14**
 - Gauss, **26**
 - for twisted polynomials, **104**
 - p*-adic, **26**
 - spectral, 95
 - supremum, **17**
 - t*-adic, **25**
 - trivial, **25**
 - normal (matrix), **60**
 - normed differential module, **96**
 - off-center Frobenius antecedents and
 - descendants, 181, 196
 - operator norm, 56, **94**, 121
 - opposite
 - difference ring/module, **245**
 - ring, 86
 - optimal basis, **198**
 - order of log-growth, **308**
 - Ostrowski’s theorem, **17**, **27**
 - outer provinces, rebellious, 300
 - overconvergent *F*-isocrystal (on the affine line), **352**
 - p*-adic
 - Cauchy theorem, **154**, 302
 - Fuchs theorem, **221**, **231**, 302
 - Hodge theory, 357
 - integers, ring of, **26**
 - Liouville number, **219**
 - local monodromy theorem, 2, 114, 234, 279, **327**
 - unit-root, **318**
 - norm, **26**
 - numbers, field of, **26**
 - Turrittin theorem, 338

- p*-divisible group, 285, 336, 364
- p*-power roots of unity, 226
- perfection, **249**, 339
- (ϕ, Γ) -module, 359
- Picard–Fuchs
 - equation, 7
 - module, 88, 122, 294, **347**
- piecewise affine function, **136**
- Poincaré pairing, 353
- potentially semistable representation, 336, 363
- prepared eigenvalues, **123**
- principal minor (of a matrix), **61**
- product formula, **26**
- product seminorm, **23**
- projective module, **84**
- pseudoconvergent sequence, **30**
- pseudolimit, **30**
- pure
 - difference module, **250**, **277**
 - differential module, **100**
- quasicontant differential module, **327**
- quasiunipotent
 - differential module, **327**
 - monodromy, **122**
- quotient seminorm, **16**
- radii of optimal convergence, **198**
- radius of convergence (of a differential module), 154
- ramification numbers, **211**
- Rapoport–Zink spaces, 285
- rational type, **164**
- real
 - semivaluation, **16**
 - valuation, **16**
- rebellious outer provinces quelled by Frobenius, 300
- refined decomposition theorem
 - strong, **179**, 214
 - visible, **113**
- refined differential module, **100**, 342
- regular differential module, **119**, 234, 347
- residue field (of a nonarchimedean field), **17**
- rigid analytic geometry, 8
- rigid cohomology, 353
- Robba condition, **223**
- Robba ring, **264**, 360
 - bounded, 217, **263**, 359
 - extended, **282**
- Schauder basis, **32**
- semiabsolute value, *see also* seminorm
- semicontinuity theorem for Newton polygons, **266**, 280, 311, 329
- seminorm, **13**
 - compatible, **17**
 - Gauss, **26**
 - metrically equivalent, **14**
 - multiplicative, **15**
 - product, **23**
 - quotient, **16**
 - submultiplicative, **15**
 - supremum, **17**
 - topologically equivalent, **14**
- semistable
 - difference module, **281**
- semivaluation, **16**
 - Gauss, **26**
- separated abelian group, **14**
- shearing transformation, **125**, 226, 227, 234, 310
- Shilov boundary, 149
- singular value, 95
 - archimedean, **56**
 - decomposition, **56**
 - nonarchimedean, **63**
- slope (of a polynomial), **36**
- slope filtration theorem, **279**, **298**, 340, 364
- Smith normal form, **63**
- solvable differential module, **211**, 294, 302
- special Newton polygon, **266**, **280**
- spectral norm, 95
- spectral radius, **94**
- spectrum
 - full, **100**
 - visible, **105**
- spherically complete metric space, **28**
- standard Frobenius lift, **297**
- strong decomposition theorem, **177**
- strong triangle inequality, **16**
- strongly difference-closed difference field, **247**
- subharmonicity, 190
- submultiplicative seminorm, **15**
- subsidiary (generic) radii (of convergence), **162**
- supersingular elliptic curve, 6
- supremum (semi)norm, **17**
- supremum-equivalent seminorm, **18**
- t*-adic valuation, **25**
- tame degree (of an extension), **49**
- tamely ramified extension, **49**, 101, 113, 179, 323
- Taylor series, **90**, 159, 173, 226, 296
- Teichmüller lift, **7**, **259**, 358
- tensor product
 - of difference modules, **244**
 - of differential modules, **81**
- topologically equivalent seminorms, **14**

- totally ramified extension, **48**
 totally wildly ramified extension, **49**
 trace (element or component), **82**
 trace-zero component, **82**
 transfer theorem, **158**
 regular singular, **234**, **295**
 triangle inequality, **14**
 strong, **16**
 trianguline representation, **364**
 trivial
 difference module, **244**
 differential module, **78**
 Turrittin–Levelt–Hukuhara decomposition
 theorem, **114**, **128**, **338**
 twisted polynomial
 over a difference ring, **246**
 over a differential ring, **85**
 type (of a field element), **218**
- ultrametric seminorm, **16**
 unipotent differential module,
 78, **153**
 unit-root
 difference module, **250**, **298**,
 315, **318**
 p -adic local monodromy theorem, **318**
 unramified extension, **47**
 upper numbering filtration, **52**, **321**
- valuation, **16**
 Gauss, **26**
 for twisted polynomials, **104**
 t -adic, **25**
 valuation ring (of a nonarchimedean field), **17**
 value group, **16**
 variation of Hodge structures, **335**
 visible decomposition theorem, **107**, **176**
 refined, **113**
 visible spectrum, **105**
- weak equivalence (of p -adic exponents), **224**
 weak topology, **358**
 weakly difference-closed difference field, **247**
 weakly prepared eigenvalues, **123**, **221**
 wedge power, *see also* exterior power
 Weierstrass preparation, **140**
 Weil conjectures, **4**
 well-ordered set, **31**, **282**
 width (of a polynomial), **35**
 wild inertia subgroup, **49**
 wildly ramified extension, **49**
 Witt vectors, **258**, **293**, **358**
- Yoneda extension group, **82**, **245**
- zero-centered Frobenius lift, **266**
 zeta function, **4**, **349**, **354**