

Cambridge University Press
0521772907 - Simulating Hamiltonian Dynamics
Benedict Leimkuhler and Sebastian Reich
Frontmatter
[More information](#)

**CAMBRIDGE MONOGRAPHS ON
APPLIED AND COMPUTATIONAL
MATHEMATICS**

Series Editors

P. G. CIARLET, A. ISERLES, R. V. KOHN, M. H. WRIGHT

14 Simulating Hamiltonian Dynamics

The *Cambridge Monographs on Applied and Computational Mathematics* reflects the crucial role of mathematical and computational techniques in contemporary science. The series publishes expositions on all aspects of applicable and numerical mathematics, with an emphasis on new developments in this fast-moving area of research.

State-of-the-art methods and algorithms as well as modern mathematical descriptions of physical and mechanical ideas are presented in a manner suited to graduate research students and professionals alike. Sound pedagogical presentation is a prerequisite. It is intended that books in the series will serve to inform a new generation of researchers.

Also in this series:

1. A Practical Guide to Pseudospectral Methods, *Bengt Fornberg*
2. Dynamical Systems and Numerical Analysis, *A. M. Stuart and A. R. Humphries*
3. Level Set Methods and Fast Marching Methods, *J. A. Sethian*
4. The Numerical Solution of Integral Equations of the Second Kind, *Kendall E. Atkinson*
5. Orthogonal Rational Functions, *Adhemar Bultheel, Pablo González-Vera, Erik Hendiksen, and Olav Njåstad*
6. The Theory of Composites, *Graeme W. Milton*
7. Geometry and Topology for Mesh Generation *Herbert Edelsbrunner*
8. Schwarz–Christoffel Mapping *Tofin A. Driscoll and Lloyd N. Trefethen*
9. High-Order Methods for Incompressible Fluid Flow, *M. O. Deville, P. F. Fischer and E. H. Mund*
10. Practical Extrapolation Methods, *Avram Sidi*
11. Generalized Riemann Problems in Computational Fluid Dynamics, *Matania Ben-Artzi and Joseph Falcovitz*
12. Radial Basis Functions, *Martin D. Buhmann*
13. Iterative Krylov Methods for Large Linear Systems, *Henk A. van der Vorst*.
15. Collocation Methods for Volterra Integral and Related Functional Equations, *Hermann Brunner*.

Cambridge University Press
0521772907 - Simulating Hamiltonian Dynamics
Benedict Leimkuhler and Sebastian Reich
Frontmatter
[More information](#)

Simulating Hamiltonian Dynamics

BENEDICT LEIMKUHLER
University of Leicester

SEBASTIAN REICH
Imperial College, London



Cambridge University Press
 0521772907 - Simulating Hamiltonian Dynamics
 Benedict Leimkuhler and Sebastian Reich
 Frontmatter
[More information](#)

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
 The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS
 The Edinburgh Building, Cambridge CB2 2RU, UK
 40 West 20th Street, New York, NY 10011–4211, USA
 477 Williamstown Road, Port Melbourne, VIC 3207, Australia
 Ruiz de Alarcón 13, 28014 Madrid, Spain
 Dock House, The Waterfront, Cape Town 8001, South Africa

<http://www.cambridge.org>

© Cambridge University Press 2004

This book is in copyright. Subject to statutory exception
 and to the provisions of relevant collective licensing agreements,
 no reproduction of any part may take place without
 the written permission of Cambridge University Press.

First published 2004

Printed in the United Kingdom at the University Press, Cambridge

Typeface CM bright 11/13.5 pt. *System* L^AT_EX 2 ϵ [TB]

A catalog record for this book is available from the British Library

Library of Congress Cataloging in Publication data

Leimkuhler, B.

Simulating Hamiltonian dynamics / Benedict Leimkuhler, Sebastian Reich.

p. cm. – (Cambridge monographs on applied and computational mathematics ; 14)

Includes bibliographical references and index.

ISBN 0 521 77290 7

1. Hamiltonian systems. I. Reich, Sebastian. II. Title. III. Series.

QA614.83.L45 2004

515'.39 – dc22 2004045685

ISBN 0 521 77290 7 hardback

The publisher has used its best endeavors to ensure that the URLs for external websites referred to in this book are correct and active at the time of going to press. However, the publisher has no responsibility for the websites and can make no guarantee that a site will remain live or that the content is or will remain appropriate.

Contents

Preface		ix
Acknowledgements		xvi
1 Introduction		1
1.1 <i>N</i> -body problems		2
1.2 Problems and applications		3
1.3 Constrained dynamics		6
1.4 Exercises		8
2 Numerical methods		11
2.1 One-step methods		13
2.2 Numerical example: the Lennard–Jones oscillator		18
2.3 Higher-order methods		20
2.4 Runge–Kutta methods		22
2.5 Partitioned Runge–Kutta methods		25
2.6 Stability and eigenvalues		27
2.7 Exercises		32
3 Hamiltonian mechanics		36
3.1 Canonical and noncanonical Hamiltonian systems		36
3.2 Examples of Hamiltonian systems		39
3.3 First integrals		44
3.4 The flow map and variational equations		48
3.5 Symplectic maps and Hamiltonian flow maps		52
3.6 Differential forms and the wedge product		61
3.7 Exercises		66
4 Geometric integrators		70
4.1 Symplectic maps and methods		74

4.2	Construction of symplectic methods by Hamiltonian splitting . . .	76
4.3	Time-reversal symmetry and reversible discretizations	81
4.4	First integrals	87
4.5	Case studies	91
4.6	Exercises	99
5	The modified equations	105
5.1	Forward v. backward error analysis	107
5.2	The modified equations	117
5.3	Geometric integration and modified equations	129
5.4	Modified equations for composition methods	133
5.5	Exercises	139
6	Higher-order methods	142
6.1	Construction of higher-order methods	143
6.2	Composition methods	144
6.3	Runge–Kutta methods	149
6.4	Generating functions	159
6.5	Numerical experiments	161
6.6	Exercises	165
7	Constrained mechanical systems	169
7.1	N -body systems with holonomic constraints	170
7.2	Numerical methods for constraints	173
7.3	Transition to Hamiltonian mechanics	184
7.4	The symplectic structure with constraints	186
7.5	Direct symplectic discretization	188
7.6	Alternative approaches to constrained integration	191
7.7	Exercises	195
8	Rigid body dynamics	199
8.1	Rigid bodies as constrained systems	201
8.2	Angular momentum and the inertia tensor	210
8.3	The Euler equations of rigid body motion	212
8.4	Order 4 and order 6 variants of RATTLE for the free rigid body	223
8.5	Freely moving rigid bodies	224
8.6	Other formulations for rigid body motion	228
8.7	Exercises	230
9	Adaptive geometric integrators	234
9.1	Sundman and Poincaré transformations	235

<i>Contents</i>	vii
9.2 Reversible variable stepsize integration	238
9.3 Sundman transformations	246
9.4 Backward error analysis	249
9.5 Generalized reversible adaptive methods	251
9.6 Poincaré transformations	253
9.7 Exercises	255
10 Highly oscillatory problems	257
10.1 Large timestep methods	258
10.2 Averaging and reduced equations	269
10.3 The reversible averaging (RA) method	271
10.4 The mollified impulse (MOLLY) method	276
10.5 Multiple frequency systems	279
10.6 Exercises	280
11 Molecular dynamics	287
11.1 From liquids to biopolymers	290
11.2 Statistical mechanics from MD trajectories	296
11.3 Dynamical formulations for the NVT, NPT and other ensembles	299
11.4 A symplectic approach: the Nosé–Poincaré method	305
11.5 Exercises	313
12 Hamiltonian PDEs	316
12.1 Examples of Hamiltonian PDEs	316
12.2 Symplectic discretizations	325
12.3 Multi-symplectic PDEs	335
12.4 Exercises	352
<i>References</i>	357
<i>Index</i>	374

Preface

About geometric integration

This book is about simulating dynamical systems, especially conservative systems such as arise in celestial mechanics and molecular models. We think of the *integrator* as the beating heart of any dynamical simulation, the scheme which replaces a differential equation in continuous time by a difference equation defining approximate snapshots of the solution at discrete timesteps. As computers grow in power, approximate solutions are computed over ever-longer time intervals, and the integrator may be iterated many millions or even billions of times; in such cases, the qualitative properties of the integrator itself can become critical to the success of a simulation. *Geometric integrators* are methods that exactly (i.e. up to rounding errors) conserve qualitative properties associated to the solutions of the dynamical system under study.

The increase in the use of simulation in applications has mirrored rising interest in the theory of dynamical systems. Many of the recent developments in mathematics have followed from the appreciation of the fundamentally chaotic nature of physical systems, a consequence of nonlinearities present in even the simplest useful models. In a chaotic system the individual trajectories are by definition inherently unpredictable in the exact sense: solutions *depend sensitively on the initial data*. In some ways, this observation has limited the scope and usefulness of results obtainable from mathematical theory. Most of the common techniques rely on local approximation and perturbation expansions, methods best suited for understanding problems which are “almost linear,” while the new mathematics that would be needed to answer even the most basic questions regarding chaotic systems is still in its infancy. In the absence of a useful general theoretical method for analyzing complex nonlinear phenomena, simulation is increasingly pushed to the fore. It provides one of the few broadly applicable and practical means of shedding light on the behavior of complex nonlinear systems, and is now a standard tool in everything from materials modeling to bioengineering, from atomic theory to cosmology.

As models grow in complexity and dimension, and the demands placed on simulation have risen, the need for more sophisticated numerical methods and analytic techniques also grows. Longer time interval simulations require more stable methods. Larger problems call for more efficient schemes, tailored to a particular application or family of applications. And more intricate modeling of delicate behaviors or properties requires corresponding improvements in the resolution of those properties during simulation.

In writing this book for a broad audience of scientists, we have attempted to limit the introduction of technical detail, but in some places this cannot be avoided. The calculations are generally included for the benefit of students. We hope that appreciation of the general principles will not be lost in following the details of arguments. In the words of John Von Neumann, “One expects a mathematical theorem or a mathematical theory not only to describe and classify in a simple and elegant way numerous and a priori disparate special cases. One also expects elegance in its architectural, structural make-up. . . . If the deductions are lengthy or complicated there should be some simple general principle involved, which explains the complications and details, reduces the apparent arbitrariness to a few simple guiding motivations.” If there is one such guiding principle underlying our work it is this: classical mechanics – on which all physical models are based – also provides the proper foundation for numerical simulation of those systems. We will attempt to show in this book that practical, efficient methods for simulating conservative systems can be realized by making judicious use of the methods of classical mechanics.

An emphasis on methods

In this book we address ourselves primarily to the following pair of questions:

Which properties should be fundamental to an integration method for a (conservative) model?

How can we design and implement schemes that respect physical principles regardless of timestep or traditional accuracy considerations?

Although our interest is always ultimately in the methods themselves and in quantifying the relative differences among them, we will find that in attempting to answer the above questions, we are drawn far afield from the usual domain of the numerical analyst. The first question will lead us into the field of mechanics so that we may appreciate something of the nature of those structures and symmetries that underlie physical models and contribute to their long-term evolution. The second question will take us outside even the areas that have traditionally been investigated by mathematicians, since the special forms of force

functions, presence of constraints, relationships among the variables, or efficiency considerations dictate to a large extent the features of appropriate (i.e. practical) methods used in applications.

It is important to emphasize that our treatment is not comprehensive; we have made a selection from the literature which comprises – in our view – the most important material from the standpoint of practical application.

Beginning with the idea of *splitting* we will show how many simple but effective integrators can be generated by using a few building blocks. The same techniques can be used to derive more sophisticated schemes. For example, explicit higher-order methods have a very natural derivation in the case of canonical mechanical systems developed in terms of the “kinetic+potential” form of the energy.

We survey recent work on methods for constrained systems and consider various approaches to the simulation of rigid body systems, methods which offer an efficient and – in many cases – demonstrably superior geometric alternative to more widespread schemes. Variable stepsize geometric integrators will be introduced based on a rescaling of the time variable. Methods for mixed systems possessing both rapidly and slowly varying degrees of freedom – or weak and strong forces – also call for the construction of specialized schemes. In all cases, our aim will be to present the ideas in as general a form as is prudent, highlighting instances where a given technique might be of use in other applications. Molecular dynamics applications are an important source of challenging problems for geometric integration, so we devote some time to their particular characteristics. Conservative partial differential equations introduce many new issues for the development of geometric integrators, a topic we touch on in the final chapter of the book.

How to use this book

This book is intended, first, as a text for a course in computational mechanics or as a tool for self-instruction, and, second, as a basic reference for researchers and educators – regardless of discipline – interested in using and developing geometric integrators. The book should serve as a bridge from traditional training in the sciences to the recent research literature in the field of geometric integration. By emphasizing mathematical and computational issues and illustrating the various concepts and techniques with carefully developed model problems, it is hoped that the book can appeal to a wide audience, including mathematicians unfamiliar with modeling issues, and physicists, chemists, and engineers wishing to gain a better understanding of the mathematical underpinnings of existing methods or in developing effective methods for new applications.

The book assumes only that the reader has had undergraduate coursework in linear algebra and differential equations. At several points we introduce, but do

not thoroughly develop, topics from dynamical systems. A good introductory text in dynamical systems is the book of VERHULST [197]. We develop – in Chapters 2 and 3 – most of the necessary preliminaries of numerical analysis and classical mechanics from the ground up, however the reader should be aware that the treatment provided here of the required background material is necessarily brief; only those elements that are essential to our later study of geometric integrators are given. For an introduction to numerical analysis, the reader is referred to the classic books of ATKINSON [11], BURDEN AND FAIRES [37], and DAHLQUIST AND BJÖRK [48]. The book of GEAR [70] can provide a useful introduction to the numerical solution of differential equations. The books of HAIRER, NØRSETT AND WANNER [82] and HAIRER AND WANNER [84] can serve as references for obtaining a more complete picture of the mathematical issues associated with construction of methods and error analysis for ordinary differential equations. ISERLES [91] has written an integrated text that introduces numerical methods for both ordinary and partial differential equations.

First published at the end of the 19th century, ROUTH's *Dynamics of a System of Rigid Bodies (Elementary Part)*¹ remains a marvelous introduction to classical mechanics and provides a wealth of examples and exercises for the student (many of which could now be revisited with the aid of the modern computational techniques developed in this book). For a more systematic treatment to Hamiltonian classical mechanics the reader is referred to the following texts: LANDAU AND LIFSHITZ [105], MARION [121], GOLDSTEIN [73], and ARNOLD [7], all of which are well-worn occupants of our bookshelves. These books are quite varied in their use of notation and even in the way in which they motivate and explain identical material, but we have found all of them to be helpful on various occasions. If only one book is to be consulted, the elegant book of LANCZOS [104] is remarkable both in terms of its readability and its breadth, owing partly to the absence of detailed proofs. A modern rigorous treatment of classical mechanics may be found in MARS DEN AND RATIU [124], a book which also contains a number of useful examples and notes on history and applications.

The book by SANZ-SERNA AND CALVO [172] was the first to cover symplectic integration methods and applications to classical mechanics and is still an excellent introduction to the subject. The more recent book by HAIRER, LUBICH, AND WANNER [80] covers a wide range of topics from geometric integration and should be very useful as an additional reference.

In a graduate course in applied mathematics or computational physics, it is probable that much of the material of Chapters 1–3 could be skipped or skimmed, depending on the backgrounds of the students and the interests of the teacher. Some caution should be exercised here. In particular, it is essential

¹Reprinted in 1960 as a Dover Edition.

that the student understand the concepts of convergence and order of convergence for a numerical method, the definition of the flow map, first integrals, and at least the condition for a symplectic map in terms of the Jacobian of the flow map.

Molecular dynamics provides a rich source of problems for geometric integration, and we often draw on examples from this field for motivation and for evaluation of concepts and methods. Here again, it is likely that the reader may, on occasion, wish for a more detailed description of the problems or of typical approaches used by chemists and physicists. One reference stands out in this area for clarity of presentation and breadth: ALLEN AND TILDESLEY [4]. More recent books of FRANKEL AND SMIT [66] and SCHLICK [174] help to fill in the picture.

Exercises included at the end of each chapter are intended to be demanding but not overwhelming; some of the multi-part problems could be assigned as projects, especially those involving the use of computers.

Computer software

This book primarily emphasizes the mathematical properties of algorithms for solving differential equations. In later chapters, we will often see the methods introduced and analyzed as abstract maps of phase space. This approach, while essential to understanding and generalizing the methods, has the tendency to obscure both the intuitive basis for the theory and the ultimate importance of the subject. We would like emphasize that *the student must implement and test numerical methods in order to gain a full understanding of the subject*.

While any programming language and graphics package could, in principle, be employed, the need for flexibility in the coding and testing of methods and the need to be able to work easily with scientific functions of vectors and matrices makes a specialized, interpreted language system for mathematics the best environment for problem solving.

At the time of this writing, there are several widely distributed commercial software packages that support the simplified design and testing of algorithms, including the commercial packages MATLAB, MAPLE, and MATHEMATICA. Of these, the authors prefer the user interface and programming structure of MATLAB, but this is largely a matter of taste and any of the three mentioned systems would be suitable. These packages are all available to students at heavily discounted prices and run on a variety of computer platforms.

For a student on a severe budget, there are several widely available *free* alternatives to the commercial packages. These options include, notably, OCTAVE, which is distributed under the Free Software Foundation's GNU Public License. Our experience with this software is that it is adequate for most study purposes,

although the commercial alternatives are generally superior with regard to ease-of-use, documentation, and reliability.

We will occasionally describe algorithms in this text, but we will attempt to avoid system-specific details, so the student is expected to supplement the mathematical study with study of the the user's guide for the software system they are using, in particular the appropriate sections on programming.

Notation

Let us summarize some basic notation used throughout the book. Dependent variables, such as positions \mathbf{q} and velocities \mathbf{v} are elements of a Euclidean space \mathbb{R}^d , where $d \geq 1$ is the appropriate dimension. More specifically, we will always identify dependent variables with *column vectors*. When two column vectors $\mathbf{u} \in \mathbb{R}^k$, $\mathbf{v} \in \mathbb{R}^l$ are given we may write $(\mathbf{u}, \mathbf{v})^T$ for the column vector in \mathbb{R}^{k+l} obtained by concatenating the two vectors. The transpose is there to remind us that the result is again a column vector. We will often need to refer to a set indexed by a parameter for which we write $\{a_t\}_{t \in P}$, where P is the index set. As a short-hand, we will write $\{a_t\}$ if the index set is clear from the context.

The set of $k \times k$ matrices with real coefficients is $\mathbb{R}^{k \times k}$ and capital bold-face letter are used to denote matrices, e.g., $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{k \times k}$. The k -dimensional identity is \mathbf{I}_k or \mathbf{I} as a short-hand if the dimension is clear from the context.

A vector-valued function $\mathbf{F} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ will be assumed to map column vectors of dimension n to column vectors of dimension m . The vector of partial derivatives of a scalar-valued function $f(\mathbf{q})$ is identified with a *row vector* in \mathbb{R}^d and is denoted by $f_{\mathbf{q}}(\mathbf{q})$ or, equivalently, by $\partial f / \partial \mathbf{q}(\mathbf{q})$. Hence the *Jacobian matrix* of a vector-valued function $\mathbf{F}(\mathbf{q})$ is identified with the $m \times n$ matrix $\mathbf{F}_{\mathbf{q}}(\mathbf{q})$.

The *scalar product*, *inner product*, or *dot product* of two column vectors \mathbf{a} and \mathbf{b} in \mathbb{R}^d is denoted by $\langle \mathbf{a}, \mathbf{b} \rangle$ or $\mathbf{a} \cdot \mathbf{b}$, or, simply, $\mathbf{a}^T \mathbf{b}$, where \mathbf{a}^T is the transpose of \mathbf{a} . The *cross product* of \mathbf{a} and \mathbf{b} is denoted $\mathbf{a} \times \mathbf{b}$. We will frequently use

$$\hat{\mathbf{a}}\mathbf{b} = \mathbf{a} \times \mathbf{b},$$

where $\hat{\mathbf{a}} \in \mathbb{R}^{3 \times 3}$ is a skew-symmetric matrix related to the vector $\mathbf{a} = (a_1, a_2, a_3)^T$ by

$$\hat{\mathbf{a}} = \begin{bmatrix} 0 & -a_3 & a_2 \\ a_3 & 0 & -a_1 \\ -a_2 & a_1 & 0 \end{bmatrix}.$$

The *norm* of a vector \mathbf{a} is defined by

$$\|\mathbf{a}\| = \langle \mathbf{a}, \mathbf{a} \rangle^{1/2}.$$

A real-valued function f defined on an interval I of the real line is said to be square-integrable if $\int_I f(x)^2 dx$ is bounded. In that case we say that f lies in the function space L_2 on the interval I . We say that the L_2 -norm of f is

$$\|f\|_2 = \left(\int_I f(x)^2 dx \right)^{\frac{1}{2}},$$

and we define the L_2 -inner product of two square integrable functions f and g by

$$(f, g) = \int_I f(x)g(x)dx$$

The *gradient* $\nabla_{\mathbf{q}}V(\mathbf{q})$ of a scalar-valued differentiable function $V(\mathbf{q})$ is defined by

$$\langle \nabla_{\mathbf{q}}V(\mathbf{q}), \mathbf{u} \rangle = \lim_{\varepsilon \rightarrow 0} \frac{V(\mathbf{q} + \varepsilon \mathbf{u}) - V(\mathbf{q})}{\varepsilon},$$

where the equality is to hold for all vectors \mathbf{u} of the same dimension as \mathbf{q} . This definition leads to the relation

$$\nabla_{\mathbf{q}}V(\mathbf{q}) = V_{\mathbf{q}}(\mathbf{q})^T,$$

and, hence, the gradient $\nabla_{\mathbf{q}}V(\mathbf{q})$ is a column vector.

The time derivative of a function $\mathbf{q}(t)$ will normally be denoted by $d\mathbf{q}/dt(t)$, but whenever it is more convenient we may instead use the short-hand $\dot{\mathbf{q}}(t)$. If clear from the context, we will also frequently drop the argument in functions and write, for example, $\dot{\mathbf{q}}$ instead of $\dot{\mathbf{q}}(t)$, or \mathbf{F} instead of $\mathbf{F}(\mathbf{q})$. The same conventions apply to higher-order derivatives.

If a variable u depends on several independent variables such as time t and space x , then the partial derivatives are often denoted by u_t and u_x , respectively.

Given two maps $\boldsymbol{\psi}_1 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ and $\boldsymbol{\psi}_2 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with compatible range and domain, we define their composition $\boldsymbol{\psi}_2 \circ \boldsymbol{\psi}_1$ by

$$[\boldsymbol{\psi}_2 \circ \boldsymbol{\psi}_1](z) = \boldsymbol{\psi}_2(\boldsymbol{\psi}_1(z)),$$

for all $z \in \mathbb{R}^n$. The inverse of a one-to-one map $\boldsymbol{\psi}$ is denoted by $\boldsymbol{\psi}^{-1}$. Hence

$$\boldsymbol{\psi}^{-1} \circ \boldsymbol{\psi} = \mathbf{id},$$

where $\mathbf{id}(z) = z$ the identity map.

Finally the Landau-order notation $\mathcal{O}(\Delta t)$ is used to denote a quantity that goes to zero as rapidly as Δt goes to zero. More generally, we will write $g(\Delta t) = \mathcal{O}(\Delta t^p)$ if $g(\Delta t)/\Delta t^p$ is bounded as $\Delta t \rightarrow 0$ but $g(\Delta t)/\Delta t^q$ is unbounded if $q > p$.

Acknowledgements

We have to admit that when we began this project, we had no appreciation for its ultimate complexity. The only thing that has kept the project going at some critical points was the broad support and encouragement we received from our families, our colleagues, and our students.

A key aspect of the book is the emphasis on applications. Brian Laird proposed the Lennard–Jones model discussed in Chapter 4 and made several other helpful comments. Chris Sweet contributed most of the numerical experiments in Chapter 6, including the long-term simulations of the solar system, and Eric Barth provided the alanine dipeptide simulation in Chapter 11. Jason Frank helped with many of the experiments in Chapter 12. Jason’s tireless proofreading of the entire manuscript must also be remarked!

We hope that the book will be useful for teaching upper-division undergraduate and graduate courses. In 2002, Ph.D. students from various fields and various parts of Europe – Martin Dahlberg (chemistry), Thomas Sauvaget (mathematics/physics), Marko Vranicar (physics), and Fredrick Hedman (mathematics) – attended an informal summer school in Leicester taught from an early draft of the book. Part of the book formed the basis for a series of survey lectures given by one of us in Bari, Italy in 2002. Mitch Luskin, Claudia Wulff, and Steve Bond have subjected their classes to a preliminary draft of the book for testing in this setting, and provided useful comments.

We would like also to thank Teijo Arponen, Jeff Cash, Colin Cotter, Ernst Hairer, Zhidong Jia, Per-Christian Moan, Brian Moore, and Antonella Zanna for reading various parts of the manuscript and providing feedback. We have also relied on the extraordinary patience of Cambridge University Press, particularly our editors David Tranah and Joseph Bottrill.