

Index

- academic problems 245–6
 - conduct disorder and 394
 - sex differences 105
 - treatment effects 446, 447, 453
- acetylcholine 179
- activation 179–80
 - effects of medication 187–8
 - evoked potentials 193–4, 195
 - motor, measurement 191
 - nonoptimal level 182–3, 188, 189
- activity level 265–6
 - in adults 478
 - increased, *see* overactivity
 - stability 266, 271
 - see also* motor activity
- ADD, *see* attention deficit disorder
- Adderall, in adult ADHD 481
- additive genetic influences 116, 117, 296, 299–300
- additives, food 3, 21
- ADHD, *see* attention deficit hyperactivity disorder
- ADHD rating scale 476
- adolescents
 - diagnostic aspects 145
 - electrophysiological studies 347
 - maternal relationships 396
 - persistence of hyperactivity 272–4
 - prevalence of ADHD 37–46, 47–8, 473
 - sex differences in outcome 108, 118
 - see also* adults; children
- adoptees, Romanian 419–25, 428–31
- adoption studies 294–6, 376–7, 418
- adult hyperactivity 36, 468–85
 - assessment 145, 475–80
 - classification 468–70
 - comorbidity 474–5, 482
 - epidemiology 470
 - further research 484–5
 - longitudinal cohorts 471–4
 - prevalence and distribution 50–1, 58, 473
 - treatment and management 480–4
- adults
 - distress threshold hypothesis 83–4, 85–7
 - outcome of hyperactivity 272–6
 - predictors 274–6
 - sex differences 108, 473–4
 - relationships with 246
 - strange, interactions with 197–8, 373
- affect
 - infant, maternal responsiveness to 251–2
 - regulation 247
- affective disorders
 - in adult ADHD 474
 - comorbidity rates 50, 104
 - cross-cultural/ethnic comparisons 88
 - diagnostic aspects 145
 - see also* anxiety disorder; depression
- African American children 74–5, 87, 88–9
- age
 - adult ADHD and 51, 473
 - genetic considerations 326
 - prevalence of ADHD and 47–8, 473
 - sex differences and 101–2
- aggression
 - developmental course 268
 - early antecedents 254, 266, 269–70
 - hyperactivity as risk factor for 394, 396–7
 - sex differences 111
- alarm
 - effect on brain function 388–9
 - reaction 386–7
 - state, stressed children 389–90
- alcohol abuse 273, 471
 - cross-cultural study 79
 - molecular genetics 320
- aldehyde dehydrogenase, variant allele (ALDH2) 79
- allelic heterogeneity 324
- allergy 177
 - food 3, 21
- α -antagonists 483, 484
- altropane 344
- American Academy of Child and Adolescent Psychiatry 4
- amfetamines
 - in adult ADHD 481
 - history of use 12–13, 14
 - mechanisms of action 352, 354
 - see also* dexamfetamine
- amygdala 387, 388
- animal studies, stress 384, 385, 386
- anoxia, perinatal 15
- antisocial behaviour 108
 - in adolescence/adulthood 272–4
 - hyperactivity as risk factor for 393–4
 - see also* conduct disorder; oppositional defiant disorder

492 Index

- antisocial personality disorder (ASPD) 292–3, 471–2, 474–5
- anxiety disorder 397–8
 - in adult ADHD 482
 - comorbidity rates 50, 104
 - cross-cultural/ethnic comparisons 88
 - diagnosis of hyperactivity and 139–40, 144–5, 149
 - role in ADHD 402
 - treatment efficacy and 448–9, 454, 457–8
- arithmetical ability 477
- arousal 19, 178–9, 247
 - anxiety and 402
 - effect on brain function 388–9
 - evoked potentials 193, 194–5
 - modulation in infants 249, 250, 262
 - regulation, *see* self-regulation; state regulation
 - unitary state concept 178–91
- Asians
 - diagnosis of hyperactivity 76
 - drug metabolism 89
 - factor analytic studies 67–9
 - genetic factors 80
 - symptom presentation 81–3
 - UK study 76–9
- assessment
 - adult ADHD 145, 475–80
 - first systematic tools 19
 - in MTA Cooperative Group study 442–3
 - see also* classification; diagnosis; laboratory measures; rating scales
- association studies 302–5
- assortative mating 297–8
- attachment relationships 252–8
 - development of self-regulation and 252–8, 380–1
 - disorganized subtype 257–8, 381
 - disrupted 252–8, 367, 379–81
 - institutional care and 418, 422–3, 427, 428, 431
 - parental psychopathology and 381–3
- attention 162–200
 - abilities, basic 167–73
 - biological basis 350
 - concept of state 178–91
 - deficits, *see* inattention
 - developmental aspects 253, 266–7
 - focused 169–70
 - mental retardation and 189–91
 - motor-related dysfunction and 173–8
 - psychophysiological measures 191–6
 - sustained
 - in adult ADHD 477, 478
 - development 267, 271
 - state regulation and 181–5
- attention deficit disorder (ADD) 130, 469
 - cognitive deficits 51–2
 - cross-cultural comparisons 75–6
 - diagnostic validity 147
 - with hyperactivity 130, 469
 - psychosocial factors 53
 - undifferentiated (UADD) 31
- attention deficit hyperactivity disorder (ADHD)
 - 243–4, 291, 469
 - in adulthood, *see* adult hyperactivity
 - behavioural genetics 292–302
 - categories vs. dimensions 33–4
 - combined with DAMP 146
 - cross-cultural comparisons 75–6
 - diagnosis, *see* diagnosis
 - diagnostic criteria (DSM-IV) 132–3
 - DSM-III-R 31–2, 131, 469
 - DSM-IV, *see* *Diagnostic and Statistical Manual* (DSM), fourth edition
 - epidemiology, *see* epidemiology
 - not otherwise specified 469
 - outcome, *see* outcome
 - in partial remission 469
 - sex differences, *see* sex differences
 - subthreshold cases 53–4
 - subtypes, *see* subtypes of ADHD
 - symptoms, *see* symptoms
 - treatment, *see* treatment
 - validity 145–6
- attunement, caregiver–infant 251–2
- auditory processing, central 226–7
- Australian Twin ADHD Project (ATAP) 224–5, 229
- autonomic nervous system 402
- balance beam task 171
- basal ganglia 195, 338–9
 - functional imaging 341–2
 - neuropharmacology 353
 - role in memory 387
- Behar Problem Behaviour Questionnaire 269
- behavioural genetics 292–302
- behavioural treatment
 - cross-cultural/ethnic comparisons 88
 - MTA Cooperative Group study (Beh) 440–1, 446–7, 448, 453–4, 457, 458–61
- behaviourist perspective 21
- behaviour rating scales, *see* rating scales
- benzedrine 12–13
- biological basis 336–55
 - electrophysiological studies 346–9
 - functional brain imaging studies 340–6
 - neuroanatomical studies 336–40
 - neurochemical studies 349–51
 - neuropharmacological studies 351–5
 - sex differences 117–18
 - speculative model 349, 355
- blastophoria 8
- blood oxygenation level-dependent (BOLD)
 - functional MRI 341, 343
- Bloomington Longitudinal Study 255, 262–3, 394
- Bradley, Charles 12–13, 130
- brain
 - anatomical studies 336–40
 - damage
 - history 2–3, 6–9, 11–12, 14–18, 129–30

493 Index

- see also* minimal brain damage; minimal brain dysfunction
 - experience-dependent development 385–9
 - functional imaging studies 340–6
 - impact of stressful experiences 384–90
 - plasticity 383
 - record of experiences 387–8
- Briquet's syndrome 292–3
- bromides 5
- Brown Attention Deficit Disorder Scale 480
- Cambridge Neuropsychological Test Automated Battery (CANTAB) 219
- candidate genes 306–23, 384
 - cross-cultural/ethnic studies 79–81
 - mode of inheritance 324
 - multiple markers within 323
 - nondopaminergic 321–3
- capacity, information-processing 170–3
- caregiver–infant relationships 250–61
 - attachment, *see* attachment relationships
 - in development of self-regulation 248–50
 - lack of personalized 429–30
 - synchrony/attunement aspects 251–2
 - see also* mother–child relationships; parent–child relationships
- case definition
 - in epidemiological studies 31–5
 - sex differences and 99–100
- catecholamines 349–51, 355
 - functional imaging studies 344
 - see also* dopamine; noradrenaline
- catechol O-methyltransferase (COMT) 307, 314–18
- Caucasians
 - comorbid disorders 87–8
 - drug treatment 88–9
 - factor analytic studies 67
 - genetic factors 80–1
 - recognition of problem/help-seeking 87
 - symptom presentation 81–2, 84
- caudate nucleus
 - anatomical studies 338, 339–40
 - functional imaging studies 341
 - neurochemistry 350
 - neuropharmacology 353
- CD, *see* conduct disorder
- cerebellum/cerebellar vermis 349, 355
 - anatomical studies 339, 340
 - functional imaging studies 343–4
- cerebral blood flow, regional 341
- cerebral glucose metabolism 108, 196, 341
- checklists 35
- child abuse 386–7, 401
- Child and Adolescent Psychiatric Assessment (CAPA) 36, 46, 54
- Child Behavior Checklist (CBCL) 36, 83, 101, 138, 139
- child-rearing practices, *see* parenting styles
- children
 - preschool, *see* preschool children
 - prevalence of hyperactivity 37–46, 58
 - see also* adolescents; infants; toddlers
- Children's Embedded Figure Test 174
- Children's Global Assessment Scale (CGAS) 54
- Chinese 90
 - behaviour rating scales 71–4
 - comorbid disorders 87–8
 - conduct disorder vs. hyperactivity 71
 - diagnosis of hyperactivity 76
 - external correlates 70
 - factor analytic studies 67, 69
 - genetic factors 79, 80–1
 - recognition of problem/help-seeking 85–7
 - symptom presentation 81–2, 84
 - vs. British children 76–9
- cingulate cortex 342–3, 350, 353
- classification 126–54, 214–15, 291
 - adult ADHD 468–70
 - for epidemiological studies 31–5
 - function in psychiatry 127–8
 - future directions 149–52
 - historical development 128–31
 - stability 55–7
 - structured interviews vs. checklists 35
 - subthreshold cases and boundaries 53–4
 - see also* assessment; diagnosis; *Diagnostic and Statistical Manual; International Classification of Disease*
- Clinical Evaluation of Language Fundamentals – Revised (CELF-R) 219
- Clinical Interview Schedule, revised (CIS-R) 476
- clomipramine 484
- clonidine 187–8, 484
- Clouston, T.S. 4–5
- clumsiness 163
- cocaine 353, 386
- cognitive function 51–3, 216–34
 - childhood stress and 388–9
 - comorbidity and 221–32
 - early caregiver relationships and 255–6, 257
 - future research 232–4
 - heritable subtypes and 151–2
 - objective measures 165–6, 167–78
 - output vs. input processes 173–4
 - role in ADHD 400–1
 - sex differences 107
 - task-inappropriate responding 174–8
 - task inefficiency, *see* task inefficiency
 - see also* executive function; inattention; intelligence; language disorders; learning disorders; reading disability/disorder
- Cohen, L.H. 11–12
- combined type of ADHD (ADHD/CT) 32, 144, 215
 - in adults 50–1
 - cognitive functioning 218–21, 224–5
 - epidemiological surveys 49
 - molecular genetics 310, 311
 - as refined phenotype 336

Cambridge University Press

0521789613 - Hyperactivity and Attention Disorders of Childhood, Second Edition

Edited by Seija Sandberg

Index

[More information](#)

494 Index

- combined type of ADHD (ADHD;CT) (*cont.*)
 - validity 147–8
- communication disorders 225–8, 400
 - see also* language disorders; speech problems
- community care (CC), in MTA Cooperative Group study 441–2, 446–7, 448, 452, 456–7
- comorbidity
 - in adult ADHD 474–5, 482
 - cognitive functioning and 221–32
 - cross-cultural/ethnic studies 87–8
 - diagnostic issues 139–40, 144–5, 148–9
 - epidemiological surveys 50, 58
 - future research 233–4
 - genetics 228–32, 390
 - long-term outcome 273–4
 - in MTA Cooperative Group study 439
 - psychosocial aspects 390–8
 - sex differences 104–5
 - treatment efficacy and 448–9, 457–8
 - validity of laboratory measures and 164
- complex traits, genetics of 305
- compliance
 - with caregiver requests 249, 253, 258–60, 381
 - with treatment 443–4, 450–1, 459–61
- computed tomography (CT) 339
- conduct disorder (CD) 245, 390
 - adult ADHD and 473, 474–5
 - cognitive function and 51–3, 224
 - comorbidity rates 50, 104
 - cross-cultural/ethnic studies 67, 69, 70–1, 85, 87–8
 - diagnosis of hyperactivity and 139–40, 145
 - early temperament and 266
 - genetics 293, 325
 - hyperactivity as risk factor for 393–7
 - hyperkinetic 33, 133, 145, 146–9
 - institutional care and 420–1, 427–8
 - mental retardation and 189–90
 - persistence in adolescence/adulthood 273–4
 - psychosocial factors 390–3
 - sex differences 101, 102–3, 104–5, 106, 111
 - sustained attention 183
 - treatment efficacy and 449, 454–5, 458
- conflict, parent–child
 - developmental course 268
 - temperament and 260, 264, 266
 - toddler control interactions and 258–61
- Conners' scales 138
 - in adult ADHD 479
 - epidemiological studies 36, 37
 - history 19
- Conners' Teacher Rating Scale (CTRS)
 - abbreviated version (ATRS) 73
 - cross-cultural/ethnic studies 65, 66, 67, 68, 71–4, 85–8
 - sex differences 103–4
- constitutional variability model 113–14
- Continuous Performance Task/Test (CPT) 52, 174, 181–5, 189
 - activity level during 184–5
 - in adult ADHD 477, 478
 - effects of comorbidity 225
 - evoked potentials 348
 - with fast event rate of stimuli 181–2
 - heart rate variability and 192–3
 - mental retardation and 190
 - with slow event rate of stimuli 182–4
 - state fluctuations and 198–9
- continuum of reproductive casualty 16
- control
 - external 182–3
 - parent–toddler 258–61
- corpus striatum (striatum) 349, 355
 - anatomical studies 338–9
 - functional imaging 196, 341–2, 345–6
 - neurochemical studies 195
 - neuropharmacology 353
- cortico-striatal-thalamocortical circuits 340, 349
- Covert Orienting of Visuospatial Attention Task (COVAT) 225
- CPT, *see* Continuous Performance Task/Test
- criminal offending 273–4, 471
- criticism, expressed 374, 378–9, 392, 396–7
- cross-cultural/ethnic aspects 64–92
 - comorbidity 87–8
 - conduct disorder vs. hyperactivity 70–1
 - genetic factors 78, 79–81, 90–2, 303, 384
 - Hong Kong and UK studies compared 76–9
 - management 88–9
 - model 90, 91
 - outcome 89–90
 - prevalence 65, 66
 - external validity 69–70
 - factorial validity 67–9
 - sex differences 73, 74, 102
 - using behaviour rating scales 71–4
 - using clinical diagnoses 75–6
 - problems of behaviour rating scales 74–5
 - recognition of problem/help-seeking 85–7, 141–2
 - symptom presentation 81–4
- cuing techniques 167
- cultural aspects, *see* cross-cultural/ethnic aspects
- culture-bound syndrome, hyperactivity as 65, 67, 76, 78
- CYP2D6 89
- DAMP, *see* disorder of attention, motor control and perception
- Darwinism, social 7, 9, 129
- DAT1 (dopamine transporter) gene 80, 89, 307
 - association and linkage studies 309–14
 - in cerebellar vermis 339
 - diagnostic specificity and heterogeneity 325–6
 - genetic considerations 324
 - multimarker haplotypes 323
- data analysis 444–5
- defect of moral control 2, 5–7
- defiance 260, 389–90
 - see also* oppositional defiant disorder

495 Index

Delayed Motor Reaction Time Task 177–8
delay gratification, inability to 177–8, 255–6, 381, 401
delinquent behaviour 273–4, 394
demographic factors 369–70
dendritic sprouting 386
depression
 animal studies 386
 comorbid 50
 cultural/ethnic differences 81, 88
 maternal 53, 372, 375–6
 as cause of hyperactivity 381–3
 conduct disorder and 392, 395–6
 postnatal 382–3
 rating 476
depth-of-processing task 168
desipramine 484
developing countries 88, 89–90
development 242–76
 childhood stress and 389–90
 early caregiving relationships 250–61
 early precursors of hyperactivity 248
 experience-dependent brain 385–9
 long-term outcome and predictors 272–6
 Romanian adoptees 419–20
 self-regulation
 normative 248–61
 temperament and 261–7
 state regulation 196–7
developmental course of hyperactivity 268–72
dexamfetamine 130
 in adult ADHD 481, 483
 encoding and 169
 mechanisms of action 352, 354
 sex differences in response 109–10
 state regulation and 181–2
dextro-amphetamine, *see* dexamfetamine
'dextro-amphetamine response disorder' 1
diagnosis 126–54
 adult ADHD 145, 475–6
 categorical approaches 33–4, 135–7, 139–41, 153
 cultural/ethnic differences 75–6, 141–2
 dimensions 33–4
 for epidemiological studies 31–5
 future directions 149–52
 in girls 112–13, 145
 in North America vs. Europe 126–7, 131–5
 quantitative approaches 135, 136, 137–9, 153
 reliability and validity of symptoms 141–3
 structured interviews vs. checklists 35
 thresholds 32, 33–4, 53–4, 131, 138
 validity 145–9
 see also assessment; classification; rating scales
diagnostic criteria
 ADHD (DSM-IV) 132–3
 comparisons between 143–5
 evolution of formal 130–1
 future directions 149–52, 153–4
 hyperkinetic disorder (ICD-10) 134
 see also *Diagnostic and Statistical Manual; International Classification of Disease*
Diagnostic Interview Schedule for Children (DISC)
 epidemiological studies 36, 46, 54
 MTA Cooperative Group study 438, 439, 454
Diagnostic and Statistical Manual (DSM) 3, 31, 291
 second edition (DSM-II) 130, 468–9
 third edition (DSM-III) 75–6, 130, 469
 third edition revised (DSM-III-R) 31–2, 131, 469
 fourth edition (DSM-IV) 32, 131, 291, 469
 ADHD diagnostic criteria 132–3
 vs. ICD-10 64–5, 133, 143–5, 146–9, 214–15
 see also attention deficit hyperactivity disorder
Differential Reinforcement of Low-rate responding (DRL) procedure 177, 178
difficultness (temperamental)
 additive model of risk 264–5
 developmental course 271
 mother–toddler conflict and 259–61, 264
 self-regulation and 262–5
directiveness, maternal 197, 258–9, 260
discipline, physical 375
disorder of attention, motor control and perception (DAMP) 135
 combined with ADHD 146
 diagnosis 143–5
 validity 145–6
dissociation 389
distractibility
 early caregiver relationships and 256
 laboratory studies 170, 174, 175
divorce, *see* marital discord
dopamine 179, 195–6, 349–51
 motivation and 399–400
 theory of ADHD 350–1
dopamine β -hydroxylase 307, 321
dopamine receptors 350–1
 D2 gene (*DRD2*) 80, 320–1
 D3 gene (*DRD3*) 307, 319
 D4 351
 gene, *see* *DRD4* gene
 D5 gene (*DRD5*) 318–19
dopaminergic system 306–7, 349, 350–1
 candidate genes 307–21
 cross-cultural/ethnic studies 79–81
 functional imaging studies 341–2, 344–6
 mechanisms of stimulant actions 352–4
 regulation 351
dopamine transporter 339, 355
 gene, *see* *DAT1* (dopamine transporter) gene
double-task paradigm 170–1
Douglas, V.I. 20
DRD2 gene 80, 320–1
DRD3 gene 307, 319
DRD4 gene 80–1, 307, 308
 diagnostic specificity and heterogeneity 325–6
 executive function and 326
 genetic considerations 324
 inhibitory control and 152

Cambridge University Press

0521789613 - Hyperactivity and Attention Disorders of Childhood, Second Edition

Edited by Seija Sandberg

Index

[More information](#)

496 Index

- DRD4* gene (*cont.*)
 multimarker haplotypes 323
- DRD5* gene 318–19
- drive reduction theory 399
- drug
 abuse 273, 471–2
 metabolism 89
 treatment, *see* medication
- DSM, *see* *Diagnostic and Statistical Manual*
- dual-task paradigm 170–1
- effort 179–80
 evoked potentials 193–4, 195
 heart rate variability and 192–3
 nonoptimal 182–3, 189
- effortful control, development 266–7
- electroencephalography (EEG) 346–7
 quantitative (QEEG) 346, 349
- electrophysiological studies 346–9
- emotional factors 401
- encephalitis epidemic (1917–18) 2, 9, 129–30
- encephalopathy 16
- encoding (information uptake) 168–9
- endophenotypes 326
- energetical systems 179–80
- environmental influences 417–18
 cross-cultural/ethnic studies 78
 development 243
 development of self-regulation 248, 250
 historical perspective 21–2
 nonshared (c^2) 296, 299–300
 sex differences 118
 shared (c^2) 296
 state regulation and 196–9
 twin studies 299–300
 vs. constitutional 402–3
see also psychosocial factors
- epidemiology 30–59
 adequacy of standard rating scales 57
 adult ADHD 470
 defining caseness 31–5
 measuring impairment 54–5
 stability of classification and natural history 55–7
 subthreshold cases and ADHD boundaries 53–4
 surveys from 1990–2000 36–53, 57–9
 age effects 47–8
 cognitive correlates 51–3
 comorbidity 50
 gender influences 46–7
 informant perspective 48–9
 methodology 36–7
 prevalence in children and adolescents 37–46
 prevalence/distribution in adults 50–1
 psychosocial correlates 53
 questionnaire-based 36, 38–9
 structured interview-based 36–7, 40–5
 subtypes of ADHD 49
- epilepsy 15
- epistasis 324
- ‘erethic child’ 11
- Errors and Time test 217–18
- ethnic aspects, *see* cross-cultural/ethnic aspects
- Europe, diagnostic practice 3, 126–7, 131–5
- European Society for Child and Adolescent Psychiatry 4
- evaluation system 179
- event rates 189
 basic attention abilities and 167, 168, 169, 173
 impulsivity and 187, 188
 sustained attention and 181–5
- event-related potentials (ERP) 193–6, 347–8, 349
- evoked potentials 193–4
 Mulder’s model 193, 194
 steady-state visually (SSVEP) 348
- exclusion criteria, MTA Cooperative Group study 439
- executive function 217–21
 in ADHD subtypes 218–21
 in adult ADHD 477, 478
 definition 217
 early development 266–7
 molecular genetics 326
 reading disability and 223
 sex differences 107
- experiential factors 402–3
see also psychosocial factors
- experimenters, presence or absence 182–3, 340
- explosiveness, mental 4–5
- expressed emotion, parental 374, 378–9, 392, 396–7
- externalizing behaviour problems 390
 early antecedents 253–5, 269–71
 early temperament and 262–3, 266
 mental retardation with 187
 sex differences 101, 109
see also conduct disorder; oppositional defiant disorder
- eye movements, ballistic 167
- factor analysis 138–9, 140
 cross-cultural comparisons 67–9
 subtypes of ADHD 147
- false negatives 150
- false positives 150
- family
 factors
 in conduct disorder 393, 395–7
 correlated with ADHD 53, 370–2
 maintaining ADHD 275, 373–9
 foster care 418, 425–8, 429–30
 size 370
 stability 370–1
 stress 106–7, 261, 271, 370
 studies 115–17, 292–4
- father–child relationships 373, 374, 391
- Fels Longitudinal Study 265
- Figures Embedded Test 165
- floor effect 172

497 Index

- food
 - additives 3, 21
 - allergy 3, 21
- forced-choice reaction time task 170–1
- foster care, family 418, 425–8, 429–30
- frontal lobe ablation studies 12
- frontostriatal system dysfunction 195–6
- functional brain imaging studies 340–6
- functional impairment 54–5
- Gastaut, H. 13–14
- gaze behaviour, newborn infants 197
- gender differences, *see* sex differences
- General Cognitive Index (GCI) 422
- genes
 - candidate, *see* candidate genes
 - vs. experiences 383–4
- genetic influences 417, 418
 - additive 116, 117, 296, 299–300
 - cross-cultural/ethnic aspects 78, 79–81, 90–2, 303, 384
 - nonadditive 296, 299
 - parental psychopathology 376–7
 - sex differences 115–17
- genetics 290–328
 - behavioural 292–302
 - comorbidity 228–32, 390
 - complex traits 305
 - future directions 323–7
 - methodology 228–9
 - molecular 302–28
 - phenotype definitions and 150–2
 - state regulation problem 188–9
- genome scans 306, 327
- germ corruption 8
- girls
 - clinical practice 118–19
 - diagnostic aspects 112–13, 145
 - general protection 111–12
 - see also* sex differences
- globus pallidus 338–9, 340, 349
- glucose metabolism, cerebral 108, 196, 341
- go–no-go test 185–8, 192, 341
- gratification, ability to delay 177–8, 255–6, 381, 401
- guidelines, management 4
- Gurevich, M.O. 11
- haplotype-based haplotype relative risk (HHRR) 303, 304
- Hawthorn effect 188
- head injury, severe closed 338
- heart rate deceleration 191–3
- heritability (h^2) 296, 298–9, 300
 - sex differences 116–17
 - subtypes of ADHD 150–2, 300–1
- heterozygosity 324
- hippocampus 179, 384, 387
- history 1–22, 128–31
 - early medical explanations 4–18
 - emergence of behavioural definitions 18–22
 - syndrome issue 22
- Hoff, H. 10–11
- home hyperactivity 48–9, 144
- homovanillic acid (HVA) 353–4
- homozygosity 324
- Hong Kong study 76–9, 141–2
 - see also* Chinese
- hyperactive child syndrome 3
- hyperactive/impulsive type of ADHD (ADHD;HI)
 - 32, 144, 215
 - in adults 50–1
 - cognitive functioning 224–5
 - epidemiological surveys 49
 - genetics of comorbidity 230–1
 - molecular genetics 310–11, 315, 317, 325
 - persistence into adulthood 473, 475
 - validity 147–8
- hyperactivity
 - developmental course 268–72
 - historical perspective 1–22, 128–31
 - in preschool children 269
 - as symptom, *see* overactivity
 - use of term 65, 127
 - see also* attention deficit hyperactivity disorder; hyperkinetic disorder
- hyperexcitability 4–5
- hyperkinetic behaviour syndrome 18
- hyperkinetic child syndrome 130
- hyperkinetic conduct disorder 33, 133, 145, 146–9
- hyperkinetic disease 10–11
- hyperkinetic disorder (HKD) 32–3, 64–5, 214–15, 291
 - cognitive deficits 52
 - cross-cultural comparisons 75–6
 - diagnosis 143–5
 - diagnostic criteria 134
 - diverging diagnostic practice 131–5
 - history 1
 - molecular genetics 325
 - prevalence 46
 - sex differences, *see* sex differences
 - subtype, *see* hyperkinetic conduct disorder
 - validity 145–6, 154
- hyperkinetic impulse disorder 13, 18
- hyperkinetic reaction of childhood 130, 468–9
- hypervigilance 389–90
- hypothalamus 399–400
- hypothyroidism, congenital 184
- ICD, *see* *International Classification of Disease*
- identical by descent (IBD) 303–4
- immunological function 386
- impairment, functional 54–5
- imprinting 324
- impulsivity 244–5
 - cross-cultural/ethnic studies 65, 66, 77
 - developmental course 268
 - in DSM diagnosis 32, 33, 132

498 Index

- impulsivity (*cont.*)
 - early caregiving relationships and 254, 255–6, 381
 - early temperament and 263–4
 - heart rate deceleration and 191–2
 - historical perspective 10
 - measures 477
 - mental retardation and 187, 190
 - motor control and 174–8
 - sex differences 103–4
 - stability over time 55–6
 - state regulation and 185–9
- inattention (attention deficits) 244
 - cross-cultural/ethnic studies 65, 66, 77
 - developmental course 268
 - in DSM diagnosis 32, 33, 132
 - genetic influences 298–9
 - historical perspective 3, 19–20
 - in hyperactivity 52–3
 - institutional care as risk factor 417–32
 - sex differences 103–4, 109
 - stability over time 55–6
 - state regulation and 181–5
 - see also* distractibility
- inattentive type of ADHD (ADHD/PI) 32, 144
 - in adults 50–1
 - cognitive functioning 216, 218–21, 224–5
 - genetics of comorbidity 230–1
 - molecular genetics 310–11, 325
 - prevalence 49, 58
 - validity 147–8
- incentive theory 399
- inclusion criteria, MTA Cooperative Group study 439
- infants
 - caregiver relationships, *see* caregiver–infant relationships
 - fixation of novel objects 266
 - modulation of arousal 249, 250, 262
 - self-regulation 247, 248–50
 - state regulation 196–7
 - temperament, *see* temperament
 - see also* newborn infants; preschool children; toddlers
- informants (raters)
 - in adult ADHD 51, 476–7, 479–80
 - agreement/disagreement between 163
 - cultural/ethnic variations 75
 - effects on prevalence rates 48–9
 - epidemiological studies 36, 58–9
 - reliability 142–3, 162–4
 - sex bias 101, 110–11
 - see also* under parents; teachers
- information processing
 - approach 165–6
 - capacity 170–3
 - output (motor) side 173–4
 - psychophysiological measures 191–6
 - slow-inaccurate 175, 180
 - slowness 177
 - state/process distinction 166
- information uptake (encoding) 168–9
- inheritance, mode of 324
- inhibition function 175–6
- inhibitory control 152, 244–5
 - defects, *see* impulsivity
 - development 267
- institutional care 21, 417–32
 - adoptees from Romania 419–25, 428–31
 - methodological issues 419
 - in UK 425–8, 429–31
- intellectual impairment, *see* learning disorders
- intelligence (IQ) 52, 216, 400
 - in adult ADHD 477
 - diagnostic aspects 147
 - executive functioning and 220
 - future research 233
 - institutional care and 419–20, 422, 426
 - sex differences 105, 109
 - state regulation and 187
 - see also* learning disorders; mental retardation
- interference, maternal 253, 256–8
- International Classification of Disease* (ICD) 3, 31
 - ninth edition (ICD-9) 75–6
 - tenth edition (ICD-10) 32–3, 133, 291
 - diagnostic criteria 134
 - vs. DSM-IV 64–5, 133, 143–5, 146–9, 214–15
 - see also* hyperkinetic disorder
- interviews
 - semistructured 142–3
 - structured 35, 142–3
- intrusiveness, maternal 253, 256–8, 378–9
- irritability, infant 263
- Kahn, E. 11–12
- knockout mouse studies 307, 308, 309, 319, 324
- laboratory measures
 - basic attention abilities 167–73
 - choosing 164–6
 - cross-cultural/ethnic studies 77, 84
 - impulsivity 174–8
 - psychophysiological 191–6
 - for refining phenotypes 326
 - validity 162–4
 - vs. clinical labels 199–200
- language disorders 225–8
 - early caregiver relationships and 257
 - in pervasive hyperactivity 48–9
 - sex differences 105
- latent class analysis 140, 147, 151, 325
- Laufer, M.N. 13–14, 18
- lead poisoning 12, 21
- learned ineffectiveness 21
- learning, use-dependent 388–9
- learning disorders 246
 - ADHD and 189–91, 222–5
 - in adult ADHD 477–8
 - future research 232–3
 - institutional care and 422, 429

Cambridge University Press

0521789613 - Hyperactivity and Attention Disorders of Childhood, Second Edition

Edited by Seija Sandberg

Index

[More information](#)**499 Index**

- language disorders and 227–8
- sex differences 105, 111
- sustained attention deficits 184
- terminology 222
- vs. hyperactivity 163
 - see also* intelligence; mental retardation
- letter-matching task 168
- liability, sex differences in 114–15
- linkage studies 302–5
- listening task, dichotic 167–8, 169
- locus ceruleus (LC) 340, 349, 350, 387
- low birth weight 421
- magnetic resonance imaging (MRI)
 - anatomical studies 336–7, 338, 339–40
 - functional (fMRI) 199, 341–4
- malnutrition 419, 421–2, 429
- management, *see* treatment
- marital discord 106–7, 371
 - conduct disorder and 392, 395–6
 - as maintaining factor 377
- Matching Familiar Figures (MFF) test 52, 165, 174, 217–18
 - in adult ADHD 477, 478
- mating, assortative (nonrandom) 297–8
- medication
 - in adult ADHD 480–4
 - compliance 450–1, 459–61
 - cultural/ethnic differences 88–9
 - dosing regimens 453, 456, 481–2
 - effect on parent–child relationship 377–9
 - history of use 12–14
 - impact of availability 21–2
 - monitoring/follow-up 456–7
 - MTA Cooperative Group study (MedMgt) 440, 446–7, 448, 452, 453, 454–6, 461
 - normalizing hypothesis 187–8
 - stimulus level shift hypothesis 188
 - withdrawal 483
 - see also* stimulants
- memory
 - recognition tasks 171–2
 - search paradigm 176
 - state-dependent 388–9
 - traumatic experiences 387–8
 - verbal 223
 - working 219
- Mental Deficiency (Amentia)* (Tredgold) 8
- mental explosiveness 4–5
- mental retardation
 - ADHD and 189–91
 - impulsivity and 187, 190
 - in Romanian adoptees 419–20
 - see also* intelligence; learning disorders
- mesocorticolimbic dopamine system 350, 351
- metabolism, drug 89
- 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) 196
- methylphenidate (MPH)
 - in adult ADHD 478, 481–2, 483
- cognitive task performance and 166, 168–9, 172, 173–4
 - cross-cultural/ethnic comparisons 88–9
 - effect on parent–child relationship 377–8
 - evoked potentials and 195
 - functional brain imaging and 342, 343, 344–5
 - genetics of response 318
 - impulsivity and 187–8
 - in institutionally reared children 423, 424
 - mechanisms of action 309, 351–4
 - in MTA Cooperative Group study 448
 - sex differences in response 109–10, 119
 - sustained attention and 182–3
 - tachyphylaxis 353
- metrozol 13–14
- minimal brain damage 2, 8, 14–16, 468
 - syndrome 15–16
- minimal brain dysfunction (MBD) 2–3, 16–18, 130
 - decline in popularity 20
 - defined 17
 - diverging diagnostic practice 131
 - sibling studies 293
- Minnesota High Risk Study 253–4, 256
- molecular genetics 302–28
- monoamine oxidase inhibitors (MAOIs) 483, 484
- mood disorders, *see* affective disorders
- moral control, defect of 2, 5–7
- moral deficiency 8
- moral dyscontrol 128–30
- mother–child relationships 373–9
 - affective content 374
 - conduct disorder and 392–3
 - effect of medication 377–9
 - marital discord and 377
 - maternal depression and 375–7, 381–2
 - parenting styles and 258–61, 375
 - sex differences 106–7
 - state regulation and 197–8
 - see also* caregiver–infant relationships
- mothers
 - depressed, *see* depression, maternal
 - directiveness 197, 258–9, 260
 - intrusiveness 253, 256–8, 260, 378–9
- motivational factors 179, 182, 267, 398–400
- motor activation, heart rate deceleration and 191, 192
- motor activity
 - during CPT performance 184–5
 - excessive 3, 18
 - see also* overactivity
- motor inhibition 271
- motor-related dysfunction 173–8, 180
 - evoked potentials and 195
 - neurochemical levels and 195–6
- motor restlessness syndrome 12
- MPTP (1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine) 196
- MRI, *see* magnetic resonance imaging

500 Index

- MTA Cooperative Group study 119, 378, 435–62
 - clinical impact of treatment 454–6
 - discussion 452–61
 - interpretation of findings 456–61
 - methods 438–46
 - data analysis 444–5
 - outcome measures 445–6
 - recruitment and sample characteristics 438–9
 - study protocol (treatment conditions) 439–42
 - study variables 442–4
 - primary aims and hypotheses 437–8
 - results 446–52
- Mulder's model 193, 194
- Multimodal Treatment Study of Children with Attention Deficit/Hyperactivity Disorder,
see MTA Cooperative Group study
- multiple threshold model, polygenetic 114–17
- multisecond oscillations 354–5
- myelin oligodendrocyte glycoprotein gene (MOG) 322
- natural history of hyperactivity 55–7
- negative wording, rating scale items 57
- neglect, child 385, 386–7, 401
- neonates, *see* newborn infants
- neurasthenia, hypersthenic 11
- neuroanatomy 336–40
- neurochemistry 195–6, 349–51
- neuroleptics 484
- neuropathic diathesis 7–9
- neuropharmacology 351–5
- neuropsychological function 107, 216–17
- neurotransmitters
 - in alarm reaction 386–7
 - in arousal 179
 - molecular genetics 306–23
 - see also* dopamine; noradrenaline; serotonin
- newborn infants
 - acute stress 388
 - gaze behaviour 197
- nicotinic acetylcholine $\alpha 4$ receptor (*CHRNA4*) gene 322–3
- nigrostriatal dopamine system 350, 351
- nonverbal skills 390
- noradrenaline (norepinephrine) 179, 195–6, 321, 349–50
- North America, diagnostic practice 3, 126–7, 131
- nosology, scientific 128, 149–52
- novelty-seeking personality trait 308
- nucleus accumbens 338
- objective measures, *see* laboratory measures
- observation, behavioural 144, 165–6
 - in adult ADHD 476–7
 - during CPT performance 184–5
 - institutional care and 426–7
- occupational status, parents' 53
- ODD, *see* oppositional defiant disorder
- oppositional defiant disorder (ODD) 390
- comorbidity rates 50, 104
- diagnostic issues 148
- executive functioning and 219
- hyperactivity as risk factor 394
- institutional care and 420–1
- molecular genetics 325
- subthreshold cases and boundaries 53–4
- treatment efficacy and 449, 454–5, 458
- Oral Directions and Recall of Sentences 219
- 'organic drivenness' 2, 11–12
- orientation 167
- oscillations, multisecond 354–5
- outcome 272–6
 - cultural/ethnic differences 89–90
 - MTA Cooperative Group study 445–6, 446–7
 - predictors 274–6
 - sex differences 108
 - see also* adult hyperactivity
- overactivity 244
 - anxiety and 402
 - cross-cultural/ethnic studies 65, 66, 77
 - developmental course 268
 - in DSM diagnosis 32, 33, 132
 - genetics 298–9
 - institutional care as risk factor 417–32
 - long-term outcome 272
 - motivational factors 398–9
 - objective measures 165
 - sex differences 103–4, 109
 - stability over time 55–7
- overcontrolled behaviour 82–3
- overexcitability 4–5
- Paired Associate Learning (PAL) test 52
- paper-and-pencil tasks 182
- parent–child relationships 246, 373–9
 - affective content 374
 - conduct disorder and 391–3
 - developmental course 269
 - early, *see* caregiver–infant relationships
 - effect of medication 377–9
 - long-term outcome and 274, 275
 - marital discord and 377
 - parenting styles and 375
 - sex differences 106–7
 - toddler control interactions 258–61
 - see also* father–child relationships; mother–child relationships
- parenting styles 21, 375, 401
 - conduct disorder and 392–3
 - cultural/ethnic differences 82, 84
 - toddler development and 258–61
 - treatment outcomes and 451
- parents
 - adoption studies 294–6
 - cross-cultural/ethnic comparisons 82, 84, 87
 - as informants
 - in adult ADHD 51, 479, 480
 - effects on prevalence rates 48–9
 - effects on sex ratios 101, 110–11

501 Index

- in epidemiological surveys 36, 46
 - reliability 142, 162–4
- psychopathology in 292–3, 371–2, 381–3
- satisfaction with treatment 459–60
- unemployment 370
- see also* marital discord
- peer relationships 246
 - activity level and 265–6
 - early predictors of problems 271–2
 - institutional care and 427, 428
 - sex differences 105–6, 109
- pemoline, in adult ADHD 481, 482, 483
- perinatal complications 15, 16
- perseverations
 - in ADHD subtypes 218–19
 - response 178
- persistence of hyperactivity 55–7, 471–4
 - sex differences 108
 - see also* adult hyperactivity
- pervasive hyperactivity
 - diagnostic aspects 143–4
 - vs. situational hyperactivity 48–9, 163–4
- pharmacological treatment, *see* medication
- phenotypes
 - defining 150–2
 - refining 326
- phenylketonuria (PKU), early-treated 186–7
- photometrozol stimulation test 13–14
- physical abuse 386–7, 401
- physiological bases 401–2
- pneumoencephalograms 12
- polygenetic multiple threshold model 114–17
- Porteus Mazes 52, 165, 174
- positivism 7
- positron emission tomography (PET) 341, 344, 353
- postencephalic behaviour disorder 9
- power analysis 444
- prefrontal cortex
 - anatomical studies 337, 339–40
 - functional imaging 341–2, 344, 345
 - neurochemistry 350
- pregnancy, stress during 385
- premature infants 421
- prenatal stress 385
- preschool children
 - activity levels 265–6
 - attention and effortful control 267
 - diagnostic aspects 145
 - institutional care 419–28
 - predictors of hyperactivity 268–72
 - prevalence of ADHD 47
 - self-regulation 248, 250
 - see also* children; infants; toddlers
- prevalence 37–46, 58, 291
 - in adults 50–1, 58, 473
 - age effects 47–8
 - and comorbidity 50
 - cultural/ethnic differences, *see*
 - cross-cultural/ethnic aspects, prevalence
 - informant perspective 48–9
 - sex differences, *see* sex differences, prevalence
 - subtypes of ADHD 49
- primate research 12, 385
- principle components analysis (PCA) 445
- problem-solving skills, development 253
- problem suppression-facilitation model (Weisz)
 - 82–3, 84
- processes 166
- psychiatric disorders
 - comorbidity, *see* comorbidity
 - hyperactivity as risk factor 394–5
 - in parents 292–3, 371–2, 381–3
 - prevalence 37–46
 - in relatives 292–4
- psychoanalytical perspective 21
- psychometric tests 65, 477–9
- psychopathic diathesis 8
- psychopathology, *see* psychiatric disorders
- psychophysiological measures 191–6
- psychosis, stimulant-induced 482–3
- psychosocial factors 367–403, 417
 - causative 379–83
 - comorbidity and 390–8
 - correlated 53, 369–71
 - maintaining 373–9
 - mechanisms of effects on brain 384–90
 - theoretical basis 398–402
 - vs. genes 383–4
 - see also* institutional care
- psychostimulants, *see* stimulants
- putamen 338, 341, 353
- raclopride 346
- random regression (RR) techniques 444–5
- raters, *see* informants
- rating scales 137–9
 - adequacy of standard 57, 142–3
 - adult ADHD 476
 - cross-cultural/ethnic studies 71–5, 84
 - structured interviews vs. checklists 35, 142–3
- reaction times (RT) 167, 173–4
- reading disability/disorder (RD) 52, 222–5
 - diagnosis of hyperactivity and 140
 - executive functioning and 219
 - genetics of comorbidity 229–31
 - language disorders and 227–8
 - sex differences 105
 - treatment effects 453
- referrals, sex differences 100–1, 110–11
- reinforcement, withholding responding to receive 177–8
- remission 55, 485
- reorientation 167–8
- reproductive casualty, continuum of 16
- respondents, *see* informants
- response
 - bias condition 176
 - competition 174
 - inhibition deficit 175–7, 188, 218

502 Index

- response (*cont.*)
 - perseveration task 178
 - withholding, to receive reinforcement 177–8
- restlessness
 - in adults 272, 478
 - see also* overactivity
- reticular activating system 13–14
- reward seeking 401
- Romanian adoptees 419–25, 428–31
- rule-governed behaviour, deficit in 247
- rural areas 74
- Rutter Parent and Teacher Questionnaire 138
 - cross-cultural/ethnic studies 67–8, 69
 - institutional care and 420, 421, 426
 - sex differences 101, 103–4
- sample size 444
- Sanders model 179–80
- Schedule for Affective Disorders and Schizophrenia for School-Age Children (K-SADS) 36
- schizophrenia 81, 89–90, 339
- school
 - based treatment 441
 - cultural/ethnic differences 82, 89
 - hyperactivity 48–9, 144, 421
- schooling, years completed 472
- scientific nosology 128, 149–52
- selective serotonin reuptake inhibitors 484
- selegiline 484
- self-control, normative development 249–50
- self-esteem 268
- self-regulation
 - construct 247
 - development 248–61
 - early caregiving relationships and 250–61, 380–1
 - phases (Kopp) 248–50
 - hyperactivity as disorder of 246–7, 401
 - individual differences in competence 250
 - temperament and 261–7
 - see also* state regulation
- self-report, adult ADHD 50–1, 479–80
- self-stimulation
 - during CPT performance 185
 - hypothesis 170
- sensation seeking 189, 401
- separation anxiety 224
- serotonin 179, 196
 - system, candidate genes 321–2
- serotonin receptors
 - 1B gene (*HTR1B*) 322
 - 2A gene (*HTR2A*) 321–2
- services, child mental health 88–9
- sex differences 99–119
 - adolescent and adult outcome 108, 473–4
 - adult ADHD 51
 - in clinical practice 118–19
 - comorbidity 104–5
 - core symptoms 103–4
 - early developmental antecedents 264, 270–1
 - electrophysiological studies 347
 - extent of 109–10
 - future research 117–18
 - institutional care effects 427
 - intellectual ability/academic performance 105
 - molecular genetics 318, 325–6
 - neuropsychological function 107
 - parent–child relationship 106–7
 - peer relationships 105–6
 - presentation 103–10
 - prevalence 46–7, 100–3
 - aetiology 113–15
 - constitutional variability model 113–14
 - cultural/ethnic differences 73, 74, 102
 - familial and genetic clustering 115–17
 - nature of 110–13
 - polygenic multiple threshold model 114–17
 - subtypes of ADHD 49
- sexual abuse 386–7
- sexual problems 471
- siblings
 - interactions with 246, 299–300, 373, 392
 - psychopathology in 293–4
 - sex differences and 116
- single-photon emission computed tomography (SPECT) 309, 341, 344–5, 353
- situational hyperactivity 368–9
 - vs. pervasive hyperactivity 48–9, 144, 163–4
- SNAP-25 gene 322
- SNAP rating scale (Swanson) 445, 451–2
- social class 53, 370, 401
- social Darwinism 7, 9, 129
- social disadvantage 370
- social factors 369–70
- social interactions
 - state regulation and 197–8
 - see also* parent–child relationships; peer relationships
- social maladjustment 21, 246
 - developmental course 269, 271–2
 - institutional care and 423–5, 427, 428, 431
 - long-term outcome 272–6
 - sex differences 118
- social skills 445
- social support 275
- socioeconomic status 53
 - in adulthood 472
 - cross-cultural/ethnic studies 74
 - treatment efficacy and 448–9
- sociopathy, *see* antisocial personality disorder
- somatization disorder 292–3
- Soviet Union 11, 15–16
- speech problems 52, 224
 - historical perspective 10
 - subtypes of ADHD and 224
 - see also* language disorders
- state 178–91
 - process distinction 166
 - state regulation 199

503 Index

- deficit 200, 336
- development 196–7
- effects of medication 187–8
- environmental factors and 196–9
- experimental testing and 198–9
- heart rate variability and 192–3
- impulsiveness and 185–9
- Sanders model 179–80
- social interactions and 197–8
- sustained attention and 181–5
 - see also* self-regulation
- Still, Sir George Frederick 2, 4, 5–7, 100, 129
- stimulants
 - in adult ADHD 481–3
 - anxiety disorder and 398
 - cognition and 234
 - contraindications in adults 482
 - cultural/ethnic differences 89
 - history of use 12–14, 130
 - in institutionally reared children 423, 424, 429
 - mechanisms of action 307, 309, 351–5
 - sex differences 109–10, 119
 - side-effects in adults 482–3
 - see also* amfetamines; dexamfetamine; methylphenidate
- stimulation
 - lack of 400–1
 - see also* self-stimulation
- stimulus degradation task 168
- stimulus-onset asynchrony (SOA) 167
- stop signal paradigm 175–7, 188–9, 191–2
- Stop Task 342
- Strange Situation paradigm 252
- Strauss, A.A. 14–15
- Strengths and Weaknesses of ADHD-symptoms and Normal-behaviors (SWAN) rating scale 57
- stress
 - family 106–7, 261, 271, 370
 - impact on brain 384–90
 - neonatal 388
 - prenatal 385
 - response 386–7
 - see also* traumatic experiences
- striatum, *see* corpus striatum
- Stroop test 52, 165, 174, 216, 217–18
 - in ADHD subtypes 220
 - in adult ADHD 477, 478
 - functional imaging studies 342–3
- structure-seeking/providing behaviour 197–8
- substance abuse 273, 386
 - in adult ADHD 471–2, 474, 482
 - sex differences 108
- substantia nigra 338–9, 349, 351
- subtypes of ADHD 32, 144, 215, 291
 - in adults 50–1
 - age and prevalence rates 47
 - cognitive functioning and 218–21, 224–5
 - epidemiological surveys 49
 - heritability 150–2, 300–1
 - molecular genetics 310–11, 325
 - validity 146–9
- symptoms 243–6
 - associated 245–6
 - cultural/ethnic differences 81–4
 - diagnostic thresholds 32, 33–4
 - functional impairment and 54–5
 - laboratory measures, *see* laboratory measures
 - MTA Cooperative Group study 445, 446–7, 452
 - primary (core) 244–5
 - reliability and validity 141–3
 - retrospective assessment, in adulthood 479–80
 - sex differences 103–4
 - stability over time 55–7
 - subthreshold 53–4
 - see also* impulsivity; inattention; overactivity
- synapses 386
- synchrony, caregiver–infant 251–2
- syndrome, issue of 22
- tachyphylaxis, methylphenidate 353
- task-inappropriate responding 174–8
- task inefficiency 167, 173, 174, 179–80
- taxonomies 127–8
- Teacher Report Form (TRF) 36, 69, 83
- teachers
 - cross-cultural/ethnic comparisons 85–7
 - as informants 36, 37
 - effects on prevalence rates 48–9
 - effects on sex ratios 101, 110–11
 - reliability 162–4
 - relationships with 246
- temperament 250, 256–7
 - cultural/ethnic studies 81–2
 - genetic influences 298
 - parent–toddler conflict and 260, 264, 266
 - self-regulation and 261–7
 - stability 264
 - see also* difficultness
- temporal insula 353
- Thailand 82–3
- tic disorder (TD) 186
- toddlers
 - compliance 253, 258–60, 381
 - development of self-regulation 258–61
 - see also* infants; preschool children
- tomoxetine 484
- Tourette syndrome 107
 - diagnostic issues 149
 - molecular genetics 320
 - neuroanatomical studies 338
- Tower of Hanoi (TOH) 217–18
- traffic task 171
- Trail-making Test 216, 217–18, 477, 478
- transactional relationships 259–60
- Transmission Disequilibrium Test (TDT) 304–5
- traumatic experiences 386–90
 - developmental consequences 389–90
 - effects on brain function 387–9
 - see also* stress

504 Index

treatment 435–62
 in adult ADHD 480–4
 cross-cultural/ethnic comparisons 88–9
 history 5, 12–14
 MTA Cooperative Group study, *see* MTA Cooperative Group study
 multimodal (Comb) 441, 446–7, 448–9, 452–6, 460–1
 sex differences 118–19
 see also medication; stimulants
Tredgold, Alfred F. 2, 4, 7–8, 129
tricyclic antidepressants 483–4
twin studies 296–301
 comorbidity 224–5, 229–30
 methodological issues 117, 296–8
 sex differences 115, 116, 117
 state regulation 188–9
tyrosine hydroxylase 307
 gene (*TH*) 323

undercontrolled behaviour 82–3
unemployment, parental 370
‘uninhibited child with choreatic symptoms’ 16
United Kingdom (UK)
 cross-cultural/ethnic study 76–9
 diagnostic practice 75–6, 127, 131–5
 institutional care 425–8, 429–31

 sex differences in prevalence 100–1
United States of America (USA)
 diagnostic practice 75–6, 126–7, 131
 sex differences in prevalence 100–1
University of Massachusetts Medical Center
 Protocol 480
urban areas 74

ventral tegmental area (VTA) 340, 349, 351
Verbal Fluency test 216
verbal memory 223
vigilance 181–5
visual behaviour, during CPT performance 184–5

Wechsler Adult Intelligence Scale (WAIS) 477
Wechsler Intelligence Scale for Children 426
 Revised (WISC-R) 52
 third edition (WISC-III) 219
Wender Utah Rating Scale (WURS) 479–80
Werner, H. 14–15
Wide Range Achievement Test (WRAT) 477
Wisconsin Card Sorting Test 174, 217–19, 220
 in adult ADHD 477, 478
working memory tasks 219

Youth Self-report Form (YSF) 36