

# Part 1

## Arrangements in Dimension Two

# 1

## Congruent Domains in the Euclidean Plane

Let  $K$  be a convex domain. According to the classical result of L. Fejes Tóth [FTL1950], the density of a packing of congruent copies of  $K$  in a hexagon cannot be denser than the density of  $K$  inside the circumscribed hexagon with minimal area. Besides this statement, we verify that the same density estimate holds for any convex container provided the number of copies is high enough. In addition, we show that if  $K$  is a centrally symmetric domain then the inradius and circumradius of the optimal convex container cannot be too different. Following L. Fejes Tóth [FTL1950] in case of coverings, the analogous density estimate is verified under the “noncrossing” assumption, which essentially says that the boundaries of any two congruent copies intersect in two points. In case of both packings and coverings, congruent copies can be replaced by similar copies of not too different sizes. Finally, we verify the hexagon bound for coverings by congruent fat ellipses even without the noncrossing assumption, a result due to A. Heppes.

Concerning the perimeter, we show that the convex domain of minimal perimeter containing  $n$  nonoverlapping congruent copies of  $K$  gets arbitrarily close to being a circular disc for large  $n$ . However, if the perimeter of the compact convex set  $D$  covered by  $n$  congruent copies of  $K$  is maximal then  $D$  is close to being a segment for large  $n$ .

### 1.1. Periodic and Finite Arrangements

Let  $K$  be a convex domain. Given an arrangement of congruent copies of  $K$  that is periodic with respect to some lattice  $\Lambda$  (see Section A.13) and given  $m$  equivalence classes, it is natural to call  $m \cdot A(K)/\det \Lambda$  the density of the arrangement. We define the *packing density*  $\delta(K)$  to be the supremum of the densities of periodic packings of congruent copies of  $K$  and the *covering density* to be the infimum of the densities of periodic coverings by congruent copies of  $K$ . In addition, we define  $\Delta(K) = A(K)/\delta(K)$  and  $\Theta(K) = A(K)/\vartheta(K)$ .

It is not hard to show that optimal clusters asymptotically provide the same densities as periodic arrangements (see Lemma 1.1.2). Our main result is that, in the planar case, finite packings are not denser (asymptotically) than periodic packings, and the analogous statement holds for coverings. We note that this is a planar phenomenon: Say, if  $d \geq 3$  and  $K$  is a right cylinder whose base is a  $(d - 1)$ -ball, then linear arrangements are of density one, whereas any periodic packing is of density at most  $\delta'$  for some  $\delta' < 1$ , and any periodic covering is of density at least  $\vartheta'$  for some  $\vartheta' > 1$  (see Lemma 7.2.5).

**Theorem 1.1.1.** *Let  $K$  be a convex domain, and let  $n$  tend to infinity.*

- (i) *If  $D_n$  is a convex domain of minimal area containing  $n$  nonoverlapping congruent copies of  $K$  then  $A(D_n) \sim n \cdot \Delta(K)$ .*
- (ii) *If  $\tilde{D}_n$  is a convex domain of maximal area that can be covered by  $n$  congruent copies of  $K$  then  $A(\tilde{D}_n) \sim n \cdot \Theta(K)$ .*

Since periodic arrangements correspond canonically to finite arrangements on tori (see Section A.13),  $\Delta(K)$  is the infimum of  $V(T)/m$  over all tori  $T$  and integers  $m$  such that there exists a packing of  $m$  embedded copies of  $K$  on  $T$ , and  $\Theta(K)$  is the supremum of  $V(T)/m$  over all tori  $T$  and integers  $m$  such that there exists a covering of  $T$  by  $m$  embedded copies of  $K$ . The first step towards verifying Theorem 1.1.1 is the case of clusters.

**Lemma 1.1.2.** *Given convex domains  $K$  and  $D$  with  $r(D) > R(K)$ , let  $N$  be the maximal number of nonoverlapping congruent copies of  $K$  inside  $D$ , and let  $M$  be minimal number of congruent copies of  $K$  that cover  $D$ . Then*

- (i)  $\left(1 + \frac{R(K)}{r(D)}\right)^2 \cdot A(D) \geq N \cdot \Delta(K) \geq \left(1 - \frac{R(K)}{r(D)}\right)^2 \cdot A(D);$
- (ii)  $\left(1 + \frac{R(K)}{r(D)}\right)^2 \cdot A(D) \geq M \cdot \Theta(K) \geq \left(1 - \frac{R(K)}{r(D)}\right)^2 \cdot A(D).$

**Remark.** Instead of the upper bound in (i), we actually prove the stronger estimate  $A(D + R(K)B^2) \geq N \cdot \Delta(K)$ .

*Proof.* We place  $K$  and  $D$  in a way that  $K \subset R(K)B^2 \subset D$ . In particular, assuming that  $K'$  is congruent to  $K$ , if the circumcentre  $c$  of  $K'$  lies outside  $D + R(K)B^2$  then  $K'$  avoids  $D$ , and if  $c \in (1 - R(K)/r(D))D$  then  $K' \subset D$ . Given a torus  $T$ , we write the same symbol to denote a convex domain in  $\mathbb{R}^2$  and its embedded image on  $T$ .

We present the proof only for packings because the case of coverings is completely analogous. Let  $T = \mathbb{R}^2/\Lambda$  be any torus satisfying that

$C = D + R(K)B^2$  embeds isometrically into  $T$ , and let  $K_1, \dots, K_m$  be the maximal number of nonoverlapping embedded copies of  $K$  on  $T$ . Writing  $x_i$  to denote the circumcentre of  $K_i$ , we have (see (A.50))

$$\int_T \#((C + x) \cap \{x_1, \dots, x_m\}) dx = m \cdot A(C). \tag{1.1}$$

Thus there exists a translate  $C + x$  that contains at most  $k \leq m \cdot A(C)/A(T)$  points out of  $x_1, \dots, x_m$ , say, the points  $x_{i_1}, \dots, x_{i_k}$ . After replacing  $K_{i_1}, \dots, K_{i_k}$  by the  $N$  nonoverlapping embedded copies of  $K$  contained in  $x + D$ , we obtain a packing of  $m - k + N$  embedded copies of  $K$  on  $T$ . In particular,  $N \leq k$  follows by the maximality of  $m$ . We conclude

$$A(D + R(K)B^2) \geq N \cdot \Delta(K),$$

which in turn yields the upper bound in (i).

Turning to (ii), we let  $\lambda < 1$  satisfy  $\lambda \cdot A(C)/\Delta(K) > \lceil A(C)/\Delta(K) \rceil - 1$ . It follows by the definition of  $\Delta(K)$  that there exist a torus  $T = \mathbb{R}^2/\Delta$  and  $m$  nonoverlapping embedded copies  $K_1, \dots, K_m$  of  $K$  on  $T$  satisfying  $A(T) < \lambda^{-1}m\Delta(K)$ , and  $D$  embeds isometrically into  $T$ . We define  $C = (1 - R(K)/r(D))D$ ; hence, (1.1) yields that some translate  $C + x$  contains at least  $m \cdot A(C)/A(T)$  points out of the circumcentres of  $K_1, \dots, K_m$ . We may assume that these points are the circumcentres of  $K_1, \dots, K_l$ ; therefore,  $l \geq \lambda \cdot A(C)/\Delta(K)$  and  $K_1, \dots, K_l$  are contained in  $D + x$ . Thus  $N \geq l \geq A(C)/\Delta(K)$  by the definition of  $\lambda$ , completing the proof of Lemma 1.1.2. □

*Proof of Theorem 1.1.1.* We present the argument only for packings because just the obvious changes are needed for the case of coverings. In the following the implied constant in  $O(\cdot)$  always depends only on  $K$ .

Theorem 1.1.1 for packings follows from the following statement: If  $\varepsilon > 0$  is small, and  $n > 1/\varepsilon^5$  then

$$A(D_n) = (1 + O(\varepsilon)) \cdot n\Delta(K). \tag{1.2}$$

Dense clusters show (see Lemma 1.1.2) that

$$A(D_n) \leq (1 + O(\varepsilon)) \cdot n\Delta(K).$$

Therefore, it is sufficient to verify that

$$A(D_n) \geq (1 - O(\varepsilon)) \cdot n\Delta(K). \tag{1.3}$$

If  $r(D_n) > 1/\varepsilon$  then (1.3) follows from Lemma 1.1.2. Thus we assume that  $r(D_n) \leq 1/\varepsilon$ , a case that requires a more involved argument. We actually prove that there exists a rectangle  $R$  that contains certain  $N$  congruent copies

of  $K$ , where

$$\frac{A(R)}{N} \leq (1 + O(\varepsilon)) \cdot \frac{A(D_n)}{n}. \tag{1.4}$$

Since the minimal width  $w$  of  $D_n$  is at most  $3/\varepsilon$  according to the Steinhagen inequality (Theorem A.8.2), there exists a rectangle  $\tilde{R}$  such that its sides touch  $D_n$ , and two parallel sides of  $\tilde{R}$  are of length  $w$ . We say that these sides are vertical; hence,  $D_n$  has a vertical section of length  $w$ . Writing  $l$  to denote the length of the horizontal sides, we have  $A(D_n) \geq wl/2$ . For  $k = \lceil 1/\varepsilon \rceil$ , we decompose  $\tilde{R}$  into  $k^3$  congruent rectangles  $R_1, \dots, R_{k^3}$  in this order, where the vertical sides of  $R_i$  are of length  $w$  and the horizontal sides are of length  $l/k^3$ .

Out of the circumcentres of the  $n$  nonoverlapping congruent copies of  $K$  that lie in  $D_n$ , let  $n_i$  be contained in  $R_i$ . Now the total area of  $R_1, \dots, R_{k^2+1}$  and of  $R_{k^3-k^2}, \dots, R_{k^3}$  is

$$\left(1 + \frac{1}{k^2}\right) \frac{2wl}{k} \leq (1 + O(\varepsilon)) 4\varepsilon A(D_n) \leq (1 + O(\varepsilon)) 4\Delta(K) \cdot \varepsilon n,$$

and hence  $\sum_{i=k^2+1}^{k^3-k^2} n_i \geq (1 - O(\varepsilon))n$ . In particular, there exists some index  $j$  such that  $k^2 + 1 \leq j \leq k^3 - k^2$  and

$$\frac{A(R_j \cap D_n)}{n_j} \leq (1 + O(\varepsilon)) \cdot \frac{A(D_n)}{n}. \tag{1.5}$$

Let  $R'$  be the rectangle whose sides are vertical and horizontal, with each touching  $R_j \cap D_n$ . We write  $a$  to denote the common length of the vertical sides of  $R'$ , which readily satisfies  $a \geq 2r(K)$ . Since  $w/k^2 < 4\varepsilon$ , we deduce that  $R_j \cap D_n$  contains a rectangle whose horizontal side is of length  $l/k^3$ , and the vertical side is of length  $a - 8\varepsilon$ . In particular,  $A(R')$  is at most  $(1 + O(\varepsilon))A(R_j \cap D_n)$ . Finally, the rectangle  $R$  whose horizontal sides are of length  $l/k^3 + 2R(K)$  and vertical sides are of length  $a$  contains  $N = n_j$  nonoverlapping congruent copies of  $K$ . Now  $3l/\varepsilon \geq n A(K)$  yields  $l/k^3 \geq 1/[4A(K)\varepsilon]$ . Thus we conclude (1.4) by (1.5).

Since the arrangement in  $R$  induces a periodic packing of  $K$ , (1.4) readily yields (1.3) and hence Theorem 1.1.1 as well.  $\square$

**Remark 1.1.3.** Given a strictly convex domain  $K$ , if  $D_n$  is a convex domain with minimal area that contains  $n$  nonoverlapping congruent copies of  $K$  then  $r(D_n)$  tends to infinity.

We sketch the argument for Remark 1.1.3: We suppose indirectly that there exists a subsequence of  $\{r(D_n)\}$  that is bounded by some  $\omega > 0$ . For any  $\varepsilon > 0$ , the proof of Theorem 1.1.1 yields a parallel strip  $\Sigma_\varepsilon$  and a packing

$\Xi$  of congruent copies of  $K$  inside  $\Sigma_\varepsilon$  such that the packing is periodic with respect to a vector parallel to  $\Sigma_\varepsilon$ , the width of  $\Sigma_\varepsilon$  is at most  $3\omega$ , and the density of the packing  $\Xi$  inside  $\Sigma_\varepsilon$  is at least  $(1 - \varepsilon)\delta(K)$ . We reflect this arrangement through one of the lines bounding  $\Sigma_\varepsilon$  and write  $\Xi'$  to denote the image of  $\Xi$ . Because  $K$  is strictly convex, there exist positive  $v_1$  and  $v_2$  depending only on  $K$  with the following property: Translating the packing  $\Xi'$  first parallel to  $\Sigma_\varepsilon$  by a vector of length  $v_1$ , then towards  $\Sigma_\varepsilon$  orthogonally by a vector of length  $v_2$ , we obtain an arrangement  $\Xi''$  such that the union of  $\Xi$  and  $\Xi''$  forms a packing. If  $\varepsilon < v_2/(2\omega)$  then the union of  $\Xi$  and  $\Xi''$  determines a periodic packing in the plane whose density is larger than  $\delta(K)$ . This contradiction verifies that  $r(D_n)$  tends to infinity.

### Open Problems.

- (i) Let  $K$  be a convex domain that is not a parallelogram. We write  $D_n$  ( $\tilde{D}_n$ ) to denote a convex domain with minimal (maximal) area that contains  $n$  nonoverlapping congruent copies of  $K$  (i.e., is covered by  $n$  congruent copies of  $K$ ). Is

$$r(D_n), r(\tilde{D}_n) > c\sqrt{n}$$

for a suitable positive constant  $c$  depending on  $K$ ? If the answer is yes then the ratio  $R(D_n)/r(D_n)$  stays bounded as  $n$  tends to infinity, and a similar property holds for  $\tilde{D}_n$ .

For packings, various partial results support an affirmative answer: The statement holds if  $K$  is centrally symmetric (see Corollary 1.4.3) or the packing is translative (see Theorem 2.4.1). Strengthening the method of Remark 1.1.3 yields that  $r(D_n) > c\sqrt[3]{n}$  holds if  $K$  is any strictly convex domain. For coverings, the statement holds if  $K$  is a fat ellipse (see Theorem 1.7.1) or if  $K$  is centrally symmetric and only translative coverings are allowed (see Corollary 2.8.2).

- (ii) Is  $\vartheta(K) \leq 2\pi/\sqrt{27} = 1.2091\dots$  for any convex domain  $K$ ; namely, is the covering density maximal for circular discs (see Theorem 1.7.1)?  
 D. Ismailescu [Ism1998] proved  $\vartheta(K) \leq 1.2281\dots$  for any convex domain  $K$ . However,  $\vartheta(K) \leq 2\pi/\sqrt{27}$  if  $K$  is centrally symmetric (see L. Fejes Tóth [FTL1972]).
- (iii) Does it hold for any convex domain that there exist a periodic packing whose density is the packing density and a periodic covering whose density is the covering density? It is known that there exist no optimal lattice arrangement for the typical convex domain (see G. Fejes Tóth and T. Zamfirescu [FTZ1994] and G. Fejes Tóth [FTG1995a]).

**Comments.** The packing and covering densities were originally introduced in the framework of infinite packing and covering of the space (see G. Fejes Tóth and W. Kuperberg [FTK1993]). Readily,  $\Theta(K) \leq A(K) \leq \Delta(K)$ . W.M. Schmidt [Sch1961] proved that  $\Theta(K) = A(K)$  or  $\Delta(K) = A(K)$  if and only if some congruent copies of  $K$  tile the plane (see also Lemma 7.2.5).

According to the hexagon bound of L. Fejes Tóth [FTL1950] (see Theorem 1.3.1),  $\Delta(K)$  is at least the minimal area of circumscribed hexagons for any convex domain  $K$ , where equality holds if  $K$  is centrally symmetric. Concerning absolute lower bounds on the packing density, G. Kuperberg and W. Kuperberg [KuK1990] verified that  $\delta(K) > \sqrt{3}/2 = 0.8660\dots$  holds for any convex domain  $K$ . In addition a beautiful little theorem of W. Kuperberg [Kup1987] states that  $\delta(K)/\vartheta(K) \geq 3/4$ , where equality holds for circular discs. It is probably surprising but the packing density,  $\pi/\sqrt{12} = 0.9068\dots$  of the unit disc is not minimal among centrally symmetric convex domains, which is shown say by the regular octagon. By rounding off the corners of the regular octagon, K. Reinhardt [Rei1934] and K. Mahler [Mah1947] proposed a possible minimal shape whose density is  $0.9024\dots$ . P. Tammela [Tam1970] proved that  $\delta(K) > 0.8926$  for any centrally symmetric convex domain  $K$ .

Concerning coverings, D. Ismailescu [Ism1998] proved  $\vartheta(K) \leq 1.2281\dots$  for any convex domain  $K$ . For very long the only convex domains with known covering densities were the tiles (when the covering density is one), and circular discs (when the covering density is  $2\pi/\sqrt{27}$  according to R. Kershner [Ker1939]; see also Corollary 5.1.2). Recently A. Heppes [Hep2003] showed that the covering density of any “fat ellipse” (when the ratio of the smaller axis to the greater axis is at least 0.86) is  $2\pi/\sqrt{27}$  (see also Theorem 1.7.1). A substantial improvement is due to G. Fejes Tóth [FTG?b]: On the one hand [FTG?b] generalized A. Heppes’ theorem to ellipses when the ratio of the smaller axis to the greater axis is at least 0.741. On the other hand if  $K$  is a centrally symmetric convex domain and  $r(K)/R(K) \geq 0.933$  then [FTG?b] proves that  $\Theta(K)$  is the maximal area of polygons with at most six sides inscribed into  $K$ . Readily if  $K$  is either type of the convex domains considered in [FTG?b], and  $C \subset K$  is a convex domain that contains a centrally symmetric hexagon of area  $\Theta(K)$  then  $\Theta(C) = \Theta(K)$ .

## 1.2. The Hexagon Bound for Packings Inside an Octagon

Given a convex domain  $K$ , we write  $H(K)$  to denote a circumscribed convex polygon with at most six sides of minimal area. The aim of this section is to verify the *hexagon bound* for packings of congruent copies of  $K$

1.2. The Hexagon Bound for Packings Inside an Octagon 9

inside a hexagon; namely, the density is at most  $A(K)/A(H(K))$ . Later we will prove the hexagon bound with respect to any convex container (see Theorem 1.4.1).

**Theorem 1.2.1.** *If a polygon  $D$  of at most eight sides contains  $n \geq 2$  congruent copies of a given convex domain  $K$  then*

$$A(D) \geq n \cdot A(H(K)).$$

The main idea of the proof Theorem 1.2.1 is to define a cell decomposition of  $D$  into convex cells in a way such that each cell contains exactly one of the congruent copies of  $K$ ; hence, the average number of sides of the cells is at most six according to the Euler formula. Then we verify that the minimal areas of circumscribed  $k$ -gons are convex functions of  $k$  (see Corollary 1.2.4), and we deduce that the average area of a cell is at least  $A(H(K))$ . Unfortunately, we cannot proceed exactly like this because no suitable cell decomposition of  $D$  may exist. In spite of this we can still save the essential properties of a cell decomposition (see Lemma 1.2.2) and verify the hexagon bound. Lemma 1.2.2 is presented in a rather general setting because of later applications.

**Lemma 1.2.2.** *Let  $D$  be a convex domain that contains the nonoverlapping convex domains  $K_1, \dots, K_n$ ,  $n \geq 2$ . Then there exist nonoverlapping convex domains  $\Pi_1, \dots, \Pi_n \subset D$  satisfying the following properties:*

- (i)  $K_i \subset \Pi_i$ .
- (ii)  $\Pi_1, \dots, \Pi_n$  cover  $\partial D$ .
- (iii)  $\Pi_i$  is bounded by  $k_i \geq 2$  convex arcs that we call edges. The edges intersecting  $\text{int } D$  are segments, and the rest of the edges are the maximal convex arcs of  $\partial D \cap \Pi_i$ .
- (iv) The number  $b$  of edges contained in  $\partial D$  satisfy

$$\sum_{i=1}^n (6 - k_i) \geq b + 6.$$

*In addition, if  $D$  is a polygon of at most eight sides and  $k_i^*$  denotes the number of sides of  $\Pi_i$  then  $\sum_{i=1}^n (6 - k_i^*) \geq 0$ .*

*Proof.* Let  $\Pi_1, \dots, \Pi_n$  be nonoverlapping convex domains such that  $K_i \subset \Pi_i \subset D$  and the total area covered by the convex domains  $\Pi_1, \dots, \Pi_n$  is maximal under these conditions. Since two nonoverlapping convex sets can be separated by a line, each  $\Pi_i$  is the intersection of a polygon  $P_i$  and  $D$ .



Now  $\text{int } P_i \cap \partial D$  consists of finitely many convex arcs whose closures we call edges of  $\Pi_i$ . The rest of the edges of  $\Pi_i$  are the segments of the form  $s \cap D$ , where  $s$  is a side of  $P_i$  that intersects  $\text{int } D$ , and the vertices of  $\Pi_i$  are the endpoints of the edges.

It may happen that  $\Pi_1, \dots, \Pi_n$  do not cover  $D$ , and we call the closure of a connected component of  $\text{int } D \setminus \cup_{i=1}^n \Pi_i$  a hole. Let  $Q$  be a hole. Then there exists an edge  $e_1$  of some  $\Pi_{i_1}$  such that  $e_1$  intersects  $\text{int } D$ , and  $e_1 \cap \partial Q$  contains a segment  $s_1$ , where we assume that  $s_1$  is a maximal segment in  $e_1 \cap \partial Q$ . Since  $\Pi_{i_1}$  cannot be extended because of the maximality of  $\sum A(\Pi_j)$ , one endpoint  $v_2$  of  $s_1$  is contained in the relative interior of  $e_1$ ; hence,  $v_2 \in \text{int } D$ . Therefore,  $v_2$  is the endpoint of an edge  $e_2$  of some  $\Pi_{i_2}$  such that  $e_2 \cap \partial Q$  contains a maximal segment  $s_2$ . Continuing this way we obtain that  $\partial Q$  is the union of segments  $s_1, \dots, s_k$  with the following properties (where  $s_0 = s_k$ ):  $s_j$  is contained in an edge  $e_j$  of some  $\Pi_{i_j}$ , and  $s_j \cap s_{j-1}$  is a common endpoint  $v_j \in \text{int } D$  that is an endpoint of  $e_j$  and not of  $e_{j-1}$  for any  $j = 1, \dots, k$ ; moreover, different  $s_i$  and  $s_j$  do not intersect otherwise. We deduce that  $Q$  is a convex polygon and  $Q \subset \text{int } D$ .

Now (ii) readily follows; namely,  $\Pi_1, \dots, \Pi_n$  cover  $\partial D$ . Next, we construct a related cell decomposition  $\Sigma$  of  $D$  by cells  $\tilde{\Pi}_1, \dots, \tilde{\Pi}_n$ . If there exists no hole then  $\tilde{\Pi}_i = \Pi_i$ . Otherwise, let  $\{Q_1, \dots, Q_m\}$  be the set of holes, let  $q_j \in \text{int } Q_j$ , and we define  $\tilde{\Pi}_i$  to be the union of  $\Pi_i$  and all triangles of the form  $\text{conv}\{q_j, s\}$  such that  $s$  is a side of  $Q_j$  and  $s \subset \Pi_i$ . In particular, the number of edges of  $\Sigma$  contained in  $\tilde{\Pi}_i$  is at least  $k_i$ ; hence  $\sum (6 - k_i) \geq b + 6$  is a consequence of Lemma A.5.9. If, in addition,  $D$  is a polygon of at most eight sides then  $\sum k_i^* \leq 8 + \sum k_i$ ; thus  $b \geq 2$  completes the proof of Lemma 1.2.2.  $\square$

Given the convex domain  $K$ , let  $t_K(m)$  denote the minimal area of a circumscribed polygon of at most  $m$  sides for any  $m \geq 3$ . Next, we show that  $t_K(m)$  is a convex function of  $m$ , more precisely, that  $t_K(m)$  is even strictly convex if  $K$  is strictly convex.

**Lemma 1.2.3.** *If  $K$  is a strictly convex domain and  $m \geq 4$  then*

$$t_K(m - 1) + t_K(m + 1) > 2t_K(m).$$

*Proof.* For any  $m \geq 3$ , we choose a circumscribed polygon  $\Pi_m$  of minimal area among the circumscribed polygons of at most  $m$  sides. Since  $K$  is strictly convex,  $\Pi_m$  is actually an  $m$ -gon, and each side of  $\Pi_m$  touches  $K$  at the midpoint of the side.

1.2. The Hexagon Bound for Packings Inside an Octagon 11

Let  $m \geq 4$ , and let  $3 \leq k \leq l$  satisfy that  $A(\Pi_k) + A(\Pi_l)$  is minimal under the condition  $k + l = 2m$ . We suppose that  $k < m$  and seek a contradiction. The idea is to decrease the total area of  $\Pi_k$  and  $\Pi_l$  by interchanging certain sides. We write  $p_1, \dots, p_k$  and  $q_1, \dots, q_l$  to denote the midpoints of the sides of  $\Pi_k$  and  $\Pi_l$ , respectively, according to the clockwise orientation, and we write  $e_i$  and  $f_j$  to denote the side of  $\Pi_k$  and  $\Pi_l$  containing  $p_i$  and  $q_j$ , respectively. For  $p, q \in \partial K$ , let  $[p, q)$  denote the semi open arc of  $\partial K$ , which starts at  $p$  and terminates at  $q$  according to the clockwise orientation, and the arc contains  $p$  and does not contain  $q$ . The  $k$  semi open convex arcs  $[p_{i-1}, p_i)$  on  $\partial K$  (with  $p_0 = p_k$ ) contain the  $l \geq k + 2$  midpoints for  $\Pi_l$ , and hence either there exists  $[p_{i-1}, p_i)$ , which contains say  $q_1, q_2, q_3$ , or there exist two semi open arcs of the form  $[p_{i-1}, p_i)$  such that each contains two midpoints from  $\Pi_l$ . In the first case, let  $\Pi'_{k+1}$  be obtained from  $\Pi_k$  by cutting off the vertex  $e_{i-1} \cap e_i$  by  $\text{aff } f_2$ , and let  $\Pi''_{l-1}$  be obtained from  $\Pi_l$  by removing the side  $f_2$ , and hence  $\text{aff } f_1 \cap \text{aff } f_3$  is the new vertex of  $\Pi''_{l-1}$ . Then  $\Pi'_{k+1}$  and  $\Pi''_{l-1}$  have  $k + 1$  and  $l - 1$  sides, respectively, and  $\Pi''_{l-1} \setminus \Pi_l$  is strictly contained in  $\Pi'_{k+1} \setminus \Pi_k$ . Therefore,

$$A(\Pi'_{k+1}) + A(\Pi''_{l-1}) < A(\Pi_k) + A(\Pi_l).$$

This is absurd, and hence we may assume that  $q_1, q_2 \in [p_1, p_2)$  and  $q_{j-1}, q_j \in [p_{i-1}, p_i)$  for  $i \neq 2$ . In this case let  $\Pi'_{k'}$  be the circumscribed  $k'$ -gon defined by affine hulls of

$$e_1, f_2, \dots, f_{j-1}, e_i, \dots, e_k.$$

In addition, let  $\Pi''_{l'}$  be the circumscribed  $l'$ -gon defined by affine hulls of

$$f_1, e_2, \dots, e_{i-1}, f_j, \dots, f_l;$$

thus  $k' + l' = 2m$ . When constructing  $\Pi'_{k'}$  and  $\Pi''_{l'}$ , we remove the part of  $\Pi_k$  at the corner enclosed by  $e_1$  and  $e_2$  and cut off by  $\partial \Pi_l$ , and we add two nonoverlapping domains contained in this part (where one of the domains degenerates if  $q_1 = p_1$ ). Because the situation is analogous at the corner of  $\Pi_k$  enclosed by  $e_{i-1}$  and  $e_i$ , we deduce that

$$A(\Pi'_{k'}) + A(\Pi''_{l'}) \leq A(\Pi_k) + A(\Pi_l).$$

The polygons were constructed in a way that  $f_{j-2} \cap f_{j-1}$  is a common vertex for  $\Pi'_{k'}$  and  $\Pi_l$ , whereas  $f_j \cap f_{j-1}$  is a vertex for  $\Pi_l$  but not for  $\Pi'_{k'}$ , and hence  $q_{j-1}$  is not the midpoint of the side of  $\Pi'_{k'}$  containing it. Therefore, there exists a circumscribed  $k'$ -gon whose area is less than  $A(\Pi'_{k'})$ , which contradicts the minimality of  $A(\Pi_k) + A(\Pi_l)$ .  $\square$