

Ravi P. Agarwal
National University of Singapore

Maria Meehan
Dublin City University

Donal O'Regan
National University of Ireland, Galway

Fixed Point Theory and Applications

 **CAMBRIDGE**
UNIVERSITY PRESS

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS
The Edinburgh Building, Cambridge CB2 2RU, UK
40 West 20th Street, New York, NY 10011-4211, USA
10 Stamford Road, Oakleigh, VIC 3166, Australia
Ruiz de Alarcón 13, 28014, Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

<http://www.cambridge.org>

© Cambridge University Press 2001

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2001

Printed in the United Kingdom at the University Press, Cambridge

Typeface Computer Modern 10/13pt. *System* L^AT_EX 2_ε [DBD]

A catalogue record of this book is available from the British Library

ISBN 0 521 80250 4 hardback

Contents

<i>Preface</i>	<i>page</i>	vii
1	Contractions	1
2	Nonexpansive Maps	12
3	Continuation Methods for Contractive and Nonexpansive Mappings	19
4	The Theorems of Brouwer, Schauder and Mönch	28
5	Nonlinear Alternatives of Leray–Schauder Type	48
6	Continuation Principles for Condensing Maps	65
7	Fixed Point Theorems in Conical Shells	78
8	Fixed Point Theory in Hausdorff Locally Convex Linear Topological Spaces	94
9	Contractive and Nonexpansive Multivalued Maps	112
10	Multivalued Maps with Continuous Selections	120
11	Multivalued Maps with Closed Graph	130
12	Degree Theory	142
<i>Bibliography</i>		159
<i>Index</i>		169

1

Contractions

Let (X, d) be a metric space. A map $F : X \rightarrow X$ is said to be *Lipschitzian* if there exists a constant $\alpha \geq 0$ with

$$(1.1) \quad d(F(x), F(y)) \leq \alpha d(x, y) \text{ for all } x, y \in X.$$

Notice that a Lipschitzian map is necessarily continuous. The smallest α for which (1.1) holds is said to be the *Lipschitz constant* for F and is denoted by L . If $L < 1$ we say that F is a *contraction*, whereas if $L = 1$, we say that F is *nonexpansive*.

For notational purposes we define $F^n(x)$, $x \in X$ and $n \in \{0, 1, 2, \dots\}$, inductively by $F^0(x) = x$ and $F^{n+1}(x) = F(F^n(x))$.

The first result in this chapter is known as Banach's contraction principle.

Theorem 1.1 *Let (X, d) be a complete metric space and let $F : X \rightarrow X$ be a contraction with Lipschitzian constant L . Then F has a unique fixed point $u \in X$. Furthermore, for any $x \in X$ we have*

$$\lim_{n \rightarrow \infty} F^n(x) = u$$

with

$$d(F^n(x), u) \leq \frac{L^n}{1 - L} d(x, F(x)).$$

Proof We first show uniqueness. Suppose there exist $x, y \in X$ with $x = F(x)$ and $y = F(y)$. Then

$$d(x, y) = d(F(x), F(y)) \leq L d(x, y),$$

therefore $d(x, y) = 0$.

To show existence select $x \in X$. We first show that $\{F^n(x)\}$ is a Cauchy sequence. Notice for $n \in \{0, 1, \dots\}$ that

$$d(F^n(x), F^{n+1}(x)) \leq L d(F^{n-1}(x), F^n(x)) \leq \dots \leq L^n d(x, F(x)).$$

Thus for $m > n$ where $n \in \{0, 1, \dots\}$,

$$\begin{aligned} d(F^n(x), F^m(x)) &\leq d(F^n(x), F^{n+1}(x)) + d(F^{n+1}(x), F^{n+2}(x)) \\ &\quad + \dots + d(F^{m-1}(x), F^m(x)) \\ &\leq L^n d(x, F(x)) + \dots + L^{m-1} d(x, F(x)) \\ &\leq L^n d(x, F(x)) [1 + L + L^2 + \dots] \\ &= \frac{L^n}{1 - L} d(x, F(x)). \end{aligned}$$

That is for $m > n$, $n \in \{0, 1, \dots\}$,

$$(1.2) \quad d(F^n(x), F^m(x)) \leq \frac{L^n}{1 - L} d(x, F(x)).$$

This shows that $\{F^n(x)\}$ is a Cauchy sequence and since X is complete there exists $u \in X$ with $\lim_{n \rightarrow \infty} F^n(x) = u$. Moreover the continuity of F yields

$$u = \lim_{n \rightarrow \infty} F^{n+1}(x) = \lim_{n \rightarrow \infty} F(F^n(x)) = F(u),$$

therefore u is a fixed point of F . Finally letting $m \rightarrow \infty$ in (1.2) yields

$$d(F^n(x), u) \leq \frac{L^n}{1 - L} d(x, F(x)). \quad \square$$

Remark 1.1 Theorem 1.1 requires that $L < 1$. If $L = 1$ then F need not have a fixed point as the example $F(x) = x + 1$ for $x \in \mathbf{R}$ shows. We will discuss the case when $L = 1$ in more detail in Chapter 2.

Another natural attempt to extend Theorem 1.1 would be to suppose that $d(F(x), F(y)) < d(x, y)$ for $x, y \in X$ with $x \neq y$. Again F need not have a fixed point as the example $F(x) = \ln(1 + e^x)$ for $x \in \mathbf{R}$ shows. However there is a positive result along these lines in the following theorem of Edelstein.

Theorem 1.2 *Let (X, d) be a compact metric space with $F : X \rightarrow X$ satisfying*

$$d(F(x), F(y)) < d(x, y) \text{ for } x, y \in X \text{ and } x \neq y.$$

Then F has a unique fixed point in X .

Proof The uniqueness part is easy. To show existence, notice the map $x \mapsto d(x, F(x))$ attains its minimum, say at $x_0 \in X$. We have $x_0 = F(x_0)$ since otherwise

$$d(F(F(x_0)), F(x_0)) < d(F(x_0), x_0)$$

– a contradiction. \square

We next present a local version of Banach's contraction principle. This result will be needed in Chapter 3.

Theorem 1.3 *Let (X, d) be a complete metric space and let*

$$B(x_0, r) = \{x \in X : d(x, x_0) < r\}, \text{ where } x_0 \in X \text{ and } r > 0.$$

Suppose $F : B(x_0, r) \rightarrow X$ is a contraction (that is, $d(F(x), F(y)) \leq Ld(x, y)$ for all $x, y \in B(x_0, r)$ with $0 \leq L < 1$) with

$$d(F(x_0), x_0) < (1 - L)r.$$

Then F has a unique fixed point in $B(x_0, r)$.

Proof There exists r_0 with $0 \leq r_0 < r$ with $d(F(x_0), x_0) \leq (1 - L)r_0$. We will show that $F : \overline{B(x_0, r_0)} \rightarrow \overline{B(x_0, r_0)}$. To see this note that if $x \in \overline{B(x_0, r_0)}$ then

$$\begin{aligned} d(F(x), x_0) &\leq d(F(x), F(x_0)) + d(F(x_0), x_0) \\ &\leq Ld(x, x_0) + (1 - L)r_0 \leq r_0. \end{aligned}$$

We can now apply Theorem 1.1 to deduce that F has a unique fixed point in $\overline{B(x_0, r_0)} \subset B(x_0, r)$. Again it is easy to see that F has only one fixed point in $B(x_0, r)$. \square

Next we examine briefly the behaviour of a contractive map defined on $\overline{B}_r = \overline{B(0, r)}$ (the closed ball of radius r with centre 0) with values in a Banach space E . More general results will be presented in Chapter 3.

Theorem 1.4 *Let \overline{B}_r be the closed ball of radius $r > 0$, centred at zero, in a Banach space E with $F : \overline{B}_r \rightarrow E$ a contraction and $F(\partial\overline{B}_r) \subseteq \overline{B}_r$. Then F has a unique fixed point in \overline{B}_r .*

Proof Consider

$$G(x) = \frac{x + F(x)}{2}.$$

We first show that $G : \overline{B}_r \rightarrow \overline{B}_r$. To see this let

$$x^* = r \frac{x}{\|x\|} \text{ where } x \in \overline{B}_r \text{ and } x \neq 0.$$

Now if $x \in \overline{B}_r$ and $x \neq 0$,

$$\|F(x) - F(x^*)\| \leq L \|x - x^*\| = L(r - \|x\|),$$

since $x - x^* = \frac{x}{\|x\|}(\|x\| - r)$, and as a result

$$\begin{aligned} \|F(x)\| &\leq \|F(x^*)\| + \|F(x) - F(x^*)\| \\ &\leq r + L(r - \|x\|) \leq 2r - \|x\|. \end{aligned}$$

Then for $x \in \overline{B}_r$ and $x \neq 0$

$$\|G(x)\| = \left\| \frac{x + F(x)}{2} \right\| \leq \frac{\|x\| + \|F(x)\|}{2} \leq r.$$

In fact by continuity we also have

$$\|G(0)\| \leq r,$$

and consequently $G : \overline{B}_r \rightarrow \overline{B}_r$. Moreover $G : \overline{B}_r \rightarrow \overline{B}_r$ is a contraction since

$$\|G(x) - G(y)\| \leq \frac{\|x - y\| + L\|x - y\|}{2} = \frac{[1 + L]}{2} \|x - y\|.$$

Theorem 1.1 implies that G has a unique fixed point $u \in \overline{B}_r$. Of course if $u = G(u)$ then $u = F(u)$. \square

Over the last fifty years or so, many authors have given generalisations of Banach's contraction principle. Here for completeness we give one such result. Its proof relies on the following technical result.

Theorem 1.5 *Let (X, d) be a complete metric space and $F : X \rightarrow X$ a map (not necessarily continuous). Suppose the following condition holds:*

$$(1.3) \quad \left\{ \begin{array}{l} \text{for each } \epsilon > 0 \text{ there is a } \delta(\epsilon) > 0 \text{ such that if} \\ d(x, F(x)) < \delta(\epsilon), \text{ then } F(B(x, \epsilon)) \subseteq B(x, \epsilon); \\ \text{here } B(x, \epsilon) = \{y \in X : d(x, y) < \epsilon\}. \end{array} \right.$$

If for some $u \in X$ we have

$$\lim_{n \rightarrow \infty} d(F^n(u), F^{n+1}(u)) = 0,$$

then the sequence $\{F^n(u)\}$ converges to a fixed point of F .

Proof Let u be as described above and let $u_n = F^n(u)$. We claim that $\{u_n\}$ is a Cauchy sequence.

Let $\epsilon > 0$ be given. Choose $\delta(\epsilon)$ as in (1.3). We can choose N large enough so that

$$d(u_n, u_{n+1}) < \delta(\epsilon) \text{ for all } n \geq N.$$

Now since $d(u_N, F(u_N)) < \delta(\epsilon)$, then (1.3) guarantees that

$$F(B(u_N, \epsilon)) \subseteq B(u_N, \epsilon),$$

and so $F(u_N) = u_{N+1} \in B(u_N, \epsilon)$. Now by induction

$$F^k(u_N) = u_{N+k} \in B(u_N, \epsilon) \text{ for all } k \in \{0, 1, 2, \dots\}.$$

Thus

$$d(u_k, u_l) \leq d(u_k, u_N) + d(u_N, u_l) < 2\epsilon \text{ for all } k, l \geq N,$$

and therefore $\{u_n\}$ is a Cauchy sequence. In addition there exists $y \in X$ with $\lim_{n \rightarrow \infty} u_n = y$.

We now claim that y is a fixed point of F . Suppose it is not. Then

$$d(y, F(y)) = \gamma > 0.$$

We can now choose (and fix) a $u_n \in B(y, \gamma/3)$ with

$$d(u_n, u_{n+1}) < \delta(\gamma/3).$$

Now (1.3) guarantees that

$$F(B(u_n, \gamma/3)) \subseteq B(u_n, \gamma/3),$$

and consequently $F(y) \in B(u_n, \gamma/3)$. This is a contradiction since

$$d(F(y), u_n) \geq d(F(y), y) - d(u_n, y) > \gamma - \frac{\gamma}{3} = \frac{2\gamma}{3}.$$

Thus $d(y, F(y)) = 0$. □

Theorem 1.6 *Let (X, d) be a complete metric space and let*

$$d(F(x), F(y)) \leq \phi(d(x, y)) \text{ for all } x, y \in X;$$

here $\phi : [0, \infty) \rightarrow [0, \infty)$ is any monotonic, nondecreasing (not necessarily continuous) function with $\lim_{n \rightarrow \infty} \phi^n(t) = 0$ for any fixed $t > 0$. Then F has a unique fixed point $u \in X$ with

$$\lim_{n \rightarrow \infty} F^n(x) = u \text{ for each } x \in X.$$

Proof Suppose $t \leq \phi(t)$ for some $t > 0$. Then $\phi(t) \leq \phi(\phi(t))$ and therefore $t \leq \phi^2(t)$. By induction, $t \leq \phi^n(t)$ for $n \in \{1, 2, \dots\}$. This is a contradiction. Thus $\phi(t) < t$ for each $t > 0$.

In addition,

$$d(F^n(x), F^{n+1}(x)) \leq \phi^n(d(x, F(x))) \text{ for } x \in X,$$

and therefore

$$\lim_{n \rightarrow \infty} d(F^n(x), F^{n+1}(x)) = 0 \text{ for each } x \in X.$$

Let $\epsilon > 0$ and choose $\delta(\epsilon) = \epsilon - \phi(\epsilon)$. If $d(x, F(x)) < \delta(\epsilon)$, then for any $z \in B(x, \epsilon) = \{y \in X : d(x, y) < \epsilon\}$ we have

$$\begin{aligned} d(F(z), x) &\leq d(F(z), F(x)) + d(F(x), x) \leq \phi(d(z, x)) + d(F(x), x) \\ &< \phi(d(z, x)) + \delta(\epsilon) \leq \phi(\epsilon) + (\epsilon - \phi(\epsilon)) = \epsilon, \end{aligned}$$

and therefore $F(z) \in B(x, \epsilon)$. Theorem 1.5 guarantees that F has a fixed point u with $\lim_{n \rightarrow \infty} F^n(x) = u$ for each $x \in X$. Finally it is easy to see that F has only one fixed point in X . \square

Remark 1.2 Note that Theorem 1.1 follows as a special case of Theorem 1.6 if we choose $\phi(t) = Lt$ with $0 \leq L < 1$.

It is natural to begin our applications of fixed point methods with existence and uniqueness of solutions of certain first order initial value problems. In particular we seek solutions to

$$(1.4) \quad \begin{cases} y'(t) = f(t, y(t)), \\ y(0) = y_0, \end{cases}$$

where $f : I \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ and $I = [0, b]$. Notice that (1.4) is a system of first order equations because f takes values in \mathbf{R}^n .

We begin our analysis of (1.4) by assuming that $f : I \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ is continuous. Then, evidently, $y \in C^1(I)$ (the Banach space of functions u whose first derivative is continuous on I and equipped with the norm $\|u\|_1 = \max\{\sup_{t \in I} |u(t)|, \sup_{t \in I} |u'(t)|\}$) solves (1.4) if and only if $y \in C(I)$ (the Banach space of functions u , continuous on I and equipped with the norm $\|u\|_0 = \sup_{t \in I} |u(t)|$) solves

$$(1.5) \quad y(t) = y_0 + \int_0^t f(s, y(s)) ds.$$

Define an integral operator $T : C(I) \rightarrow C(I)$ by

$$Ty(t) = y_0 + \int_0^t f(s, y(s)) ds.$$

Then the equivalence above is expressed briefly by

$$y \text{ solves (1.4) if and only if } y = Ty, \quad T : C(I) \rightarrow C(I).$$

In other words, classical solutions to (1.4) are fixed points of the integral operator T . We now present a result known as the Picard–Lindelöf theorem.

Theorem 1.7 *Let $f : I \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ be continuous and Lipschitz in y ; that is, there exists $\alpha \geq 0$ such that*

$$|f(t, y) - f(t, z)| \leq \alpha |y - z| \text{ for all } y, z \in \mathbf{R}^n.$$

Then there exists a unique $y \in C^1(I)$ that solves (1.4).

Proof We will apply Theorem 1.1 to show that T has a unique fixed point. At first glance it seems natural to use the maximum norm on $C(I)$, but this choice would lead us only to a local solution defined on a subinterval of I . The trick is to use the weighted maximum norm

$$\|y\|_\alpha = |e^{-\alpha t} y(t)|_0$$

on $C(I)$. Observe that $C(I)$ is a Banach space with this norm since it is equivalent to the maximum norm, that is,

$$e^{-\alpha b} |y|_0 \leq \|y\|_\alpha \leq |y|_0.$$

We now show that T is a contraction on $(C(I), \|\cdot\|_\alpha)$. To see this let $y, z \in C(I)$ and notice

$$Ty(t) - Tz(t) = \int_0^t [f(s, y(s)) - f(s, z(s))] ds \text{ for } t \in I.$$

Thus for $t \in I$,

$$\begin{aligned} e^{-\alpha t} |(Ty - Tz)(t)| &\leq e^{-\alpha t} \int_0^t \alpha e^{\alpha s} e^{-\alpha s} |y(s) - z(s)| ds \\ &\leq e^{-\alpha t} \left(\int_0^t \alpha e^{\alpha s} ds \right) \|y - z\|_\alpha \\ &\leq e^{-\alpha t} (e^{\alpha t} - 1) \|y - z\|_\alpha \\ &\leq (1 - e^{-\alpha b}) \|y - z\|_\alpha, \end{aligned}$$

and therefore

$$\|Ty - Tz\|_\alpha \leq (1 - e^{-\alpha b}) \|y - z\|_\alpha.$$

Since $1 - e^{-\alpha b} < 1$, the Banach contraction principle implies that there is a unique $y \in C(I)$ with $y = Ty$; equivalently (1.4) has a unique solution $y \in C^1(I)$. \square

Now we relax the continuity assumption on f and extend the notion of a solution of (1.4) accordingly. We want to do this in a way that preserves the natural equivalence between (1.4) and the equation $y = Ty$, which was obtained by integrating. To this end we follow the ideas of Carathéodory and make the following definitions.

Definition 1.1 A function $y \in W^{1,p}(I)$ is an L^p -Carathéodory solution of (1.4) if y solves (1.4) in the almost everywhere sense on I ; here $W^{1,p}(I)$ is the Sobolev class of functions u , with u absolutely continuous and $u' \in L^p(I)$.

Definition 1.2 A function $f : I \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ is an L^p -Carathéodory function if it satisfies the following conditions:

- (c1) the map $y \mapsto f(t, y)$ is continuous for almost every $t \in I$;
- (c2) the map $t \mapsto f(t, y)$ is measurable for all $y \in \mathbf{R}^n$;
- (c3) for every $c > 0$ there exists $h_c \in L^p(I)$ such that $|y| \leq c$ implies that $|f(t, y)| \leq h_c(t)$ for almost every $t \in I$.

If f is an L^p -Carathéodory function, then $y \in W^{1,p}(I)$ solves (1.4) if and only if

$$y \in C(I) \text{ and } y(t) = y_0 + \int_0^t f(s, y(s)) ds.$$

In fact (c1) and (c2) imply that the integrand on the right is measurable for any measurable y , and (c3) guarantees that it is integrable for any bounded measurable y . The stated equivalence now is clear. Therefore just as in the continuous case,

$$(1.4) \text{ has a solution } y \text{ if and only if } y = Ty, \quad T : C(I) \rightarrow C(I).$$

Theorem 1.8 Let $f : I \times \mathbf{R}^n \rightarrow \mathbf{R}^n$ be an L^p -Carathéodory function and L^p -Lipschitz in y ; that is, there exists $\alpha \in L^p(I)$ with

$$|f(t, y) - f(t, z)| \leq \alpha(t)|y - z| \text{ for all } y, z \in \mathbf{R}^n.$$

Then there exists a unique $y \in W^{1,p}(I)$ that solves (1.4).

Proof The proof is similar to Theorem 1.7 and will only be sketched here. Let

$$A(t) = \int_0^t \alpha(s) ds.$$

Then $A'(t) = \alpha(t)$ for a.e. t . Define

$$\|y\|_A = \left| e^{-A(t)} y(t) \right|_0.$$

The norm is equivalent to the maximum norm because

$$e^{-\|\alpha\|_1} |y|_0 \leq \|y\|_A \leq |y|_0, \text{ where } \|\alpha\|_1 = \int_0^b |\alpha(t)| dt.$$

Thus $(C(I), \|\cdot\|_A)$ is a Banach space and use of the Banach contraction principle, essentially as in the proof of Theorem 1.7, implies that there exists a unique $y \in C(I)$ with $y = Ty$. It follows that (1.4) has a unique L^p -Carathéodory solution on I . \square

Notes Most of the results in Chapter 1 may be found in the classical books of Dugundji and Granas [55], Goebel and Kirk [77] and Zeidler [191].

Exercises

- 1.1 Show that a contraction F from an incomplete metric space into itself need not have a fixed point.
- 1.2 Let (X, d) be a complete metric space and let $F : X \rightarrow X$ be such that $F^N : X \rightarrow X$ is a contraction for some positive integer N . Show that F has a unique fixed point $u \in X$ and that for each $x \in X$, $\lim_{n \rightarrow \infty} F^n(x) = u$.
- 1.3 Using the result obtained in Exercise 1.2, give an alternative proof for the Picard–Lindelöf theorem (Theorem 1.7).
- 1.4 Let \overline{B}_r be the closed ball of radius $r > 0$, centred at zero, in a Banach space E with $F : \overline{B}_r \rightarrow E$ a contraction and $F(-x) = -F(x)$ for $x \in \partial \overline{B}_r$. Show F has a fixed point in \overline{B}_r .
- 1.5 Let U be an open subset of a Banach space E and let $F : U \rightarrow E$ be a contraction. Show that $(I - F)(U)$ is open.

1.6 Let (X, d) be a complete metric space, P a topological space and $F : X \times P \rightarrow X$. Suppose F is a contraction uniformly over P (that is, for each $x, y \in X$, $d(F(x, p), F(y, p)) \leq L d(x, y)$ for all $p \in P$) and is continuous in p for each fixed $x \in X$. Let x_p be the unique fixed point of $F_p : X \rightarrow X$, where $F_p(x) = F(x, p)$. Show that $p \mapsto x_p$ is continuous.

1.7 Let $k : [0, 1] \times [0, 1] \times \mathbf{R} \rightarrow \mathbf{R}$ be continuous with

$$|k(t, s, x) - k(t, s, y)| \leq L|x - y|$$

for all $(t, s) \in [0, 1] \times [0, 1]$ and $x, y \in \mathbf{R}$ (here $L \geq 0$ is a constant) and $v \in C[0, 1]$.

(a) Show that

$$u(t) = v(t) + \int_0^t k(t, s, u(s)) ds, \quad 0 \leq t \leq 1,$$

has a unique solution $u \in C[0, 1]$.

(b) Choose $u_0 \in C[0, 1]$ and define a sequence of functions $\{u_n\}$ inductively by

$$u_{n+1}(t) = v(t) + \int_0^t k(t, s, u_n(s)) ds, \quad n = 0, 1, \dots$$

Show that the sequence $\{u_n\}$ converges uniformly on $[0, 1]$ to the unique solution $u \in C[0, 1]$.

1.8 Let (X, d) be a complete metric space and let $\phi : X \rightarrow [0, \infty)$ be a map (not necessarily continuous). Suppose

$$\inf\{\phi(x) + \phi(y) : d(x, y) \geq \gamma\} = \mu(\gamma) > 0 \text{ for all } \gamma > 0.$$

Show that each sequence $\{x_n\}$ in X , for which $\lim_{n \rightarrow \infty} \phi(x_n) = 0$, converges to one and only one point $u \in X$.

1.9 Let (X, d) be a complete metric space and let $F : X \rightarrow X$ be continuous. Suppose $\phi(x) = d(x, F(x))$ satisfies

$$\inf\{\phi(x) + \phi(y) : d(x, y) \geq \gamma\} = \mu(\gamma) > 0 \text{ for all } \gamma > 0,$$

and that $\inf_{x \in X} d(x, F(x)) = 0$. Show that F has a unique fixed point.

1.10 If in Theorem 1.6 the assumptions on ϕ are replaced by $\phi : [0, \infty) \rightarrow [0, \infty)$ is upper semicontinuous from the right on $[0, \infty)$ (that is, $\limsup_{s \rightarrow t^+} \phi(s) \leq \phi(t)$ for $t \in [0, \infty)$) and satisfies

$\phi(t) < t$ for $t > 0$. Show that F has a unique fixed point $u \in X$ with $\lim_{n \rightarrow \infty} F^n(x) = u$ for each $x \in X$.

- 1.11** Let T be a map of the metric space (X, ρ) into itself such that, for a fixed positive integer n ,

$$\rho(T^n x, T^n y) \leq \alpha^n \rho(x, y) \text{ for } x, y \in X;$$

here α is a positive real number. Show that the function σ defined by

$$\sigma(x, y) := \rho(x, y) + \frac{1}{\alpha} \rho(Tx, Ty) + \cdots + \frac{1}{\alpha^{n-1}} \rho(T^{n-1}x, T^{n-1}y)$$

is a metric on X and T satisfies

$$\sigma(Tx, Ty) \leq \alpha \sigma(x, y) \text{ for } x, y \in X.$$