

## INDEX

- acceptance region
  - similar of type  $\alpha$ , 134
- adjoint identities, 72
- adjoint of a linear transformation
  - definition of, 38, 39
  - dependence on the inner product, 72
  - for a nonorthogonal projection, 142
    - formula for, given a basis for the range space, 145
    - “reverse and perp” rule, 142
  - for an orthogonal projection, 39
  - for an orthogonal transformation, 38
  - properties of, 38–40
- admissible estimator, 97
- affine estimator, 77
- analysis of covariance
  - $A$  and  $B$  transformations, 147
  - estimation of  $\sigma^2$ , 152–153
  - $F$ -test of  $H: \mu \in M_0 + N$ , 156
    - degrees of freedom for, 156
    - noncentrality parameter for, 158
    - power of, 158
  - $F$ -test of  $H: P_M \mu \in M_0$ , 157
  - formulation of, 146–147
  - Gauss-Markov estimators, 147–149
  - one-way layout with one covariate
    - estimation of  $\sigma^2$ , 153
    - $F$ -test of the hypothesis of equal adjusted group means, 157, 158
    - $F$ -test of the hypothesis of equal unadjusted group means, 158
    - Gauss-Markov estimators, 149
    - introduction to, 146
    - simultaneous confidence intervals for all contrasts in the adjusted group means, 155
    - variances of GMEs, 152
  - Scheffé confidence intervals
    - for linear functionals of  $\mu_M$ , 153–154
    - for linear functionals of  $\mu_N$ , 155
  - variances and covariances of GMEs, 150–151
- analysis of variance
  - Latin square design, 159–163
  - one-way layout
    - ANOVA table for, 25
    - estimable parametric functionals, 79–80
    - estimation of  $\sigma^2$ , 71
    - Gauss-Markov estimation in, 63
    - Gauss-Markov estimator of an estimable parametric functional, 81
    - projections in, 15
    - simultaneous confidence intervals for all contrasts in the group means, 135
    - testing the hypothesis that the group means are equal, 111, 116
    - with a covariate, *see* analysis of covariance, one-way layout with one covariate
  - two-way additive layout
    - ANOVA table for, 25
    - differential effect, 65
    - estimable parametric functionals, 80
    - estimation of  $\sigma^2$ , 72
    - Gauss-Markov estimation in, 65, 69
    - Gauss-Markov estimator of an estimable parametric functional, 82
    - grand mean, 65
    - projections in, 17, 24
    - with a missing observation, *see* incomplete observations model, two-way additive layout with a missing observation
    - with replicated observations, 187
  - analysis of variance table
    - for a Latin square design, 162
    - for a Tjur system, 25
    - for the two-way layout, 25
  - ANCOVA, *see* analysis of covariance
  - ANOVA, *see* analysis of variance
- Baranchik’s theorem, 107
- bases
  - canonical, *see* canonical bases for a pair of subspaces

- basis, 42
  - coordinates with respect to, 42
  - orthogonal, 9
  - usual coordinate basis for  $\mathbb{R}^n$ , 42
- Bayes estimator
  - as the mean of the posterior distribution of the parameter given the data, 94
  - definition of, 92
  - extended, 92
  - formal, 108
  - generalized, 108
  - improper, 108
  - proper, 108
  - within  $\epsilon$ , 92
- best linear unbiased estimator, 64
- bilinear functionals
  - definition of, 37
  - positive-definite, 39
  - representation theorem for, 38
  - symmetric, 38
- BLUE, *see* best linear unbiased estimator
- book orthogonal, *see* subspaces, book orthogonal
- Borel  $\sigma$ -field, 45
  
- canonical bases for a pair of subspaces, 86
- canonical transformation, 75
- Cauchy-Schwarz inequality, 10
- characteristic function
  - of a normal random vector, 53
  - of a random vector, 53
- chisquare distribution
  - central, 54
  - moment generating function of, 56
  - noncentral, 54, 55
  - Poisson representation of, 56–57
- Cleveland's identity, 40
- closed form of a residual squared length, 71
- Cochran's theorem
  - algebraic form of, 20–21
  - for quadratic forms in normally distributed random variables, 58
- coefficient vector, 36
  - confirmation method for, 37
  - direct-construction method for, 37
  - effect of covariates on, 148
  - effect of missing observations on, 181
  - for a coordinate functional, 37
    - in the case of an orthogonal basis, 62
    - obtained by partial orthogonalization, 61–62
    - properties of, 62
  - obtained by projection, 60
- collinearity
  - effect on the variance of a GME, 64
- complete statistic
  - condition for, 91
  - definition of, 91
  - in the GLM, 91
- composition of linear transformations, 43
- confidence intervals
  - for the value of a linear functional  $\psi$  of  $\mu$ , 128
  - relation to the  $F$ -test of  $H: \psi(\mu) = 0$ , 128
  - simultaneous, *see* simultaneous confidence intervals
- confidence region procedure, 134
- contrasts, 134
  - spanning set for a space of contrasts, 135
- coordinate functional, 37, 43, 61
- covariance matrix, 47
- covariance operator, 47–48
  
- degrees of freedom
  - of a chisquare distribution, 54
  - of a squared length, 25
  - of a weakly spherical random vector, 51
  - of an  $\mathcal{F}$  distribution, 57
- design matrix, 1
- design of experiments, 26
- differential effect
  - in a Latin square design, 160
  - in the two-way additive layout, 65
- dimension
  - of a vector space, 42
- direct sum, *see* subspaces, direct sum of
- disjoint subspaces, 43
- dispersion matrix, 32, 49
- dispersion operator ( $\Sigma$ ), 49–50
  - of the projection of a weakly spherical random vector onto a subspace, 51
- dot-product, 6
  - convention about  $\mathbb{R}^n$ , 6, 45
  - weighted, 39
  - used with uncorrelated random variables, 52
- dual space, 43
  
- Efron-Morris estimator of a mean vector
  - assumed to lie in the space spanned by the equiangular line, 104
- eigenmanifold, 34
- eigenvalue, 34
- eigenvector, 34
- $\epsilon$ -Bayes estimator, 92
- equiangular vector ( $e$ ), 3
- equicorrelated, 73
- equivalence relation, 27
- estimable parametric functional, 78–85
  - and the covariance of Gauss-Markov estimators, 82
  - characterization of, 78–79
  - subject to constraints, 84–85
- Gauss-Markov estimator of, 80–81
  - subject to constraints, 85
- in a Tjur design, 82–84

## estimation

- of a mean, *see* estimation assuming normality, Gauss-Markov estimator, James-Stein estimator of  $\mu$
  - of a variance, *see* estimation of  $\sigma^2$
  - estimation assuming normality, 89–109
  - Bayes estimator of  $\mu$  versus the prior  $N_M(0, \lambda I_M)$ , 96
    - with a hyper-prior on  $\lambda$ , 108
  - complete sufficient statistic for  $\mu$  and  $\sigma^2$ , 91
  - conditional distribution of  $\Theta$  given  $Y = y$ , when the marginal distribution of  $\Theta$  is  $N_M(0, \lambda I_M)$  and the conditional distribution of  $Y$  given  $\Theta = \mu$  is  $N_V(\mu, I_V)$ , 96
  - desirable properties of the GME of  $\mu$ 
    - extended Bayes, 97
    - maximum likelihood estimator, 89
    - minimax for mean square error, 97
    - minimum dispersion in the class of all unbiased estimators, 90
  - Efron-Morris estimator of  $\mu$ , 104
  - maximum likelihood estimators of  $\mu$  and  $\sigma^2$ , 89–90
    - joint distribution thereof, 90
  - minimum dispersion unbiased estimators of  $\mu$  and  $\sigma^2$ , 90
  - smoothness with respect to  $\mu$  and  $\sigma^2$  of the expected value of a function of  $Y$ , 92
  - undesirable properties of the GME of  $\mu$ 
    - inadmissibility with respect to mean square error when  $\dim(M) \geq 3$ ., 102
  - see also* James-Stein estimator of  $\mu$ , James-Stein type estimators of  $\mu$  of the form  $\phi(S)X$ , Scheffé confidence intervals
- estimation of  $\sigma^2$
- effect of covariates, 152–153
  - effect of extra observations, 186
  - effect of missing observations, 176–177
  - in a Tjur system of subspaces, 71
  - unbiased, 70–72
    - potential bias in, 71
  - under normality, *see* estimation assuming normality
  - see also* specific models, such as Latin square designs and simple linear regression
- expectation
- as a nondecreasing function of a parameter, 119
  - of a random vector, 46–47
  - of the squared length of a weakly spherical random vector, 52
- experimental units, 26
- exponential family
- and completeness, 91
  - and MLR, 117

extended Bayes estimator, 92

extra observations model, 184–187

 $\mathcal{F}$  distribution, 57

central, 57

noncentral, 57

MLR property of, 118

Poisson representation of, 57

unnormalized ( $\mathcal{F}^*$ ), 57 $F$ -test of  $H: \mu \in M_0$ 

as a likelihood ratio test, 112–113

as a similar test with maximum average power over the surface of certain spheres, 137

as a uniformly most powerful invariant test, 127

when  $\sigma^2$  is known, 127as a uniformly most powerful test whose power depends only on the distance from  $\mu$  to  $M_0$  in  $\sigma$  units, 140

definition of, 113

effect of covariates, *see* analysis of covariance

effect of extra observations, 186

effect of missing observations, *see* incomplete observations model

noncentrality parameter for, 115

power of, 115

as an increasing function of the noncentrality parameter, 117, 120

simple interpretations of, 113–114

sum of squares for error ( $SS_e$ ), 113sum of squares for testing  $H$  ( $SS_H$ ), 113

unbiasedness of, 120

when  $M_0$  is the null space of a collection of linear functionals of  $\mu$ 

related confidence intervals, 133

when  $M_0$  is the null space of a linear functional of  $\mu$ , 116

related confidence intervals, 128

when  $M_0$  is the null space of a set of contrasts, 135

factor, 26

balanced, 26

blocks of, 26

equivalence relation for, 27

levels of, 26

orthogonal projection associated with, 30

subspace associated with, 29

sum of squares associated with, 30

trivial, 26

units, 26

factors

cross-classification of ( $\times$ ), 27, 29maximum of ( $\vee$ ), 27minimum of ( $\wedge$ ), 28

nestedness of, 27

- factors (*cont'd*)  
 orthogonal, 30  
   condition of proportional cell counts for, 31  
 partial ordering ( $\leq$ ) of, 27  
 Tjur design of, *see* Tjur design
- fitted values, 68
- flat, 43
- four-penny problem, 65–66
- Gauss-Markov estimator  
 of a linear functional of  $\mu$   
   center of a confidence interval, 128  
   closure under linear combinations, 63  
   covariance between the two such estimators, 64  
   definition of, 62  
   effect of covariates, *see* analysis of covariance  
   effect of extra observations, 186  
   effect of missing observations, *see* incomplete observations model  
   estimated standard deviation of, 70  
   guessing method for, 63  
   invariance under linear transformations, 75  
   mean and variance of, 63–64  
   minimum variance among linear unbiased estimators, 64  
   significantly different from zero, 133  
   used to estimate a mean vector, 69–70
- of an estimable parametric functional  
 with constraints, 85  
 without constraints, 80–81
- of  $\mu$   
 as a unique invariant estimator, 77  
 as possibly distinct from the least squares estimator, 72  
 definition of, 67  
 effect of covariates, *see* analysis of covariance  
 effect of extra observations, 186  
 effect of missing observations, *see* incomplete observations model  
 invariance under linear transformations, 74–75  
 methods for finding, 68, 75  
 minimum dispersion among linear unbiased estimators, 67  
 minimum MSE among affine estimators having bounded MSE, 76  
 risk under squared error, 92  
 used to estimate linear functionals, 69  
*see also* estimation assuming normality, James-Stein estimator of  $\mu$ , using the wrong inner product, as well as specific models, such as the Latin square design and simple linear regression
- Gauss-Markov theorem, 2  
 for estimable parametric functionals, 80, 85  
 for linear functionals of  $\mu$ , 64  
 for linear transformations of  $\mu$ , 67  
 for  $\mu$ , 67  
   when there are missing observations, 167  
 quantification of, 85–88
- general linear model  
 basis case, *see* multiple linear regression  
 geometric formulation of, 2, 60  
 matricial formulation of, 1  
 reduction to canonical form, 75
- generalized Bayes estimator, 108
- GLM, *see* general linear model
- GME, *see* Gauss-Markov estimator
- goodness of fit statistic  
 asymptotic distribution of, 55
- Gram-Schmidt orthogonalization, 9  
 in the one-sample problem, 10
- grand mean  
 in a Latin square design, 160  
 in the two-way additive layout, 65
- Hsu's theorem, 140
- hypothesis test  
 admissible, 140  
 invariant, 122  
 power function of, 125, 136  
 randomized, 136  
 similar of size  $\alpha$ , 137  
 unbiased of level  $\alpha$ , 137  
 uniformly most powerful, 125  
 uniformly most powerful invariant, 125  
*see also*  $F$ -test of  $H: \mu \in M_0$
- hypothesis testing assuming normality  
 $F$ -test, *see*  $F$ -test of  $H: \mu \in M_0$   
 general formulation of the problem, 110–111, 133  
 likelihood ratio test of  $H: \mu \in M_0$ , 112  
*see also* confidence intervals, simultaneous confidence intervals, estimation assuming normality, invariance in hypothesis testing, as well as specific models, such as the Latin square design and simple linear regression
- idempotent, *see* linear transformations, idempotent
- identity transformation, 44
- improper Bayes estimator, 108
- inadmissible estimator, 97
- incomplete observations model  
 $A$  and  $B$  transformations, 168  
 diagrams for, 168–169  
 estimation of  $\sigma^2$ , 176–177  
 $F$ -test of  $H: \mu \in M_0$ , 177–178  
 degrees of freedom for, 178  
 noncentrality parameter for, 178  
 Yates procedure, 178, 180

- incomplete observations model (*cont'd*)  
 formulation of, 165–169  
 Gauss-Markov estimator of a linear functional of  $\mu$ , 181–182  
 Gauss-Markov estimator of  $\mu$   
 ANCOVA method for, 173–174  
 consistency equation for, 169–171  
 definition of, 167  
 quadratic function method for, 172–173  
 Gauss-Markov theorem for, 167, 182  
 Scheffé confidence intervals, 183  
 two-way additive layout with a missing observation  
 $F$ -test of the null hypothesis of no row effects, 178–181  
 Gauss-Markov estimator of  $\mu$ , 171–175  
 GMEs of  $\nu$ ,  $\alpha_i$ , and  $\beta_j$ , 172  
 introduction to, 164–165  
 variances of the GMEs of  $\nu$ ,  $\alpha_i$ , and  $\beta_j$ , 182–183
- indicator function of a set, 29
- inner product  
 between two sets of vectors, 11  
 of linear functionals, 61  
 of vectors, 6
- inner product space, 6
- integrability of random variables  
 facts about, 106
- intersection of subspaces, 43  
 and projections, 10
- invariance in hypothesis testing, 121–127  
 and the  $F$ -test, 127  
 constancy on orbits, 122  
 distinguish between orbits, 122  
 group of transformations leaving the testing problem invariant, 121–122  
 invariance reduction, 125  
 and UMP tests, 125  
 invariant test, 122  
 maximal invariant, 122  
 orbits, 122  
 principle of invariance, 122  
 reduced problem, 125
- inverse of a linear transformation, 44
- isomorphism of vector spaces, 43
- James-Stein estimator of  $\mu$   
 Bayesian motivation for, 97–98  
 definition of, 98  
 improvement in Bayes risk, 102  
 inadmissibility of, 102  
 positive part thereof, 103  
 regression motivation for, 98  
 risk of, 99  
 compared to the risk of other estimators, 103  
 upper bound on, 101  
 shrinkage toward a subspace, 104  
 when  $\sigma^2$  is unknown, 104
- James-Stein theorem, 98
- James-Stein type estimators of  $\mu$  of the form  $\phi(S)X$ , 97–104  
 admissible minimax generalized Bayes, 109  
 as spherically symmetric estimators, 105  
 condition to be minimax, 107  
 motivation for, 105  
 risk of, 100  
 compared to the risk of  $\phi^+(S)X$ , 101  
 unbiased estimator for, 106
- Latin square, 159
- Latin square design, 159–163
- least squares, 8
- least squares estimator of  $\mu$ , 72
- Lebesgue measure, 53
- Lehmann-Scheffé theorem  
 for complete sufficient statistics, 91
- length of a vector, 7
- likelihood ratio test of  $H: \mu \in M_0$ , 112
- linear algebra, review of, 41–44  
 basis, 42  
 composition of two linear transformations, 43  
 coordinate functional, 43  
 dimension, 42  
 direct sum of subspaces, 43  
 disjoint subspaces, 43  
 dual space, 43  
 flat, 43  
 idempotent linear transformation, 43  
 identity transformation, 44  
 intersection of subspaces, 43  
 inverse of a linear transformation, 44  
 isomorphism, 43  
 linear dependence, 42  
 linear functional, 43  
 linear independence, 42  
 linear transformation, 43  
 manifold, 43  
 matrix of a linear transformation, 44  
 nonsingular linear transformation, 44  
 null space of a linear transformation, 44  
 range of a linear transformation, 44  
 rank of a linear transformation, 44  
 span, 42  
 subspaces, 43  
 sum of subspaces, 43  
 transpose  
 of a matrix, 44  
 of a vector, 42  
 vector space, 42
- linear dependence of vectors, 42

- linear functional
- coefficient vector of, *see* coefficient vector
  - confidence interval for, *see* confidence interval for the value of a linear functional  $\psi$  of  $\mu$
  - coordinate functional, 37, 43
  - definition of, 36, 43
  - Gauss-Markov estimator of, *see* Gauss-Markov estimator of a linear functional of  $\mu$
  - inner product for, 61
  - norm of, 61
  - representation theorem for, 36
  - testing an hypothesis about, 116
- linear independence of vectors, 42
- linear transformation
- adjoint of, *see* adjoint of a linear transformation
  - composition of, 43
  - definition of, 43
  - eigenmanifolds of, 34
  - eigenvalues of, 34
  - eigenvectors of, 34
  - idempotent, 12, 43
  - identity, 44
  - inverse of, 44
  - matrix of, 33, 44
    - notation for, 44
  - nonsingular, 44
  - null space of ( $\mathcal{N}$ ), 44
  - orthogonal, *see* orthogonal, transformation
  - orthogonal projection, *see* projections, orthogonal
  - range of ( $\mathcal{R}$ ), 44
  - rank of ( $\rho$ ), 44
  - self-adjoint, *see* self-adjoint linear transformation
  - uniqueness of
    - in terms of inner products, 8
- LSD, *see* Latin square design
- manifold
- eigen, 34
  - regression, 60
  - see also* subspaces
- matrix
- of a linear transformation, 33
  - of a quadratic form, 57
  - of the adjoint of a linear transformation, 38
  - symmetric
    - diagonalization of, 36
- maximal invariant, 122
- maximum likelihood estimators of  $\mu$  and  $\sigma^2$ , assuming normality, 89–90
- mean
- grand
    - in the two-way additive layout, 65
  - of a random vector, 46–47
- mean square error
- definition of, 76
  - formula for, 76, 92
  - normalized, 97
- method of  $\text{su}_b^{\text{per}}$  scripts, 24
- metric induced by an inner product, 10
- minimax estimator
- as a constant risk, extended-Bayes estimator, 92
  - definition of, 92
- minimum mean square error predictor, 11
- Minkowski's inequality, 10
- missing observations model, 166
- see also* incomplete observations model
- mixed-up observations model, 166
- see also* incomplete observations model
- MLR, *see* monotone likelihood ratio
- Möbius function, 25
- Möbius inversion formula, 25
- models
- analysis of covariance
    - one-way layout with one covariate, *see* analysis of covariance
  - analysis of variance
    - one-way layout, *see* analysis of variance
    - two-way layout, *see* analysis of variance
  - extra observations, *see* extra observations model
  - four-penny problem, 65–66
  - general linear, *see* general linear model
  - incomplete observations, *see* incomplete observations model
  - Latin square design, 159–163
  - missing observations, *see* incomplete observations model
  - mixed-up observations, *see* incomplete observations model
  - multiple linear regression, *see* multiple linear regression
  - one-sample problem, *see* one-sample problem
  - replicated observations model, *see* extra observations model
  - simple linear regression, *see* simple linear regression
  - split plot design, *see* split plot design
  - triangle problem, *see* triangle problem
  - two-sample problem, *see* two-sample problem
- monotone likelihood ratio
- and exponential families, 117
  - and hypothesis testing, 126
  - and the expectation of a nondecreasing function, 119
  - closure under composition, 118
  - definition of, 117
  - examples of
    - exponential families, 117
    - noncentral  $\mathcal{F}$  family, 118

- monotone likelihood ratio (*cont'd*)  
 examples of (*cont'd*)  
   Poisson family, 118  
   unnormalized  $\mathcal{F}$  family, 118  
   unnormalized  $t$  family, 120  
 stochastic monotonicity, 119
- MSE, *see* mean square error
- multiple linear regression  
 coordinate functionals in, 61  
 estimable parametric functionals, 79  
 estimation of  $\sigma^2$ , 71  
 expected response at the mean values of  
   the independent variables  
   Gauss-Markov estimation of, 65  
 regression coefficients  
   covariances of, 66  
   expectations of, 66  
   formula for, 66  
   normal equations for, 63  
   variances of, 64
- noncentrality parameter  
 of a chisquare distribution, 54  
 of an  $\mathcal{F}$  distribution, 57
- nonestimable parametric functional, 79  
 completely undetermined by  $\mu$ , 79  
 no unbiased estimator for, 81
- nonsingular distribution, 50–51
- nonsingular linear transformation, 44
- norm  
 of a linear functional, 61  
 of a vector, 7
- normal distribution, 53  
 and conditional distributions, 94–95  
   conditional distribution of  $\Theta$  given  $Y = y$ ,  
   when the marginal distribution of  $\Theta$   
   is  $N_M(0, \lambda I_M)$  and the conditional  
   distribution of  $Y$  given  $\Theta = \mu$  is  
    $N_V(\mu, I_V)$ , 96  
 and marginal distributions, 94–95  
 and quadratic forms, 57–59  
 characteristic function for, 53  
 density of, 53  
 in a lower dimensional subspace, 54
- normal equations  
 abstract formulation of, 40  
 and estimable parametric functionals, 81,  
   85  
 and Gauss-Markov estimation, 68  
 basis formulation of, 10, 69  
 obtained via Gauss-Markov estimation, 63  
 normalized mean square error, 97
- notational conventions  
 difficulty level of exercises, 5  
 inner product between  
   a single vector and a set of vectors, 11  
   two sets of vectors, 11  
 markers for the end of a proof, example,  
   exercise, and part of a problem set, 5
- notational conventions (*cont'd*)  
 matrices versus linear transformations, 5  
 numbering scheme, 4  
 summation over nonmissing subscripts ( $\oplus$ ),  
   171
- null space of a linear transformation, 44
- observed values, 68
- one-sample problem  
 and Gram-Schmidt orthogonalization, 10  
 condition for the least squares and  
   Gauss-Markov estimators to be the  
   same, 73  
 equicorrelated case, 73  
 heteroscedastic case, 52  
 independence of the sample mean and  
   variance, 2  
 projections in, 9–10  
 re Cochran's theorem, 59
- one-way layout, *see* analysis of variance,  
 analysis of covariance
- open form of a residual squared length, 71
- orbit, 122
- orthogonal  
 basis, 9  
   and Gauss-Markov estimation, 68  
 complement of a subspace, 12  
   properties of, 12  
 complement of one subspace within  
   another, 13  
   and spanning vectors, 13  
 decompositions, 19  
 factors, *see* factors, orthogonal  
 polynomials  
   derived from 1,  $i$ , and  $i^2$ , 7  
 projection, *see* projections, orthogonal  
 subspaces, 16  
 transformation, 7  
   adjoint of, 38  
   inverse of, 38  
   reflection through a subspace, 13  
   used to characterize strict sphericity, 53  
   used to characterize weak sphericity, 51  
 vectors, 7
- parameter vector, 1
- parametric functional, 78  
 estimable, *see* estimable parametric  
   functional  
 nonestimable, *see* nonestimable parametric  
   functional
- partial order relation ( $\leq$ ) on  
 factors, 27  
 self-adjoint linear transformations, 13

- partial orthogonalization
  - in the analysis of covariance, 149
- partially ordered set, 23
  - greatest lower bound ( $\wedge$ ), 28
  - least upper bound ( $\vee$ ), 27
  - minimal element, 23
  - Möbius function for, 25
  - Möbius inversion formula for, 25
  - principle of induction for, 23
  - zeta function for, 26
- perpendicularity
  - of subspaces, 16
  - of vectors, 7
- positive definiteness, 6, 39
- power function, 125
  - of the  $F$ -test, 115
  - effect of covariates, 158
  - effect of extra observations, 187
  - effect of missing observations, 178
- prediction
  - mean square error of, 11
- principle of invariance
  - in estimation, 77
  - in hypothesis testing, 122
- prior, 92
- projections
  - nonorthogonal
    - adjoints of, 142
    - as idempotent linear transformations, 142
    - definition of, 141
    - formula for the adjoint, given a basis for the range space, 145
    - formula for, given a basis for the range space, 144
    - partial inverses to, 143
  - orthogonal
    - adjoints of, 39
    - and minimum distance, 8, *see also* Cleveland's identity
    - and residuals, 8
    - as idempotent self-adjoint linear transformations, 12–13
    - definition of, 8
    - differences of, 14–15
    - matrix for, 11
    - obtained by making a linear transformation of the problem, 74
    - onto  $\mathcal{O}M$  for an orthogonal transformation  $\mathcal{O}$ , 38
    - onto a one-dimensional space, 9, 13
    - onto a subspace of a subspace, 14–15
    - onto a subspace that is an element of a Tjur system, 23
    - onto a sum of mutually orthogonal subspaces, 16
    - onto arbitrary subspaces, 10
    - onto the intersection of two subspaces, 18
- projections (*cont'd*)
  - orthogonal (*cont'd*)
    - onto the orthogonal complement of the null space of a set of contrasts, 135
    - products of, 17–18
    - sums of, 16–17
    - used in establishing the distributions of quadratic forms, 58
    - used to find Gauss-Markov estimators, 68
- proper Bayes estimator, 108
- proportional cell counts
  - condition for orthogonal factors, 31
- Pythagorean theorem, 7
- quadratic form
  - definition of, 39
  - of a normal random vector
    - distribution of, 57–59
    - on an inner product space, 57
    - representation theorem for, 39
- random vectors
  - characteristic function for, 53
  - covariance operators for, 47–48
  - definition of, 45
  - dispersion operators of, 49–50
    - spectral representation of, 50
  - distribution of projections of
    - under first and second moment conditions, 54
    - under normality assumptions, 54
  - expectation of, 46–47
  - mean of, 46–47
  - normally distributed, 53
    - conditional distribution, 94–95
    - marginal distribution, 94–95
    - quadratic forms in, 57–59
  - spherically distributed, 53
  - uncorrelated, 48
  - weakly spherical, 51–52
    - variance parameter of, 51
  - with nonsingular distributions, 50–51
  - with singular distributions, 50–51
- range of a linear transformation, 44
- rank of a linear transformation, 44
- reduction to canonical form, 75
- regression, *see* multiple linear regression, simple linear regression
- regression coefficients, 66
  - dependence on the number of explanatory variables, 67
- regression manifold, 60
- regression matrix, 1
- relative orthogonal complement, 13
  - and spanning vectors, 13
- replicated observations model, 184



- representation theorem
  - for bilinear functionals, 38
  - for linear functionals, 36
  - for quadratic forms, 39
- residual squared length
  - closed form of, 71
  - open form of, 71
- residuals, 8
  - inner products of, 145
  - length of, 8
  - uncorrelated with fitted values, 68
- risk, 92
  - of the GME of  $\mu$  relative to mean square error, 92
- see also* James-Stein estimator of  $\mu$ ,  
James-Stein type estimators of  $\mu$   
of the form  $\phi(S)X$
- $R^2$ 
  - as the ratio of explained variation to total variation, 68
  - as the square of the correlation coefficient between the observed and fitted values, 68
- Scheffé confidence intervals
  - for all contrasts in the group means in one-way ANOVA, 135
  - for linear functionals of  $\mu$ , 129–131
    - effect of covariates, *see* analysis of covariance
    - effect of missing observations, *see* incomplete observations model
  - for linear functionals of the component of  $\mu$  orthogonal to  $M_0$ , 134
  - relation to the  $F$ -test, 133
- Scheffé multiplier, 131
- SDFZ, *see* significantly different from zero
- self-adjoint linear transformation
  - as a covariance operator, 33
  - definition of, 12
  - extremal characterization of the eigenvalues thereof, 34
  - matrix for, 33
  - partial ordering ( $\leq$ ) of, 13
  - positive semi-definite, 36
    - square root of, 36
  - positive-definite, 39
  - range and null space of, 13
  - spectral theorem for, 33–36
  - trace of, 58
    - and preservation of order ( $\leq$ ), 76
- shrinkage estimator, 104
  - see also* James-Stein estimator of  $\mu$ ,  
James-Stein type estimators of  $\mu$  of  
the form  $\phi(S)X$
- significantly different from zero, 133
- simple linear regression
  - coefficient vector for the slope functional, 62
  - confidence interval(s)
    - for a single predicted value, 129
    - for all predicted values, 131
    - for predicted values when the predictand ranges over an interval, 132
  - definition of, 1
  - estimated SD of
    - the response at the mean of the predictor variable, 71
    - the slope coefficient, 71
  - estimator of  $\sigma^2$ , 70
  - Gauss-Markov estimate of the slope, 63
  - projections in, 10, 11
  - testing for zero slope, 110, 116
  - variance of the GME of the slope coefficient, 64
- simultaneous confidence intervals
  - for a line segment of linear functionals, 131
  - for the values of  $\psi(\mu)$  for a linear space of linear functionals  $\psi$ , *see* Scheffé confidence intervals
- singular distribution, 50–51
- span of vectors, 42
- spectral theorem, 33
- spherical distribution, 53
- spherically symmetric estimator, 105
- split plot design, 31
  - estimable parametric functionals, 84
- Stein's lemma, 105
- Stein-Efron-Morris theorem, 105
- stochastic monotonicity, 119
- Strawderman-Berger theorem, 109
- subspaces
  - book orthogonal, 17–18
    - see also* Tjur system
  - definition of, 43
  - direct sum ( $\oplus$ ) of, 19, 43
  - disjoint, 43
  - eigenmanifolds, 34
  - intersection ( $\cap$ ) of, 43
  - nonorthogonal
    - measuring the degree of, 85
  - orthogonal ( $\perp$ ), 16
  - orthogonal complements ( $\cdot^\perp$ ) of, 12
    - properties of, 12
  - orthogonal decompositions of, 19
  - perpendicular ( $\perp$ ), 16
  - relative orthogonal complements of, 13
    - and spanning vectors, 13
  - shifted, 43
  - sum ( $+$ ) of, 43
  - Tjur system of, *see* Tjur system of subspaces

- sufficient statistic  
 definition of, 90  
 factorization criterion for, 91  
 for  $\mu$  and  $\sigma^2$ , 91  
 having minimum dispersion in the class of  
 all unbiased estimators, 91  
 sum of squares for error ( $SS_e$ ), 113  
 sum of squares for testing  $H$  ( $SS_H$ ), 113  
 summation over nonmissing subscripts ( $\oplus$ ),  
 171
- $t$  distribution  
 unnormalized noncentral  
 definition of, 120  
 MLR of, 120
- test, *see*  $F$ -test of  $H: \mu \in M_0$ , hypothesis test
- Tjur design, 31  
 estimable parametric functional, 82–84  
 factor structure diagram for, 32  
 Gauss-Markov estimator of an estimable  
 parametric functional, 83–84
- Tjur system of subspaces, 21–26  
 analysis of variance table for, 25  
 and estimation of  $\sigma^2$ , 71  
 and Gauss-Markov estimation, 69  
 method of  $\text{sup}_b^{\text{per}}$  scripts for, 24  
 Möbius inversion formulas for, 26  
 partial order on, 21  
 representation of, 21  
 structure diagram for, 24
- Tjur's theorem, 21–23
- Tjur, Tue, 21
- trace  
 of a linear transformation, 76  
 of a self-adjoint linear transformation, 58,  
 76
- transformation, *see* linear transformation
- triangle problem  
 coefficient vectors in, 61  
 covariances of GMEs, 64  
 Gauss-Markov estimation in, 63  
 testing the equilateral hypothesis, 110, 115
- trivial factor, 26
- two-sample problem, 1–2
- two-way layout, *see* analysis of variance
- UMP, *see* uniformly most powerful
- UMPI, *see* uniformly most powerful invariant
- unbiased  
 estimator, 63  
 hypothesis test, 137
- uncorrelated random vectors, 48
- uniformly most powerful, 125
- uniformly most powerful invariant, 125
- units factor, 26
- using the wrong inner product  
 conditions for false and true estimates to  
 be the same  
 for linear functionals of  $\mu$ , 74  
 for  $\sigma^2$ , 74  
 for the mean vector  $\mu$ , 72–73  
 difference between the false and true GMEs  
 of  $\mu$ , 87  
 effect on residual squared length, *see*  
 Cleveland's identity  
 ratio of the variances of the false and true  
 GMEs of a linear functional of  $\mu$ ,  
 87–88
- value of a statistical game, 92
- variance parameter  
 estimation of, *see* estimation of  $\sigma^2$   
 of a weakly spherical random vector  
 assumptions about, 60  
 definition of, 51
- variance-covariance matrix, 49
- variation  
 addition rule for, 68  
 explained, 68  
 residual, 68  
 total, 68
- vector space, 42  
 basis for, 42  
 dimension of, 42  
 dual of, 43  
 finite dimensional, 42  
 isomorphism of, 43  
 manifolds of, 43  
 subspaces of, 43
- vectors  
 angle between, 7  
 equiangular ( $e$ ), 3  
 length of, 7  
 linear dependence of, 42  
 linear independence of, 42  
 norm of, 7  
 orthogonal, 7  
 orthogonal projection of, 8  
 span of, 42  
 uniqueness of  
 in terms of inner products, 8
- Wald's theorem, 137  
 corollaries to, 140  
 reformulation of, 137
- weak sphericity, 51–52  
 achieved by changing the inner product, 52  
 achieved by transformation, 52
- weighing design, 66
- Yates procedure, 178
- zeta function, 26