



## INDEX

- absorption bands, 27
- acceleration
  - at a fixed point in a fluid, 56
  - following a particle of fluid, 56
  - geostrophic, 54
  - relative to Earth, 54
- accessory pigments, 133
- Acoustic Doppler Current Profiler (ADCP), 16–17, 19, 242
- acoustics
  - observing plankton, 14, 19
- active fluorescence, 136
- advection-diffusion equation, 108–9
  - for Fickian diffusion, 111
- ageostrophic flow, 247
- albedo, 27
- Amazon ROFI, 275
- ammonium, 13, 144, 146, 197
  - preferential uptake by phytoplankton, 146
- Arctic shelf
  - freshwater supply, 359
  - future challenges, 358–60
  - sea ice, 359
- Argo floats, 18
- atmospheric absorption, 27, 29
- attenuation coefficient
  - PAR, 132
- attenuation of motion with depth
  - surface waves, 92
- attenuation, light, 132–3
- autonomous underwater vehicles (AUVs), 19
- autotrophs, 129, 135
  
- back radiation, 29–30
- bacteria, 146, 197, 299
  - heterotrophic, 160–1
  - recycling organic matter, 155, 160
- baroclinic flow, 63. *See also* barotropic flow
- baroclinic instability, 245
  - eddy scale, 246
  - energy source, 246
  - in frontal regions, 245–7
  - of coastal currents, 270
- barotropic flow, 63. *See also* baroclinic flow
- basking sharks, 256, 257, 344
- Bay of Fundy tides, 85
- bed stress
  - boundary condition in model, 206
  - drag coefficient, 179
  - effect on along-slope flow, 316
  - quadratic drag law, 179, 206
- Benguela current, 313
- Benguela upwelling system, 334
- Bering Sea
  - downwelling and primary production, 338
  - fisheries, 343
  - Green Belt, 338, 339
- Bernoulli equation, 92
- Bigelow, Henry, 7
- biological pump, 157, 163
- black body, 26, 29
- blooms, 140, 154, 169. *See also* spring bloom
  - harmful, 226
  - timing in Region of Freshwater-Influence, 295
- bottom Ekman layer, 70–2
- boundary layers in shelf seas and deep ocean, 305
- Boussinesq approximation, 60
- Bowden, Kenneth, 8, 9, 87, 114
- brine rejection, 360
- Bumpus, Dean, 9
- buoyancy
  - changes due to heating, 35–6
  - buoyancy force, 68
    - from heat input, 178
  - buoyancy frequency, 68, 118
    - limit on internal wave frequency, 104
  - buoyancy inputs, 173
  - buoyancy of water particles, 26
  
- caesium tracer distribution, 86, 298
- California current, 313, 337
- California upwelling, 333, 335, 345
  - hypoxia, 336
- canyons
  - carbon export, 345, 348
  - effects on downslope flow, 318
- carbohydrates, 131
- carbon
  - supply by rivers, 298
- carbon export, 155–8, 171, 337, 344–5
  - biological pump, 157
  - by slope current, 347
  - canyons, 348
  - continental shelf pump, 348

## Index

- carbon export (cont.)
  - DIC, 345
  - downwelling, 345–6
  - mesozooplankton role, 163, 171
  - North Sea, 364
  - phytoplankton cell size, dependence on, 156
  - role of the thermocline, 345–7
  - shelf seas' role, 157
  - sinking phytoplankton, 156
  - upwelling systems, 345
- carbon fixation, 131–3, 140. *See also* primary production
  - global map, 3
  - in the dark, 140
  - measurement using  $^{14}\text{C}$ , 134–5
  - rates during spring bloom, 192
- carbon supply by rivers, 298
- cascades, 317–20
  - carbon export, 348
  - force balance in down-slope flow, 317
  - global impact, 320
  - Gulf of Lions, 348
  - Malin shelf edge, 318
- cascading
  - desalination plants, 320
- case 1 water, 136
- case 2 water, 136
- CDOM. *See* gelbstoff
- cell quota, 148
  - maximum quota, 150
  - subsistence (minimum) quota, 150
- Celtic Sea, 7, 166
  - frontal jet, 242
  - internal tide, 322, 327, 339
  - internal tide and phytoplankton community, 341
  - internal tide and primary production, 341
  - macronutrient profiles, 144
  - shelf edge and fisheries, 163, 343–4
  - shelf edge fisheries, 343–4
  - shelf edge primary production, 193
  - spring-neap cycle, 196
  - subsurface chlorophyll maximum, 218
  - vertical profile of phytoplankton species, 227
- centre of gravity of the Earth-Moon system, 41
- centrifugal force, 40, 42
- Chesapeake Bay
  - phytoplankton transport by residual flow, 293–5
  - residual flow, 275
- chlorophyll *a*, 12, 133
  - fluorometer, 13
  - measurement from satellites, 20, 136
  - measurement using fluorescence, 133
  - tidal mixing fronts, 249
- ciliates, 161, 197
- climate change, 368–9
  - ecosystem changes, 369
  - impacts of, 196
  - ROFIs, 369
  - stratification, 368
  - tidal energy dissipation, 368
  - warming of shelf seas, 369
- CO<sub>2</sub>
  - air-sea flux, 136, 155
  - air-sea flux in upwelling systems, 345
  - fixing to organic compounds, 131
  - inorganic carbon release to the atmosphere, 161
  - mixed water column, 346
  - phytoplankton and climate, 130
  - source to atmosphere, 298
  - trapping by thermocline, 347
  - uncertainties in fluxes, 363–4
  - uptake by cooler water, 348, 349
- coastal buoyancy current, 270–2
  - $^{137}\text{Cs}$  transport around the UK, 298
  - Norwegian Coastal Current, 270
  - Rhine, 270
- coccolithophores, 153–4
  - Emiliana huxleyi*, 153
- compensation depth, 140–2
- compensation light intensity. *See* compensation depth
- conservation rules, 52
- continental shelf pump, 348, 368
- continuity
  - continuity statements, 51
  - fluid, 50
  - volume continuity, 50
- convective chimneys, 52
- convective mixing, 191, 360
  - due to tidal straining, 281, 283
  - phytoplankton growth in winter, 191
- convective overturning. *See* convective mixing
- convectively unstable, 173
- copepod, 161, 162, 165, 169, 195, 333
  - egg production, 169, 255, 334
  - nauplii, 161, 165
  - response to phytoplankton bloom, 169
- co-range lines, 80
- Coriolis force, 54–5
  - derivation for a particle moving east, 55
  - horizontal components, 54
  - vertical component, 55
- Coriolis parameter, 54
- co-tidal lines, 80
- critical depth, 142, 143, 191, 192, 198
- critical mixing, 143, 198
- cross-isobath flow, 307
- cross-slope transport, 313–20
  - bottom boundary layer, 316–17
  - filaments, 314, 337

- net exchange in boundary layers, 317
- surface boundary layer, 314–16
- CTD, 10–11, 19
  - accuracy, 11
  - rosette, 11
- currents
  - measurement, 15–18
- Cushing, David, 196
- cyanobacteria, 152, 154, 161, 227, 343
  - Trichodesmium*, 146
- D'Alembert's principle, 54
- Dalton number, 30
- dark reactions. *See* photosynthesis – light-independent stage
- deep water waves, 93
  - group velocity, 96
- Delaware Bay
  - coastal current, 289
  - transport of crab larvae, 292
- denitrification, 147
- density, calculation of, 11
- density-driven circulation, 267. *See also* estuarine circulation
  - crab larvae, Delaware Bay, 292
  - dependence on stratification, 287
  - dinoflagellate transport, Chesapeake Bay, 295
  - Heaps solution with rotation, 273
  - transport of organisms, 292–5
- desalination plants, 320
- diatoms, 145, 154, 155, 167, 193, 226
  - importance to food chain, 195
  - shelf edge, 341
  - sinking, 154, 195
  - storage of nitrate, 148
  - upwelling regions, 332
- diazotrophs, 146
- diffuse attenuation coefficient, 28
- diffusion by random walk, 112–13
- diffusivity
  - for random walk, 112
  - increase with scale, 113
  - Osborn relation, 121
  - vertical mixing time scale, 114
- dinoflagellates, 152–3, 155, 161, 194, 225, 227
  - food for fish larvae, 335
  - harmful algal blooms, 299
  - swimming speeds, 225
  - transport by density-driven circulation, 293–5
  - upwelling regions, 333
- dissipation in shelf seas and deep ocean, 46
- dissipation of turbulent kinetic energy (TKE)
  - conversion to heat, 120
- dissolved inorganic carbon (DIC), 130, 155
- dissolved organic carbon (DOC), 160
- dissolved organic matter (DOM), 160
- dissolved organic nitrogen (DON), 160, 161
- dissolved organic phosphorus (DOP), 161
- dissolved oxygen, 131, 134, 161, 335
  - measurement, 13
  - saturated concentration, 131
  - super-saturation, 336
- Dohrn, Anton, 6
- Doppler effect, 16. *See also* Acoustic Doppler Current Profiler
- downward flux of the total radiant energy, 28
- downwelling
  - carbon export, 345–6
  - Mid Atlantic Bight, 338
  - shelf edge primary production, 338
- drifters
  - current measurement, 17–18
  - frontal jet, 244
  - gyre circulation, Irish Sea, 259
  - slope current, 308
- dye tracer experiments on cross-frontal flow, 247
- East Australian Current
  - upwelling, 337
- ecosystem models, 355
  - emergent properties of communities, 356
  - new approaches, 354–6
- eddies
  - along-slope current, 320–1
  - baroclinic instability at fronts, 245–7
  - in upwelling regions, 331
- eddy diffusivity, 111
  - Osborn relation, 211
- eddy viscosity, 69, 110
- efficiency of mixing, 179
  - by tides, 181
  - by wind, 184
  - estimates for tide and wind, 185
  - reduction by stratification, 175
- Ekman depth, 69
- Ekman drain, 316
- Ekman dynamics, 68–70
- Ekman layer, 69
  - at the seabed, 70–2
- Ekman number, 273
- Ekman spiral, 70
- Ekman transport, 70, 314
  - at a boundary, 72–5
    - baroclinic coastal response, 74–5
    - barotropic coastal response, 72–4
- Ekman veering, 72
- Ekman, V. W., 15
- El Niño (ENSO), 334–5, 337
- energy dissipation
  - in wave breaking, 96
- energy flux exponential decay, 28
- energy input at the top of the atmosphere, 26

- Energy sources, 25–6  
 entrainment velocity, 175  
 equations of motion, 59–60  
 estuarine circulation, 267–9. *See also*  
   density-driven circulation  
   components due to density gradient and river flow, 269  
   enhancement by tidal straining, 288  
   velocity profile, 269  
   versus stirring in an estuary, 275–7  
 estuarine stratification  
   criterion, 276  
 eukaryotes, 152  
 Eulerian current, 17  
 Eulerian velocity, 56–7  
 eutrophication, 146, 161, 299  
 evaporation, 38  
 export production, 156  
   nutrient requirements, 330  
   nutrient requirements in shelf seas, 330  
 export ratio, 156
- fast repetition rate fluorescence. *See* active fluorescence  
 fast repetition rate fluorometer. *See* active fluorescence  
 Fick's law, 106, 110  
 Fickian diffusion, 110–13  
 filaments  
   carbon export, 345  
   East Australian Current, 337  
   upwelling regions, 314, 331, 337  
 finite amplitude waves, 93–5  
 fish larvae, 169  
   diet, 344  
   food requirements, 164  
   shelf edge, 344  
   subsurface chlorophyll maximum, 335  
   transport by frontal jet, 260  
 fisheries, 195, 363  
   shelf edge, 163, 343–4  
   tidal mixing fronts, 256–7, 262  
 flagellates, 152–3, 161, 227  
   upwelling regions, 333  
 flow components  
   mean and turbulent, 107–8  
 flow in horizontal plane, simplification, 52  
 flow instability  
   in slope currents, 321  
 fluorescence, 133. *See also* active fluorescence  
 fluorometer, 133  
 force due to frictional stress, 57, 59  
 force due to pressure gradient, 58  
 forcing by atmospheric pressure, 40  
 forcing by wind stress, 39–40  
 $f$ -ratio, 155, 157  
   at a tidal mixing front, 251  
   shelf edge, 341  
   spring bloom, 193  
   summer surface layer, 197  
 freshwater, 26  
   buoyancy source, 36–9  
 freshwater buoyancy  
   contrast with buoyancy from heating, 266–7  
 frictional and pressure forces, 57–9  
 frictional stress, 48, 58  
   restraining torque on the Earth, 48  
 front. *See* tidal mixing (TM) front  
 frontal convergence, 52  
 frontal jet, 241  
   baroclinic instability, 245  
   contribution to summer circulation, 245  
   retention of larvae, 260  
   transport of fish larvae, 260  
 frontal position  
   condition including wind stress, 184  
   tidal mixing only, 180
- Gaussian function  
   solution for 1D diffusion, 111  
 gelbstoff, 20  
 Georges Bank, 9, 200  
   fish stocks, 260–2  
   frontal jet, 244  
   subsurface chlorophyll maximum, 218  
   tidal mixing front, 239, 244, 247  
   tidal mixing front and primary production, 251  
   tidal mixing front and zooplankton, 255  
 geostrophic acceleration, 54  
 geostrophic balance, 61–2  
   tidal mixing (TM) front, 241–2  
 geostrophic flow  
   control by topography, 306–7  
   direction, 61  
   equations, 61  
   level of no motion, 65  
   thermal wind  
     shear, 65  
   thermal wind equations, 65  
   thermal wind relation at a front, 240  
   velocity profile, 64  
 gliders, 19  
 global distribution of freshwater input, 37–8  
 global overturning circulation, 360  
 gradient equation, 64  
 grazers  
   effect on spring bloom, 192  
 group velocity, 96  
   internal waves, 103  
 Gulf of Maine, 7, 196, 229  
 Gulf of Thailand, 37, 277, 362

## Index

- Gulf Stream  
influence on Irish Sea, 87
- Haackel, Ernst, 6
- Hansen and Rattray  
estuarine circulation theory, 267  
estuarine classification scheme, 269
- harmful algal blooms (HABs), 226, 299
- Harvey, H. W., 12
- heat exchange  
back radiation, 29–30  
by conduction, 30–1  
by evaporation, 30  
solar radiation, 26–9  
with sediments, 32
- heat flux  
for onset of stratification, 185  
variation with latitude, 34–5
- heat input to surface layer, 177
- heat storage and rate of surface heating, 34
- heat stored in the water column, 32
- heat transport by currents, 32
- heating and cooling  
two mixed layer model, 186
- Hensen, Victor, 6
- heterotrophs, 129, 135, 158–64
- history of shelf sea science, 6–9
- holographic analysis of particles, 14
- holoplankton, 162
- holoplanktonic, 162
- horizontal density gradients, 26
- hydrostatic approximation, 60
- hydrostatics, 59–60
- hypoxia, 4, 336  
bottom waters of ROFI, 299–301  
Louisiana dead zone, 299  
upwelling regions, 335–6
- Iberian upwelling system, 331, 345
- ICES, 7
- inertial frequency  
limit on internal wave frequency, 104
- inertial oscillations, 66–7, 353  
contribution to pycnocline mixing, 215  
mixing, 353
- inertial period, 66
- infinitesimal waves, 92
- internal Rossby radius, 75, 246
- internal tide, 321–6  
breaking, 324  
Celtic Sea, 339  
energy propagation onto the shelf, 327  
examples, 322–4  
generation over slope, 321–2  
intermittent character, 324  
nitrate flux at the shelf edge, 339–40  
phytoplankton community, 341–3  
propagation and solibore development, 322–6  
shelf edge primary production, 341
- internal waves  
allowed frequency range, 104  
bands of surface material, 102  
continuous stratification, 103–4  
contribution to pycnocline mixing, 214–15  
convergence and divergence, 102  
energy propagation, 103  
finite amplitude, 101  
group velocity, 103  
importance, 104–5  
in models, 216  
kinetic energy, 103  
mode 1, 104  
modelling, 353, 354  
packets in breaking internal tide, 324  
parameterisation of pycnocline mixing, 216  
particle orbits, 100–1  
potential energy, 103  
residual transport, 102  
streamlines of flow, 102  
two-layer stratification, 99–103  
velocity components, 100–1  
velocity potential, 99–100
- Irish Sea, 7, 8, 86, 200  
attenuation of PAR, 200  
gyre and *nephrops norvegicus*, 257–60  
modelled turbulence, 209  
sediment nutrient flux, 146  
Sellafeld discharge, 298  
tidal mixing front, 181, 240, 241  
tidal mixing front and phytoplankton, 253  
tidal mixing front and seabirds, 256  
tidal mixing front and zooplankton, 255
- irrotational, 90
- island mixing front, 251
- isobar  
definition, 61  
slope, 64
- JEBAR  
defined, 311  
forcing mechanism, 311–13  
theory, 311–13
- Kelvin wave, 79  
direction, 78  
energy loss, 80  
equations, 78  
incident and reflected, 79  
reflection at the head of a gulf, 81  
tide in Rhine ROFI, 286
- kinematics, 50–2

## Index

- kinetic energy
  - internal waves, 103
  - surface waves, 95
- Knudsen, Martin, 8
- Kolmogorov, Andrey, 121, 124
- Kuroshio
  - deflection into Sea of Japan, 321
- laboratory experiments
  - baroclinic instability in fronts, 246
  - estuarine circulation and mixing, 279–81
  - stratification and mixing, 174
- Lagrangian
  - current, 17
  - currents in tidal mixing front, 244
  - model of phytoplankton trajectories, 201, 225
  - velocity, 56–7
- laminar flow, 105
- Langmuir circulation, 97
- Laplace, Pierre-Simon, 44
  - Laplace's equation, 91
- laser particle size analyser, 14
- Lasker Event, 335
- Lasker, Reuben, 228
- latent heat transfer, 30
- Liebig's Law of the Minimum, 150
- linearising approximation, 57
- Liverpool Bay ROFI
  - biological response, 296
  - semi-diurnal stratification, 281
  - standing wave tide, 283
  - stratification time scales, 277
  - tidal straining, 283
- lock-exchange experiment, 279
- long wave
  - amplitude change across the shelf edge, 79
  - energy flux, 79
  - equation, 76
  - internal, 100
  - particle velocity, 76
  - reflection, 76
  - with rotation, 77–9
  - without rotation, 75–7
- Louisiana Dead Zone, 299–301, 335
- macronutrients, 330
  - vertical profile, Celtic Sea, 144
- Marine Biological Association, Plymouth, UK, 7
- Massachusetts Bay, 198
- match-mismatch hypothesis, 195–6, 256
- Mekong River, 37
- Meridional Overturning Circulation, 367.
  - See also* Global Overturning Circulation
- meroplankton, 162
- mesozooplankton, 162–4, 195
  - swimming, 162
  - swimming in a tidal environment, 228
  - vertical migration, 162
- microbial loop, 160, 161, 169
- microscale
  - in shelf seas, 124
- microzooplankton, 161–2, 195
- mixed layer development
  - laboratory analogue, 174–6
- mixing in pycnoclines
  - candidate mechanisms, 214–16
- mixotrophs, 161
- models. *See also* ecosystem models
  - challenges in linking to observations, 357
  - importance in oceanography, 21–2
  - Occam's razor, 22
  - potential energy anomaly model of heating-stirring, 179
  - ROFIs, 290–1
  - turbulence closure (TC) model, 205–9
  - two mixed layer (TML) model, 185–90
  - unstructured grids, 354
- molecular diffusion, 106
- molecular diffusivity, 106
- molecular viscosity, 106
- momentum flux, 109
- Monod growth curve, 139
- Monterey Bay, 337
  - phytoplankton thin layer, 218, 226
  - zooplankton layers, 228
- negative buoyancy
  - from winter cooling, 317
- net transport
  - due to large internal waves, 324
- new production, 155
  - shelf seas, 158
- New Zealand
  - internal tide, 322, 324
  - mixing by tropical cyclone, 198
  - sub-surface chlorophyll maximum, 218
  - upwelling, 314
- Newton, Isaac, 43
  - second law in a rotating frame of reference, 54–6
  - second law of motion, 52
- nitracline, 218
- nitrate, 13, 144, 145, 197
  - atmospheric supply, 146
  - conversion to ammonium by
    - phytoplankton, 146
  - demand at a tidal mixing front, 251
  - distribution at a tidal mixing front, 249
  - flux from deep ocean, 145
  - flux from sediments, 146
  - flux through thermocline, 155, 157, 221
  - measurement on a mooring, 198
  - supply by rivers, 330

## Index

- nitrate flux  
 comparison between shelf sea and ocean, 343  
 control on thermocline primary production, 223  
 internal tide, 339–40  
 into the thermocline, 221, 224
- nitrification, 146
- nitrite, 146
- nitrogen  
 atmospheric gas, 146  
 fixation by diazotrophs, 146  
 flux from rivers, 145
- nitrogen cycle, 147
- nitrogen fixation, 146–7
- nodes of wave motion, 77
- non-linear terms, 57
- North Sea, 6, 36, 86, 196, 298  
 carbon export, 346–7, 364  
 fisheries, 4  
 primary production, 223  
 salinity and Rhine ROFI, 270  
 spring bloom, 192
- Norwegian Coastal Current, 270
- nutrient uptake, 148–9  
 cell size, role of, 149, 167, 197  
 half-saturation concentration for nitrate uptake, 149  
 measurement, 148–9  
 Michaelis-Menten uptake kinetics, 148  
 nitrate uptake rate, 149  
 nutrient quota, 148  
 specific nutrient uptake rate, 149
- nutrients. *See also* nitrate, phosphate, silicate, ammonium  
 depletion by spring bloom, 192  
 inorganic nitrogen, 145. *See also* nitrate  
 iron, 144, 339  
 limiting nutrient, 144–5, 197  
 macronutrients, 13, 144  
 measurement, 13  
 micronutrients, 144  
 nitrogen, 146  
 nutrient quota, 148  
 phytoplankton requirements, 143–4  
 pre-formed, 156  
 requirements for phytoplankton, 143  
 sources, 145–7  
 storage in phytoplankton cells, 148  
 supply by upwelling, 75  
 units of concentration, 145
- Okubo, Akira  
 diffusion diagrams, 113
- ontogenic, 163
- Optical Plankton Counter (OPC), 14
- organic matter  
 consumption by heterotrophs, 155, 158  
 export. *See* carbon export  
 fate of, 155
- Osborn relation for vertical diffusivity, 121, 211
- PAR. *See* photosynthetically available radiation (PAR).
- particulate organic carbon (POC), 156
- Péclet number, 152, 225, 228
- Peru upwelling system, 332, 335
- phase speed  
 internal waves, 100  
 Kelvin wave, 78  
 long wave, 76  
 surface waves, 92
- phosphate  
 flux from rivers, 145
- phosphate, 13, 144, 145  
 flux from deep ocean, 145
- photic zone, 137
- photoacclimation, 140, 355  
 mixed water column, 200–2
- photoautotrophs, 130, 135
- photoinhibition, 140
- photosynthesis, 131–3, 134  
 energy required, 131  
 light-dependent stage, 131  
 light-independent stage, 131  
 maximum light utilisation coefficient, 139  
 non-photochemical quenching, 133, 140, 192  
 photosynthesis-PAR curve, 138–40  
 photosystems I and II, 131  
 rate of, 139  
 reaction centre, 131, 140  
 saturation of, 133  
 summary of reactions, 131
- photosynthetic quotient, 134
- photosynthetically available radiation (PAR), 27, 131  
 units, 140–1
- phytoplankton, 130  
 community, 135–6, 154–5, 158  
 community, the role of grazers, 167–9, 194  
 functional types (PFTs), 355  
 groups, 133, 150–4  
 internal tide and phytoplankton community, 341–3  
 sinking, 156  
 species. *See* phytoplankton, groups  
 spring bloom succession of groups, 194  
 support of fisheries, 363  
 swimming, 152–3, 218, 225
- Planck radiation law, 26, 29
- pollutants  
 agricultural and wastewater run-off, 4, 145, 298, 330  
 nutrients supplied via rivers, 298–301

## Index

- potential energy
  - increase due to mixing, 116
  - internal waves, 103
  - surface waves, 95
  - water column, 177
- potential energy anomaly  $\Phi$ , 176
  - contribution due to estuarine circulation, 275–7
  - contribution from tidal straining, 282–3
  - decrease due to cooling, 179
  - decrease due to tidal stirring, 179
  - effect of wind stress, 184
  - equation
    - for heating versus stirring, 179
  - increase due to heating, 178
  - integrating forward in time, 184
  - regression analysis for mixing efficiencies, 185
- potential function, 91
- practical salinity scale, 11
- practical salinity unit (psu), 11
- Prandtl number, 118
- predator-prey interaction, 164–5
  - turbulence and encounter rates, 165
  - viscosity, 164–5
- pressure, 57
- pressure force, 57
- pressure gradient
  - in estuarine flow, 268
- primary production, 130
  - cycles of stratification in ROFIs, 295–8
  - downwelling shelf edge, 338
  - effects of spring-neap cycle, 222–3
  - export production, 156
  - gross, 133
  - gross community, 134
  - in upwelling regions, 331–3
  - limitation by light or nutrients, 150
  - measurement of, 134–5, 140
  - measurement using  $^{14}\text{C}$ , 134–5, 138–40
  - measurement using active fluorescence, 136–7
  - measurement using oxygen, 134
  - measurement using satellites, 3, 136
  - mixed water column, 199–202
  - net, 133
  - net community, 134
  - new production, 155
  - regenerated production, 155, 197
  - semi-diurnal tidal cycle in a ROFI, 296–8
  - shelf edge, 338, 341
  - simulated *in situ*, 137–8
  - spring bloom, 192–3
  - spring-neap adjustment of a tidal
    - mixing front, 252
  - spring-neap cycles in ROFIs, 296
  - sub-surface chlorophyll maximum, 223
  - tidal mixing front, 251
  - triggered by a storm, 198
  - turbulent environment, 136, 142, 199–202
- progressive wave, 77
- prokaryotes, 152
- pycnocline. *See* thermocline
- rainfall, 38
- Redfield ratio, 150
  - carbon export, 156–7
- reduced gravity, 74, 100
- reference frame
  - rotating and non-rotating, 54
- regenerated production, 155
- Region Of Freshwater Influence (ROFI),
  - 36, 266
  - Arctic shelf, 359
  - bloom timing, 295
  - climate change, 369
  - contrasts with estuaries, 266–7
  - density-driven circulation, 272–5
  - effects of riverine nutrients, 298–301
  - effects of wind stress, 288–90
  - hypoxia in bottom water, 299–301
  - modulation of residual circulation, 287–8
  - numerical models, 290–1
  - primary production over semi-diurnal tidal
    - cycle, 296–8
  - spring-neap stratification and primary
    - production, 296
  - spring-neaps cycle in stratification, 279
  - stratification, 277–9
  - time scales of mixing and stratification, 277
  - transport of organisms by density-driven
    - circulation, 292–5
  - tropical regions, 360–2
- remote sensing. *See* satellite remote sensing
- research vessels, 19–20
- residence time
  - phytoplankton above critical depth, 143
- residual circulation, 86–7, 287
  - residual flow model, 86
- residual currents
  - from tracer distributions, 86
- residual transport
  - surface waves, 94
- resonance, 85
- respiration
  - cellular, 133
  - net community, 134
- Reynolds number, 105
  - at microscale, 123
  - fish, 165
  - for plankton, 106
  - mesozooplankton, 165
- Reynolds stress
  - components, 109
  - formal derivation of components, 110

- relating to mean flow, 110
- tensor, 109
- Reynolds, Osborne, 107
  - pipe flow studies, 105
- Rhine ROFI, 270–2
  - semi-diurnal behaviour, 286
  - buoyancy input to the North Sea, 36
  - coastal buoyancy current, 270
  - nitrate, 299
  - progressive wave tide, 283
  - river discharge rate, 36
  - salinity and nutrients, 271
  - sub-tidal flow, 273
- Richardson number, 176
  - control of stability functions, 208
  - derivation, 115–18
  - flux form, 118, 121
  - gradient form, 118
  - parameterisations of eddy coefficients, 125
- Riley, Gordon, 9, 12, 191
- river plumes. *See* Region of Freshwater
  - Influence (ROFI), coastal buoyancy current
- ROFI. *See* Region Of Freshwater
  - Influence (ROFI)
- Rossby number
  - criterion for geostrophy, 321
- Rossby radius of deformation, 73, 78
- salinity
  - measurement, 10, 11
  - units, 11
- satellite remote sensing, 20–1
  - chlorophyll, 339
  - chlorophyll *a*, 249
  - chlorophyll in upwelling regions, 331
  - infra-red mapping of sea surface temperature, 30
  - ocean colour and chlorophyll, 20
  - primary production, 3, 136
  - SAR images of internal waves, 103
  - synthetic aperture radar (SAR), 21, 324
  - temperature, 20
  - tidal mixing fronts, 249
- sea surface micro-layer, 30
- seabirds
  - at tidal mixing fronts, 256
- seasonal cycle
  - freshwater input, 38
  - heat budget, 33
  - heat fluxes and budget, 32–4
  - heating and cooling, 26–36
  - heat storage, 189
  - heating and cooling, 25
  - potential energy anomaly model, 179
  - temperature in mixed region, 187
  - temperature in stratified region, 187
  - two mixed layer model of heating-stirring, 185–90
- SEEP (Shelf Edge Exchange Processes experiment), 344
- sensible heat, 30
- shear dispersion, 113–15
  - horizontal diffusivity, 114
  - maximum horizontal diffusivity, 115
- shear profilers, 210–11
- shelf edge
  - band of cooler water, 327
  - band of increased chlorophyll, 329
  - biogeochemical fluxes, 330
  - carbon export, 344–5
  - cold water coral, 344
  - cross-slope transport mechanisms, 313–21
  - enhanced mixing, 326–9
  - fisheries, 338, 343–4
  - future challenges, 364–6
  - isopycnals spread due enhanced mixing, 327
  - nutrient supplies to support shelf sea export production, 330
  - organic matter export, 364
  - phytoplankton community, 341–3
  - primary production, 338
- shelf seas
  - contrasts with open ocean, 1–4, 304–6
  - contribution to global carbon export, 4
  - contribution to global fish catch, 4
  - contribution to global primary production, 3
  - definition, 1
  - economic value, 4–5
  - fishing pressure, 4–5, 6–7
  - future challenges, 23
  - pressures from humans, 4–5
  - resources, 4
- silicate, 13, 144, 145, 147
  - source to the ocean, 330
  - supply by rivers, 298
- Simpson-Hunter parameter (*SH*)
  - choice of current speed, 180–1
  - frontal position, 180
  - mapping in different regions, 182
  - test using frontal positions, 181–3
- slope current, 307–8. *See also* JEBAR
  - carbon export, 347
  - continuity, 308
  - deflection in bottom boundary layer, 316
  - eddies, 320–1
  - forcing, 311–13
  - Hebridean shelf edge, 308
  - joint effect of baroclinicity and relief (JEBAR), 311
  - measurements with drogue buoys, 308
  - seasonal change, 313
  - spatial consistency, 308
  - steadiness, 308
  - wind stress forcing, 313

- solar energy input
  - average attenuation coefficient, 28
  - total energy absorption, 28
- solar heating, 26–9
- spectral energy distribution, 26
- Spencer Gulf, S. Australia, 38
  - draining of high salinity water, 320
- spring bloom, 9, 191–3
  - control by spring-neap tidal cycle, 196
  - diatoms, 194
  - importance to annual primary production, 217
  - match-mismatch hypothesis, 195–6
  - mechanism, 191–2
  - observations, example, 192
  - phytoplankton species, 193–5
  - rates of primary production, 192–3
  - timing, 195–7
- spring-neap cycle, 46
  - contribution to pycnocline mixing, 215
  - internal tide and nitrate flux, 340
  - mixing at the thermocline, 222
  - spring bloom timing, 196
  - variation in tidal mixing, 235
- stability functions
  - for local equilibrium, 208
- stability of the water column, 68
- standing wave, 77
- Stanton number, 30
- Stefan's law, 29
- stirring and mixing, 106
- stirring by wind and tide
  - two mixed layer model, 186
- stirring power
  - variation over spring-neaps cycle, 235
- Stokes drift, 94
- Stokes wave, 94
- Stommel, Henry, 9
- strain induced periodic stratification (SIPS), 281
  - condition for periodic stratification, 282
- stratification
  - barrier to bottom water oxygen replenishment, 299, 346
  - bottom layer inorganic carbon, 346
  - climate change, 368
  - criterion for seasonal stratification, 180
  - cycle due to tidal straining, 283
  - effect on density-driven circulation, 287
  - estuarine, 275–7
  - induced by estuarine circulation, 269
  - inhibition of mixing, 175, 207, 238
  - primary production in ROFIs, 295–8
  - spring onset, 191
  - suppression of vertical motion, 52
  - trigger for phytoplankton growth, 192
- subsurface chlorophyll maximum (SCM)
  - contrasts with open ocean, 224
  - effects of spring-neap cycle, 222–3
  - examples, 218
  - importance of interior mixing, 219
  - importance to annual primary production, 223–4
  - light, 223
  - nitrate flux, 221
  - phytoplankton species, 226–7
  - possible mechanisms, 218
  - primary production rates, 223
  - role of phytoplankton swimming, 225–6
  - shelf edge, 341
  - targeting by zooplankton, 228–9
- surface boundary layer
  - cross-slope transport, 314–16
- surface microlayer, 32
- surface waves
  - boundary conditions, 91
  - energy propagation, 95–6
  - finite amplitude, 93–5
  - kinetic energy, 95
  - orbital motion, 93
  - potential energy, 95
  - pressure variation, 92
  - residual transport, 94
  - velocity components, 92
  - velocity potential, 90–2
  - wave breaking, 96–9
  - whitecapping, 96
- Sverdrup, Harald, 142, 191
- Synthetic Aperture Radar
  - internal wave observations, 324
- Taylor hypothesis, 121, 124
- Taylor, Geoffrey, 8, 82, 113
- Taylor-Proudman theorem, 306–7
- Thailand, Gulf of, 361
- thermal wind, 65–6. *See also* geostrophic flow
- thermocline. *See also* stratification
  - carbon export, 345–7
  - inhibition of turbulence, 192, 346
- thin layers
  - phytoplankton, 226
  - zooplankton, 228
- tidal ellipse
  - in Rhine ROFI, 285
  - response to changes in stratification, 285
- shear dispersion, 115
- variability in Rhine ROFI, 283–7
- tidal forcing, 25
- tidal mixing (TM) front
  - baroclinic instability, 245–7
  - basking sharks, 256
  - change in density gradients with depth, 241
  - chlorophyll distribution, 249
  - condition for location, 180
  - difference in sea surface elevation, 241
  - enhanced primary production, 251
  - equilibrium adjustment over spring-neap cycle, 235–7

- fisheries, 256–62
- flow direction, 241
- frontal jet, 241
- geostrophic balance, 241–2
- gyre circulation and larval retention, 257–60
- horizontal temperature gradients, 234
- infra-red imaging of temperature, 238, 245, 249
- larger marine animals, 256
- measurements of residual velocity, 242
- mechanisms for enhanced chlorophyll, 251
- mechanisms for primary production, 251–4
- mixing efficiency and spring-neap adjustment, 238
- nitrate distribution, 249
- non-equilibrium adjustment, 238
- ocean colour imaging of chlorophyll *a*, 249
- positions from satellite imagery, 233–4
- seabirds, 256
- seasonal adjustment, 239
- since the last ice age, 368
- spring-neap adjustment and primary production, 252
- spring-neaps adjustment, 235–9
- stored buoyancy and spring-neap adjustment, 237
- transverse circulation, 247–8
- velocity from density section, 241
- zooplankton, 254–6, 257
- tidal rectification, 245, 260
- tidal stirring
  - for  $M_2$  constituent, 179
- tidal straining, 281
  - contribution to the potential energy anomaly, 282
  - convective mixing, 281, 283
  - enhancement of estuarine circulation, 288
  - impact on sediment transport, 288
  - Liverpool Bay ROFI, 283
  - mechanism, 281–3
  - over-straining, 283
  - primary production, 296–8
  - Rhine ROFI, 283–7
- tides
  - amphidrome, 80
  - amphidromic systems, 80–3
  - amplitude at the head of a gulf, 85
  - angular momentum of Earth-Moon system, 48
  - centre of gravity of the Earth-Moon system, 40
  - centrifugal force, 42
  - co-range lines, 80
  - current amplification at the shelf edge, 79
  - degenerate amphidrome, 81
  - difference between the gravitational and centrifugal forces, 42
  - diurnal and semi-diurnal constituents, 45
  - effect of the sun, 40
  - equilibrium theory of the tides, 43
  - equilibrium tide range, 43
  - frictional losses in a gulf, 80
  - global distribution tidal energy dissipation, 46
  - harmonic method of tidal analysis, 45
  - Moon's gravitational attraction, 40, 42
  - Moon's orbital acceleration, 48
  - neap tides, 46
  - power supply from Moon and Sun tides, 46
  - slope forcing in model, 206
  - spring tides, 46
  - tidal constituents, 44–6
  - tidal ellipses, 82
  - tidal energy, 25
  - tidal energy dissipation
    - in the past, 366–8
  - tidal energy supply to shelf seas, 46–7
  - tidal forcing, 40–6
  - tidal resonance in a gulf, 84–6
  - tidal species or bands, 44
  - tidal wave behaviour crossing the shelf edge, 79–80
  - tidal waves, 46, 77
  - tide generating force (TGF), 40, 42
  - tide generating force horizontal component, 43
  - tide generating force variation around globe, 43
  - tide generating force vertical component, 43
  - wave amplification across the shelf edge, 79
  - wave amplification and reflection, 79–84
  - wave reflection at the head of a gulf, 81
- top-down control, 168
- total derivative, 57
- towed vehicles, 11
- trophic levels, 151
- tsunami waves, 77
- turbulence
  - billows in the atmosphere and ocean, 119
  - criterion for development from
    - a laminar flow, 118
  - criterion for maintenance, 118
  - development of billows in a shear flow, 118
  - diffusivity dependence on scale, 113
  - dispersive tendency, 106
  - energetics, 115–20
  - energy dissipation, 106–7
  - intensity, 207
  - Kolmogorov constant, 123
  - Kolmogorov microscales, 123
  - laboratory experiments on billows, 118
  - length scale, 207, 208
  - local equilibrium, 121
  - mean and turbulent flow components, 107–8
  - momentum flux component, 109
  - Ozmidov length, 125
  - power available in a shear flow, 117
  - predator–prey encounter rates, 165, 262
  - properties and relation to mixing, 105–7
  - Reynolds and Richardson numbers, 120
  - scales of turbulence, 121–5
  - spectrum, 121
  - Taylor hypothesis, frozen field, 121, 124

- turbulence (cont.)
  - Thorpe scale, 125
  - turbulent Prandtl number, 118
  - vertical mixing time scale, 114
- turbulence closure, 125–6
  - length scale, 125
  - stability functions, 126
- turbulence closure (TC) model, 205–9
  - eddy viscosity and diffusivity, 207
  - missing mixing, 213–14
  - tidal forcing, 208
- turbulence closure scheme, 207–8
  - feedback mechanisms, 208
- turbulent dissipation
  - delay of maximum with height
    - above seabed, 212
  - in the thermocline, 213
  - levels in shelf seas, 213
  - model-observation comparison, 210–13
  - quarter diurnal variation, 212
  - relation to velocity shear, 210
  - shear profiler measurement, 210
  - tidal cycle in a ROFI, 283
- turbulent flow, 105
- turbulent fluxes
  - of momentum, 109–10
  - of scalar properties, 107–8
- turbulent kinetic energy (TKE), 120
  - diffusion, 120
  - dissipation, 120
  - energy cascade, 123
  - equation, 120–1, 207
  - equilibrium spectrum, 122–4
  - inertial sub-range, 123
  - inertial sub-range in spectrum, 123
  - production, 117
  - scalar energy spectrum, 122
  - total water column production, 178
  - universal equilibrium spectrum, 124
- turbulent stress, 69
- Turner, J. S.
  - tank experiment and grid turbulence, 175
- two mixed layer (TML) model, 185–90
- upwelling
  - at shelf edge, 314–15
  - biogeochemical response, 331–3
  - carbon export, 345
  - filaments and eddies, 314, 331, 337
  - fisheries, 334–5
  - front, 314
  - frontal instabilities, 314
  - hypoxia in bottom waters, 335–6
  - in Rhine ROFI, 286
  - nitrate supply, 330, 331–3
  - nutrient supply, 75
  - sections of temperature, nitrate
    - and chlorophyll, 331
  - surface chlorophyll, 331
  - surface temperature, 331
  - time scale of phytoplankton response, 332
  - time scale of zooplankton response, 333
  - under western boundary current, 337
  - zooplankton response, 333–4
- upwelling/dwelling, 74–5
- Ushant tidal mixing front, 245
- variable-efficiency model
  - S–N frontal adjustment, 239
- velocity potential
  - internal waves, 99–100
  - surface waves, 90–2
- vertical migration (of zooplankton), 162, 228
- vertical mixing
  - by waves in ROFIs, 290
  - induced by air bubbles in laboratory
    - experiments, 280
- vertical motions in the ocean, 52
- vertical oscillations, 67–8
- vorticity, 91
- waves. *See also* deep water waves, internal waves, Kelvin waves, long waves, surface waves
  - angular frequency, 92
  - group velocity definition, 96
  - phase speed, 92
  - wavenumber, 92
- western boundary current
  - trapping on continental slope, 311
- whitecapping, 96. *See* wave breaking
- wind stirring
  - contribution in seasonally stratifying regions, 183–5
  - contribution to mixing in ROFIs, 285
- wind stress, 25, 39
  - boundary condition in model, 206
  - drag coefficient, 39, 184
  - drag coefficient variation with stratification and
    - wind speed, 40
  - drag law, 39
  - effects in ROFIs, 288–90
  - primary production after a storm, 198
  - rate of working, 183
  - variability and primary production, 221
- Woods Hole Oceanographic Institute, 6, 7
- Yellow Sea
  - co-tidal charts, 82
- zooplankton
  - at a tidal mixing front, 254–6
  - dispersion and life history, 257, 333
  - patchiness, 257