

INDEX

- AAP. *See* American Academy of Pediatrics
AAPCC. *See* American Association of Poison Control Centers
abdomen, evaluation of, 10–11. *See also*
 umbilical area, evaluation of
 for kidneys, 10
 for liver, 10
 for spleen, 10
 for umbilical area, 10–11
 Eagle-Barrett triad in, 11
 hernias in, 11
 hygiene for, 10
 omphalitis in, 10–11
 patent urachus in, 11
 Prune Belly Syndrome in, 11
abdominal breathing, 9
absence seizures, 142
acetaminophen, 28
 recommended dosing for, 28, 296
 as toxic, 296–297
 NAC for, 296–297
 treatment for, 296–297
N-acetylcysteine (NAC), 296–297
acne, neonatal, 183–184
acrocyanosis, 14
 of infancy, 190
Acropustulosis of infancy, 184
ACS. *See* acute chest syndrome
activated charcoal, 295
 for salicylate poisoning, 298
acute chest syndrome (ACS), 240–241
 definition of, 240
 management of, 241
 mortality rate for, 240
 pathophysiology for, 240
 symptoms of, 240–241
acute gastroenteritis (AGE), 215–217
 antibiotic therapy for, 218
 complications from, 217
 dehydration from, 215–216
 oral replacement therapy for, 218
 rehydration for, 216, 217
acute mastoiditis (AM), 89
acute otitis media (AOM), 83–88
 common pathogens for, 83
 PCV-7 and, 83
 differential diagnosis of, 85, 87
 excessive crying in infants from, 71
 follow-up for, 88
 lab work for, 85
 management of, 85–88
 HD amoxicillin in, 86–87
 infant age as factor in, 85–86
 maculopapular allergies from, 88
 pain, 88
 primary goals of, 85
 treatment failure in, 87–88
 “watchful waiting” strategy in, 87
 OE and, 89
 otorrhea and, 88
 radiology for, 85
 symptoms of, 83–85
 middle ear complications in, 85
 otalgia as, 83
 SMAOM guidelines for, 83–84, 87
 TM examination for, 84–85
 tympanostomy tubes and, 88
 URIs and, 83
 viral causes of, 83
acute vasoocclusive crisis. *See* vasoocclusive crisis, acute
acyanotic heart disease, 165–166, 167
 airway management in, 174
 ECG for, 167
adrenal insufficiency, 259–260
 CAH and, 12, 260
 symptoms of, 260
 differential diagnosis for, 260
 lab work for, 260
 symptoms of, 260
 treatment of, 260
 with corticosteroids, 261
AGE. *See* acute gastroenteritis
AHA. *See* American Heart Association
airway management, pediatric, 33–39. *See also*
 laryngeal mask airway; rapid-sequence intubation, pediatric
anatomy and, 33
 adult v. children, 33
 bag-mask-valve ventilation in, 33–34
 American Heart Association guidelines for, 34
 complications of, 34
 sizing for, 33–34
 for burn injuries, 203
 clinical scenarios for, 34
 failed intubation in, 38
 LMA and, 38
 with foreign bodies, 37–38
 Heimlich maneuver for, 38
 indications for, 34
 needle cricothyrotomy in, 39
 complications of, 39
 membrane identification in, 39
 ventilation methods for, 39
 with rapid-sequence intubation, 34–37
 assessment of, 36–37
 medications for, 37
 nasotracheal, 36
 patient weight and, 37
 preoxygenation for, 34
 procedures for, 37
 tube sizing in, 34–36
 suggested equipment for, 37
alagille syndrome, 134
ALCAPA syndrome. *See* anomalous origin of the left coronary artery syndrome
Alert/Verbal/Painful/Unresponsive (AVPU) scale, 293
ALTE. *See* apparent life threatening events
AM. *See* acute mastoiditis
ambiguous genitalia, 12
 females with, 12
 CAH and, 12
 males with, 12
American Academy of Pediatrics (AAP)
 Committee on Fetus and Newborn, 18
 minor head injury guidelines of, 323
 Policy Statement on Prevention and Management of Pain in the Neonate, 26
 SIDS recommendations by, 52–53, 61
 toxicologic emergency guidelines for, 295
American Academy of Toxicologists, 295
American Association of Poison Control Centers (AAPCC), 291
American College of Emergency Physicians, 113
American Heart Association (AHA)
 bag/mask ventilation guidelines of, 34
 pediatric resuscitation guidelines of, 40, 44
amnesia. *See* post-traumatic amnesia
analgesia, 26–29
 definition of, 27
 PSA and, 30
analgesics, 28–29
 non-opioid, 28–29
 acetaminophen, 28
 NSAIDs, 28–29
 opioid, 29
 dosing guidelines for, 29

- analgesics (*cont.*)
 meperidine, 29
 morphine, 29
 anomalous origin of the left coronary artery
 (ALCAPA) syndrome, 169
 anterior temporal lobectomy, 154
 antibiotics
 for AGE, 218
 for bone/joint infections, 115
 for cervical lymphadenitis, 123
 for infant fevers, 111
 for meningitis, 114
 for neonatal fevers, 109–110
 for pertussis, 285
 anticholinergic medications, 78
 anticholinergics, 286–287
 antifibrinolytic therapy, 250
 AOI. *See* apnea of prematurity
 AOM. *See* acute otitis media
 aortic stenosis (AS), 163, 169
 Apert's syndrome, 6
 aphthous stomatitis, 183
 aplastic crisis, 237
 apnea, 9. *See also* apnea of prematurity;
 apparent life threatening events
 ALTE and, 61–67
 definition of, 61
 diagnoses of, 63–64
 for central apnea, 63
 for obstructive apnea, 63
 GERD and, 63
 SIDS and, 17–18
 apnea monitors, 17–18
 ALTE and, 18, 67
 American Academy of Pediatrics Committee
 on Fetus and Newborn on, 18
 Collaborative Home Infant Monitoring
 Evaluation for, 18
 apnea of prematurity (AOI), 17–18. *See also*
 apnea monitors
 ALTE from, 66
 devices for, 17–18
 epidemiology of, 17
 management of, 17–18
 with monitors, 17–18, 67
 methylxanthine therapy in, 17
 with caffeine, 17
 pathophysiology of, 17
 for brainstem development, 17
 for respiratory system, 17
 SIDS and, 17–18
 apparent life threatening events (ALTE), 18,
 61–67
 from AOI, 66
 home monitors for, 18, 67
 apnea and, 61–67
 definition of, 61
 diagnoses of, 63–64
 cardiac causes of, 65
 HRV issues, 65
 from child abuse, 64
 Münchausen by proxy, 64
 definition of, 61
 GERD and, 62–63
 hematological causes of, 65
 from IEM, 65
 incidence rates of, 61
 from infections, 64–65
 bordetella pertussis, 65
 RSV, 64–65
 lab evaluation of, 66
 management of, 66–67
 from MAO, 65–66
 neurological causes of, 63–64
 BHS, 64
 intracranial pressure, 64
 seizures, 63–64
 from non-accidental trauma, 66
 presentation of, 62
 radiology evaluation of, 66
 SIDS v., 61
 from toxins, 66
 appendicitis, 212–213
 lab work for, 212
 physical examinations for, 212
 radiography for, 212–213
 symptoms of, 212
 treatment for, 213
 Apt-Downey test, 213–214
 arthritis. *See* septic arthritis
 AS. *See* aortic stenosis
 ASD. *See* atrial septic defects
 asystole, 45
 definition of, 45
 management of, 45
 atonic seizures, 142
 atopic dermatitis, 193–195
 pityriasis alba and, 184
 atrial septic defects (ASD), 166–167
 atrioventricular septal defects (AVSDs),
 10
 Down syndrome and, 10
 AVPU scale. *See*
 Alert/Verbal/Painful/Unresponsive
 scale
 AVSDs. *See* atrioventricular septal defects
 bacterial tracheitis, 272–273
 complications from, 273
 diagnosis of, 273
 lab work for, 273
 radiography for, 273
 symptoms of, 273
 treatment of, 273
 BAER. *See* brainstem auditory evoked
 responses
 bag-mask-valve ventilation, 33–34
 American Heart Association guidelines for,
 34
 complications of, 34
 sizing for, 33–34
 balanitis, 222
 balanoposthitis, 222
 barbiturates, 30
 Barlow tests, for DDH, 343
 Beckwith-Wiedemann syndrome, 4
 benign familial neonatal convulsions (BFNC),
 145
 benign idiopathic neonatal convulsions
 (BINC), 145
 benzodiazepines, 30
 diazepam, 144, 149
 lorazepam, 149
 midazolam, 149
 BFNC. *See* benign familial neonatal
 convulsions
 BHS. *See* breath holding spells
 bicarbonate therapy, for DKA, 266
 biliary atresia, 134, 207–209
 diagnosis of, 208–209
 with ultrasonography, 209
 forms of, 207–208
 incidence rates for, 207
 symptoms of, 208
 bilirubin
 increased indirect, 132
 unconjugated, 128–129
 BINC. *See* benign idiopathic neonatal
 convulsions
 Blalock-Taussig shunt, 175
 bleeding disorders, inherited, 242–253
 clotting factor deficiencies, 244, 247
 differential diagnoses for, 247
 factor replacement guidelines for, 250
 factor XI deficiency, 244
 hemophilia, 243–244, 245–246
 CNS and, 245
 genitourinary tract bleeding from, 246
 GI bleeding from, 246
 in head and neck, 246
 hemarthrosis, 246
 hemostasis and, 243
 incidence rates for, 243
 in intra-abdomen, 246
 posttraumatic bleeding and, 246
 by sex, 243
 skeletal muscle and, 246
 symptoms of, 243–244, 245
 lab work for, 246–248
 bleeding time in, 247
 factor level assays in, 248
 platelet count in, 247
 PT, 247
 management of, 248–251
 in ED, 248–249
 with factor treatment products, 249
 prevention of, 253
 for VWD, 253
 prophylaxis for, 253
 radiology for, 248
 treatment of, 250, 251–253
 with antifibrinolytic therapy, 250
 blood product safety and, 252
 complications of, 252–253
 with desmopressin, 249–250
 fractures and, 251
 for GI bleeding, 251
 for head and neck bleeding, 251
 for hemarthrosis, 251
 for hematomas, 251
 for hematurias, 251
 with hemostatic blood products,
 250–251
 of patients with inhibitors, 252–254
 for severe hemorrhage, 251

- surgery as, 252
 - wound management and, 251
 - vitamin K deficiency, 244–245
 - prevention of, 249
 - symptoms of, 246
 - VWD, 244
 - desmopressin for, 253
 - etiology of, 244
 - prevention of, 253
 - ristocetin cofactor assay for, 248
 - symptoms of, 245, 246
 - treatment of, 253
 - types of, 244
 - bleeding time, 247
 - blisters, friction/sucking, 178–179
 - blood products
 - hemostatic, 250–251
 - safety issues with, 252
 - blood urea nitrogen (BUN), 258
 - blue sclera, 6
 - Ehlers-Danos syndrome and, 6
 - Bohn nodules, 8
 - bone/joint infections, 114–115
 - antibiotic therapy for, 115
 - blood testing for, 115
 - causes of, 114–115
 - fevers from, 114–115
 - radiographs for, 115
 - septic arthritis and, 115, 344
 - bordetella pertussis. *See* pertussis
 - Boston criteria, 107–108
 - bowing fractures, 336
 - brachial plexus injury, 14
 - Erb palsy and, 14
 - Klumpke palsy and, 14
 - bradycardia, resuscitation and, 44–45
 - evaluation of, 44
 - management of, 45
 - brainstem auditory evoked responses (BAER), 199–200
 - branchial cleft anomalies, 8–9, 122–123
 - excision of, 123
 - Goldenhar syndrome as, 9
 - location of, 123
 - Pierre-Robin Association as, 9
 - breast enlargement, 9
 - breastfeeding
 - colic and, 70
 - crying in infants from, from toxins, 73–74
 - diet modification of mothers and, 79
 - FTT and, in management of, 102
 - hyperbilirubinemia and, 137
 - increased indirect bilirubin and, 132–133
 - for pain management, 27
 - breath holding spells (BHS)
 - as ALTE, 64
 - during seizures, 146
 - breathing. *See* abdominal breathing; periodic breathing
 - bronchiolitis, 279–281
 - causes of, 279
 - epidemiology of, 279
 - hospital admission criteria for, 281
 - lab work for, 279–280
 - prevention of, 280–281
 - from RSV, 279
 - symptoms of, 279
 - treatment for, 280
 - adjunctive therapies for, 280
 - with β_2 bronchodilators, 280
 - with hypertonic saline, 280
 - with nebulized epinephrine, 280
 - for steroids, 280
 - β_2 bronchodilators, 280
 - bruising, 77, 313
 - on back, 77
 - Brushfield spots, 6
 - Down syndrome and, 6
 - buckle fractures. *See* torus fractures
 - BUN. *See* blood urea nitrogen
 - bundling, 3
 - burn injuries, 200–204
 - classification of, 201–204
 - from child abuse, 203–204, 315–316
 - electrical, 203
 - incidence rates for, 201
 - management of, 203–204
 - burn central referral criteria in, 201
 - in ED, 202–203
 - field resuscitation in, 201–202
 - fluid resuscitation in, 203
 - with Lund and Browder chart, 201
 - prevention of, 200–204
 - types of, 201
 - caffeine, 17
 - CAH. *See* congenital adrenal hyperplasia
 - calcium channel blockers, 302–303
 - CAM. *See* complementary and alternative medicine, excessive crying in infants and
 - camphor, 304
 - Canadian Assessment of Tomography for Child hood Head Injury (CATCH), 330
 - Canadian CT Head Rule (CCHR), 329
 - candidiasis, 186
 - capnography, 36
 - caput succedaneum, 5
 - from vacuum extraction, 5
 - carbamazepine, 150
 - cardiac emergencies, in infants, 159–176.
 - See also* carditis; cyanotic heart disease
 - cardiac disease and, 159
 - cardiomyopathies and, 173
 - congested/dilated, 173
 - hypertrophic, 173
 - restrictive, 173
 - carditis in, 169–172
 - endocarditis, 171–172
 - myocarditis, 169
 - pericarditis, 169–171
 - chest X-rays for, 159–160
 - congestive heart failure and, 174
 - treatment of, 174
 - cyanotic heart disease in, 162–163, 169
 - AS and, 163, 169
 - acyanotic v., 165–166, 174
 - airway management for, 173
 - ALCAPA syndrome and, 169
 - ASD and, 166–167
 - cardiogenic shock and, 163
 - coarctation of the aorta and, 163, 168
 - ductal dependent lesions and, 173–174
 - endocardial cushion defect and, 167–168
 - HLHS and, 163, 168–169
 - Hypoplastic Right Heart Syndrome and, 163
 - hypoxemic episodes for, 163
 - pathophysiology of, 162–163
 - PDA and, 167
 - prophylaxis for, 172
 - pulmonary atresia and, 163
 - TA and, 163, 165
 - TAPVR and, 163, 164–165
 - TGA and, 163, 164
 - TOF from, 163–164
 - truncus arteriosus and, 163, 165
 - VSD and, 167
 - ECG for, 159–160
 - heart chamber size and, 161–162
 - heart rate and, 160
 - Kawasaki's disease in, 172–173
 - presentation of, 173
 - treatment of, 174–175
 - PR intervals during, 161
 - QRS duration and, 161
 - surgery for, 174, 175–176
 - Blalock-Taussig shunt, 175
 - Fontan operation, 175
 - Mustard and Senning operation, 175
 - Norwood operation, 175
 - post-operative complications of, 175–176
 - Rashkind procedure, 175
 - Rastelli procedure, 175
 - T wave morphology in, 161
 - ventricle axis dominance and, 161
- cardiac pulmonary resuscitation (CPR), 40
- AHA guidelines for, 44
- cardiac tamponade, 170–171
- cardiomyopathies, 173
 - congested/dilated, 173
 - hypertrophic, 173
 - ECG for, 173
 - restrictive, 173
- cardiovascular systems, evaluation of, 9–10.
 - See also* asystole; murmurs; pulseless electrical activity; tachycardia; tachydysrhythmias
- murmurs in, 10
 - AVSD and, 10
 - holosystolic, 10
 - PDA, 10
 - sound of, in diagnosis of, 10
- in pediatric resuscitation, 43–48
 - for advanced support, 44–48
 - for asystole, 45
 - basic support for, 43–44
 - for bradycardia, 44–45
 - for PEA, 45
 - for tachycardia, 45, 46–47
 - for tachydysrhythmias, 47–48
 - for ventricular fibrillation, 45–46
- post-birth transition for, typical, 9–10
- vascular access in, 351–354
 - complications of, 352

- cardiovascular systems (*cont.*)
 - for femoral vein, 352
 - for internal jugular vein, 352
 - for intraosseous line, 352–353, 354
 - peripheral, 351–352
 - saphenous vein cutdown in, 352
 - for subclavian vein, 352
 - for umbilical vein, 354
- carditis, 169–172
 - endocarditis, 171–172
 - categorization of, 172
 - causes of, 171
 - Duke criteria for, 171
 - ECG for, 171
 - lesions from, 191
 - management of, 171–172
 - prophylaxis for, 172
 - myocarditis, 169
 - causes of, 169
 - symptoms of, 169
 - pericarditis, 169–171
 - cardiac tamponade and, 170–171
 - cause of, 169
 - presentation of, 169–170
- casein hydrolysate formula, 79
- cataracts, congenital, 6
- CATCH. *See* Canadian Assessment of Tomography for Child hood Head Injury
- CCHR. *See* Canadian CT Head Rule
- CDC. *See* Centers for Disease Control
- CDH. *See* congenital diaphragmatic hernia
- cellulitis, 24
- Centers for Disease Control (CDC), 93
 - on OTC medications poisonings, 301
- central apnea, 63
- central hypotonia, 13
 - Down syndrome and, 13
 - Prader-Willi Syndrome and, 13
- central nervous system (CNS)
 - hemophilia and, 245
 - trauma to, 331
- central pontine myelinolysis, 267–268
- cephalhematomas, 5
- cerebral edema, 266–267
- cerebrospinal fluid (CSF) shunts, 18–21
 - components of, 18
 - infection from, 20–21
 - etiology of, 20
 - lab work for, 20–21
 - radiology for, 21
 - symptoms of, 20
 - treatment therapies for, 21
 - leaks in, 21
 - malfunction of, 18–20
 - assessment of, 19–20
 - NCCT and, 19–20
 - shunt tapping and, 20
 - overdrainage by, 21
 - pseudocyst formation from, 21
 - purpose of, 18
 - seizures from, 21
 - ventricular/vascular, 21
 - visceral damage from, 21
- cerebrovascular accidents, 239
 - SCI, 239
 - strokes, 239–240
 - management of, 240
- cervical lymphadenitis, 123–124
 - antibiotic therapy for, 123
 - causes of, 123
 - bacterial infection as, 123
 - lymphadenopathy and, 123
- CF. *See* cystic fibrosis
- CHALICE. *See* Children’s Head Injury Algorithm for the Prediction of Important Clinical Events
- CHD. *See* congenital heart disease
- chest, evaluation of, 9
 - breast enlargement and, 9
 - clavicle fractures and, 9
 - mastitis in, 9
- chest X-rays, for infant cardiac emergencies, 159–160
- chicken pox. *See* varicella
- child abuse. *See also* shaken baby syndrome
 - ALTEs and, 64
 - burn injuries from, 203–204, 315–316
 - excessive crying in infants and, 77–78
 - in drowning/submersion injuries, 198
 - excessive crying in infants from, 77–78
 - bruising and, 77, 313
 - burn injuries and, 77–78
 - fractures and, 78
 - head/scalp injuries and, 78
 - fractures from, 313–314
 - excessive crying in infants and, 78
 - Münchausen by proxy as, 64
 - respiratory emergencies as result of, 271
 - sexual abuse as, 315–318
 - diagnosis of, 315–316
 - evaluation of, 318
 - evidence testing for, 316–317
 - radiographic evidence for, 318
 - victim history in, 317–318
 - SIDS as, 52, 319
 - infant/neonatal mortality and, 52
 - trauma from, 307–319
 - abdominal/thoracic injuries as, 312–313
 - age as factor in, 308–309
 - burns as, 315–316
 - caregivers’ role in, 309
 - categories of, 307–308
 - cutaneous manifestations of, 315–316
 - death review teams for, 319
 - from falls, 314–315
 - family history characteristics and, 307–309, 311
 - incidence rates of, 307
 - iTBI as, 309
 - morbidity rates of, 308
 - mortality rates of, 308
 - physical examinations for, 312
 - radiographic evaluation of, 315
 - SBS as, 309–311
 - SIDS as, 52
- Children’s Head Injury Algorithm for the Prediction of Important Clinical Events (CHALICE), 330
- CHIP study. *See* CT in head injury patients study
- chlamydia, 283–284
 - conjunctivitis from, 7
 - epidemiology of, 283
 - management of, 284
 - presentation of, 283
 - radiography for, 283–284
- chloral hydrate, 30
- choledochal cysts, 134
- circumcision, complications from, 223
 - meatal stenosis from, 223
 - Plastibell method of, 223
- cleaning agents, poisoning from, 301–302
 - epidemiology of, 301
 - under PPA, 301
 - symptoms of, 302
- cleft lip/palate, 8
- clonazepam, 152
- clonidine, 303
- clotting factor deficiencies, 244, 247
- clubfoot, 12
- CMT. *See* congenital muscular torticollis
- CNS. *See* central nervous system
- coagulation disorders. *See* clotting factor deficiencies
- coarctation of the aorta, 163, 168
 - PDA and, 168
- coin rubbing, lesions from, 192
- colic, 70–71, 79–80. *See also* crying in infants, excessive
 - breastfeeding and, 70
 - from excessive colonic gas, 70
 - from GERD, 70
 - maternal behavior and, 70–71
 - presentation of, 71
- Collaborative Home Infant Monitoring Evaluation, 18
- complementary and alternative medicine (CAM), excessive crying in infants and, 79
- complete blood counts (CBC), for excessive crying in infants, 78
- computed tomography (CT)
 - for head and neck emergencies, 122
 - for hemangiomas, 125
 - for meningitis, 113–114
 - for minor head injuries, 325–326
 - risk factors for, 326
 - NCCT, 19–20
 - for neonatal seizures, 145
- congenital adrenal hyperplasia (CAH), 12
 - adrenal insufficiency and, 260
 - symptoms of, 260
- congenital diaphragmatic hernia (CDH), 11, 278–279
 - complications from, 279
 - epidemiology for, 275–278
 - lab work for, 278
 - management of, 278
 - presentation of, 278
- congenital heart disease (CHD), excessive crying in infants and, 74
- congenital hypothyroidism, 132

- congenital muscular torticollis (CMT), 8.
See also neonatal torticollis
- congenital subglottic stenosis, 270–271
- congested/dilated cardiomyopathy, 173
- congestive heart failure, 174
treatment of, 174
- conjugated hyperbilirubinemia, 133
causes of, 130
ED evaluation of, 135–136
- conjunctivitis, 7
bacterial causes of, 7
chemical causes of, 7
chlamydia and, 7
from gonorrhea, 7
Herpes simplex and, 7
- constipation, 217–219
causes of, 217–219
definition of, 217
NAPSGHAN guidelines in, 217
differential diagnosis for, 219
evaluation for, 219
treatment for, 219
- contact diaper dermatitis, 73, 183, 185–186
- corpus callostomy, 154
- Coxsackie virus, 182–183
- CPR. *See* cardiac pulmonary resuscitation
- cranial nerves, 12, 13
- craniosynostosis, 5
Apert’s syndrome and, 6
Crouzon’s disease and, 6
FGFR and, 6
as inherited, 5–6
- crawling reflex, 13
- Crigler-Najjar syndrome, 132
type I, 132
type II, 132
- croup, 273–275
cause of, 273–274
complications of, 275
lab work for, 274
severe, 275
symptoms of, 274
treatment for, 274–275
with glucocorticoids, 274
with mist therapy, 275
with racemic epinephrine, 274–275
- Crouzon’s disease, 6
- crying in infants, excessive, 69–80
child abuse and, 77–78
bruising and, 77, 313
burn injuries and, 77–78
fractures and, 78
head/scalp injuries and, 78
colic and, 70–71, 79–80
breastfeeding and, 70
from excessive colonic gas, 70
from GERD, 70
maternal behavior and, 70–71
presentation of, 71
differential diagnoses for, 71–72, 78
CHD in, 74
eye examinations in, 75
feeding types in, 74
immunizations in, 74
infection and, 71–73
insect bites and, 74–75
maternal medications in, 74
medications as factor in, 74
for metabolic disorders, 76–77
for normal v. excessive, 71
for pathological crying, 71
pneumonic in, 72
strangulation in, 76
surgical issues for, 75–76
toxins in, 73–74
trauma in, 73
epidemiology of, 69
Hirschsprung’s disease and, 76
hypoglycemia and, 76–77
incarcerated inguinal hernias and, 76
lab work for, 78
CBC in, 78
urine analysis in, 78
management of, 78–79
with anticholinergic medications, 78
with behavioral interventions, 79
with breastfeeding modifications, 79
with CAM, 79
with defoaming agents, 78
with formula modifications, 78–79
with sucrose, 78
midgut volvulus and, 76
from otitis media, 71
pathophysiology of, 69
stress from, 69
testicular torsion and, 76
- CSF shunts. *See* cerebrospinal fluid shunts
- CT. *See* computed tomography
- CT in head injury patients (CHIP) study, 329
- cuffed tubes
endotracheal, 42
tracheostomy, 22
- cultural beliefs, about grieving, 54, 55
- cupping, lesions from, 192
- cutis marmorata, 14
- cyanotic heart disease, 162–163, 169
AS and, 163, 169
acyanotic v., 165–166, 167
airway management in, 174
ECG for, 167
airway management for, 173
ALCAPA syndrome and, 169
ASD and, 166–167
cardiogenic shock and, 163
coarctation of the aorta and, 163, 168
ductal dependent lesions and, 173–174
ECG findings for, 163
endocardial cushion defect and, 167–168
HLHS and, 163, 168–169
Glenn operation for, 175
Hypoplastic Right Heart Syndrome and, 163
Glenn operation for, 175
hypoxemic episodes for, 163
pathophysiology of, 162–163
assessment of, 162
etiologies of, 162
infant age as factor for, 162–163
presentation of, 162
PDA and, 167
prophylaxis for, 172
pulmonary atresia and, 163
TA and, 163, 165
TAPVR and, 163, 164–165
TGA and, 163, 164
Mustard and Senning operation for, 175
TOF from, 163–164
Blalock-Taussig shunt for, 175
Rastelli procedure for, 175
truncus arteriosus and, 163, 165
VSD and, 167
- cystic fibrosis (CF), 134–135
- cystic hygromas, 124
- cystic lesions, 122
- cysts
choledochal, 134
cystic hygromas, 124
cystic lesions, 122
dermoid, 124
pseudocyst formation, from CSF shunts, 21
- cytomegalovirus, 133–134
- dactylitis, 238–239
- DDH. *See* developmental dysplasia of the hip
- death practices, by religion, 56
- defoaming agents, 78
- dehydration, 215–216
from AGE, 215–216
diarrhea with, 217
oral replacement therapy for, 218
- dermatitis
atopic, 193–195
contact diaper, 73, 183, 185–186
from GTs, 24
seborrheic, 193
- dermatologic conditions, in infants/neonates, 178–195. *See also* herpes simplex virus; lesions; varicella syndrome; Zoster
- aphthous stomatitis, 183
- dermatitis, 24
contact diaper, 183, 185–186
initial evaluation of, 178
lesions, 184–193
erythematous maculopapular, 185, 190
erzematous, 193–195
Mongolian spots, 185
papulosquamous, 192–193
purpuric, 191
vascular, 190–192
physical exams for, 178
pustular, 183–184
Acropustulosis of infancy, 184
erythema toxicum, 183
neonatal acne, 183–184
scabies, 184
TNPM, 183
treatment of, 193
with Fingertip Method, 178
- vesiculobullous, 178–183
Coxsackie virus, 182–183
friction/sucking blisters, 178–179
HSV, 133–134, 179–180
impetigo contagiosa, 180
Small pox, 182
SSSS, 180

- dermatologic conditions (*cont.*)
 - varicella syndrome, 180–182
 - Zoster, 182
- dermoid cysts, 124
- epidermoid, 124
- midline, 124
- presentation of, 124
- desmopressin, 249–250
- for VWD, 253
- developmental dysplasia of the hip (DDH), 343
 - Barlow/Ortolani tests for, 343
 - symptoms of, 343
 - treatment for, 343
- devices, for premature infants, 17–24. *See also*
 - cerebrospinal fluid shunts; feeding tubes
 - with apnea of prematurity, 17–18
- CSF shunts, 18–21
 - components of, 18, .
 - infection from, 20–21
 - leaks in, 21
 - malfunction of, 18–20
 - overdrainage by, 21
 - pseudocyst formation from, 21
 - purpose of, 18
 - seizures from, 21
 - ventricular/vascular, 21
 - visceral damage from, 21
- feeding tubes, 22–24
 - GTs, 23–24
 - NGTs, 22–23
- patient history with,
- tracheostomy tubes, 21–22
 - cuffed, 22
 - dislodgement/obstruction of, 22
 - infections from, 22
- diabetes, LGA and, 4
- diabetic ketoacidosis (DKA), 265–267
 - cerebral edema and, 266–267
 - definition of, 265
 - lab work for, 265
 - pathophysiology of, 265
 - treatment for, 265–267
 - bicarbonate therapy in, 266
 - insulin therapy in, 266
 - potassium levels management in, 266
- diarrhea, 216–217
 - with dehydration, 217
- diazepam, for febrile seizures, 144, 149
- DIC. *See* disseminated intravascular coagulation
- digital hair tourniquet, 76
- disseminated gonorrhea, 191
- disseminated intravascular coagulation (DIC), 192
- DKA. *See* diabetic ketoacidosis
- Doll’s eyes reflex, 13
- Doppler imaging, for hemangiomas, 125
- Down syndrome, 6
 - AVSDs and, 10
 - central hypotonia and, 13
 - palmar creases and, 12
- drowning/submersion injuries, 197–200
 - in bathtubs, 197
 - caregiver/parental counseling for, 199
 - child abuse as factor in, 198
 - definition of, 197
 - in ED, 198–200
 - epidemiology of, 197–198
 - management of, 198
 - mortality rates for, 197–198
 - pathophysiology of, 198
 - prevention of, 200
 - prognosis for, 199–200
 - BAER in, 199–200
 - risk factors for, 199
- drugs. *See* over the counter medications, poisoning from; prescribed medications, poisoning from
- ductal dependent lesions, 173–174
- Duke criteria, for endocarditis, 171
- duodenal atresia, 205–206
- dysconjugate gaze, 6
- Eagle-Barrett triad, 11
- ears. *See also* acute otitis media
 - AOM in, 71, 83–88
 - common pathogens for, 83
 - differential diagnosis of, 85
 - excessive crying in infants from, 71
 - follow-up for, 88
 - lab work for, 85
 - management of, 85–88
 - otorrhea and, 88
 - radiology for, 85
 - symptoms of, 83–85
 - tympanostomy tubes and, 88
 - URIs and, 83
 - viral causes of, 83
 - evaluation of, 8
 - mastoiditis in, 88–89
 - AM, 89
 - common pathogens for, 89
 - treatment for, 89
 - OE in, 89
 - AOM and, 89
 - treatment for, 84, 89
 - pain in, for infants/newborns, 83–89
 - pre-auricular pitting of, 8
- ECG. *See* electrocardiograms
- E coli* bacteremia, infant fevers from, 111
- eczema. *See also* atopic dermatitis
 - nummular, 195
- ED. *See* emergency department
- EEG. *See* electroencephalography, for seizures
- Ehlers-Danos syndrome, 6
- elbows, orthopedic injuries to, 336, 340
 - radial head subluxation, 341–342
 - hyperpronation method for, 342
 - supination/flexion technique for, 342
- electrical burn injuries, 203
- electrocardiograms (ECG)
 - for acyanotic heart disease, 167
 - for cyanotic heart disease, 163
 - for endocarditis, 171
 - for hypertrophic cardiomyopathy, 173
 - for infant cardiac emergencies, 159–160
- electroencephalography (EEG), for seizures, 148
- elliptocytosis, 131–132
- EM. *See* erythema multiforma
- emergency department (ED)
 - burn injuries in, 202–203
 - endocrine disorders in, 256–257
 - FTT diagnoses in, 93
 - hyperbilirubinemia evaluation in, 135–138
 - for conjugated, 135–136
 - kernicterus and, 136
 - management and, 136–138
 - for UTIs, 135
 - infant/neonatal mortality in, 51–56
 - differential diagnosis for, 52
 - educational programs about, 56
 - epidemiology of, 51
 - GREIV’ING mnemonic for, 56
 - management of, 53–56
 - National Association of Social Workers’ guidelines for, 53, 54
 - patient history in, 52
 - physical examination for, 52
 - presentation of, 51
 - resource/support organizations for, 56
 - SIDS as cause, 52–53
 - inherited bleeding disorders in, 248–249
 - metabolic disorders in, 256–257
 - sickle cell emergencies in, 235
- EMLA. *See* eutectic mixture of local anesthetics
- encephaloceles, 124
 - nasal, 124
- endocardial cushion defect, 167–168
- endocarditis, 171–172
 - categorization of, 172
 - causes of, 171
 - Duke criteria for, 171
 - ECG for, 171
 - lesions from, 191
 - management of, 171–172
 - prophylaxis for, 172
 - for gastrointestinal/genitourinary procedures, 172
- endocrine disorders, 256–268. *See also* adrenal insufficiency
 - adrenal insufficiency, 259–260
 - CAH and, 12
 - differential diagnosis for, 260
 - lab work for, 260
 - symptoms of, 260
 - treatment of, 260
 - diagnostic testing for, 256–257
 - in ED, 256–257
 - physical exams for, 256
 - symptoms of, 256
- endoscopic retrograde cholangiopancreatography (ERCP), 209
- environmental emergencies, 197–200. *See also*
 - burn injuries; drowning/submersion injuries
 - burn injuries, 200–204
 - classification of, 201–204
 - incidence rates for, 201
 - management of, 203–204
 - prevention of, 200–204
 - types of, 201

- drowning/submersion injuries, 197–200
 - in bathtubs, 197
 - caregiver/parental counseling for, 199
 - child abuse as factor in, 198
 - definition of, 197
 - in ED, 198–200
 - epidemiology of, 197–198
 - management of, 198
 - mortality rates for, 197–198
 - pathophysiology of, 198
 - prevention of, 200
 - prognosis for, 199–200
- epidermoid cysts, 124
- epididymitis, 225
 - causes of, 225
 - symptoms of, 225
 - treatment for, 225
- epiglottitis, 275–276
 - causes of, 275
 - complications from, 276
 - incidence rates for, 275
 - lab work for, 276
 - management of, 276
 - symptoms of, 275–276
- epilepsy
 - antiepileptic drug therapy for, 144
 - seizures and, 141
 - syndromes of, 142–143
 - juvenile myoclonic epilepsy of Janz syndrome, 143
 - Lennox-Gastaut syndrome, 142–143
 - in nighttime, 143
 - West’s syndrome, 142
- Epstein’s pearls, 8
- Erb palsy, 14
- ERCP. *See* endoscopic retrograde cholangiopancreatography
- erythema infectiosum, 188–189
- erythema multiforma (EM), 186
- erythema nodosum, 186
- erythema toxicum, 183
- ethosuximide, 152
- etomidate, 31
- European Association of Poison Centers and Clinical Toxicologists, 295
- eutectic mixture of local anesthetics (EMLA), 27–28
- examinations, approaches to, 14. *See also*
 - abdomen, evaluation of; cardiovascular systems, evaluation of; extremities, evaluation of; eyes, evaluation of; genitourinary tract, evaluation of; neck, evaluation of; neurological system, evaluation of; skull, evaluation of; vital signs, for neonates
- of abdomen, 10–11
 - for kidneys, 10
 - for liver, 10
 - for spleen, 10
 - for umbilical area, 10–11
- of cardiovascular systems, 9–10
 - murmurs in, 10
 - post-birth transition for, typical, 9–10
- of chest, 9
 - breast enlargement and, 9
 - clavicle fractures and, 9
 - mastitis in, 9
 - of common primitive reflexes, 13
 - of ears, 8
 - pre-auricular pitting of, 8
 - of extremities, 12
 - feet, 12
 - neurological system and, 12
 - palms, 12
 - polydactyly in, 12
 - of eyes, 6–8
 - blue sclera in, 6
 - Brushfield spots in, 6
 - congenital cataracts in, 6
 - conjunctivitis in, 7
 - ductal obstructions in, 7–8
 - dysconjugate gaze in, 6
 - iris of, 6
 - leukocoria in, 6
 - nystagmus in, 7
 - red reflexes in, 6
 - retinoblastomas in, 6
 - subconjunctival hemorrhages in, 6–7
 - of genitourinary tract, 11–12
 - ambiguous genitalia in, 12
 - in male infants, 11–12
 - of lungs, 9
 - abdominal breathing in, 9
 - apnea and, 9
 - periodic breathing in, 9
 - TNN and, 9
 - of mouth, 8
 - Bohn nodules in, 8
 - cleft lip/palate in, 8
 - Epstein’s pearls in, 8
 - of neck, 8–9
 - branchial cleft anomalies in, 8–9
 - CMT in, 8
 - of neurological system, 12–14
 - for brachial plexus injury, 14
 - for cranial nerves, 12, 13
 - extremities and, movement of, 12
 - hypotonia and, 12–14
 - for sensation, 12
 - Werdnig-Hoffman syndrome and, 14
 - of nose, 8
 - of pharynx, 8
 - of skin, 14
 - for acrocyanosis, 14
 - for cutis marmorata, 14
 - of skull, 4–6
 - caput succedaneum and, 5
 - cephalhematomas and, 5
 - craniosynostosis in, 5
 - fontanel size and, 6
 - molding in, 4–5
 - plagiocephaly in, 5
 - subgaleal bleeds in, 5
 - of vital signs, 3–4
 - fevers, 3
 - weight, 3–4
- exanthem subitum. *See* roseolla infantum
- extremities, evaluation of, 12
 - feet, 12
 - clubfoot in, 12
 - neurological system and, 12
 - palms and, inspection of, 12
 - Down syndrome and, 12
 - polydactyly in, 12
 - eyes, evaluation of, 6–8
 - blue sclera in, 6
 - Ehlers-Danos syndrome and, 6
 - Brushfield spots in, 6
 - Down syndrome and, 6
 - congenital cataracts in, 6
 - conjunctivitis in, 7
 - bacterial causes of, 7
 - chemical causes of, 7
 - chlamydia and, 7
 - from gonorrhea, 7
 - Herpes simplex and, 7
 - corneal abrasions, 75
 - ductal obstructions in, 7–8
 - dysconjugate gaze in, 6
 - for excessive crying in infants, 75
 - iris of, 6
 - leukocoria in, 6
 - nystagmus in, 7
 - red reflexes in, 6
 - colobomas and, 6
 - retinoblastomas in, 6
 - subconjunctival hemorrhages in, 6–7
 - factor XI deficiency, 244
 - failure to thrive (FTT), 92–103
 - cases of, 92–93
 - differential diagnosis for, 95–97
 - for chronic diseases, 96
 - environmental/social factors in, 96–97
 - in ED, 93
 - ICD-9 code for, 93
 - lab tests for, 97–98
 - screening tests, 97–98
 - management of, 99–103
 - breastfeeding patterns in, 102
 - hospitalization for, 102
 - medical instability in, 102–103
 - patient history in, 101–102
 - physical examinations for, 102
 - social instability in, 102
 - pathophysiology of, 93–95
 - anthropometric measurements in, 95
 - weight measurement in, 93–95
 - radiological studies for, 98–99
 - falls, from child abuse, 314–315
 - FDA. *See* Food and Drug Administration
 - feeding tubes, 22–24. *See also* gastronomy tubes
 - GTs, 23–24
 - characteristics of, 23
 - complications of, 23
 - dislodgment of, 23
 - leaks with, 24
 - obstruction of, 23
 - stomal complications with, 24
 - NGTs, 22–23
 - feet
 - clubfoot and, 12
 - digital hair tourniquet, 76
 - plantar reflexes for, 13

- felbamate, 152
- femoral vein, 352
- fever(s), 3, 105–116
 - from bundling, 3
 - FWS, 105, 106
 - hypothermia and, 3
 - in infants, 110–112
 - antibiotics for, 111
 - from bone/joint infections, 114–115
 - definition of, 110
 - E coli* bacteremia and, 111
 - evaluation of, 110
 - general bacteremia in, 110–112
 - from HSV, 114
 - meningitis and, 113–114
 - meningococcal bacteremia and, 111
 - pneumococcal bacteremia and, 110–111, 112–113
 - from UTIs, 112
 - from meningitis, 3
 - in neonates, 107–110
 - antibiotics for, 109–110
 - from bacterial infections, 107
 - Boston criteria for, 107–108
 - evaluation of, 108–109
 - lumbar punctures for, 108
 - Philadelphia criteria for, 108
 - Rochester criteria for, 107
 - testing for, 108
 - treatment for, 109–110
 - UTI testing for, 108
 - WBC testing for, 108
 - from osteomyelitis, 3
 - patient history for, 105–107
 - immunization history in, 105
 - physical examinations for, 105–107
 - bacteremia in, 105
 - infections in, 107
 - seizures from, 143, 145
 - antiepileptic drug therapy for, 144
 - complications from, 144
 - imaging for, 144
 - incidence rates for, 143
 - lumbar punctures for, 144
 - pathophysiology of, 144
 - from sepsis, 3
 - from UTIs, 3
- fever without obvious source (FWS), 105, 106
- FGFR genes. *See* Fibroblast Growth Factor Receptor genes
- Fibroblast Growth Factor Receptor (FGFR) genes, 6
- Fifth's disease. *See* erythema infectiosum
- Fingertip Method, 178
- Fontan operation, 175
- Food and Drug Administration (FDA), 272
 - on OTC medications poisonings, 301
- foreign bodies
 - female genitourinary emergencies from, 229
 - pediatric airway management with, 37–38
 - pediatric resuscitation from, 41
 - removal of, 350
 - respiratory emergencies from, in infants/neonates, 271–272
 - epidemiology of, 271
 - management of, 271–272
 - radiography for, 271
 - symptoms of, 271
- formula, for infants, 78–79
 - casein hydrolysate, 79
 - soy-based, 79
 - whey hydrolysate, 79
- fosphenytoin, 149
- fractures, pediatric, 335–337, 339, 340, 341
 - bowing, 336
 - from child abuse, 78, 313–314
 - of skull, 311–312
 - of clavicle, 9, 339–340
 - greenstick, 336
 - inherited bleeding disorders and, 251
 - metaphyseal, 337, 338
 - physeal, 336
 - Salter-Harris classification of, 336, 337
 - of skull, 73, 324–325
 - supracondylar, 340–341
 - imaging for, 340–341
 - toddler's fracture, 344–345
 - torus, 336, 338
- friction blisters, 178–179
- FTT. *See* failure to thrive
- FWS. *See* fever without obvious source
- gabapentin, 152–153
- gastroenteritis. *See* acute gastroenteritis
- gastroesophageal reflux disease (GERD) as ALTE, 62–63
 - apnea and, 63
 - colic from, 70
- gastrointestinal (GI) bleeding, 213–215
 - causes of, 213
 - from hemophilia, 246
 - treatment for, 251
 - lab work for, 213
 - Apt-Downey test in, 213–214
 - with guiac test, 213
 - management for, 214–215
 - Meckel's diverticulum and, 214
 - radiography for, 214
- gastrointestinal emergencies, 205–219. *See also* acute gastroenteritis; gastrointestinal bleeding
 - AGE, 215–217
 - antibiotic therapy for, 218
 - complications from, 217
 - dehydration from, 215–216
 - rehydration for, 216
 - appendicitis, 212–213
 - lab work for, 212
 - physical examinations for, 212
 - radiography for, 212–213
 - symptoms of, 212
 - treatment for, 213
 - biliary atresia, 134, 207–209
 - diagnosis of, 208–209
 - forms of, 207–208
 - incidence rates for, 207
 - symptoms of, 208
 - causes of, 205, 215
 - constipation, 217–219
 - causes of, 217–219
 - definition of, 217
 - differential diagnosis for, 219
 - evaluation for, 219
 - treatment for, 219
 - diarrhea in, 216–217
 - with dehydration, 217
 - GI bleeding, 213–215
 - causes of, 213
 - lab work for, 213
 - management for, 214–215
 - Meckel's diverticulum and, 214
 - radiography for, 214
 - HPS, 206–207
 - examinations for, 206–207
 - incidence rates for, 206
 - lab work for, 207
 - presentation of, 206
 - radiography for, 207
 - treatment for, 207
 - during immediate newborn period, 205–206
 - duodenal atresia, 205–206
 - NEC, 206
 - intussusception, 210–212
 - causes of, 211
 - lab work for, 211
 - physical exams for, 211
 - radiography for, 211
 - symptoms of, 211
 - treatment for, 211–212
 - malrotation/volvulus of intestines, 209–210
 - embryonic development of, 209
 - incidence rates for, 209
 - lab work for, 209
 - radiography for, 209–210
 - symptoms of, 209
 - treatment of, 210
 - patient anxiety for, 205
- gastronomy tubes (GTs), 23–24
 - characteristics of, 23
 - complications of, 23
 - dislodgment of, 23
 - leaks with, 24
 - obstruction of, 23
 - of outlets, 24
 - stomal complications with, 24
 - allergic/chemical dermatitis, 24
 - cellulitis, 24
 - granulomas, 24
- genital bleeding, 227–228
- genitalia. *See also* genital bleeding;
 - genitourinary emergencies, female;
 - genitourinary emergencies, male;
 - genitourinary tract, evaluation of
- ambiguous, 12
 - females with, 12
 - males with, 12
 - examination of, in females, 227
- genitourinary emergencies, female, 227–232
 - ambiguous genitalia and, 12
 - from foreign bodies, 229
 - genital bleeding, 227–228
 - hemophilia and, 246

- imperforate hymen, 229
- labial adhesions, 231–232
 - treatment for, 231–232
- lichen sclerosus, 229–230
- ovarian torsion, 232
- physical examinations with, 227
- straddle injuries, 231
- urethral prolapse, 228–229
 - symptoms of, 228–229
 - treatment for, 229
- vulvovaginitis, 230–231
 - risk factors for, 230
 - symptoms of, 230
 - treatment for, 230–231
- genitourinary emergencies, male, 11–12, 221–227
 - ambiguous genitalia and, 12
 - balanitis/balanoposthitis, 222
 - circumcision complications as, 223
 - meatal stenosis from, 223
 - phimosis and, 223
 - Plastibell method of, 223
 - entrapment injuries as, 223
 - epididymitis, 225
 - causes of, 225
 - symptoms of, 225
 - treatment for, 225
 - hemophilia and, 246
 - hydroceles and, 11–12, 224
 - idiopathic scrotal edema, 224–225
 - HSP and, 224–225
 - inguinal hernias, 12, 226–227
 - orchitis, 225
 - paraphimosis, 221
 - phimosis, 221
 - priapism, 222–223
 - causes of, 222
 - complications from, 222
 - hospitalization for, 223
 - treatment for, 222–223
 - testicular masses, 227
 - testicular torsion, 225–226
 - treatment for, 226
 - torsion of testicular appendage, 226
 - trauma in, 227
 - varicoceles and, 11, 224
- genitourinary tract, evaluation of, 11–12
 - ambiguous genitalia in, 12
 - females with, 12
 - males with, 12
 - in male infants, 11–12
 - ambiguous genitalia in, 12
 - hydroceles in, 11–12
 - inguinal hernias in, 12
 - varicoceles in, 11
- GERD. *See* gastroesophageal reflux disease
- German measles. *See* rubella
- GI bleeding. *See* gastrointestinal bleeding
- Glasgow Coma Scale, 293
 - for minor head injuries, 323
- Glenn operation, 175
- gliomas, 124–125
- glucocorticoids, 274
- Goldenhar syndrome, 9
- gonorrhea, conjunctivitis from, 7
- granuloma annulare, 193
- granulomas, 24
- greenstick fractures, 336
- GREIV_ING mnemonic, for infant/neonatal mortality, 56
- grieving, cultural concepts of, 55
- GTs. *See* gastrostomy tubes
- guaiac test, 213
- Hand-Foot-Mouth disease. *See* Coxsackie virus
- hand-foot syndrome. *See* dactylitis
- hand-to-mouth reflex, 13
- HD amoxicillin. *See* high-dose amoxicillin
- headaches, 328
- head and neck, emergencies with, 122–126.
 - See also* head injuries, minor; neck, evaluation of
 - cervical lymphadenitis, 123–124
 - antibiotic therapy for, 123
 - causes of, 123
 - lymphadenopathy and, 123
 - cystic hygromas, 124
 - dermoid cysts, 124
 - epidermoid, 124
 - midline, 124
 - presentation of, 124
 - encephaloceles, 124
 - nasal, 124
 - evaluation of, 122
 - with CT, 122
 - symptoms in, 122
 - ultrasonography for, 122
 - gliomas, 124–125
 - hemangiomas, 125
 - CT for, 125
 - dermatologic conditions v., 125
 - Doppler imaging for, 125
 - incidence rates for, 125
 - presentation of, 125
 - thrombocytopenia and, 125
 - treatment therapies for, 125
 - hemophilia as, 246
 - treatment for, 251
 - for lesions, 122
 - cystic, 122, 123
 - inflammatory, 122
 - neck
 - branchial cleft anomalies in, 8–9, 122–123
 - CMT in, 8
 - evaluation of, 8–9
 - tonic neck reflex for, 13
 - neonatal torticollis, 122, 125–126
 - orthopedic injuries in, 337–339
 - pseudosubluxation, 338–339
 - pre-auricular pits, 8, 126
 - Sturge-Weber syndrome, 126
- head injuries, minor, 322–332
 - brain injury from, 331
 - CNS trauma in, 331
 - clinical decision rules for, 329–330
 - under CATCH, 330
 - under CCHR, 329
 - under CHALICE, 330
 - under CHIP, 329
 - under NEXUS II, 329
 - under PECARN study, 330
 - University of California-Davis rule, 329
 - CT scans for, 325–326
 - risk factors for, 326
 - definition of, 323–324
 - AAP guidelines for, 323
 - Glasgow Coma scale for, 323
 - evaluation of, 324
 - clinical variables in, 326–327
 - headaches from, 328
 - heterogeneity of studies for, 328
 - ICI, 324–330
 - clinical indicators for, 328
 - incidence rates for, 322
 - infant v. adult brain and, 322–323
 - injury mechanism differences and, 323
 - LOC from, 327
 - long-term psycho-social outcomes of, 330
 - MRI for, 326
 - in neurologically normal infants, 323
 - post-traumatic amnesia from, 328
 - scalp hematomas, 327
 - seizures from, 327
 - skull radiographs for, 324–325
 - for skull fractures, 324–325
 - ultrasound for, 331–332
 - limitations of, 331–332
 - vomiting as result of, 328
- heart rate variability (HRV), 65
- HEI. *See* hemorrhagic edema of infancy
- Heimlich maneuver, 38
- hemangiomas, 125
 - CT for, 125
 - dermatologic conditions v., 125
 - Doppler imaging for, 125
 - incidence rates for, 125
 - lesions from, 190
 - presentation of, 125
 - thrombocytopenia and, 125
 - treatment therapies for, 125
- hemarthrosis, 246
 - treatment for, 251
- hematology. *See* bleeding disorders, inherited; sickle cell, emergencies with
- hematomas
 - scalp, 327
 - treatment for, 251
- hematurias, treatment for, 251
- hemodialysis, for salicylate poisoning, 298
- hemolytic disease, 130
- hemolytic transfusion reaction (HTR), 241
- hemophilia, 243–244, 245–246
 - CNS and, 245
 - genitourinary tract bleeding from, 246
 - GI bleeding from, 246
 - in head and neck, 246
 - hemarthrosis, 246
 - hemostasis and, 243
 - incidence rates for, 243
 - in intra-abdomen, 246
 - posttraumatic bleeding and, 246
 - by sex, 243
 - skeletal muscle and, 246
 - symptoms of, 243–244, 245

- hemorrhagic edema of infancy (HEI), 191
- hemostasis, hemophilia and, 243
- Henoch-Schoenlein purpura (HSP), 173
 - idiopathic scrotal edema and, 224–225
 - lesions from, 191
- hereditary spherocytosis (HS), 131–132
- hernias, 11
 - CDH, 11
 - incarcerated inguinal, 76
 - inguinal, 12
- herpes simplex virus (HSV), 133–134, 179–180
 - causes of, 179
 - conjunctivitis and, 7
 - diagnosis of, 179
 - fevers from, 114
 - genital, 180
 - lesions with, 179
 - presentation of, 179
 - treatment for, 179–180
- high-dose (HD) amoxicillin, for AOM, 86–87
- hip, injuries to, 343–344
 - DDH, 343
 - Barlow/Ortolani tests for, 343
 - symptoms of, 343
 - treatment for, 343
 - osteomyelitis, 344
 - causes of, 344
 - diagnosis of, 344
 - fever from, 3
 - radiography for, 344
 - symptoms of, 344
 - treatment for, 344
 - septic arthritis, 115, 344
 - causes of, 344
 - diagnosis of, 344
 - radiography for, 344
 - symptoms of, 344
 - treatment for, 344
 - transient synovitis, 343–344
 - lab testing for, 343–344
- Hirschsprung’s disease, 76
- HLHS. *See* Hypoplastic Left Heart Syndrome
- holosystolic murmurs, 10
 - VSD as cause of, 10
- HPS. *See* hypertrophic pyloric stenosis
- HRV. *See* heart rate variability
- HS. *See* hereditary spherocytosis
- HSP. *See* Henoch-Schoenlein purpura
- HSV. *See* herpes simplex virus
- HTR. *See* hemolytic transfusion reaction
- human milk hyperbilirubinemia, 132–133
- hydroceles, 11–12, 224
- hydroxyurea, 238
- hyperbilirubinemia, 128–138. *See also*
 - kernicterus; neonatal hepatitis syndrome
 - complications of, 128
 - kernicterus as, 128, 136
 - conjugated, 133
 - causes of, 130
 - ED evaluation of, 135–136
 - differential diagnoses for, 130–135
 - for blood group incompatibility, 130–131
 - for congenital hypothyroidism, 132
 - for Crigler-Najjar syndrome, 132
 - duration in, 133
 - for elliptocytosis/ovalocytosis, 131–132
 - for enzyme deficiencies, 131
 - for hemolytic disease, 130
 - for HS, 131–132
 - for human milk hyperbilirubinemia, 132–133
 - for increased indirect bilirubin, 132–133
 - for NHS, 133–134
 - for PKD, 131
 - ED evaluation of, 135–138
 - for conjugated, 135–136
 - management and, 136–138
 - patient admission in, 138
 - for UTIs, 135
 - epidemiology of, 128
 - family history for, 129
 - pathophysiology of, 128–129
 - causes in, 129
 - unconjugated bilirubin in, 128–129
 - patient birth history and, 129–130
 - LGA and, 129–130
 - SGA and, 129–130
 - physical assessment for, 129
 - prognosis for, 138
 - risk factors for, 129, 131
 - ethnicity as, 129
 - maternal medications as, 130
 - symptoms of, 129–130
 - treatment for, 137–138
 - breastfeeding in, 137
 - with exchange transfusions, 138
 - pharmacologic therapy in, 138
 - with phototherapy, 137–138
 - unconjugated, 130
 - management of, 133
- hyperglycemia, 264–267
 - definition of, 264
 - differential diagnosis of, 265
 - IDDM and, 264–265
 - diagnosis of, 265
 - lab work for, 265
 - hyperpronation method, 342
 - hyperthyroidism, 261–262
 - symptoms of, 261
 - hypertonic saline, 267
 - for bronchiolitis, 280
 - central pontine myelinolysis and, 267–268
 - hypertrophic cardiomyopathy, 173
 - ECG for, 173
 - hypertrophic pyloric stenosis (HPS), 206–207
 - examinations for, 206–207
 - incidence rates for, 206
 - lab work for, 207
 - presentation of, 206
 - radiography for, 207
 - treatment for, 207
 - hypoglycemia, 263–264
 - causes of, 264
 - definition of, 263
 - differential diagnosis of, 263
 - excessive crying in infants and, 76–77
 - glucose levels and, 263
 - symptoms of, 263
 - testing for, 263
 - treatment of, 263–264
 - hyponatremia, 267–268
 - differential diagnosis of, 267
 - lab work for, 267
 - symptoms of, 267
 - treatment of, 267–268
 - with hypertonic saline, 267
 - Hypoplastic Left Heart Syndrome (HLHS), 163, 168–169
 - Glenn operation for, 175
 - Hypoplastic Right Heart Syndrome, 163
 - Glenn operation for, 175
 - hypothermia, fevers and, 3
 - hypothyroidism, 261
 - hypotonia, 12–14
 - central, 13
 - Down syndrome and, 13
 - Prader-Willi Syndrome and, 13
 - peripheral, 13–14
 - severe, 13
- ICD-9. *See* International Classification of Diseases
- ICI. *See* intra-cranial injuries
- IDDM. *See* insulin-dependent diabetes mellitus
- idiopathic scrotal edema, 224–225
 - HSP and, 224–225
- idiopathic thrombocytopenic purpura (ITP), 191
- IEM. *See* inborn errors of metabolism
- immunizations, 348
 - contraindication factors for, 349
 - excessive crying in infants and, 74
- imperforate hymen, 229
- Impetigo contagiosa, 180
- IMR. *See* infant mortality rate
- inborn errors of metabolism (IEM), 65, 257–259
 - differential diagnosis for, 258
 - excessive crying in infants and, 76
 - incidence rates for, 257
 - lab work for, 258–259
 - with urinalysis, 259
 - symptoms of, 257–258
 - urea cycle defects in, 257–258
 - treatment for, 259
- incarcerated inguinal hernia, 76
- increased indirect bilirubin, 132–133
 - breastfeeding and, 132–133
- infant mortality rate (IMR), 51
- infant/neonatal mortality, in ED, 51–56
 - differential diagnosis for, 52
 - educational programs about, 56
 - epidemiology of, 51
 - IMR in, 51
 - PMR in, 51
 - GREIV_ING pneumonic for, 56
 - management of, 53–56
 - cultural/religious beliefs and, 54, 55, 56
 - law enforcement and, 54
 - National Association of Social Workers’ guidelines for, 53, 54
 - patient history in, 52

- physical examination for, 52
- presentation of, 51
- resource/support organizations for, 56
- SIDS as cause, 52–53
 - AAP on, 52–53
 - child abuse v., 52
 - demography for, 52
 - diagnosis of, 52–53
 - lab studies for, 53
 - postmortem examinations for, 53
 - radiology studies for, 53
 - sleeping position and, 52
- infants. *See also* burn injuries; cardiac emergencies, in infants; crying in infants, excessive; dermatologic conditions, in infants/neonates; devices, for premature infants; drowning/submersion injuries; head injuries, minor; infant/neonatal mortality, in ED; orthopedic injuries, in infants; premature infants; respiratory emergencies, in infants/neonates; Sudden Infant Death Syndrome
- cardiac emergencies in, 159–176
- cardiac disease and, 159
- cardiomyopathies and, 173
- carditis in, 169–172
- chest X-rays for, 159–160
- congestive heart failure and, 174
- cyanotic heart disease in, 162–163, 169
- ECG for, 159–160
- heart chamber size and, 161–162
- heart rate and, 160
- Kawasaki’s disease in, 172–173
- PR intervals during, 161
- QRS duration and, 161
- surgery for, 174, 175–176
- T wave morphology in, 161
- ventricle axis dominance and, 161
- dermatologic conditions in, 178–195
 - aphthous stomatitis, 183
 - dermatitis, 24
 - initial evaluation of, 178
 - lesions, 184–193
 - physical exams for, 178
 - pustular, 183–184
 - treatment of, 193
 - vesiculobullous, 178–183
- fevers in, 110–112
 - antibiotics for, 111
 - from bone/joint infections, 114–115
 - definition of, 110
 - E coli* bacteremia and, 111
 - evaluation of, 110
 - general bacteremia in, 110–112
 - from HSV, 114
 - meningitis and, 113–114
 - meningococcal bacteremia and, 111
 - pneumococcal bacteremia and, 110–111, 112–113
 - from UTIs, 112
- formula for, 78–79
 - casein hydrolysate, 79
 - soy-based, 79
 - whey hydrolysate, 79
- IMR, 51
- lesions in, 184–193
 - erythematous maculopapular, 185–190
 - erzematous, 193–195
 - from malaria rubia, 186
 - Mongolian spots, 185
 - papulosquamous, 192–193
 - from tinea versicolor, 192–193
 - vascular, 190–192
- minor head injuries in, 322–332
 - clinical decision rules for, 329–330
 - CT scans for, 325–326
 - definition of, 323–324
 - evaluation of, 324
 - headaches from, 328
 - heterogenicity of studies for, 328
 - ICL, 324–330
 - incidence rates for, 322
 - infant v. adult brain and, 322–323
 - LOC from, 327
 - long-term psycho-social outcomes of, 330
 - MRI for, 326
 - in neurologically normal infants, 323
 - post-traumatic amnesia from, 328
 - scalp hematomas, 327
 - seizures from, 327
 - skull radiographs for, 324–325
 - ultrasound for, 331–332
 - vomiting as result of, 328
- mortality, in ED, 51–56
 - differential diagnosis for, 52
 - educational programs about, 56
 - epidemiology of, 51
 - GREIV’ING pneumonic for, 56
 - management of, 53–56
 - National Association of Social Workers’ guidelines for, 53, 54
 - patient history in, 52
 - physical examination for, 52
 - presentation of, 51
 - resource/support organizations for, 56
 - SIDS as cause, 52–53
- orthopedic injuries in, 335–345
 - common terms in, 335
 - fractures, 335–337
 - of hip, 343–344
 - of lower extremities, 344–345
 - of neck, 337–339
 - physical examinations for, 335
 - radiographic examinations for, 335
 - of upper extremities, 339
- infections
 - ALTEs from, 64–65
 - bordetella pertussis, 65
 - RSV, 64–65
 - bone/joint, 114–115
 - fevers from, 114–115
 - cervical lymphadenitis from, 123
 - from CSF shunts, 20–21
 - etiology of, 20
 - lab work for, 20–21
 - radiology for, 21
 - symptoms of, 20
 - treatment therapies for, 21
 - excessive crying in infants from, 71–73
 - in bones/joints, 72–73
 - otitis media as cause of, 71
 - otoscopy use for, 71
 - fevers from, 107
 - in neonates, 107
 - respiratory emergencies from, 272–278
 - from bacterial tracheitis, 272–273
 - from croup, 273–275
 - from epiglottitis, 275–276
 - FDA recommendations for, 272
 - from retropharyngeal abscess, 276–278
 - from URIs, 83, 272
 - sickle cell and, 236
 - from tracheostomy tubes, 22
 - URIs, 83
 - UTIs, 3
 - ALTEs from, 65
 - hyperbilirubinemia from, 135
- infectious mononucleosis. *See* mononucleosis, infectious
- inflicted traumatic brain injury (iTBI), 309
- inguinal hernias, 12, 226–227
 - excessive crying in infants and, 76
- inherited bleeding disorders. *See* bleeding disorders, inherited
- insect bites
 - excessive crying in infants from, 74
 - lesions from, 184–185
- insulin-dependent diabetes mellitus (IDDM), 264–265
 - diagnosis of, 265
- insulin therapy, for DKA, 266
- International Association for the Study of Pain, 26
- International Classification of Diseases (ICD-9), 93
- intra-cranial injuries (ICI), 324–330
- intraosseous line, 352–353, 354
- intussusception, 210–212
 - causes of, 211
 - lab work for, 211
 - physical exams for, 211
 - radiography for, 211
 - symptoms of, 211
 - treatment for, 211–212
- iris, 6
- iron, ingestion of, 298–299
 - clinical stages of, 299
 - lab testing for, 299
 - treatment for, 299
- iTBI. *See* inflicted traumatic brain injury
- ITP. *See* idiopathic thrombocytopenic purpura
- jaundice. *See* hyperbilirubinemia
- jet ventilation. *See* needle cricothyrotomy
- jock itch. *See* tinea cruris
- jugular vein, 352
- Juvenile myoclonic epilepsy of Janz syndrome, 143
- Kawasaki’s disease, 172–173
 - lesions from, 189–190
 - presentation of, 173
 - treatment of, 174–175

- keratosis pilaris, 193
- kernicterus, 128
 - brain damage from, 136
 - ED evaluation of, 136
 - in Pilot Kernicterus Registry, 128
 - symptoms of, 136
- ketamine, 31
- ketogenic diet, 153–154
- kidneys, 10
- Klumpke palsy, 14
- labial adhesions, 231–232
 - treatment for, 231–232
- lamotrigine, 152
- large gestational age (LGA), 4
 - Beckwith-Wiedemann syndrome and, 4
 - diabetes and, 4
 - hyperbilirubinemia and, 129–130
 - Simpson-Golabi-Behmel syndrome and, 4
 - Soto syndrome and, 4
- laryngeal mask airway (LMA), 38
 - complications of, 38
 - placement techniques for, 38
 - replacement of, 38
 - sizing for, 38
- laryngomalacia, 270
- laryngoscopes, 36
- lead, poisoning from, 299–300
 - management of, 300
 - sources of, 300
 - symptoms of, 299–300
 - testing for, 300
- Lennox-Gastaut syndrome, 142–143
- lesions
 - cystic hygromas, 124
 - ductal dependent, 173–174
 - gliomas, 124–125
 - in head and neck, 122
 - head and neck, emergencies with
 - cystic, 122, 123
 - inflammatory, 122
 - from herpes simplex, 179
 - infant/neonatal, 184–193
 - from acrocyanosis of infancy, 190
 - from atopic dermatitis, 193–195
 - from candidiasis, 186
 - from coin rubbing/cupping, 192
 - from contact diaper dermatitis, 185–186
 - from DIC, 192
 - from disseminated gonorrhea, 191
 - from drug reactions, 185
 - from EM, 186
 - from endocarditis, 191
 - from erythema infectiosum, 188–189
 - from erythema nodosum, 186
 - from granuloma annulare, 193
 - harlequin color change in, 190
 - from HEI, 191
 - from hemangiomas, 190
 - from HSP, 191
 - from infectious mononucleosis, 189
 - from insect bites, 184–185
 - from ITP, 191
 - from Kawasaki's disease, 189–190
 - from keratosis pilaris, 193
 - from Lyme disease, 190
 - from measles, 188
 - from meningococemia, 192
 - from milaria rubra, 184, 186
 - from milia, 184
 - from molluscum contagiosum, 184
 - from Mongolian spots, 185
 - from neonatal sebaceous gland hyperplasia, 184
 - from nummular eczema, 195
 - pityriasis rosea, 192–193
 - from pityriasis alba, 184
 - from psoriasis, 192
 - purpuric, 191
 - from pyogenic granulomas, 190–191
 - from RMSF, 192
 - from roseolla infantum, 187
 - from rubella, 187–188
 - from scarlet fever, 189
 - from seborrheic dermatitis, 193
 - from Steven-Johnson syndrome, 186
 - from stork bites, 190
 - from subcutaneous fat necrosis of the newborn, 185
 - from syphilis, 191–192
 - from tinea capitis, 193
 - from tinea corporis, 192
 - from tinea cruris, 193
 - from tinea pedis, 193
 - from tinea versicolor, 184, 192–193
 - from toxic shock, 190
 - from TTP, 191
 - from urticaria, 186
 - from viral exanthem, 186–187, 194
 - leukocoria, 6
 - levetiracetam, 153
 - LGA. *See* large gestational age
 - lichen sclerosis, 229–230
 - liver, 10
 - LMA. *See* laryngeal mask airway
 - LOC. *See* loss of consciousness
 - lorazepam, 149
 - loss of consciousness (LOC), 327
 - lumbar punctures, 108
 - for meningitis, 114
 - for pediatric seizures, from fevers, 144
 - for seizures, 147
 - Lund and Browder chart, for burn injuries, 201
 - lungs, evaluation of, 9
 - abdominal breathing in, 9
 - apnea and, 9
 - periodic breathing in, 9
 - TNN and, 9
 - Lyme disease, 190
 - lymphadenopathy, 123
 - macrocephaly, 4
 - magnetic resonance imaging (MRI)
 - for minor head injuries, 326
 - for neonatal seizures, 145
 - malrotation/volvulus of intestines, 209–210
 - embryonic development of, 209
 - incidence rates for, 209
 - lab work for, 209
 - radiography for, 209–210
 - symptoms of, 209
 - treatment of, 210
 - MAO. *See* mechanical airway obstruction
 - mastitis, 9
 - mastoiditis, 88–89
 - AM, 89
 - common pathogens for, 89
 - treatment for, 89
 - measles, 188
 - meatal stenosis, 223
 - mechanical airway obstruction (MAO), 65–66
 - Meckel's diverticulum, 214
 - medications. *See* over the counter medications, poisoning from; prescribed medications, poisoning from
 - meningitis
 - causes of, 113
 - diagnosis of, 113
 - fever from, 3
 - antibiotic therapy for, 114
 - in infants, 113–114
 - testing for, 113–114
 - with blood cultures, 113
 - with CT, 113–114
 - with lumbar puncture, 114
 - meningococcal bacteremia, infant fevers from, 111
 - meningococemia, 192
 - meperidine, 29
 - metabolic acidosis, 262–263
 - causes of, 262
 - differential diagnosis of, 262
 - symptoms of, 262
 - treatment of, 262–263
 - metabolic disorders, 256–268. *See also* diabetic ketoacidosis; inborn errors of metabolism
 - diagnostic testing for, 256–257
 - DKA, 265–267
 - cerebral edema and, 266–267
 - definition of, 265
 - lab work for, 265
 - pathophysiology of, 265
 - treatment for, 265–267
 - in ED, 256–257
 - excessive crying in infants from, 76–77
 - hospitalization for, 257
 - hyperglycemia, 264–267
 - definition of, 264
 - differential diagnosis of, 265
 - IDDM and, 264–265
 - lab work for, 265
 - hypoglycemia, 263–264
 - causes of, 264
 - definition of, 263
 - differential diagnosis of, 263
 - excessive crying in infants and, 76–77
 - glucose levels and, 263
 - symptoms of, 263
 - testing for, 263
 - treatment of, 263–264
 - hyponatremia, 267–268

- IEM, 65, 257–259
differential diagnosis for, 258
excessive crying in infants and, 76
incidence rates for, 257
lab work for, 258–259
symptoms of, 257–258
treatment for, 259
metabolic acidosis, 262–263
causes of, 262
differential diagnosis of, 262
symptoms of, 262
treatment of, 262–263
in NHS, 134
physical exams for, 256
symptoms of, 256
of thyroid, 261–262
differential diagnosis for, 261–262
hyperthyroidism, 261–262
hypothyroidism, 261
lab work for, 262
treatment for, 262
metaphyseal fractures, 337, 338
methemoglobinemia, 300–301
congenital forms of, 300
external causes of, 300
lab testing for, 300–301
symptoms of, 300
treatment of, 301
methemoglobin inducing substances, 300–301.
See also methemoglobinemia
methemoglobinemia and, 300–301
methylxanthine therapy, 17
with caffeine, 17
microcephaly, 4
midazolam, 149
midgut volvulus, 76
midline dermoid cysts, 124
malaria rubra, 184, 186
milia, 184
minor head injuries. *See* head injuries, minor
mist therapy, 275
molluscum contagiosum, 184
Mongolian spots, 185
mononucleosis, infectious, 189
Moro reflex, 13
morphine, 29
mortality. *See* infant/neonatal mortality,
in ED
mouth, evaluation of, 8
Bohn nodules in, 8
cleft lip/palate in, 8
Epstein's pearls in, 8
MRI. *See* magnetic resonance imaging
mucocutaneous lymph node syndrome. *See*
Kawasaki's disease
Münchhausen by proxy, 64
murmurs, 10
AVSD and, 10
Down syndrome and, 10
holosystolic, 10
VSD as cause of, 10
PDA, 10
sound of, in diagnosis of, 10
Mustard and Senning operation,
175
myocarditis, 169
causes of, 169
symptoms of, 169
myoclonic seizures, 142
NAC. *See* N-acetylcysteine
narcolepsy, 146
nasogastric tubes (NGTs), 22–23
nasotracheal intubation, 36
NASPGHAN. *See* North American Society of
Pediatric Gastroenterology Hepatology
and Nutrition
National Association of Social Workers, on
infant/neonatal mortality, bereavement
guidelines for, 53, 54
National Center for Health Statistics Growth
Charts (NCHS), 93
National Emergency X-Radiography
Utilization Study II (NEXUS II),
329
NCCT. *See* non-contrast computed
tomography, for CSF shunts
NCHS. *See* National Center for Health
Statistics Growth Charts
nebulized epinephrine, 280
NEC. *See* necrotizing enterocolitis
neck, evaluation of, 8–9
branchial cleft anomalies in, 8–9, 122–123
excision of, 123
Goldenhar syndrome as, 9
location of, 123
Pierre-Robin Association as, 9
CMT in, 8
orthopedic injuries and, 337–339
pseudosubluxation, 338–339
tonic neck reflex for, 13
necrotizing enterocolitis (NEC), 206
needle cricothyrotomy, 39
complications of, 39
membrane identification in, 39
ventilation methods for, 39
needle thoracostomy, 350–351
diagnosis for, 350
neonatal acne. *See* acne, neonatal
neonatal hepatitis syndrome (NHS),
133–134
alagille syndrome and, 134
biliary atresia and, 134
CF and, 134–135
choledochal cysts and, 134
conjugated hyperbilirubinemia and,
133
metabolic disorders in, 134
ToRCH infections and, 133–134
neonatal sebaceous gland hyperplasia, 184
neonatal torticollis, 122, 125–126
neonates. *See also* dermatologic conditions, in
infants/neonates; pain management,
for neonates; respiratory emergencies,
in infants/neonates; sedation, for
neonates; vital signs, for neonates;
weight, of neonates
dermatologic conditions in, 178–195
aphthous stomatitis, 183
dermatitis, 24
initial evaluation of, 178
lesions, 184–193
physical exams for, 178
pustular, 183–184
treatment of, 193
vesiculobullous, 178–183
fevers in, 107–110
antibiotics for, 109–110
from bacterial infections, 107
Boston criteria for, 107–108
evaluation of, 108–109
lumbar punctures for, 108
Philadelphia criteria for, 108
Rochester criteria for, 107
testing for, 108
treatment for, 109–110
UTI testing for, 108
WBC testing for, 108
lesions in, 184–193
erythematous maculopapular, 185–190
erzematous, 193–195
Mongolian spots, 185
papulovesicular, 192–193
vascular, 190–192
pain management for, 26–31
AAP Policy Statement on Prevention and
Management of Pain in the Neonate, 26
analgesia in, 26–29
analgesics, 28–29
animal/human studies on, 26
for AOM, 88
combination therapies for, 27
infiltrative medications for, 28
International Association for the Study of
Pain on, 26
nonpharmacologic, 27
sedation as part of, 29–31
topical medications for, 27–28
sedation for, 29–31
barbiturates for, 30
benzodiazepines for, 30
chloral hydrate for, 30
dosing guidelines for, 31
etomidate for, 31
ketamine for, 31
propofol for, 31
types of, 29–30
seizures in, 145
BFNC and, 145
BINC and, 145
causes of, 145
complications of, 145
imaging for, 145
symptoms of, 145
treatment of, 145
vital signs for, 3–4
fever in, 3
weight in, 3–4
weight of, 3–4
average gains in, 4
LGA and, 4
macrocephaly and, 4
microcephaly and, 4
SGA and, 4
standardized growth charts for, 4

- neurological system, evaluation of, 12–14. *See also* hypotonia
 - for brachial plexus injury, 14
 - Erb palsy and, 14
 - Klumpke palsy and, 14
 - for cranial nerves, 12, 13
 - extremities and, movement of, 12
 - hypotonia and, 12–14
 - central, 13
 - peripheral, 13–14
 - severe, 13
 - for sensation, 12
 - Werdnig-Hoffman syndrome and, 14
- NEXUS. *See* National Emergency X-Radiography Utilization Study II
- NGTs. *See* nasogastric tubes (NGTs)
- NHS. *See* neonatal hepatitis syndrome
- night terrors, 146
- non-contrast computed tomography (NCCT),
 - for CSF shunts, 19–20
- non-opioid analgesics, 28–29
 - acetaminophen, 28
 - recommended dosing for, 28
 - NSAIDs, 28–29
- non-steroidal anti-inflammatory drugs (NSAIDs), 28–29
- North American Society of Pediatric Gastroenterology Hepatology and Nutrition (NASPGHAN), 217
- Norwood operation, 175
- nose, evaluation of, 8
- NSAIDs. *See* non-steroidal anti-inflammatory drugs
- nursemaid’s elbow. *See* radial head subluxation
- nystagmus, 7
- obstructive apnea, 63
- OE. *See* otitis externa
- OI. *See* osteogenesis imperfecta
- omphalitis, 10–11
- opioid analgesics, 29
 - dosing guidelines for, 29
 - meperidine, 29
 - morphine, 29
 - toxic poisoning from, 303
- orchitis, 225
- orthopedic injuries, in infants, 335–345. *See also* fractures, pediatric; hip, injuries to; osteomyelitis; radial head subluxation; septic arthritis
 - common terms in, 335
 - fractures, 335–337, 339, 340, 341
 - bowing, 336
 - from child abuse, 78, 313–314
 - of clavicle, 9, 339–340
 - greenstick, 336
 - inherited bleeding disorders and, 251
 - metaphyseal, 337
 - physeal, 336
 - Salter-Harris classification of, 336, 337
 - of skull, 73, 324–325
 - toddler’s fracture, 344–345
 - torus, 336
 - of hip, 343–344
 - DDH, 343
 - osteomyelitis, 344
 - septic arthritis, 115, 344
 - transient synovitis, 343–344
- of lower extremities, 344–345
 - OI, 345
 - toddler’s fracture, 344–345
- of neck, 337–339
 - pseudosubluxation, 338–339
- physical examinations for, 335
- radiographic examinations for, 335
 - common ossification centers in, 335
- of upper extremities, 339
 - clavicle fractures, 339–340
 - of elbow, 336, 340
 - radial head subluxation, 341, 342
 - supracondylar fractures, 340–341
- Ortolani tests, for DDH, 343
- osteogenesis imperfecta (OI), 345
- osteomyelitis, 344
 - causes of, 344
 - diagnosis of, 344
 - fever from, 3
 - radiography for, 344
 - symptoms of, 344
 - treatment for, 344
- OTC medications. *See* over the counter medications, poisoning from
- otitis externa (OE), 89
 - AOM and, 89
 - treatment for, 84, 89
- otorrhea, 88
- otoscopy, 71
- ovalocytosis, 131–132
- ovarian torsion, 232
- over the counter (OTC) medications,
 - poisoning from, 301
 - camphor, 304
 - CDC and, 301
 - FDA and, 301
- oxcarbazepine, 153
- pain, definition of, 26–27
- pain management, for neonates, 26–31. *See also* analgesics; sedation, for neonates
 - AAP Policy Statement on Prevention and Management of Pain in the Neonate, 26
 - analgesia in, 26–29
 - definition of, 27
 - analgesics, 28–29
 - non-opioid, 28–29
 - opioid, 29
 - animal/human studies on, 26
 - for AOM, 88
 - combination therapies for, 27
 - infiltrative medications for, 28
 - International Association for the Study of Pain on, 26
 - nonpharmacologic, 27
 - breastfeeding as, 27
 - environmental interventions as, 27
 - sucrose administration as, 27
 - sedation as part of, 29–31
 - barbiturates for, 30
 - benzodiazepines for, 30
 - chloral hydrate for, 30
 - dosing guidelines for, 31
 - etomidate for, 31
 - ketamine for, 31
 - propofol for, 31
 - types of, 29–30
 - topical medications for, 27–28
 - EMLA, 27–28
 - palmar reflex, 13
 - palms, inspection of, 12
 - palmar reflexes in, 13
 - paracetamol. *See* acetaminophen
 - paralytic medications, for rapid-sequence intubation, 35
 - paraphimosis, 221
 - partial prothrombin time (PTT), 247
 - patent ductus arteriosus (PDA) murmur, 10
 - coarctation of the aorta and, 168
 - cyanotic disease and, 167
 - patent urachus, 11
 - VCUG for, 11
 - PCV-7. *See* pneumococcal conjugate vaccine
 - PDA murmur. *See* patent ductus arteriosus murmur
 - PEA. *See* pulseless electrical activity
 - PECARN study, 330
 - pediatric airway management. *See* airway management, pediatric
 - pediatric fractures. *See* fractures, pediatric
 - pediatric rapid-sequence intubation. *See* rapid-sequence intubation, pediatric
 - pediatric resuscitation, 40–49
 - pediatric seizures. *See* seizures, pediatric
 - percutaneous chest tubes, 351
 - complications of, 351
 - pericarditis, 169–171
 - cardiac tamponade and, 170–171
 - cause of, 169
 - presentation of, 169–170
 - perinatal mortality rate (PMR), 51
 - periodic breathing, 9
 - peripheral hypotonia, 13–14
 - pertussis, 65, 284–285
 - antibiotic therapy for, 285
 - clinical features of, 284
 - complications from, 284–285
 - epidemiology of, 284
 - lab work for, 284
 - management of, 284
 - petit mal seizures. *See* absence seizures
 - pharynx, evaluation of, 8
 - phenobarbital, for febrile seizures, 144, 149, 150–152
 - phenytoin, 150
 - Philadelphia criteria, 108
 - phimosis, 221
 - circumcision and, 223
 - phototherapy, 137–138
 - physeal fractures, 336
 - Salter-Harris classification of, 336
 - Pierre-Robin Association, 9
 - Pilot Kernicterus Registry, 128
 - pityriasis rosea, 192–193
 - pityriasis alba, 184
 - atopic dermatitis and, 184
 - PKD. *See* pyruvate kinase deficiency

- plagiocephaly, 5
- plantar reflex, 13
- Plastibell method, for circumcision, 223
- PMR. *See* perinatal mortality rate
- pneumococcal bacteremia
 - infant fevers from, 110–111, 112–113
 - radiology studies for, 113
 - risk factors for, 349
- pneumococcal conjugate vaccine (PCV-7), 83
- pneumonia, 281–283
 - complications from, 283
 - epidemiology of, 281
 - etiology of, 281
 - lab work for, 282–283
 - management of, 283
 - presentation of, 281–282
 - radiography for, 283–285
 - treatment for, by age and type, 282–283
- Poison Prevention Act (PPA) (1970), 292
 - cleaning agent poisoning and, 301
- polydactyly, 12
- post-traumatic amnesia, 328
- PPA. *See* Poison Prevention Act
- Prader-Willi Syndrome, central hypotonia and, 13
- pre-auricular pitting, 8, 126
 - treatment for, 126
- premature infants, 16–24. *See also* apnea of prematurity; devices, for premature infants
 - apnea of prematurity in, 17–18
 - epidemiology of, 17
 - methylxanthine therapy in, 17
 - monitoring of, 17–18
 - pathophysiology of, 17
 - SIDS and, 17–18
 - assessment of, 16–17
 - definition of, 16
 - devices for, 17–24
 - for apnea of prematurity, 17–18
 - CSF shunts, 18–21
 - feeding tubes, 22–24
 - history for, 21–22
 - tracheostomy tubes, 21–22
 - mortality rates for, 16
 - terms related to, 16
- preoxygenation, 34
- prescribed medications, poisoning from, 302–303, 304. *See also* over the counter medications, poisoning from
 - calcium channel blockers, 302–303
 - from clonidine, 303
 - opioids, 303
 - sulfonylureas, 303
 - TCAs, 302
 - treatment of, 302
- priapism, 222–223
 - causes of, 222
 - complications from, 222
 - hospitalization for, 223
 - treatment for, 222–223
- primidone, 152
- primitive reflexes. *See* reflexes, primitive
- PR intervals, 161
- procedural sedation and analgesia (PSA), 30
- propacetamol. *See* acetaminophen
- propofol, 31
- protective reflex, 13
- prothrombin time (PT), 247
 - PTT, 247
- Prune Belly Syndrome, 11
- PSA. *See* procedural sedation and analgesia
- pseudocyst formation, from CSF shunts, 21
- pseudosubluxation, 338–339
- psoriasis, 192
- PT. *See* prothrombin time
- PTT. *See* partial prothrombin time
- pulmonary atresia, 163
- pulseless electrical activity (PEA), 45
 - definition of, 45
 - management of, 45
- pyogenic granulomas, 190–191
- pyruvate kinase deficiency (PKD), 131
- QRS duration, 161
- racemic epinephrine, 274–275
- radial head subluxation, 341, 342
 - hyperpronation method for, 342
 - supination/flexion technique for, 342
- radiology
 - for ALTE, 66
 - for AOM, 85
 - for bone/joint infections, 115
 - for CSF shunts, 21
 - FTT studies with, 98–99
 - for inherited bleeding disorders, 248
 - for pneumococcal bacteremia, 113
 - for SIDS, 53
- rapid-sequence intubation, pediatric, 34–37
 - assessment of, 36–37
 - with capnography, 36
 - medications for, 37
 - induction agents, 35
 - paralytics, 35
 - pre-treatment, 35
 - nasotracheal, 36
 - patient weight and, 37
 - preoxygenation for, 34
 - procedures for, 37
 - ventilator settings in, 37
 - tube sizing in, 34–36
 - complications from, 36
 - of laryngoscopes, 36
- Rashkind procedure, 175
- Rastelli procedure, 175
- reactive airways diseases, 285–287
 - anticholinergics, 286–287
 - complications from, 287
 - lab work for, 285
 - SABA for, 285, 286
 - steroids for, 287
 - supplemental oxygen for, 286
 - symptoms of, 285
 - systemic therapies for, 287
 - treatment for, 285–287
- red reflexes, in eyes, 6
 - colobomas and, 6
- reflexes, primitive, 13
 - crawling, 13
 - Doll’s eyes reflex, 13
 - hand-to-mouth reflex, 13
 - Moro reflex, 13
 - palmar, 13
 - plantar, 13
 - protective, 13
 - red reflexes, 6
 - colobomas and, 6
 - rooting reflex, 13
 - Swimmer’s response, 13
 - tonic neck reflex, 13
 - walking reflex, 13
- Regional Poison Control Centers, 292
- rehydration, 216, 217
- religious beliefs, death practices by, 56
- respiratory emergencies, in infants/neonates, 270–287. *See also* bacterial tracheitis; bronchiolitis; chlamydia; congenital diaphragmatic hernia; croup; epiglottitis; pertussis; pneumonia; retropharyngeal abscess; upper respiratory infections
 - dosing reference guide for, 286
 - from foreign bodies, 271–272
 - epidemiology of, 271
 - management of, 271–272
 - radiography for, 271
 - symptoms of, 271
 - from infections, 272–278
 - bacterial tracheitis, 272–273
 - croup, 273–275
 - epiglottitis, 275–276
 - FDA recommendations for, 272
 - retropharyngeal abscess, 276–278
 - URIs, 83, 272
 - in lower airway, 278–285
 - bronchiolitis, 279–281
 - CDH, 11, 278–279
 - chlamydia, 283–284
 - pertussis, 65, 284–285
 - pneumonia, 281–283
 - pathophysiologies of, 270
 - reactive airways diseases in, 285–287
 - anticholinergics, 286–287
 - complications from, 287
 - lab work for, 285
 - SABA for, 285, 286
 - steroids for, 287
 - supplemental oxygen for, 286
 - symptoms of, 285
 - systemic therapies for, 287
 - treatment for, 285–287
 - from thermal burns, 272
 - from trauma, 271
 - child abuse as, 271
 - upper airway anomalies, 270–271
 - congenital subglottic stenosis, 270–271
 - laryngomalacia, 270
 - vocal cord paralysis, 271
- respiratory syncytial virus (RSV), 64–65
 - bronchiolitis from, 279
- restrictive cardiomyopathy, 173

- resuscitation, pediatric, 40–49. *See also*
 - bradycardia
 - for advanced support, 42
 - breathing assessment in, 42
 - circulation assessment in, 44–48
 - cuffed endotracheal tubes in, 42
 - American Heart Association guidelines for, 40
 - for basic support, 40–42
 - airway patency evaluation in, 40
 - breathing assessment in, 42
 - circulation assessment in, 43–44
 - foreign bodies and, 41
 - opening the airway in, 40–41
 - respiration restoration in, 41–42
 - for burn injuries, 201–202
 - for fluids, 203
 - circulation and, 43–48
 - in advanced support, 44–48
 - for asystole, 45
 - in basic support, 43–44
 - for bradycardia, 44–45
 - for neonates, 48
 - for PEA, 45
 - for tachycardia, 45, 46–47
 - for tachydysrhythmias, 47–48
 - for ventricular fibrillation, 45–46
 - CPR and, 40
 - goals for, 40
 - for neonates, 42–43, 49
 - circulation and, 48
 - post-resuscitative care, 49
- retinal hemorrhages, 75, 310, 312, 313
- retinoblastomas, 6
- retropharyngeal abscess, 276–278
 - complications from, 278
 - epidemiology of, 276–277
 - lab work for, 277
 - management of, 277–278
 - symptoms of, 277
- ringworm. *See* tinea corporis
- ristocetin cofactor assay, 248
- RMSF. *See* Rocky Mountain Spotted Fever
- Rochester criteria, 107
- Rocky Mountain Spotted Fever (RMSF), 192
- rooting reflex, 13
- roseolla infantum, 187
- RSV. *See* respiratory syncytial virus
- rubella, 133–134, 187–188
- rubeola. *See* measles
- SABA. *See* short acting beta antagonists
- salicylate, poisoning from, 297–298
 - incidence rates for, 297
 - pharmacologic effects of, 297–298
 - symptoms of, 298
 - treatment for, 298
 - with activated charcoal, 298
 - with hemodialysis, 298
 - lab testing in, 298
- Salter-Harris classification, of fractures, 336, 337
- Sandifer’s syndrome, 146
- SBS. *See* shaken baby syndrome
- scabies, 184
- scalp hematomas, 327
- scalpringworm. *See* tinea capitis
- scarlet fever, 189
- SCI. *See* silent cerebral infarcts
- SCIWORA. *See* spinal cord injury without radiographic abnormality
- seborrheic dermatitis, 193
- sedation, for neonates, 29–31
 - barbiturates for, 30
 - benzodiazepines for, 30
 - chloral hydrate for, 30
 - dosing guidelines for, 31
 - etomidate for, 31
 - ketamine for, 31
 - propofol for, 31
 - types of, 29–30
 - deep, 30
 - minimal, 29
 - moderate, 29
 - PSA, 30
- seizures, pediatric, 141–154, 155
 - as ALTE, 63–64
 - from CSF shunts, 21
 - definition of, 141
 - differential diagnosis for, 146
 - BHS and, 146
 - dystonic reactions in, 146
 - for narcolepsy, 146
 - for night terrors, 146
 - Sandifer’s syndrome in, 146
 - shuddering attacks in, 146
 - for somnambulism, 146
 - syncopal events in, 146
 - tics in, 146
 - epilepsy and, 141
 - syndromes of, 142–143
 - etiology of, 141
 - febrile, 143, 145
 - antiepileptic drug therapy for, 144
 - complications from, 144
 - imaging for, 144
 - incidence rates for, 143
 - lumbar punctures for, 144
 - pathophysiology of, 144
 - incidence rates for, 141
 - lab testing for, 147
 - lumbar punctures for, 147
 - metabolic demands during, 141–142
 - from minor head injuries, 327
 - neonatal, 145
 - BFNC and, 145
 - BINC and, 145
 - causes of, 145
 - complications of, 145
 - imaging for, 145
 - symptoms of, 145
 - treatment of, 145
 - neuroimaging for, 147–148
 - with EEG, 148
 - patient history and, 147
 - physical exams for, 147
 - treatment of, 148–154
 - with airway stabilization, 148–149
 - with anterior temporal lobectomy, 154
 - with anticonvulsant agents, 151
 - with carbamazepine, 150
 - with clonazepam, 152
 - with corpus callostomy, 154
 - with diazepam, 144, 149
 - with ethosuximide, 152
 - with felbamate, 152
 - with fosphenytoin, 149
 - with gabapentin, 152–153
 - with ketogenic diet, 153–154
 - with lamotrigine, 152
 - with levetiracetam, 153
 - with lifestyle modifications, 154
 - long-term, 150–154
 - with lorazepam, 149
 - with midazolam, 149
 - with oxcarbazepine, 153
 - with phenobarbital, 144, 149, 150–152
 - with phenytoin, 150
 - with primidone, 152
 - with tiagabine, 153
 - with topiramate, 153
 - with vagal nerve stimulation therapy, 154
 - with valproate, 152
 - with valproic acid, 144, 149
 - with vigabatrin, 153
 - with zonisamide, 153
 - types of, 142
 - absence, 142
 - atonic, 142
 - generalized, 142
 - myoclonic, 142
- sensation, evaluation of, 12
- sepsis, 235–237
 - fever from, 3
 - infections from, 236
 - management of, 236, 237
 - streptococcus pneumonia and, 235–236
 - vaccinations and, 236
- septic arthritis, 115, 344
 - causes of, 344
 - diagnosis of, 344
 - radiography for, 344
 - symptoms of, 344
 - treatment for, 344
- severe croup, 275
- severe hypotonia, 13
- sexual abuse, 315–318
 - diagnosis of, 315–316
 - evaluation of, 318
 - evidence testing for, 316–317
 - radiographic evidence for, 318
 - victim history in, 317–318
- SGA. *See* small gestational age
- shaken baby syndrome (SBS), 309–311
 - diagnosis of, 310, 313
 - morbidity/mortality rates of, 310
 - ophthalmic findings with, 310–311
 - retinal hemorrhages in, 75, 310, 312, 313
 - presentation of, 310
- short acting beta antagonists (SABA), 285, 286
- shunts. *See also* cerebrospinal fluid shunts
 - CSF, 18–21
 - components of, 18, .
 - infection from, 20–21
 - leaks in, 21

- malfunction of, 18–20
 - overdrainage by, 21
 - pseudocyst formation from, 21
 - purpose of, 18
 - seizures from, 21
 - ventricular/vascular, 21
 - visceral damage from, 21
- shunt tapping, 20
- sickle cell, emergencies with, 235–241
- ACS and, 240–241
 - definition of, 240
 - management of, 241
 - mortality rate for, 240
 - pathophysiology for, 240
 - symptoms of, 240–241
- acute splenic sequestration, 237
- acute VOC, 237–238
- hydroxyurea and, 238
 - pain in, 237–238
- aplastic crisis and, 237
- causes of, 235
- cerebrovascular accidents and, 239
- SCI, 239
 - strokes, 239–240
- dactylitis, 238–239
- diagnosis of, 235
- ED evaluation of, 235
- HTR, 241
- incidence rates for, 235
- infections and, 236
- race as factor for, 235
- sepsis and, 235–237
- fever from, 3
 - infections from, 236
 - management of, 236, 237
 - streptococcus pneumonia and, 235–236
 - vaccinations and, 236
- symptoms of, 235
- syndromes associated with, 236
- SIDS. *See* Sudden Infant Death Syndrome
- silent cerebral infarcts (SCI), 239
- Simpson-Golabi-Behmel syndrome, 4
- skin, evaluation of, 14
- for acrocyanosis, 14
 - for cutis marmorata, 14
- skull, evaluation of, 4–6
- caput succedaneum and, 5
 - from vacuum extraction, 5
 - cephalhematomas and, 5
 - craniosynostosis in, 5
 - Apert’s syndrome and, 6
 - Crouzon’s disease and, 6
 - FGFR and, 6
 - as inherited, 5–6
 - fontanel size and, 6
 - for fractures, 73, 324–325
 - molding in, 4–5
 - plagiocephaly in, 5
 - with radiographs, 324–325
 - subgaleal bleeds in, 5
- small gestational age (SGA), 4
- hyperbilirubinemia and, 129–130
- Small pox, 182
- SMAOM. *See* Subcommittee on Management of Acute Otitis Media
- somnambulism, 146
- Soto syndrome, 4
- soy-based formula, 79
- spinal cord injury without radiographic abnormality (SCIWORA), 339
- spinal muscular atrophy type 1. *See* Werdnig-Hoffman syndrome
- spleen, 10
- splenic sequestration, acute, 237
- SSSS. *See* staphylococcal scalded skin syndrome
- standardized growth charts, 4
- staphylococcal scalded skin syndrome (SSSS), 180
- steroids
- for bronchiolitis, 280
 - for reactive airway diseases, 287
- Steven-Johnson syndrome, 186
- stork bites, 190
- straddle injuries, 231
- strangulation, 76
- streptococcus pneumonia, 235–236
- strokes, 239–240. *See also* cerebrovascular accidents
- management of, 240
- Sturge-Weber syndrome, 126
- subclavian vein, 352
- Subcommittee on Management of Acute Otitis Media (SMAOM), 83–84, 87
- subconjunctival hemorrhages, 6–7
- subcutaneous fat necrosis of the newborn, 185
- subgaleal bleeds, 5
- subluxation. *See* pseudosubluxation; radial head subluxation
- submersion injuries. *See* drowning/submersion injuries
- sucking blisters, 178–179
- sucrose
- for excessive crying in infants, 78
 - for pain management, 27
- Sudden Infant Death Syndrome (SIDS), 52–53
- AAP recommendations for, 52–53, 61
 - ALTEs v., 61
 - apnea and, 17–18
 - child abuse and, 52, 319
 - demography for, 52
 - diagnosis of, 52–53
 - lab studies for, 53
 - postmortem examinations for, 53
 - radiology studies for, 53
 - sleeping position and, 52
- sulfonylureas, 303
- supination/flexion technique, 342
- supracondylar fractures, 340–341
- imaging for, 340–341
- supraventricular tachycardia, 46–47
- Swimmer’s response reflex, 13
- syncopal events, 146
- syphilis, lesions from, 191–192
- TA. *See* tricuspid atresia
- tachycardia, 45
- algorithm, 47
 - supraventricular, 46–47
 - ventricular, 46
- tachydysrhythmias, 47–48
- pharmacologic therapies for, 47–48
 - tachycardia algorithm, 47
- talipes equinovarus. *See* clubfoot
- TAPVR. *See* total anomalous pulmonary venous return
- TCAs. *See* tricyclic antidepressants
- testicular masses, 227
- testicular torsion, 76, 225–226
- Tetralogy of Fallot (TOF), 163–164
- Blalock-Taussig shunt for, 175
 - Rastelli procedure for, 175
- TGA. *See* Transposition of the Great Arteries
- thermal burns, 272
- thrombocytopenia, 125
- thrombotic thrombocytopenic purpura (TTP), 191
- thyroid disorders, 261–262
- differential diagnosis for, 261–262
 - hyperthyroidism, 261–262
 - hypothyroidism, 261
 - lab work for, 262
 - treatment for, 262
- tiagabine, 153
- tics, 146
- tinea capitis, 193
- tinea corporis, 192
- tinea cruris, 193
- tinea pedis, 193
- tinea versicolor, 184, 192–193
- TM. *See* tympanic membrane
- TNN. *See* transient tachypnea of the newborn
- TNPM. *See* transient neonatal pustular melanosis
- toddler’s fracture, 344–345
- TOF. *See* Tetralogy of Fallot
- tonic neck reflex, 13
- topiramate, 153
- torsion of testicular appendage, 226
- torus fractures, 336, 338
- total anomalous pulmonary venous return (TAPVR), 163, 164–165
- toxicologic emergencies, 291–304. *See also* methemoglobinemia; prescribed medications, poisoning from; salicylate, poisoning from
- AAPCC data for, 291
 - from acetaminophen, 296–297
 - NAC for, 296–297
 - treatment for, 296–297
- American Academy of Toxicologists and, 295
- ancillary evaluation in, 294–295
 - from cleaning agents, 301–302
 - epidemiology of, 301
 - under PPA, 301
 - symptoms of, 302
- European Association of Poison Centers and Clinical Toxicologists and, 295
- gastrointestinal decontamination in, 295
 - AAP guidelines for, 295
 - activated charcoal for, 295
 - incidence rates for, 291
 - intervention programs for, 292
 - through legislation, 292

- toxicologic emergencies (*cont.*)
 - through Regional Poison Control Centers, 292
 - from iron ingestion, 298–299
 - clinical stages of, 299
 - lab testing for, 299
 - treatment for, 299
 - from lead ingestion, 299–300
 - management of, 300
 - sources of, 300
 - symptoms of, 299–300
 - testing for, 300
 - main causes of, 291, 292
 - from methemoglobin inducing substances, 300–301
 - methemoglobinemia and, 300–301
 - mortality rates in, 292
 - from OTC medications, 301
 - camphor, 304
 - CDC and, 301
 - FDA and, 301
 - from prescribed medications, 302–303, 304
 - calcium channel blockers, 302–303
 - clonidine, 303
 - opioids, 303
 - sulfonylureas, 303
 - TCAs, 302
 - primary survey of, 293
 - AVPU scale in, 293
 - decontamination during, 293
 - Glasgow Coma Scale in, 293
 - from salicylate, 297–298
 - incidence rates for, 297
 - pharmacologic effects of, 297–298
 - symptoms of, 298
 - treatment for, 298
 - secondary survey of, 293–294
 - physical examination in, 294
 - toxidromes in, 294
- toxic shock, 190
- toxidromes, 294
- toxins
 - ALTEs from, 66
 - excessive crying in infants from, 73–74
 - breastfeeding modifications for, 79
- toxoplasmosis, 133–134
- tracheostomy tubes, 21–22
 - cuffed, 22
 - dislodgement/obstruction of, 22
 - infections from, 22
- transfusions, for hyperbilirubinemia, 138
- transient neonatal pustular melanosis (TNPM), 183
- transient synovitis, 343–344
 - lab testing for, 343–344
- transient tachypnea of the newborn (TNN), 9
- Transposition of the Great Arteries (TGA), 163, 164
 - Mustard and Senning operation for, 175
- trauma, 307–319. *See also* shaken baby syndrome
 - ALTE from, 66
 - anatomic difference for pediatric patient and, 307
 - from child abuse, 307–319
 - abdominal/thoracic injuries as, 312–313
 - age as factor in, 308–309
 - burns as, 315–316
 - caregivers’ role in, 309
 - categories of, 307–308
 - cutaneous manifestations of, 315–316
 - death review teams for, 319
 - excessive crying in infants and, 78
 - from falls, 314–315
 - family history characteristics and, 307–309, 311
 - fractures and, 73, 311–312, 314
 - incidence rates of, 307
 - iTBI as, 309
 - morbidity rates of, 308
 - mortality rates of, 308
 - physical examinations for, 312
 - radiographic evaluation of, 315
 - SBS as, 309–311
 - sexual abuse as, 315–318
 - SIDS as, 319
 - socioeconomic status and, 308–309
 - swelling/bruising injuries as, 77, 313
 - to CNS, 331
 - excessive crying in infants from, 73
 - incidence rates for, 307
 - in male genitourinary emergencies, 227
 - respiratory emergencies from, 271
- tricuspid atresia (TA), 163, 165
- tricyclic antidepressants (TCAs), 302
 - treatment of, 302
- truncus arteriosus, 163, 165
- TTP. *See* thrombotic thrombocytopenic purpura
- T wave morphology, 161
- tympanic membrane (TM), 84–85
- tympanostomy tubes, 88
- ultrasonography
 - for biliary atresia, 209
 - for head and neck emergencies, 122
 - for minor head injuries, 331–332
 - limitations of, 331–332
- ultrasound or voiding cystourethrogram (VCUG), 11
- umbilical area, evaluation of, 10–11
 - Eagle-Barrett triad in, 11
 - hernias in, 11
 - CDHs, 11
 - hygiene for, 10
 - omphalitis in, 10–11
 - patent urachus in, 11
 - VCUG for, 11
 - Prune Belly Syndrome in, 11
- umbilical vein, 354
 - complications of, 354
- unconjugated bilirubin, 128–129
- unconjugated hyperbilirubinemia, 130
 - management of, 133
- University of California-Davis rule, 329
- upper respiratory infections (URIs)
 - AOM and, 83
 - over-the-counter medications and, 272
 - as respiratory emergency, 272
- urea cycle defects, 257–258
- BUN in, 258
- urethral prolapse, 228–229
 - symptoms of, 228–229
 - treatment for, 229
- urinalysis
 - for excessive crying in infants, 78
 - for IEM, 259
- urinary tract infections (UTIs)
 - ALTEs from, 65
 - fevers from, 3
 - in infants, 112
 - testing for, 108
 - hyperbilirubinemia and, 135
- URIs. *See* upper respiratory infections; upper respiratory infections (URIs)
- urticaria, 186
- UTIs. *See* urinary tract infections
- vaccines, 349
 - sepsis and, 236
 - for varicella, 181
- vagal nerve stimulation therapy, 154
- valproate, 152
- valproic acid, for febrile seizures, 144, 149
 - valproate and, 152
- varicella syndrome, 180–182
 - cause of, 181
 - complications of, 181
 - congenital, 181
 - diagnosis of, 181
 - presentation of, 181
 - treatment for, 181–182
 - vaccine for, 181
 - zoster and, 182
- varicoceles, 11, 224
- vasoocclusive crisis (VOC), acute, 237–238
 - hydroxyurea and, 238
 - pain in, 237–238
 - treatment for, 238
- VCUG. *See* ultrasound or voiding cystourethrogram
- veins
 - femoral, 352
 - jugular, 352
 - subclavian, 352
 - umbilical, 354
 - complications of, 354
- ventricular septal defects (VSD), 10, 167
- ventricular tachycardia, 46
- vigabatrin, 153
- viral exanthem, 186–187, 194
- vital signs, for neonates, 3–4. *See also* weight, of neonates
 - fever in, 3
 - from bundling, 3
 - hypothermia and, 3
 - from meningitis, 3
 - from osteomyelitis, 3
 - from sepsis, 3
 - from UTIs, 3
 - weight in, 3–4
 - average gains in, 4
 - LGA and, 4
 - macrocephaly and, 4
 - microcephaly and, 4

Index		385
SGA and, 4	vulvovaginitis, 230–231	SGA and, 4
standardized growth charts for, 4	risk factors for, 230	standardized growth charts for, 4
vitamin K deficiency, 244–245	symptoms of, 230	Werdnig-Hoffman syndrome, 14
prevention of, 249	treatment for, 230–231	West’s syndrome, 142
symptoms of, 246	VWD. <i>See</i> von Willebrand disease	whey hydrolysate formula, 79
vocal cords, paralysis of, 271		white blood cell (WBC) testing, for fevers, in
vomiting, 328	walking reflex, 13	neonates, 108
von Willebrand disease (VWD), 244	WBC testing. <i>See</i> white blood cell testing	white reflexes. <i>See</i> leukocoria
clotting factor deficiencies and,	weight, of neonates, 3–4	whooping cough. <i>See</i> pertussis
244	average gains in, 4	
desmopressin for, 253	LGA and, 4	X-rays, for infant cardiac emergencies,
etiology of, 244	Beckwith-Wiedemann syndrome and, 4	159–160
prevention of, 253	diabetes and, 4	
ristocetin cofactor assay for, 248	Simpson-Golabi-Behmel syndrome and,	zonisamide, 153
symptoms of, 245, 246	4	zoster, 182
treatment of, 253	Soto syndrome and, 4	complications of, 182
types of, 244	macrocephaly and, 4	diagnosis of, 182
VSD. <i>See</i> ventricular septal defects	microcephaly and, 4	presentation of, 182