

## Index

- abiotic environment, biophysical model  
 adult fish, 96  
 fish eggs and larval models, 89
- acidification  
 climate change effects, phyto- and zooplankton, 328  
 habitat effects, 37–8
- Adriatic Sea  
 fishing collapse analysis, 163–4  
 catch ratios and abundance models, 166–9  
 climate change effects, 181–2  
 effects on top predators, 160, 172  
 flows to detritus model, 172–4f, 175t  
 gelatinous zooplankton simulations, 169–72  
 mixed trophic impact, 174, 175t  
 small pelagic catch reduction and predatory fish enhancement, 180–1  
 trophic levels, 168f  
 stock assessment and management, European anchovy, 201–2
- adult pelagic populations, biophysical models, 95–102  
 abiotic/biotic environment, growth, and mortality, 96  
 climate change, 103  
 full life cycle models, 98–102  
 geographic shifts, 104  
 movement and behavior, 96  
 overview, 88–9  
 physics and physics-related variables, 97–8
- advective transport, biophysical models  
 fish egg and larval dynamics, 90  
 movement and behavior in adult fish, 96
- Aegean Sea, stock assessment and management, European anchovy, 201–2
- aerial fish spotting data, stock assessment and management, 217
- age-dependent fecundity, low-frequency population fluctuations, 291t
- Age-Structured Assessment Program (ASAP), modeling and harvest strategies, California stock, 218–19
- age-structured Sequential Population Analysis, modeling and harvest strategies  
 Benguela operational management, 219–20  
 Humboldt stock, 219
- Agulhas Current  
 Benguela Current ecosystem, 76–7
- biophysical models, climate change, 103–4  
 habitat characteristics, 22  
 pelagic regime shifts and, 78  
 pelagic spawning patterns, 77–8  
 sardine/anchovy habitats, 22–6
- Agulhas Retroflection, 22
- Agulhas Return Current, 22
- albacore, climate-based fishery fluctuations, California ecosystem model, 268
- Aleutian Low pressure system  
 California Current ecosystem regime shifts, 68–70  
 climate change and ocean surface layer stratification, 315  
 decadal-scale population variability, climate-based regime shifts, 82  
 trophic dynamics and climate change in, 139–41
- Allowable Biological Catches (ABC)  
 limitations of, 236  
 modeling and harvest strategies, Japanese stocks, 217–18
- Alosa sapidissima*  
 global harvesting, production, and trade practices, 257–61f  
 trophic dynamics, California Current habitat, 118
- Anchoa mitchilli*, hydrodynamic water quality model, 99–100
- anchovy (anchoveta). *See also* specific species, e. g. *Engraulis* and *Anchoa* spp.  
 Benguela Current habitat, 22–6  
 northern Benguela ecosystem, 77  
 regime shift, 78  
 southern Benguela ecosystem, 77–8  
 spawning changes in, 324
- biophysical models  
 Benguela Current upwelling system, 91–3  
 mortality models, 101  
 NEMURO.SAN simulation model, 102  
 upwelling systems, 93–5
- California Current ecosystem, regime shifts, 70
- Canary Current ecosystem  
 biota changes and population dynamics, 78–9  
 regime shift and geographic distribution, 79–80
- cannibalization of eggs by, 20
- Catalan and Adriatic fisheries, 163–4
- climate-based fishery fluctuations  
 California ecosystem model, 268  
 circulation models, 326  
 distributional dynamics, 321–2  
 decadal-scale population variability  
 Humboldt Current ecosystem regime shift, 75  
 regime shifts, 65, 66f  
 research overview, 64–5
- feeding and food environment, 19–20
- fishing pressures  
 Benguela upwelling region, 161–2  
 predator triggering events, 302
- global harvesting, production, and trade practices, 257–61f
- habitat characteristics, 12–14
- Humboldt Current habitat, 18–20
- inter-species relationships, scale deposition rates and, 58–9
- Kuroshio Current ecosystem, climate-based decadal-scale population variability, 72–4
- low-frequency population fluctuations  
 circumstantial basis for sardine–anchovy alternation  
 nearshore habitat restriction, 289  
 theoretical background, 287
- niche replacement by, 305–6
- Northeast Atlantic habitat, 26–7
- Peruvian fisheries, 7  
 trophic dynamics and climate change, 163
- population changes and habitat characteristics, 34
- as prey species, 20
- sardine–anchovy alternation theory, 7  
 fishing pressures and, 302
- scale deposition rates, 57f, 59f
- site-record relationships of scale depositions for, 57–8
- stock assessment and management, 192–202  
 Benguela anchovy, 199–200  
 Black Sea anchovy, 202  
 California anchovy, 193–8  
 egg production surveys, 213–14  
 European anchovy–Bay of Biscay, 200–1  
 European anchovy–Mediterranean Sea, 201–2  
 Humboldt anchovy, 198–9

- Japanese anchovy, 192–3  
 trophic dynamics  
   Benguela Current, 113–18f  
   climate change and, 147  
   Humboldt Current species, 123–6  
   Northeast Atlantic/Mediterranean systems, 128–33  
   resource partitioning of zooplankton, 137–8  
   summary of research, 146–7  
   wind-driven upwelling and, 37
- Angola–Benguela front  
   Benguela Current ecosystem biota, 76–7  
   habitat characteristics, 22  
   sardine/anchovy habitats, 22–6
- Annual Egg Production (AEP), stock  
   assessment and management, 213–15
- aquaculture  
   climate change effects, California  
   ecosystem management, 270  
   small pelagic fisheries, growth trends for, 274–262
- aquafeeds, global harvesting, production, and trade practices, small pelagic fisheries, 261
- archeological records, pelagic fish population variability, 48
- Arcto-Norwegian cod, biophysical models, 102–3
- Arcto-Norwegian herring stock, modeling and harvest strategies, 232
- artisanal fisheries, global SPF production and economics, 275–6  
   scale issues, 276–8
- Assessment Model Combining Information (AMCI), European sardine, 224
- Atlantic herring, stock assessment and management  
   Baltic Sea, 210–1  
   North Sea autumn spawning, 208–10  
   Norwegian spring spawning stock, 210
- Atlantic Multidecadal Oscillation (AMO), Canary Current ecosystem, climate change-based decadal-scale population dynamics, 80, 81f
- atmospheric forcing, climate change and, 314f
- Bakun's triad  
   climate change and, 325–6  
   empirical models, 326
- Balistes carolinensis*, Canary Current ecosystem populations, 79
- Baltic Sea  
   stock assessment and management  
   Atlantic herring, 210–1, 233–4  
   European sprat, 212–13, 232–3
- hydroacoustic surveys, 215–17
- trophic dynamics, 135–7, 136–42f,  
   140–4f, 146f  
   climate change and, 145  
   physical forcing, 142f
- basic-scale circulation  
   decadal-scale population variability, 64–5  
   climate-based regime shifts, 82  
   synchronies and teleconnections, 82–3  
   research gaps and limitations, 329
- basin hypothesis  
   of pelagic population dynamics, 1–2  
   population expansion, contraction, and refugia, 35  
   site-record relationships of scale  
   depositions and, 57–8
- “basin model” of habitat selection, low-frequency population fluctuations, 293–4
- bay anchovy, hydrodynamic water quality model, 99–100
- Bay of Biscay anchovy  
   modeling and harvest strategies, 221–4  
   Northeast Atlantic habitat, 27–30  
   stock assessment and management, 200–1  
   egg production survey, 213–14  
   hydroacoustic surveys, 215–17
- bearded goby, Northern Benguela ecosystem, 77, 78
- behavioral mechanisms  
   biophysical models  
   adult fish, 96  
   fish egg/larval environment, 91  
   low-frequency population fluctuations, 292–4  
   “basin model” of habitat selection, 293–4  
   entrainment hypothesis, 293  
   fisheries' behavior, 296  
   imprinting, 292–3  
   mixed-species schooling and “school trap” 293  
   school-mix feedback, 293, 301–2
- Benguela anchovy  
   Benguela Current habitat, 22–6  
   biophysical models  
   Benguela Current upwelling system, 91–3  
   miscellaneous upwelling systems, 93  
   Canary Current ecosystem biota, 78–9  
   Catalan and Adriatic fisheries, 163–4  
   modeling and harvest strategies, operational management approach, 219–20  
   mortality models, 101  
   Northeast Atlantic habitat, 27–30  
   stock assessment and management, 199–200  
   egg production survey, 213–14  
   trophic dynamics
- Benguela Current, 113–18  
   Northeast Atlantic/Mediterranean systems, 128–33
- Benguela Current (BC)  
   acidification effects, 37–8  
   anchovy spawning ground changes in, 324  
   biophysical models  
   anchovy/sardine models, upwelling system, 91–3  
   OSMOSE model, 100–1  
   climate change effects on circulation in, 326  
   decadal-scale population dynamics, 76–8  
   biota changes, 76–7  
   northern Benguela, 77  
   regime shifts, 78  
   southern Benguela, 77–8  
   fishery fluctuations, 9  
   fishing collapse analysis, 161–2  
   benthopelagic coupling, 179–80  
   catch ratios and abundances, 166–9f  
   effects on small pelagics, 176–8  
   effects on top predators, 160, 172, 179  
   flows to detritus model, 172–4f, 175t  
   mixed trophic impact, 174, 175t  
   small pelagic catch reduction and predatory fish enhancement, 180–1  
   trophic flow control models, 174–6  
   trophic levels, 168f  
   trophic pathways disruption, 182  
   upwelling intensification and climate change, 183  
   geography and hydrography, 34–5  
   habitat characteristics, 22–6  
   sardine/anchovy habitats, 22–6  
   transport and migration patterns, 35  
   trophic dynamics  
   biomass variability, 138–9  
   decadal-scale population dynamics, 81  
   regime shifts, 78  
   summary of findings, 140–4t  
   upwelling system, 113–18f, 115–16f
- Benguela Jet, 22
- Benguela Niño  
   climate change and fishing collapse, 183  
   pelagic regime shift, 78
- Benguela sardine  
   modeling and harvest strategies, operational management approach, 219–20  
   stock assessment and management, 205–7  
   hydroacoustic surveys, 215–17
- benthopelagic coupling, fishing collapse analysis  
   flow to detritus models, 179–80  
   species proliferation, 178–9  
   trophic pathways disruption, 182

- bioeconomic modeling
  - climate-related fisheries fluctuations in
    - developing coastal countries, 270–1
  - global SPF production and economics, 277–8
- bioenergetics sub-models
  - Baltic sea system, herring and sprat diet, 137
  - fish egg/larval growth, 90
  - physics/physics-related variables, adult fish models, 97–8
- biological overfishing, climate-based fishery fluctuations and, 265
- biological time series, decadal-scale variability research, 64–5
- biomass variability
  - Benguela Current
    - ecosystem, 76–7
    - population dynamics, 26
  - biophysical models, fish egg/larval growth, 90
  - California Current ecosystem, biota changes in, 66–8
  - Canary Current ecosystem, 78–9
  - decadal-scale regime shifts, 65
  - fish scale deposition and, 56
  - full life cycle models, adult pelagic fish, 98–102
  - Humboldt Current ecosystem, 74–5
  - Kuroshio Current ecosystem, 71
  - pelagic:demersal ratios and, 169, 170f
  - phenology and, 37
  - stock assessment and management, 192, 196t
- biophysical models
  - adult pelagic populations, 95–102
    - abiotic/biotic environment, growth, and mortality, 96
    - full life cycle models, 98–102
    - movement and behavior, 96
    - physics and physics-related variables, 97–8
  - climate change and, 102–5
    - research gaps and limitations, 329–30
  - early life stages, 89–91
    - behavior, 90–1
    - Benguela upwelling system anchovy/sardine case study, 91–3
    - fish egg and larval dynamics, 89–90
    - fish egg and larval environment, 89
    - growth patterns, 90
    - mortality, 90–1
    - “non-SPACC” small pelagic fish case study, 95
    - upwelling systems, anchovy/sardine models, 93–5
  - overview and research background, 88–9
  - summary of applications, 105
- biotic environment
  - Benguela Current, ecosystem, 76–7
  - biophysical model
    - adult fish, 96
    - fish eggs and larval models, 89
  - California Current ecosystem, biota changes in, 66–8
  - Canary Current ecosystem, 78–9
  - decadal-scale regime shifts, 65
  - Humboldt Current ecosystem, 74–5
  - Kuroshio Current ecosystem, 71
- bioturbation, chronostratigraphy limitations and, 51
- Black Sea
  - jellyfish predation in, 302
  - modeling and harvest strategies, 234–5
  - stock assessment and management
    - anchovy, 202
    - European sprat, 211
    - hydroacoustic surveys, 215–17
    - trophic flow control and fishing collapse in, 174–6
- blue shrimp, biophysical models, upwelling systems, 93
- BM2 models, adult pelagic fish, 99
- boarfishes, fishing collapse analysis and proliferation of, 178–9
- bottom temperature, Bay of Biscay anchovy spawning, 29
- bottom trawl surveys, stock assessment and management, 217
- bottom-up analysis
  - climate change, trophic productivity and composition predictions, 316–20
  - decadal-scale population variability, 64–5
  - fishing collapse, effects on small pelagics, 176–8
  - future research application, 308
  - low-frequency population fluctuations, 294
  - pelagic regime shifts, 159
  - wasp-waist ecosystem research, 301
- box cores, scale deposition rates and scale shedding, 51–3
- Brazilian sardine, stock assessment and management, 205
- breakout thresholds, fishing collapse research, 300–1
- Brevoortia tyrannus*
  - global harvesting, production, and trade practices, 257–61f
  - physics/physics-related variables, adult fish models, 97–8
  - trophic dynamics, Northwestern Atlantic system, 133–5f, 140–4t
- brown pelicans, climate-based fishery fluctuations, ecosystem models, 266–70
- brown shrimp, biophysical models, upwelling systems, 93
- caducous species, scale deposition rates and scale shedding, 51–3
- California anchovy, 7
  - California Current habitat, 14
  - diet, 119
  - scale deposition rate variability, 45
  - stock assessment and management, 193–8
    - egg production surveys, 213–14
    - modeling and harvest strategies, 218–19
- California Commercial Fish Landing Records (CACom), climate-based fishery fluctuations, 266–70f
- California Cooperative Oceanic Fisheries Investigations (CalCOFI) program
  - egg production surveys, sardine stock, 214–15
  - zooplankton displacement research, 68
- California Current (CC)
  - biophysical models, 94–5
  - climate change effects
    - food web models, trophic dynamics and, 318–19
    - ocean surface layer stratification, 315–16
    - recruitment patterns, 326–7
    - wind intensification, 314
  - decadal-scale population variability
    - biota changes in, 66–8
    - climate-based changes, 70–1
    - regime shifts in, 67, 68–70f
  - expansion, contraction and refugia in, 35
  - geography and hydrography, 34–5
  - habitat characteristics, 14–18
  - hydrology, 37
  - low-frequency population fluctuations
    - cyclonic eddies, 290
    - indeterminate fecundity and forage access, 291
    - latitudinal source water shifts, 290–1
    - life history plasticity, 291–2t
    - sardine upstream movement and offshore habitat utilization, 288
  - phenology, 37
  - physical characteristics, 14
  - population expansion, contraction, and refugia, 35
  - presence/absence of scales in, 56–7
  - sardine population variability and, 47f
  - scale deposition variability in, 55, 60–1
  - site-record relationships of scale depositions in, 57–8

- transport and migration patterns, 35  
 trophic dynamics  
   summary of findings, 140–4t  
   upwelling system, 118–23, 119–20f, 119–20t
- California fisheries  
 climate-based fluctuations, ecosystem models, economic and management implications, 266–70f  
 equilibrium theory and competition models, 7
- California sardine  
 climate change and recruitment patterns, 326–7  
 low-frequency population fluctuations, life history plasticity, 292  
 stock assessment and management, 203  
   modeling and harvest strategies, 218–19  
 upstream movement and offshore habitat utilization, 288
- California Undercurrent, California Current and, 14
- Canary Current  
 climate change and wind intensification in, 314  
 decadal-scale population dynamics, 78–81  
   biota changes, 78–9  
   climate-based mechanisms, 80–1  
   geographic distribution and regime shifts, 79–80  
 European Atlantic sardine habitat, 31  
 latitudinal shifts in, 322–3
- canned and preserved small pelagic fish, global harvesting, production, and trade practices, 260f, 261f
- cannibalization  
 anchovy mortality and, 20, 35–6  
 trophic dynamics and, 143
- Cape gannets, sardine occurrence in, 77–8
- Cape hake, mortality models, 101
- Capros aper*, fishing collapse analysis and proliferation of, 178–9
- capture-based aquaculture, SPF demand and, 274–262
- Carbon-14 values, chronostratigraphy interference, 49–51
- carbon budget  
 fish scale geochemistry, 54, 55–6t  
 trophic dynamics  
   Benguela current pelagics, 117  
   resource partitioning and, 137–8
- carbon dioxide effects  
 acidification and UV radiation, 328  
 climate change and wind intensification, 314
- Carcharodon carcharias*, climate-based fishery fluctuations, ecosystem models, 266–70
- Carcinus maenas*, biophysical models, upwelling systems, 93
- Catalan Sea  
 fishing collapse analysis, 163–4  
 catch ratios and abundance models, 166–9f  
 climate change effects, 181–2  
 effects on top predators, 160, 172, 179  
 flows to detritus model, 172–4f, 175t  
 gelatinous zooplankton simulations, 169–72  
 mixed trophic impact, 174, 175t  
 small pelagic catch reduction and predatory fish enhancement, 180–1  
 species proliferation, 178–9  
 trophic flow control, 174–6  
 trophic levels, 168f  
 trophic pathways disruption, 182  
 stock assessment and management, European anchovy, 201–2
- catastrophe theory, climate-related fisheries fluctuations and, 8–9
- catch and abundance data series  
 fishing collapse analysis and, 164  
 small pelagic fish results, 166–9  
 population variability and, 45  
 stock assessment and management, Brazilian sardine (*brasilensis*), 205
- catch-at-age analysis for science (CANSAR), modeling and harvest strategies, California stock, 218–19
- catching technologies, global SPF production and economics, 280–2
- catch-per-unit-effort (CPUE)  
 climate-related fisheries fluctuations and, 8–9  
 limitations of, 236  
 modeling and harvest strategies, Humboldt stock, 219  
 Pacific sardine, 323f  
 stock assessment and management, 199  
   Chilean common sardine, 205
- catch statistics  
 modeling and harvest strategies, Japanese stocks, 217–18  
 stock assessment and management, 192  
   Benguela sardine, 205–7  
   Chilean common sardine, 205  
   European anchovy, 200–1  
   Japanese sardine, 202–4
- Catch-Survey Analysis (CSA), North Sea sprat, 232
- Chaetoceros*, trophic dynamics, Humboldt Current dietary studies, 118–23
- Chilean common sardine  
 stock assessment and management, 205  
 upstream movement and offshore habitat utilization, 288
- Chilean fisheries, global harvesting, production, and trade practices and, 257–61f  
 human interaction and governance issues, 279
- Chincha period, archeological records of fish populations, 48
- chlorophyll-*a*, dissolved inorganic nitrogen and, 99
- chronostratigraphy  
 fish scale deposition rates and, 49–51  
 surface sediment recovery, 49–51
- circulation effects of climate change, recruitment patterns and, 325–7
- classical period of pelagic fisheries, 6–7
- Climate and Eastern Ocean Systems (CEOS) project, international research co-operation, 1–4
- climate change  
 acidification, 37–8, 328  
 aquaculture growth and, 262  
 Bakun's triad and, 325–6  
 biophysical models, pelagic fish  
   current research, 102–5  
   overview, 88–9  
   research gaps and limitations, 329–30  
 characteristics of, 64–5  
 decadal-scale population dynamics  
   California Current ecosystem, 70–1  
   Canary Current ecosystem, 80–1  
   Humboldt Current ecosystem, 75–6  
   Kuroshio Current ecosystem, 72–4  
   regime shifts, 81–2  
 disease rates and temperature increases, 328  
 distributional changes, 320–5  
 eco-dynamic complexity, 308  
 ecosystem stresses and, 331  
 event-driven, 37  
 fishing collapse and effects of, 181–2  
   individual-based models, 183  
   modeling techniques, 182–3  
   “what-if” climate model, 183  
 global production and economics of small pelagic fish, 262–71  
 in developing coastal countries, 270–1  
 ecosystem economics and management issues, 266–70  
 fisheries collapse and regime shifts, 263–4  
 human dimensions of, 280  
 optimal time path of resource exploitation, discount rates and, 264–6  
 research overview, 256–7  
 sharing of fish stocks and global warming, 262

- climate change (*cont.*)  
 habitat and, 12, 36  
 herring and sprat diet, Baltic Sea system, 136  
 hydrology, 37  
 impact on human communities, 281  
 models of pelagic collapse and, 158–9  
 ocean surface layer stratification, 37, 315–16  
 ocean temperature, 36  
 phenology, 37  
 physical oceanographic future scenarios, 313–16f  
 prediction techniques, research gaps in, 330–1  
 recruitment patterns and, 325–7  
 research gaps concerning, 328–31  
 research summary, 332–3  
 riverine input, storm frequency and intensity and, 327–8  
 SPACC research overview, 312–13  
 spatio-temporal complexity, 330  
 stock assessment and management, 241  
 regime shifts and, 241  
 trophic dynamics and, 139–41  
   Baltic herring/sprat recruitment and, 145  
   menhaden distribution and, 145–6  
   productivity and composition  
     predictions, 316–20  
     summary of research, 147  
 winds and, 37
- Clupea harengus*, stock assessment and management, North Sea spawning stock, 208–10
- clupeoids, 6–7. *See also* herring; sprats  
 early life hydrodynamic models, 95  
 feedback instability and regime shifts and, 302  
 NEMURO.FISH model, 101–2  
 trophic dynamics, 112–13  
   Baltic Sea system, 135–7  
   Humboldt Current dietary studies, 124
- coastal communities  
 climate change impact on, 281  
 developing coastal countries, climate-related fisheries fluctuations in, 270–1  
 global SPF production and economics, human dimensions of, 276–8
- coastal sea levels, low-frequency population fluctuations, synchronous interdecadal mechanisms, 287–8
- cod predators  
 Baltic Sea herring stock assessment and management, 211  
 feedback instability and, 302
- cod stocks, distributional change and climate effects, 319, 321–2f
- cold coastal water (CCW), Humboldt Current ecosystem, climate-based decadal-scale population dynamics, 76
- collapse. *See* fishing collapse analysis
- Cololabis saira*  
 Kuroshio Current ecosystem, biota changes, 71  
 NEMURO.FISH model, 101–2
- Commission for the Conservation of Antarctic Living Marine Resources (CCAMLR), 242
- commodity production, global harvesting, production, and trade practices, small pelagic fisheries, 259f
- “common sense” ideology, future research applications, 300–1
- community characteristics  
 climate change impact on, 281  
   acidification and UV radiation, 328  
   plankton and fish communities, 319–20  
 global SPF production and economics  
   historical background, 278–9  
   scale mismatching, 276–8
- competition models  
 Baltic Sea system, herring and sprat feeding, 136  
 California fisheries, 7  
 fishing collapse analysis, species proliferation, 178–9  
 low-frequency population fluctuations  
   cyclic dominance model, 294–5  
   interspecies competition, 294
- connectivity mechanisms, climate change and circulation patterns, 327
- Continuous Plankton Recorder data, European latitudinal shifts, 323–4
- Continuous Underway Fish Egg Sampler (CUFES), Bay of Biscay anchovy spawning, 29
- contrast harvest rate strategies (CHRS), stock assessment and management and, 240
- copepods  
 in anchovy diet, 119, 123–6, 126–8, 133  
 Baltic sea system, herring and sprat diet, 135–7  
 California Current ecosystem, 68  
 decadal-scale population dynamics, trophodynamic aspects, 81  
 food web models, trophic dynamics and climate change, 318–19  
 Humboldt Current ecosystem, 76  
 Kuroshio Current Extension, 71, 72–4  
 northern anchovy spawning patterns and, 14  
 northern Benguela ecosystem, 77
- in sardine diet, 119–23, 123–6, 126–8  
 sardine migration patterns and, 15–17  
 southern Benguela ecosystem, 77–8
- coring and sampling techniques, surface sediment recovery, 49
- Coscinodiscus*, trophic dynamics, 118–23
- crab, biophysical models, upwelling systems, 93
- cross-disciplinary research, global SPF production and economics, human dimensions of, 275–6
- ctenophore predation, feedback instability and, 302
- cultivation effect, wasp-waist ecosystem models, 301
- curl, wind-stress, 14, 15, 18, 34, 315, 332–3
- currents (oceanic). *See also* specific currents, e.g. California Current  
 low-frequency population fluctuations, synchronous interdecadal mechanisms, 287–8  
 oceanographic climate change, future scenarios, 313–15  
 winds and changes in, 37
- cyclic dominance model, low-frequency population fluctuations, 294–5
- cyclonic eddies, low-frequency population fluctuations, 290
- Cynoscion nebulosus*, predator–prey models, 98
- Daily Egg Production Method (DEPM). *See also* egg production surveys  
 fish recruitment mechanisms, 1–2  
 mortality of species and, 35–6  
 Southwestern Australia sardine modeling, 235  
 stock assessment and management, 213–15  
 anchovies, 213–14  
 Bay of Biscay anchovy, 221–4  
 Benguela anchovy, 199  
 modeling and harvest strategies, California stock, 218–19  
 sardine stock, 214–15
- Daily Fecundity Reduction (DFR), stock assessment and management, 213–15
- decadal-scale population dynamics  
 Benguela Current ecosystem, 76–8  
 biota changes, 76–7  
 northern Benguela, 77  
 regime shifts, 78  
 southern Benguela, 77–8
- California Current ecosystem  
 biota changes in, 66–8  
 climate-based changes, 70–1  
 regime shifts, 67, 68–70f

- Canary Current ecosystem, 78–81  
 biota changes, 78–9  
 climate-based mechanisms, 80–1  
 geographic distribution and regime shifts, 79–80  
 climate-based regime shifts, 81–2  
 climate-related fisheries fluctuations,  
 economic uncertainty linked to, 263–4  
 fishing collapse analysis, 169  
 global SPF production and economics,  
 human dimensions of, 278  
 Humboldt Current ecosystem,  
 biota changes, 74–5  
 climate change and, 75–6  
 regime shifts, 75  
 Kuroshio Current ecosystem,  
 biota changes, 71  
 climate change and, 72–4  
 regime shifts, 71–2  
 timing of events table, 73t  
 nineteenth- vs. twentieth-century  
 paradigms of, 61  
 regime shifts, 65, 66f  
 climate-based mechanisms, 81–2  
 small pelagic fish populations, research  
 overview, 64–5, 65–83  
 stock assessment and management,  
 European anchovy, 200–1  
 synchronies and teleconnections, 82–3  
 temperature effects, 81  
 trophic dynamics, 81, 139–41  
 degradation assessment, scale deposition rates  
 and, 53–5  
 demersal species. *See also* pelagic:demersal  
 fish ratio  
 Catalan and Adriatic overfishing, 163–4  
 climate change, distributional dynamics  
 and, 320–5  
 fishing pressures on  
 Benguela upwelling region, 161–2  
 small pelagic catch reduction and  
 predatory fish enhancement, 180–1  
 demographic effects of fishing, low-frequency  
 population fluctuations and, 296  
 density-dependent response  
 biophysical models, fish egg/larval  
 mortality, 90–1  
 future research in, 303–5  
 Northern Benguela ecosystem biota, 77  
 developing coastal countries, climate-related  
 fisheries fluctuations in, 270–1  
 developmental period of pelagic fisheries, 6  
 developmental time threshold, low-frequency  
 population fluctuations, 289–90  
 diatoms, trophic dynamics  
 California Current dietary studies, 119–23  
 Humboldt Current dietary studies, 118–23  
 dietary studies. *See* feeding ecology  
 dinoflagellates, in anchovy diet, 119  
 disease rates, climate change effects  
 on, 328  
 dissolved inorganic nitrogen (DIN),  
 chlorophyll-*a* and, 99  
 dissolved oxygen-related mortality, NPZ adult  
 fish model, 97–8  
 distributional dynamics  
 climate change and, 320–5f  
 future research in, 302–3  
 latitudinal shifts, 321, 322–3f  
*Doscidus gigas*, low-frequency population  
 fluctuations, predator outbreaks, 295  
 doubt period of pelagic fisheries collapse, 6,  
 7–10  
 down-scaling problems  
 climate change, research gaps concerning,  
 328–9  
 global SPF production and economics,  
 277  
 drift index, sprat recruitment and, 92  
 dynamic prey, biophysical models  
 Benguela Current upwelling system,  
 sardine/anchovy models, 91–3  
 individual-based models, 89  
 early life stages, small pelagic fish  
 biophysical models, 89–91  
 behaviour, 90–1  
 Benguela upwelling system anchovy/  
 sardine case study, 91–3  
 climate change, 102–3  
 fish egg and larval dynamics, 89–90  
 fish egg and larval environment, 89  
 growth patterns, 90  
 mortality, 90–1  
 “non-SPACC” small pelagic fish case  
 study, 95  
 overview, 88  
 spawning habitats, 104  
 upwelling systems, anchovy/sardine  
 models, 93–5  
 climate change effects, 326  
 low-frequency population fluctuations,  
 egg retention and developmental time  
 threshold, 289–90  
 trophic dynamics  
 California Current, 118–19  
 environmental factors, 141  
 Humboldt Current upwelling system, 123–6  
 Northeast Atlantic/Mediterranean  
 systems, 128–9  
 eastern boundary upwelling systems (EBUS),  
 climate change effects on circulation in,  
 322, 326f  
 economic issues  
 climate-based fishery fluctuations,  
 ecosystem models, 266–70  
 global production and trade, small pelagic  
 fish  
 aquaculture operations, 274–262  
 climate change effects, 262–71  
 in developing coastal countries, 270–1  
 ecosystem economics and management  
 issues, 266–70  
 fisheries collapse and regime shifts,  
 263–4  
 optimal time path of resource  
 exploitation, discount rates and,  
 264–6  
 sharing of fish stocks and global  
 warming, 262  
 future research issues, 271–2  
 harvesting, production and trade  
 practices, 257–61  
 human dimensions of, 278–9  
 research overview, 256–7  
 social-ecological research collaborations,  
 280–2  
 economic overfishing, climate-based fishery  
 fluctuations and, 265  
 EcoPath with EcoSim (EwE) family of models  
 adult pelagic fish, 98–102  
 fishing collapse analysis applications,  
 165–76  
 fishing mortality rates, 101  
 EcoSpace model, adult pelagic fish, 98  
 Ecosystem Approach, stock assessment and  
 management, 192, 238, 241–2  
 scientific contributions to, 238  
 ecosystem models. *See also* fishing collapse  
 analysis  
 Benguela upwelling region, 161–2  
 climate-based fishery fluctuations  
 economic and management issues,  
 266–70  
 non-market values and, 265–6  
 climate change and, 308  
 predicted avenues of, 316–28  
 SPACC research overview, 312–13  
 stress interactions, 331f  
 trophic productivity and composition,  
 316–20  
 uncertainty and research gaps  
 concerning, 330  
 collapse indicators, 184n.2  
 small pelagics as, 181  
 decadal-scale population variability, 64–5  
 fishery management systems, 10  
 future research issues, less-studied systems,  
 306  
 global SPF production and economics,  
 human response in West Africa, 279

- ecosystem models (*cont.*)  
 trophic flows, 174–6  
 wasp-waist pelagic fish models, 158–60, 301  
 overview of, 158–60
- Effingham Inlet, inter-species relationships  
 and scale deposition from, 58–9
- egg buoyancy schemes, biophysical models,  
 fish egg/larval behaviour, 91
- egg cannibalism estimates  
 development of, 1–2  
 trophic dynamics, Northeastern Atlantic/  
 Mediterranean systems, 133
- egg distributions, biophysical models, climate  
 change, 103–4
- egg production surveys. *See also* Daily Egg  
 Production Method (DEPM)  
 biophysical models  
 abiotic environment, 89  
 biotic environment, 89  
 stock assessment and management, 213–15  
 anchovy stocks, 213–14  
 sardine stocks, 214–15  
 technology innovations in, 307
- Ekman pumping, wind-driven upwelling and,  
 18, 34, 37
- Ekman transport, 22, 25, 147, 326
- El Niño. *See also* Benguela Niño  
 California Current spawning patterns,  
 17, 18  
 decadal-scale population variability,  
 64  
 trophodynamic aspects, 81  
 fishing collapse and effects of, 183  
 Humboldt Current ecosystem  
 biota changes and, 74–5  
 climate-based decadal-scale population  
 dynamics, 75–6  
 habitat characteristics, 18  
 spawning patterns and, 18–20, 21–2  
 Peruvian anchovy fishery collapse and, 7  
 stock assessment and management,  
 Peruvian anchovy, 199  
 temperature and salinity changes, 34  
 trophic dynamics  
 Humboldt Current dietary studies, 124  
 regime shifts and, 139–41
- El Niño-Southern Oscillation (ENSO)  
 ecosystem variability and human response,  
 West African case study, 279  
 future research applications, 306  
 Humboldt Current and, 18  
 ecosystem effects, 162  
 scale deposition variability and, 45–6, 47f  
 SEPODYM model, 100
- empirical orthogonal function (EOF),  
 ecosystem models, climate-based fishery  
 fluctuations, 266–70f
- energy costs of feeding, trophic dynamics,  
 Benguela current, 117
- Engraulis capensis*, 7
- Engraulis encrasicolus*  
 Benguela Current habitat, 22–6  
 biophysical models  
 Benguela Current upwelling system, 91–3  
 miscellaneous upwelling systems, 93  
 Canary Current ecosystem biota, 78–9  
 Catalan and Adriatic fisheries, 163–4  
 mortality models, 101  
 Northeast Atlantic habitat, 27–30  
 stock assessment and management,  
 199–200  
 Bay of Biscay, 200–1  
 Black Sea, 202  
 egg production survey, 213–14  
 Mediterranean Sea, 201–2  
 trophic dynamics  
 Benguela Current, 113–18  
 Northeast Atlantic/Mediterranean  
 systems, 128–33
- Engraulis japonicus*  
 biophysical models, upwelling systems, 93  
 habitat characteristics, 31–3  
 stock assessment and management, 192–3  
 modeling and harvest strategies, 217–18  
 trophic dynamics, Northwestern Pacific  
 ecosystem, 126–8
- Engraulis mordax*, 7  
 California Current habitat, 14  
 diet, 119  
 scale deposition rate variability, 45  
 stock assessment and management, 193–8  
 egg production surveys, 213–14  
 modeling and harvest strategies, 218–19
- Engraulis ringens*, 7, 13  
 biophysical models, 93  
 upwelling systems, 94  
 Humboldt Current habitat, 18–20, 123–6  
 stock assessment and management, 198–9  
 egg production survey, 213–14  
 modeling and harvest strategies, 219  
 trophic dynamics  
 Humboldt Current upwelling system,  
 123–6
- entrainment hypothesis, low-frequency  
 population fluctuations, 293
- “environmental loophole hypothesis,” low-  
 frequency population fluctuations,  
 295
- environmental processes  
 fishery collapse and, 10  
 fish recruitment mechanisms and, 1  
 global SPF production and economics,  
 future social–ecological research  
 collaborations, 280–2
- Equatorial Undercurrent, Humboldt Current  
 habitat and, 18
- equilibrium theory, California fisheries  
 management and, 7
- equity issues, global SPF production and  
 economics, 280–2
- Eschrichtius robustus*, climate-based fishery  
 fluctuations, ecosystem models, 266–70
- estuarine habitat, trophic dynamics and, 145–6
- ethical issues, global SPF production and  
 economics, 280–2
- Etrumeus teres*, trophic dynamics,  
 Northwestern Pacific ecosystem, 126–8
- Etrumeus whiteheadi*  
 modeling and harvest strategies, Benguela  
 operational management, 219–20  
 mortality models, 101  
 trophic dynamics, Benguela Current, 114
- Eulerian approach, biophysical models  
 fish egg and larval dynamics, 89–90  
 SEPODYM model, 100
- euphausiid eggs, trophic dynamics  
 California Current, pelagic dietary  
 patterns, 119–23  
 Humboldt Current dietary studies, 125–6
- EUREKA project, stock assessment and  
 management, Humboldt anchovy, 199
- European anchovy  
 Benguela Current habitat, 22–6  
 biophysical models  
 Benguela Current upwelling system,  
 91–3  
 miscellaneous upwelling systems, 93  
 biophysical models, upwelling  
 systems, 93  
 Canary Current ecosystem biota, 78–9  
 Catalan and Adriatic fisheries, 163–4  
 mortality models, 101  
 Northeast Atlantic habitat, 27–30  
 stock assessment and management,  
 199–200  
 Bay of Biscay, 200–1  
 Black Sea, 202  
 Mediterranean Sea, 201–2  
 trophic dynamics  
 Benguela Current, 113–18  
 Northeast Atlantic/Mediterranean  
 systems, 128–33
- European sardine  
 biophysical model, 93  
 Canary Current ecosystem  
 biota changes and population dynamics,  
 78–9  
 regime shift and geographic distribution,  
 79–80  
 Catalan and Adriatic seas, 163–4  
 modeling and harvest strategies, 224

- Northeast Atlantic habitat, 30–1  
 stock assessment and management,  
 207–8  
 hydroacoustic surveys, 215–17  
 trophic dynamics, Northeast Atlantic/  
 Mediterranean systems, 128–33
- event-driven climate change, habitats and, 37
- expansion–contraction model, population  
 dynamics, 324f
- expansion of population, habitat  
 characteristics and, 35
- exploitation rate/production rate ratio, low-  
 frequency population fluctuations and,  
 295–6
- export values, global harvesting, production,  
 and trade practices, small pelagic fish, ,  
 260–1f, , 265f
- Extended Survivor Analysis (XSA)  
 Baltic Sea herring, 233–4  
 Baltic Sea sprat, 232–3  
 ICES stocks, 220–1  
 North Sea autumn spawning herring stock,  
 224–32
- ex-vessel revenue, climate-based fishery  
 fluctuations, California ecosystem  
 model, 267–8, 272n.8
- Farfantepenaeus californiensis*, biophysical  
 models, upwelling systems, 93
- fatty acid availability in prey, effects on  
 predators of, 179
- feast-or-famine trophic strategy, Benguela  
 Current anchovy, 118
- fecundity rates, low-frequency population  
 fluctuations  
 age-dependent fecundity, 291t  
 indeterminate fecundity and forage access,  
 291
- Federal Exclusive Economic Zones (EEZ),  
 California fisheries and, 268–9
- feedback instability  
 climate change and, 331f  
 future research methodology, 301–2
- feeding ecology  
 fishing collapse analysis, benthopelagic  
 coupling, 179–80  
 global production and trade, small pelagic  
 fish, human–ecosystem interaction,  
 278–9  
 niche replacement and, 305–6  
 trophic dynamics  
 Baltic Sea system, 135–7  
 Benguela Current, 113–18, 115–16f  
 California Current upwelling system,  
 118–23, 119–20f, 119–20t  
 Humboldt Current upwelling system,  
 118–23
- Northeast Atlantic/Mediterranean  
 systems, 128–33
- Northwestern Atlantic system, menhaden  
 species, 134–5  
 species morphology and, 138  
 summary of research, 146–7
- filtering strategies of pelagics, trophic  
 dynamics  
 Baltic Sea system, 136  
 Benguela current, 116  
 California Current species, 121–3  
 Northeast Atlantic/Mediterranean systems,  
 128–9  
 Northwestern Atlantic menhaden species,  
 133–5  
 resource partitioning and, 138
- Fish Ecology III Conference, 1
- fishery characteristics  
 climate change and global production,  
 263–4, 331f  
 global harvesting, production, and trade  
 practices and, 257–61f  
 low-frequency population fluctuations and,  
 295–6  
 demographic and behavioral effects,  
 296  
 exploitation rate/productive rate ratio,  
 295–6  
 optimal harvesting, 296  
 stock assessment and management, 192  
 summary of research, 347–8
- fishery-independent stock surveys, 213–17  
 aerial fish spotting data, 217  
 bottom trawl surveys, 217  
 driftnet surveys, 217  
 egg production surveys, 213–15  
 anchovy stocks, 213–14  
 sardine stocks, 214–15  
 hydroacoustic surveys, 215–17  
 success of, 236  
 surface trawl surveys, 217
- fishing collapse analysis. *See also* stock  
 assessment and management  
 Atlantic herring, North Sea autumn stock,  
 210  
 basic definition, 184n.1, 184n.2  
 Benguela upwelling region, 161–2  
 benthopelagic coupling, flow to detritus  
 models, 179–80  
 Catalan and Adriatic seas, 163–4  
 catch and abundance data series, 164  
 climate change effects, 181–2  
 individual-based models, 183  
 modeling techniques, 182–3  
 “what-if” climate model, 183  
 Ecopath with Ecosim models, 165–76  
 ecosystem indicators, small pelagics as, 181
- effects on top predators, 160, 172, 179  
 future research issues, 300–1  
 gelatinous zooplankton simulations, 169–72  
 generic effects on small pelagics, 176–8  
 Humboldt Current upwelling system,  
 162–3  
 pelagic food web and, 160  
 predatory fish, small pelagic collapse  
 effects on, 160, 172  
 scale mismatching, 276–8  
 small pelagic catch reduction and predatory  
 fish enhancement, 180–1  
 species proliferation, 178–9  
 “static” (snapshot) models, 164–5  
 time series data models, 165–6  
 trophic flow control models, 174–6  
 trophic pathways disruption, 182
- Fishing-in-Balance (FiB) index  
 fishing collapse analysis, 164, 166–9
- fishmeal/fish oil exports  
 aquaculture growth and, 274–262  
 climate change effects, California  
 ecosystem management, 270  
 global harvesting, production, and trade  
 practices, 261, 272f  
 human dimensions of, 275–6
- fish scale deposition. *See* scale deposition rates
- fitness-based approach, biophysical models,  
 adult pelagic fish  
 movement and behavior models, 96  
 predator–prey models, 98
- fitted models, fishing collapse analysis, time  
 series data, 166
- flow hypotheses  
 decadal-scale population variability  
 climate-based regime shifts, 82  
 synchronies and teleconnections, 82–3  
 low-frequency population fluctuations, 286–9  
 physical variability, 287
- flow to detritus model, fishing collapse  
 analysis, 172–4f, 175t  
 benthopelagic coupling, 179–80
- fold catastrophes, climate-related fisheries  
 fluctuations and, 9
- Food and Agriculture Organization (FAO),  
 international research co-operation, 1–4
- food chain hypothesis, trophic dynamics,  
 112–13
- food supplies, mortality of species and, 35–6
- food web model  
 climate change and, research gaps and  
 limitations, 329–30  
 fishing collapse analysis, trophic flow  
 control, 174–6  
 global production and trade, small pelagic  
 fish, human–ecosystem interaction,  
 278–9

- food web model (*cont.*)  
 trophic dynamics and climate change, 318–19, 320f
- forage models  
 low-frequency population fluctuations, indeterminate fecundity and forage access, 291  
 tuna population, 100
- fossil imprints, surface sediment recovery, coring and sampling techniques, 49
- fresh, frozen, and chilled small pelagic fish, global harvesting, production, and trade practices, 260ff, 261f
- fresh water input, climate change  
 ocean surface layer stratification, 316  
 riverine input, 327–8
- full life cycle models  
 adult pelagic fish  
 definition, 95  
 EcoPath/EcoSim models, 98–102  
 climate change, 103
- Gadus morhua*, biophysical models, 102–3
- gastric evacuation modeling, Baltic sea system, herring and sprat diet, 137
- gelatinous zooplankton, fishing collapse analysis and, 169–72f, 169–72t
- generalized additive models (GAMs)  
 Bay of Biscay anchovy spawning, 29  
 Mediterranean and Black Sea pelagic stock, 234–5  
 trophic dynamics, Humboldt Current dietary studies, 125–6
- General Ocean Circulation Models (GCMs), research gaps and limitations, 329
- geochemistry, fish scales, scale deposition rates and, 55–6
- geographic shifts  
 biophysical models, 104  
 Canary Current ecosystem, pelagic regime shifts and, 79–80  
 distributional dynamics, future research in, 302–3  
 fishing collapse and predator–prey variations in, 179  
 habitat characteristics and, 13–14, 34–5
- Ghana case study, ecosystem variability and human response, 279
- gill raker gap measurements, trophic dynamics  
 Benguela current, 114  
 California Current, 118  
 Kuroshio Current/Northwestern Pacific pelagic comparisons, 127  
 Northeast Atlantic/Mediterranean systems, 128–9
- Northwestern Atlantic system, menhaden species, 133–5  
 species differences in, 138
- Global Ocean Ecosystem Dynamics (GLOBEC)  
 Baltic Sea system, herring and sprat feeding, 136  
 global SPF production and economics, human dimensions of, 275–6  
 retrospective studies, decadal-scale variability research, 65
- global SPF production and economics  
 aquaculture operations, 274–262  
 climate change effects, 262–71  
 in developing coastal countries, 270–1  
 ecosystem economics and management issues, 266–70  
 fisheries collapse and regime shifts, 263–4  
 optimal time path of resource exploitation, discount rates and, 264–6  
 sharing of fish stocks and global warming, 262
- future research issues, 271–2
- harvesting, production, and trade practices, 257–61f
- human dimensions of  
 climate impacts on human communities, 281  
 cross-disciplinary research, 275–6  
 future social–ecological research issues, 280–2  
 historical context, 278–9  
 research overview, 275  
 scale issues, 276–8  
 social–natural science research collaboration, 279–80  
 research overview, 256–7
- global warming  
 ocean surface layer stratification, 316  
 small pelagic fish stocks fluctuations, 262  
 economic issues, 263–4  
 SPACC research review, 313  
 SPF production and economics, human interaction and, 280  
 trophic productivity and composition predictions, 316–20
- goby  
 fishing collapse analysis, 169  
 small pelagic catch reduction and predatory fish enhancement, 180–1
- Northern Benguela ecosystem, 78
- governance  
 climate-based fishery fluctuations and, 263–4  
 global SPF production and economics
- human dimensions of, 275–6  
 scale mismatching, 276–8  
 stock assessment and management and, 239
- gray whales, climate-based fishery fluctuations, ecosystem models, 266–70
- “Group of Four” experts, formation of, 1
- growth algorithms, biophysical models, fish egg and larval dynamics, 90
- guano scraping, global production and trade, small pelagic fish, 278–9
- guillemot, prey quality effects on, 179
- “gulping” feeding behavior, trophic dynamics, Kushio Current/Northwestern Pacific pelagic comparisons, 127
- Gymnodinium splendens*, in anchovy diet, 119
- habitats  
 acidification effects, 37–8  
 anchovy and sardine population shifts, 34  
 anchovy and sardine taxa, 12–14  
 Bay of Biscay anchovy, 27–30  
 Benguela Current, 22–6  
 California Current, 14–18  
 climate change and, 12, 36, 264  
 European Atlantic sardine, 30–1  
 event-driven climate change, 37  
 expansion, contraction, and refugia, 35  
 food and predator characteristics and species mortality, 35–6  
 geography and hydrography of, 13–14, 34–5  
 hydrology, 37  
 Japanese anchovy, 31–3  
 Kuroshio–Oyashio current, 31–4  
 low-frequency population fluctuations  
 “basin model” of habitat selection, 293–4  
 nearshore habitat restriction, 289  
 sardine upstream movement and offshore habitat utilization, 288–9  
 mortality rates and changes in, 35–6  
 Northeast Atlantic, 26–7  
 ocean stratification and changes in, 37  
 ocean temperature and changes in, 36  
 phenology, 37  
 summary of research, 38  
 temperature and salinity properties, 35  
 transport and migration patterns, 35  
 trophic dynamic implications for, 143  
 winds and changes in, 37
- habitat suitability indices (HSI), physics/physics-related variables, adult fish models, 97
- hake, fishing pressures on  
 Benguela upwelling region, 161–2  
 catch ratios and abundance data, 166–9

- small pelagic catch reduction and predatory fish enhancement, 180–1
- Hamburg Shelf Ocean Model (HAMSOM)  
 early life models, small pelagic fish, 95  
 Lagrangian simulations, upwelling systems, 94–5  
 mackerel biophysical model, 95
- harvest control rules, modeling and harvest strategies, California stock, 218–19
- harvesting strategies and modeling  
 climate-related fisheries fluctuations in developing coastal countries, 270–1  
 global production and economics of small pelagic fish, 257–61f  
 commodity production, 259f  
 discounted present values, 265f  
 export values, 260f, 261f  
 future trends in, 271–2
- low-frequency population fluctuations and, 296
- stock assessment and management, 217–42  
 Arcto-Norwegian herring spring spawning, 232  
 Baltic Sea herring, 233–4  
 Baltic Sea sprat, 232–3  
 Bay of Biscay anchovy, 221–4  
 Benguela operational management procedure, 219–20  
 California sardine and anchovy, 218–19  
 European sardine, 224  
 Humboldt fisheries, 219  
 ICES guidelines, 220–1  
 Japanese anchovy, 217–18  
 Mediterranean and Black Sea pelagic fisheries, 234–5  
 North Sea herring, autumn spawning stock, 224–32  
 North Sea sprat, 232  
 Southwestern Australia sardine, 235
- herring. *See also* specific species  
 early life hydrodynamic models, 95  
 feedback instability and population growth of, 302  
 global harvesting, production, and trade practices, 257–61f  
 historical records of, 46  
 inter-species relationships, scale deposition rates and, 58–9  
 mortality models, 101  
 NEMURO.FISH model, 101–2  
 scale deposition rate variability, 51  
 site-record relationships of scale depositions for, 57–8  
 stock assessment and management, 208–11  
 Baltic Sea, 210–1  
 modeling and harvest strategies, 210–1, 224–32, 233–4
- North Sea autumn spawning, 208–10  
 Norwegian spring spawning, 210
- trophic dynamics, Baltic Sea system, 135–7  
 climate change and, 145
- historical records  
 environmental significance of, 56–60  
 nineteenth- vs. twentieth-century paradigms and interpretation of, 61  
 ocean variability indications from, 60–1  
 pelagic fish population variability, 46–8  
 site characteristics and, 57–8
- Hokkaido herring, 6–7
- Holocene period  
 archaeological record of fish populations during, 48  
 climate variability during, 46  
 dietary role of pelagics during, 48  
 presence/absence of scales in, 56–7
- homing instinct in fish, climate-related fisheries fluctuations and, economic uncertainty linked to, 263–4
- horse mackerel  
 Benguela Current population changes, 26, 78  
 fishing collapse analysis, 166–9  
 Northern Benguela ecosystem, 77
- human interaction  
 climate change and, 312–13  
 global SPF production and economics  
 climate impacts on human communities, 281  
 cross-disciplinary research, 275–6  
 ecosystem variability and, West African case study, 279  
 future social–ecological research issues, 280–2  
 historical context, 278–9  
 research overview, 275  
 scale issues, 276–8  
 social–natural science research collaboration, 279–80
- Humboldt anchovy  
 biophysical models, 93  
 upwelling systems, 94
- Humboldt Current habitat, 18–20, 123–6  
 stock assessment and management, 198–9  
 egg production survey, 213–14  
 modeling and harvest strategies, 219
- trophic dynamics  
 Humboldt Current upwelling system, 123–6
- Humboldt Current (HC)  
 biophysical models, upwelling systems, 93
- decadal-scale population variability, 64  
 biota changes, 74–5  
 climate change and, 75–6  
 Kuroshio Current synchronies and teleconnections, 82–3f  
 regime shifts, 75  
 trophodynamic aspects, 81
- fishing collapse analysis  
 benthopelagic coupling, 179–80  
 effects on small pelagics, 176–8  
 flows to detritus model, 172–4f, 175t  
 mixed trophic impact, 174, 175t  
 small pelagic catch reduction and predatory fish enhancement, 180–1  
 trophic flow control, 174–6  
 trophic pathways disruption, 182
- fishing collapse in, 162–3  
 catch ratios and abundance models, 166–9  
 effects on top predators, 160, 172  
 trophic levels, 168f
- geography and hydrography, 34–5  
 mackerel/jack mackerel habitat, 22  
 models of pelagic collapse in, 158–9  
 population expansion, contraction, and refugia, 35  
 regime shifts in, 67f  
 sardine habitat, 21–2  
 transport and migration patterns, 35
- trophic dynamics  
 climate change and, 139–41  
 upwelling system, 123–6f, 124–6f, 125–30t, 140–4t
- Humboldt sardine  
 Benguela Current habitat, 22–6  
 biophysical models, Benguela Current upwelling system, 91–3  
 California Current habitat, 14–17  
 Humboldt Current habitat, 21–2  
 mortality models, 101  
 scale deposition rate variability, 45, 53f  
 stock assessment and management, South Peru/North Chile, 203–5
- trophic dynamics  
 Benguela Current, 113–18  
 Humboldt Current upwelling system, 123–6
- hydroacoustic surveys  
 Bay of Biscay anchovy, 221–4  
 stock assessment and management, 215–17
- hydrodynamics  
 biophysical models  
 Benguela Current upwelling systems, sardine/anchovy models, 91–3

- hydrodynamics (*cont.*)  
 climate change and, 102–3  
 early life models, small pelagic fish, 95  
 fish egg and larval dynamics, 89–90  
 fish egg/larval abiotic environment, 89  
 MARS 3-dimensional circulation, upwelling systems 93  
 NPZ adult fish model, 97–8  
 physics/physics-related variables, adult fish models, 97–8  
 predator-prey models, 98  
 water quality models, 99–100  
 individual-based models, 89  
 three-dimensional model, Pacific Ocean, 100
- hydrography  
 habitat characteristics and, 13–14, 34–5  
 ocean stratification and, 37
- hydrologic cycle, climate change research, 36, 37
- trophic productivity and composition  
 predictions, 316–20
- Iberian Poleward Current (IPC), Northeast Atlantic habitat, 26–7
- ICES stocks, modeling and harvest strategies, 220–1
- “ideal free distribution” hypothesis, low-frequency population fluctuations, 293–4
- IGBEM model, adult pelagic fish, 99
- immigration trends, California fisheries and, 268
- imprinting, low-frequency population fluctuations, 292–3
- individual-based models (IBMs)  
 biophysical models  
 fish egg/larval mortality, 90–1  
 overview, 89  
 fish egg and larval dynamics, 89–90  
 fishing collapse and climate change effects, 183
- Individual Transferable Quota (ITQ), Southwestern Australia sardine modeling and harvest strategies, 235
- Initial Framework hypothesis, low-frequency population fluctuations, 286–9  
 “basin model” of habitat selection, 293–4  
 circumstantial sardine–anchovy alternations, 289  
 cyclic dominance model, 295  
 physical variability, 287  
 synchronous interdecadal fluctuations, ocean circulation, 287–8
- instantaneous surplus production rate (ISPR), Kuroshio Current ecosystem biota and, 71
- Instituto de Fomento Pesquero (IFOP), sardine stock egg production survey, 214–15
- Integrated-Catch-at-Age (ICA) models  
 Baltic Sea sprat, 232–3  
 Bay of Biscay anchovy, 221–4  
 European sardine, 224  
 ICES stocks, 220–1  
 North Sea autumn spawning herring stock, 224–32
- interest groups, climate-based fishery fluctuations and policy options and, 264–5
- Intergovernmental Oceanographic Commission (IOC)  
 international research co-operation, 1–4  
 resolution XI-17, 1
- international agreements  
 climate-based fishery fluctuations and, 263–4  
 in developing coastal countries, 270–1  
 future trends in, 271–2  
 stock assessment and management and, 240–2
- International Council for Exploration of the Sea (ICES) Working Group, entrainment hypothesis, low-frequency population fluctuations, 293
- International Geosphere Biosphere Program (IGBP), decadal-scale variability research, 65
- International Recruitment Project (IREP), development of, 2
- intraguild predation (IGP), trophic dynamics and, 143
- Investigaciones Mexicanas de la Corriente de California (IMECOCAL), stock assessment and management, sardine stock egg production, 214–15
- isotherm movements, distributional dynamics, future research in, 303
- jack mackerel  
 Humboldt Current habitat, 22  
 small pelagic catch reduction and predatory fish enhancement, 180–1
- Japanese anchovy  
 biophysical models, upwelling systems, 93  
 decadal-scale population variability and, 64  
 habitat characteristics, 31–3  
 Kuroshio Current ecosystem, biota changes, 69, 71f  
 stock assessment and management, 192–3  
 modeling and harvest strategies, 217–18  
 trophic dynamics, Northwestern Pacific ecosystem, 126–8
- Japanese fisheries, global harvesting, production, and trade practices and, 257–61f
- Japanese sardine  
 decadal-scale population variability, 64  
 temperature effects, 81  
 Kuroshio Current ecosystem, biota changes, 69, 71f  
 Kuroshio–Oyashio habitat characteristics, 33–4  
 low-frequency population fluctuations, life history plasticity, 292  
 regime shifts of, 67f  
 stock assessment and management, 202–4  
 modeling and harvest strategies, 217–18  
 trophic dynamics, Northwestern Pacific ecosystem, 126–8  
 upstream movement and offshore habitat utilization, 288
- Japan Sea  
 biota changes in, 71
- jellyfish predation  
 Benguela Current population changes, 26  
 feedback instability and, 302  
 fishing collapse analysis and, 169–72f, 169–72t  
 benthopelagic coupling, 179–80  
 species proliferation, 178–9  
 trophic pathways disruption, 182
- jumbo squid, low-frequency population fluctuations, predator outbreaks, 295
- Kasten cores  
 scale deposition rate variability and, 51, 52f  
 slumps and limitations of, 51  
 surface sediment recovery, 49
- kinesis  
 biophysical models, adult pelagic fish, movement and behavior models, 96  
 predator–prey models, 98
- Kuroshio Current Extension (KC Extension)  
 biota changes in, 71  
 climate-based decadal-scale population variability, 72–4  
 regime shifts, 71–2
- Kuroshio Current (KC)  
 biophysical models, 94  
 climate change, ocean surface layer stratification, 316  
 decadal-scale population dynamics  
 biota changes, 71  
 climate change and, 72–4  
 Humboldt current synchronies, 82–3f  
 regime shifts, 71–2  
 temperature effects, 81  
 timing of events table, 73t  
 low-frequency population fluctuations  
 egg retention and developmental time threshold, 289–90  
 future research issues, 296  
 meandering characteristics, 290

- research overview, 286–9
  - sardine upstream movement and offshore habitat utilization, 288
  - trophic dynamics, 126–8, 127–8f, 128–9f, 140–4t
    - climate change and, 139–41
- Kuroshio–Oyashio (KO) current. *See also* Oyashio Current
  - characteristics of, 31–4
  - decadal-scale population variability and, 64
  - geography and hydrography, 34–5
  - phenology, 37
  - population expansion, contraction, and refugia, 35
- Lagrangian approach, biophysical models
  - Benguela current, sardine/anchovy models, 92–3
  - fish egg and larval dynamics, 89–90
  - upwelling systems, , 94–5
- laminae structure, sedimentary pelagic fish records, 50f
- La Niña, trophic dynamics and, 139–41
- larval environment. *See* early life stages, small pelagic fish
  - stock assessment and management
    - Atlantic herring spawning stock, 208–10
    - time series data, 217
- “Lasker windows” model, development of, 1–2
- latitudinal shifts
  - Benguela Current, 324
  - Canary Current, climate change and, 322–3
  - distributional dynamics, future research in, 320
  - Europe, 323–4
  - generalizations concerning, 321, 324–5f
  - low-frequency population fluctuations, 290–1
- life history and reproduction, low-frequency population fluctuations, temperature optima, 292
- Limacina bulmumoides*, anchovy diet, 119
- Lima Workshop, activities of, 1–2
- Litopenaeus stylirostris*, biophysical models, upwelling systems, 93
- Lo Demás archeological site, archeological records of fish populations, 48
- low-frequency population fluctuations anchovy
  - initial framework, 287
  - nearshore habitat restriction, 289
- behavioral mechanisms, 292–4
  - “basin model” of habitat selection, 293–4
  - entrainment hypothesis, 293
  - imprinting, 292–3
- mixed-species schooling and “school trap,” 293
  - school-mix feedback, 293, 301–2
- “environmental loophole hypothesis,” 295
- fisheries’ impact on, 295–6
  - demographic and behavioral effects, 296
  - exploitation rate/productive rate ratio, 295–6
  - optimal harvesting, 296
- initial framework hypothesis, 286–9
  - physical variability, 287
  - synchronous interdecadal fluctuations, ocean circulation, 287–8
- life history and reproduction, 291–2
  - age-dependent fecundity and reproductive age structure, 291
  - indeterminate fecundity and forage access, 291
  - plasticity, 291–2
  - temperature optima, 292
- multispecies mechanisms, 294–5
  - cyclic dominance of pelagic species, 294–5
  - predator outbreaks, 295
  - predator–prey dynamics, 294
  - species competition, 294
  - trophic dynamics, 294
- over-fishing vs. natural fluctuations, 286
- physical processes, 289–91
  - cyclonic eddies, 290
  - Kuroshio Current meandering, 290
  - latitudinal shifts, source water, 290–1
  - retention and critical development time, 289–90
- sardine/anchovy populations
  - circumstantial basis for alternation of, 289
  - future research issues, 296
  - research overview, 285–6
- sardines
  - initial framework, 287
  - upstream movement and offshore habitat utilization, 288–9
- low-latitude distributions, ecosystem analysis, 306
- Lüderitz cell
  - Benguela Current ecosystem
    - biota changes, 76–7
    - upwelling region, 161–2
  - biophysical models, spawning patterns, 92–3
  - climate change effects, 326
- habitat characteristics, , 22–6
- mackerel
  - biophysical model, 95
  - mortality models, 101
  - Humboldt Current habitat, 22
- macro-electronics revolution, SPF research applications, 307
- Macrorhamphosus* species, Canary Current ecosystem populations, 79
- management theory, global SPF production and economics, social–ecological research collaborations, 280–2
- Marbled sale, hydrodynamic model, 97–8
- marine mammals, climate-based fishery fluctuations, California fisheries, 269
- Marine Stewardship Council (MSC), stock assessment and management and, 241
- MARS 3-dimensional circulation model, biophysical models, upwelling systems 93
- match-mismatch hypothesis, trophic dynamics and climate change, 318
- Maximum Likelihood Estimator (MLE), Bay of Biscay anchovy, 221–4
- Mediterranean ecosystems
  - Catalan and Adriatic seas, fishing collapse in, 163–4
  - fishing collapse analysis
    - benthopelagic coupling, 179–80
    - small pelagic catch reduction and predatory fish enhancement, 180–1
  - stock assessment and management
    - egg production survey, 213–14
    - European anchovy, 201–2
    - hydroacoustic surveys, 215–17
    - modeling and harvest strategies, 234–5
  - trophic dynamics, 128–33, 129–34f, 130–2t, 132–40t, 140–4t
    - climate change and, 141–3
    - comparison of indices, 143–6f
- menhaden
  - global harvesting, production, and trade practices, 257–61f
  - physics/physics-related variables, adult fish models, 97–8
  - trophic dynamics, Northwestern Atlantic system, 133–5, 140–4f, 145–6t
- Merluccius capensis*, mortality models, 101
- midden dating
  - archeological records of fish populations, 48
  - presence/absence of scales in, 56–7
- migration patterns
  - biophysical models
    - Benguela Current upwelling system, sardine/anchovy models, 92–3
    - fish egg and larval dynamics, 90
    - fish egg/larval behavior, 91
    - upwelling systems, 93
  - climate-related fisheries fluctuations, economic issues, 263–4
  - habitat characteristics and, 35

- migration patterns (*cont.*)  
 Northern Benguela ecosystem biota, 77  
 Pacific sardine, 15–17  
 scale deposition rates and regional fish abundance variations, 55  
 minimum number of individuals (MIN),  
 archeological records of fish populations, 48  
 mixed layer depth (MLD)  
 climate change, ocean surface layer stratification, 316  
 decadal–scale population variability, climate-based regime shifts, 82  
 Kuroshio Current ecosystem  
 climate-based decadal-scale population variability, 72–4  
 regime shifts and, 71–2  
 mixed-species schooling, low-frequency population fluctuations, 293  
 Mixed Trophic Impact (MTI) analysis  
 defined, 184n.6  
 fishing collapse, 165  
 demersal–pelagic compartments, 174, 175t  
 flows to detritus model, 172–4, 175  
*Mnemiopsis leidyi*  
 feedback instability and, 302  
 Mediterranean and Black Sea pelagic stock modeling and, 234–5  
 molecular genetics, SPF research using, 307  
 mortality of species  
 biophysical models  
 dissolved oxygen-related mortality, 97–8  
 fish egg/larval environment, 90–1  
 habitat change and, 35–6  
 movement of adult fish  
 biophysical models, 96  
 climate change and, 105  
 low-frequency population fluctuations, sardine upstream movement and offshore habitat utilization, 288–9  
 multi-core techniques, surface sediment recovery, 49  
 multidisciplinary research methodology,  
 global SPF production and economics,  
 human dimensions of, 275–6  
 MULTIFAN-CL stock assessment model, 100  
 multispecies mechanisms, low-frequency population fluctuations, 294–5  
 cyclic dominance of pelagic species, 294–5  
 predator outbreaks, 295  
 predator–prey dynamics, 294  
 species competition, 294  
 trophic dynamics, 294  
 Namibia  
 Benguela Current ecosystem  
 Benguela Niño and regime shifts near, 78  
 ecosystem biota, 76–7  
 upwelling region, 161–2  
 fishing collapse in  
 benthopelagic coupling, 179–80  
 pilchard overfishing case study, 265n.6, 272f  
 scale issues concerning, 277  
 trophic flow control, 174–6  
 trophic pathways disruption, 182  
 inter-species relationships and scale deposition off coast of, 58–9  
 pelagic fish distribution of, 25, 48  
 purse-seine fisheries, 278–9  
 sardine upstream movement and offshore habitat utilization near, 289  
 natural mortality, stock assessment and management vs., 237  
 natural science, global SPF production and economics  
 human dimensions of, 275–6  
 social science and, 279–80  
 niche replacement, future research issues, 305–6  
 nitrogen budget  
 dynamically coupled biophysical models, 99–100  
 fish scale geochemistry, 54, 55–6t  
 trophic dynamics  
 Benguela current pelagics, 117  
 resource partitioning and, 137–8  
*Nitzschia*, trophic dynamics, Humboldt Current dietary studies, 118–23  
 non-consumptive use values, climate-based fishery fluctuations and, 264, 272n.4  
 nonlinear feedback mechanisms  
 feedback instability, 301–2  
 fishing collapse research, 300–1  
 low-frequency population fluctuations,  
 school-mix feedback, 293  
 non-market values, climate-based fishery fluctuations, ecosystem models, 265–6  
 non-Small Pelagic Fish and Climate Change (non-SPACC) regions, biophysical models, 88  
 early life models, small pelagic fish, 95  
 overview, 89  
 North Atlantic Central Water (NACW),  
 Canary Current ecosystem, climate change-based decadal-scale population dynamics, 81  
 North Atlantic Oscillation (NAO)  
 Baltic Sea sprat recruitment, 232–3  
 pelagic fish population variability and historical records of, 46  
 trophic dynamics, Baltic Sea system, climate change and, 145  
 Northeast Atlantic (NEA) current  
 geography and hydrography, 34–5  
 habitat characteristics, 26–7  
 hydrology, 37  
 inter-species relationships and scale deposition from, 58–9  
 transport and migration patterns, 35  
 trophic dynamics, 128–33, 129–34f, 130–2t, 132–40t, 140–4t  
 comparison of indices, 143–6f  
 northern anchovy  
 California Current habitat, 14  
 scale deposition rate variability, 45  
 North Pacific Ecosystem Model for Understanding Regional Oceanography (NEMURO), biophysical model, adult fish, 95–6  
 NPZ models, 101–2  
 North Pacific ecosystem regime,  
 climate-based mechanisms, 81–2  
 North Sea, stock assessment and management  
 Atlantic herring, autumn spawning, 208–10  
 hydroacoustic surveys, 215–17  
 modeling and harvest strategies, 224–32  
 sprats, 211–12  
 Northwestern Atlantic system, trophic dynamics, 133–5f, 140–4t  
 Northwestern Pacific ecosystem  
 climate change and ocean surface layer stratification, 316  
 trophic dynamics, 126–8, 127–8f, 128–9f, 140–4t  
 NORWECOM hydrodynamic model  
 Benguela Current upwelling system,  
 sardine/anchovy models, 91  
 early life dynamics, small pelagic fish, 95  
 Norwegian fisheries  
 economics of migration changes and, 264  
 global harvesting, production, and trade practices and, 257–61f  
 Norwegian herring, 6–7  
 stock assessment and management  
 hydroacoustic surveys, 215–17  
 spring stock spawning, 210  
 number of identified specimens (NISP),  
 archeological records of fish populations, 48  
 nutrient–phytoplankton–zooplankton (NPZ)  
 biophysical models, 88  
 climate change, 103  
 individual-based models, 89  
 NEMURO model family, 101–2  
 OSMOSE model, 101  
 SEPODYM model, 100  
 water quality model, 99–100

- nutrient recycling dynamics, water quality model, 99–100
- oceanographic analysis  
 climate change, 313–16f  
 scale deposition rates and, 60–1  
 sedimentary pelagic fish records, 49
- Ocean Science in Relation to Living Resources Program (OSLR), formation of, 1–4
- ocean stratification  
 California Current ecosystem, climate-based decadal-scale population dynamics, 70–1  
 habitat change and, 37
- ocean temperature, habitat change and, 36
- optimal growth temperatures, distributional change and climate effects, 321–2
- optimal harvesting practices, low-frequency population fluctuations and, 296
- optimal stability window hypothesis, climate change and ocean surface layer stratification, 316
- optimal time path, climate-based fishery fluctuations and, 264–6
- OSMOSE model, adult fish biophysical models, 100–1
- overcoat sardine  
 biophysical model, 93  
 Canary Current ecosystem  
 biota changes and population dynamics, 78–9  
 regime shift and geographic distribution, 79–80  
 Catalan and Adriatic seas, 163–4  
 trophic dynamics, Northeast Atlantic/Mediterranean systems, 128–33
- overfishing hypotheses  
 biological vs. economic overfishing, 265, 272n.6  
 classical theories on pelagic collapse and, 6–7  
 climate-related fisheries fluctuations in developing coastal countries, 270–1
- oxygen rates  
 fish scale geochemistry, 55–6  
 sedimentary pelagic fish records, oceanographic settings, 49
- Oyashio Current  
 biota changes in, 71  
 regime shifts and, 72
- Pacific Decadal Oscillation (PDO)  
 decadal-scale population variability, climate-based regime shifts, 82  
 historical records, 56–60
- Pacific ecosystems  
 climate-based regime shifts, 81–2  
 global harvesting, production, and trade practices, small pelagic fisheries, 257–61f
- Pacific hake, scale deposition rate variability, 45
- Pacific sardine  
 California Current habitat, 14–17  
 decadal-scale population dynamics, temperature effects, 81  
 geographic variability, 323f  
 global production and economics, human interaction and, 280  
 low-frequency population fluctuations, upstream movement and offshore habitat utilization, 288  
 scale deposition rate variability, 45, 53f
- Pacific saury, Kuroshio Current ecosystem  
 biota changes, 71  
 regime shifts, 71–2
- paleoecological patterns, scale deposition variability in, 60
- Paleo-oceanography, California sardine biomass, 203
- particulate-based feeding  
 niche replacement and, 305  
 trophic dynamics  
 Baltic Sea system, 136  
 Benguela current, 116–17  
 California Current, 119  
 California Current species, 121–3  
 resource partitioning and, 138
- pelagic:demersal fish ratio. *See also* demersal species  
 Benguela upwelling region, 161–2  
 collapse detection, 158–9  
 fishing collapse analysis and, 166–9f, 170f  
 effects on top predators, 172  
 flows to detritus model, 172–4f, 175t  
 mixed trophic impact, 174, 175t  
 small pelagic catch reduction and predatory fish enhancement, 180–1  
 trophic pathways disruption, 182
- Pelecanus occidentalis californicus*, climate-based fishery fluctuations, ecosystem models, 266–70
- Penaeus duorarum*, predator–prey models, 98
- Peruvian Coastal Current (PCC), Humboldt Current and, 18
- Peruvian Current (PC), Humboldt Current and, 18
- Peruvian fisheries  
 decadal-scale population variability and, 64  
 density dependence and, 304–5  
 global harvesting, production, and trade practices and, 257–61f  
 historical background, 278–9  
 historic collapse of, 6  
 Humboldt Current ecosystem  
 biota changes and decline in, 74–5  
 regime shift, 75  
 trophic dynamics and climate change, 163  
 scientific management of, 7
- Peruvian–German Cooperative Program for Fisheries Investigations (PROCOPA)  
 establishment of, 1  
 Peruvian anchovy ecosystem research, 3
- phenology  
 habitat change and, 37
- phosphorus loading, dynamically coupled biophysical models, 99–100
- physics  
 biophysical models, adult pelagic fish case studies, 97–8  
 definition, 95  
 movement and behavior models, 96  
 low-frequency population fluctuations and role of, 287  
 oceanographic climate change, future scenarios, 313–16f
- phytoplankton  
 Canary Current ecosystem, climate change-based decadal-scale population dynamics, 80–1  
 climate change effects  
 acidification and UV increases, 328  
 ocean surface layer stratification, 315–16  
 decadal-scale population dynamics, trophodynamic aspects, 81  
 Humboldt Current ecosystem, 162  
 trophic dynamics, 133–5  
 climate change and, 316–20  
 community structure and climate change, 319–20  
 Humboldt Current dietary studies, 124–6  
 Kushio Current/Northwestern Pacific pelagic comparisons, 127  
 Northeast Atlantic/Mediterranean systems, 129–31  
 phenology and match–mismatch hypothesis, 318  
 research gaps and limitations, 329–30  
 research overview, 112–13  
 summary of research, 147
- pink shrimp, predator–prey models, 98
- piscivorous fish, fishing pressures on, 160

- piston cores  
 scale deposition rate variability and, 51  
 surface sediment recovery, 49
- plasticity modeling, low-frequency population fluctuations, life history plasticity, 291–2t
- Pleistocene deposits, presence/absence of scales in, 56–7
- Pleuronectes yokohamae*, hydrodynamic model, 97–8
- PLUME configuration, biophysical models, Benguela Current upwelling system, sardine/anchovy models, 91–3
- population variability of pelagics. *See also* decadal-scale population dynamics  
 adult pelagic populations, biophysical models, 95–102  
 age-structured Sequential Population Analysis, 219–20  
 anchovy (anchoveta), habitat changes and, 34  
 archeological records of, 48  
 basin hypothesis, 1–2, 35  
 in Benguela Current habitat, 26  
 catch records and, 45  
 distributional dynamics, future research in, 302–3  
 expansion and contraction, habitat change and, 35  
 expansion–contraction model, 324f  
 fishing collapse analysis and species proliferation, 178–9  
 fish scale deposition rates, 45  
 future research issues, 60  
 historical records of, 46–8  
 low-frequency population fluctuations  
 behavioral mechanisms, 292–4  
 fisheries' impact on, 295–6  
 life history and reproduction, 291–2  
 “loophole hypothesis,” 295  
 multispecies mechanisms, 294–5  
 physical processes, 289–91  
 prehistoric models, 8  
 salinity and temperature and, 34–5  
 sardine, habitat changes and, 34  
 stock assessment and management, 192  
 Virtual Population Analysis model, 6–7, 25
- predator pit loop, wasp-waist ecosystem models, 301
- predator–prey models  
 anchovy populations, 20  
 biophysical models  
 fish egg/larval biotic environment, 89  
 fish egg/larval mortality, 90–1  
 physics/physics-related variables, 98  
 climate-based fishery fluctuations  
 California fisheries, 269  
 ecosystem models, 266–70
- feedback instability, 301–2
- fishing collapse analysis  
 catch reduction and predatory fish enhancement, 180–1  
 quality of prey factors, 179  
 small pelagic fish effects, 172, 179  
 species proliferation, 178–9  
 trophic flow control models, 174–6
- food web models, trophic dynamics and climate change, 318–19f
- full life cycle models, 99
- low-frequency population fluctuations, 294
- predator outbreaks, 295
- mortality of species, 35–6
- phenology and match–mismatch hypothesis, 318
- scale deposition rates and scale shedding, 51–3
- Southern Benguela pelagic population dynamics, 77–8
- stock assessment and management, Baltic Sea herring, 211
- von Bertalanffy equation, 100
- prediction techniques  
 research gaps in, 330–1  
 trophic productivity, 316–20
- prehistoric sardine/anchovy population models, 8
- pre-season forecasting, stock assessment and management, 237
- prey populations. *See also* predator–prey models  
 biophysical models  
 fish egg/larval biotic environment, 89  
 fish egg/larval growth, 90  
 density dependence, future research in, 303–5  
 trophic dynamics  
 California Current, 119–23  
 Humboldt Current, 118–23  
 Kuroshio Current/Northwestern Pacific pelagic feeding, 126–8  
 Northeast Atlantic/Mediterranean systems, 129–31  
 resource partitioning and, 137–8
- prey-to-predator feedback loop  
 tuna population, 306  
 wasp-waist ecosystem models, 301–2
- Princeton Ocean Model (POM), early life models, small pelagic fish, 95
- production/biomass P/B ratios, full life cycle models, 99
- production statistics  
 global production small pelagic fish  
 production and trade, 257–61f  
 commodity production, 259f  
 discounted present values, 265f  
 export values, 260f, 261f
- low-frequency population fluctuations  
 and, exploitation rate/production rate ratio, 295–6
- Pseudocalanus acuspes*, herring and sprat diet, Baltic Sea system, 135–7
- pteropod, anchovy diet, trophic dynamics, California current, 119
- public held fisheries, climate-based fishery fluctuations and, 265–6
- quantitative fishery dynamics, classical theories on pelagic collapse and, 6–7
- quota systems, global harvesting, production, and trade practices and, 279
- quotient analysis, 25, 28–9
- radioisotopes, chronostratigraphy interference from, 49–51
- recruitment experiment (REX) framework, 1
- recruitment patterns  
 Benguela habitat, 25  
 full life cycle models, 99  
 Catalan and Adriatic overfishing, 163–4  
 classical theories on pelagic collapse and, 7  
 climate change and  
 circulation effects, 325–7  
 distributional dynamics, 321–2  
 environment-based analysis, 237  
 European Atlantic sardine, 30–1  
 international co-operation on research in, 1  
 low-frequency population fluctuations,  
 retention of eggs and critical development times, 289–90
- MULTIFAN-CL stock assessment model, 100
- NEMURO.FISH model, 101–2
- regulating mechanisms, 1
- stock assessment and management, 191–2  
 Baltic Sea herring, 211  
 European sardine, 207–8  
 European sprat, 212–13  
 hydroacoustic surveys, 215–17  
 Japanese sardine, 202–4  
 Southwestern Australia sardine, 208  
 survey-based analysis of, 236–7  
 trophic dynamics and, 139–41  
 Baltic Sea herring/sprat recruitment and, 145
- Recruitment-per-Spawner (RPS)  
 measurements, modeling and harvest strategies, Japanese stocks, 217–18
- red-eye round herring, trophic dynamics, Benguela Current, 114
- refugia  
 acidification effects, 37–8  
 habitat change and, 35

- regime/regime shift  
 Benguela Current ecosystem, 78  
 Canary Current ecosystem, 79–80  
 climate-related fisheries fluctuations and, 8, 10  
 aquaculture growth and, 262  
 California ecosystem management, 270  
 economic issues, 263–4  
 future trends in, 271–2  
 decadal-scale population variability, 65, 66f  
 California Current ecosystem, 68–70  
 climate-based mechanisms, 81–2  
 Humboldt Current ecosystem, 75  
 Kuroshio Current ecosystem (KCE), 71–2  
 research overview, 64  
 defined, 65–6  
 feedback instability and, 301–2  
 historical records and, 56–60  
 low-frequency population fluctuations  
 life history plasticity, 291–2  
 research overview, 753.5  
 NEMURO.FISH model, 101–2  
 stock assessment and management and, 239–40  
 regime-specific harvest rates (RSHR), stock assessment and management and, 239–40  
 Regional Ocean Model System (ROMS)  
 Benguela Current upwelling system, sardine/anchovy models, 91–3  
 biophysical models  
 climate change and, 102–3  
 upwelling systems, 93  
 velocity measurement and modeling, 93  
 regulatory policies, California fisheries and, 268–9  
 replicate sampling techniques, scale deposition rates and scale shedding, 53  
 reproductive age structures, low-frequency population fluctuations, 291  
 research methodology  
 future research trends, 300–1  
 climate change and eco-dynamics, 308  
 density dependence, 303–5  
 distributional dynamics, 302–3  
 feedback instability, 301–2  
 less-studied ecosystems, 306  
 new technologies and techniques, 306–7  
 niche replacement, 305–6  
 wasp-waist ecosystems, 301  
 international co-operation in, 1  
 resource exploitation, optimal time path, climate-based fishery fluctuations and, 264–6  
 resource partitioning  
 filter vs. particulate feeding strategies, 138  
 Northwestern Atlantic system, menhaden species, 134–5  
 prey size and, 137–8  
 trophic dynamics, summary of research, 147  
 retention of planktonic eggs, low-frequency population fluctuations, 289–90  
 rhomboidal modeling, research gaps and limitations, 329  
 riverine input, climate change effects, 327–8  
 round herring  
 fishing collapse analysis in Benguela Current, 169  
 small pelagic catch reduction and predatory fish enhancement, 180–1  
 modeling and harvest strategies, Benguela operational management, 219–20  
 trophic dynamics, Northwestern Pacific ecosystem, 126–8  
 Russia/USSR fisheries, global harvesting, production, and trade practices and, 257–61f  
 salinity. *See also* sea surface salinity (SSS)  
 herring and sprat diet, Baltic Sea system, 136, 145–6  
 pelagic fish population shifts and, 34–5  
 salmon migrations, climate change and economics of, 264  
 salp biomass, California Current ecosystem, climate-based decadal-scale population dynamics, 70–1  
 Santa Barbara Basin (SBB)  
 archeological records of fish populations, 48  
 inter-species relationships and scale deposition in, 58–9  
 scale deposition variability in, 45, 55, 60–1  
 sedimentary pelagic fish records, 49  
 site-record relationships of scale depositions in, 57–8  
*Sardina pilchardus*  
 biophysical model, 93  
 Canary Current ecosystem  
 biota changes and population dynamics, 78–9  
 regime shift and geographic distribution, 79–80  
 Catalan and Adriatic seas, 163–4  
 distributional dynamics, 321–2f  
 modeling and harvest strategies, 224  
 stock assessment and management, 207–8  
 trophic dynamics, Northeast Atlantic/Mediterranean systems, 128–33  
 Sardine–Anchovy Recruitment Program (SARP), development of, 2  
*Sardinella* spp.  
 Canary Current ecosystem  
 biota changes and population dynamics, 78–9  
 regime shift and geographic distribution, 79–80  
 habitat characteristics, 30–1  
 stock assessment and management, Brazilian sardine (*brasilensis*), 205  
 triggering mechanisms for collapse of, 302  
 sardines. *See also* specific species, e.g. *Sardinops* and *Sardina* spp.  
 Benguela Current habitat, 22–6  
 northern Benguela ecosystem, 77  
 regime shift, 78  
 southern Benguela ecosystem, 77–8  
 biophysical models  
 Benguela Current upwelling system, 91–3  
 mortality models, 101  
 NEMURO.SAN model, 102  
 in upwelling systems, 93–5  
 California Current ecosystem  
 biota changes and population fluctuations, 66–8  
 regime shifts in, 70  
 California fisheries, fluctuations in, 269  
 Canary Current ecosystem  
 biota changes and population dynamics, 78–9  
 climate change-based decadal-scale population dynamics, 80–1  
 regime shift and geographic distribution, 79–80  
 classical period of collapse of, 6–7  
 climate-based fishery fluctuations  
 circulation models, 326  
 distributional dynamics, 321–2f  
 ecosystem models, 266–70f  
 decadal-scale population variability  
 Humboldt Current ecosystem regime shift, 75  
 regime shifts, 65, 66f  
 research overview, 64–5  
 El Niño conditions and collapse, 7  
 fishing pressures  
 Benguela upwelling region, 161–2  
 global harvesting, production, and trade practices, 257–61f  
 human interaction and, 280  
 habitat characteristics, 12–14  
 historical observations of, 46–8  
 Humboldt Current ecosystem, 21–2, 74–5  
 inter-species relationships, scale deposition rates and, 58–9  
 Kuroshio Current ecosystem, climate-based decadal-scale population variability, 72–4

- sardines. (*cont.*)  
 low-frequency population fluctuations  
 circumstantial basis for sardine–anchovy alternation, 289  
 indeterminate fecundity and forage access, 291  
 theoretical background, 287  
 upstream movement and offshore habitat utilization, 288–9  
 niche replacement by, 305–6  
 Northeast Atlantic habitat, 26–7  
 population changes and habitat characteristics, 34  
 regime shifts in, 67f  
 Kuroshio Current ecosystem, 71–2, 74f  
 sardine–anchovy alternation theory, 7  
 scale deposition rates, 52, 53f, 54–9f  
 site-record relationships of scale depositions for, 57–8  
 stock assessment and management, 202–8  
 Benguela sardine, 205–7  
 Brazilian sardine, 205  
 California sardine, 203  
 Chilean common sardine, 203–5  
 egg production surveys, 214–15  
 European sardine, 207–8  
 Japanese sardine, 202–4  
 Southwestern Australia sardine, 208  
 trophic dynamics  
 Benguela Current, 113–18  
 climate change and, 139–41, 147  
 Humboldt Current species, 123–6  
 Northeast Atlantic/Mediterranean systems, 128–33  
 resource partitioning of zooplankton, 137–8  
 summary of research, 146–7  
*Sardinia pilchardus*, Northeast Atlantic habitat, 30–1  
*Sardinops melanostictus*, 6–7  
 biophysical models, 93, 94  
 Kuroshio–Oyashio habitat characteristics, 33–4  
 stock assessment and management, 202–4  
 modeling and harvest strategies, 217–18  
 trophic dynamics, Northwestern Pacific ecosystem, 126–8  
*Sardinops ocellata*, scale deposition rate variability, 53  
*Sardinops sagax*  
 Benguela Current habitat, 22–6  
 biophysical models, Benguela Current upwelling system, 91–3  
 California Current habitat, 14–17  
 climate change and recruitment patterns, 326–7  
 Humboldt Current habitat, 21–2  
 low-frequency population fluctuations, upstream movement and offshore habitat utilization, 288–9  
 mortality models, 101  
 scale deposition rate variability, 45, 53f  
 stock assessment and management  
 Benguela sardine, 205–7  
 modeling and harvest strategies, 218–19  
 South Peru/North Chile, 203–5  
 trophic dynamics  
 Benguela Current, 113–18  
 Humboldt Current upwelling system, 123–6  
*Sardinops sagax caerulea*, 6–7  
 stock assessment and management, 203  
*Sardinops sagax neopilchardus*, 208  
*Sardinops sagax ocellata*, 6–7  
*Sardinops sagax sagax*, 7  
 SARDYN (SARDine DYNamics) project, European Atlantic sardine studies, 30–1  
 scale deposition rates (SDR)  
 distributional dynamics, future research in, 302–3  
 scale deposition rates (SDRs)  
 biomass variability calibrations, 56  
 chronostratigraphies, 49–51  
 El Niño conditions and, 45–6, 47f  
 interpretation guidelines, 51–6  
 inter-species relationships, 58–9  
 Kuroshio Current ecosystem, biota changes, 71  
 ocean variability implications of, 60–1  
 pelagic fish population variability and, 45  
 population variability and, 45  
 presence/absence as variability indicator, 56–7  
 regional fish abundance variations, 55  
 scale degradation and, 53–5  
 scale geochemistry and, 55–6  
 scale shedding and, 51–3  
 sedimentary environment models, 47f  
 site-record relationships, 57–8  
 site variability of, 51  
 timescales of variability, 53, 59–60t  
 scale issues, global SPF production and economics, human dimensions of, 276–8  
 schooling behavior, low-frequency population fluctuations, 293  
 school-mix feedback hypothesis  
 low-frequency population fluctuations, 293  
 wasp-waist ecosystem models, 301–2  
 school size, density dependence and, 304  
 school “trap,” low-frequency population fluctuations, 293  
 Scientific Committee on Oceanic Research (SCOR), Working Group 67 and 1  
 scientific research  
 ecosystem approach to stock management, 238  
 global SPF production and economics  
 human dimensions of, 275–6  
 social science and, 279–80  
 Small Pelagic Fish and Climate Change (SPACC) project and, 345–7  
 stock assessment and management detached from, 235–6  
*Scomber japonicus*  
 Canary Current ecosystem populations, 79  
 mortality models, 101  
 seabird populations  
 climate-based fishery fluctuations, California fisheries, 269  
 feeding interactions of SPF, guano scraping and, 278–9  
 fishing collapse effects on, 172, 179  
 sea lions, fishing collapse effects on, 172  
 SEAPOPDM application, biophysical models, 100  
 SeaStar model  
 Arcto-Norwegian herring stock, 232  
 ICES stocks, 220–1  
 sea surface salinity (SSS)  
 Bay of Biscay anchovy spawning, 27–30  
 ocean surface layer stratification, 316  
 sea surface temperature (SST)  
 Aleutian Low pressure system, trophic dynamics and, 139–41  
 Bay of Biscay anchovy spawning, 27–30  
 California Current ecosystem regime shifts, 68  
 Canary Current ecosystem, climate change-based decadal-scale population dynamics, 81  
 decadal-scale population dynamics and, 81  
 climate-based regime shifts, 82  
 forage models, 100  
 Humboldt Current ecosystem  
 climate-based decadal-scale population dynamics, 75–6  
 fishing productivity and, 162  
 Kuroshio Current ecosystem, regime shifts and, 71–2, 76f  
 modeling and harvest strategies  
 California stock, 218–19  
 Japanese stocks, 217–18  
 ocean surface layer stratification, 316  
 trophic dynamics and sardine/anchovy regime shifts, 146  
 sedimentary environments  
 chronostratigraphies and fish scale deposition rates, 49–51

- coring and sampling techniques, 49
  - oceanographic settings, 49
  - presence/absence of scales in, 56–7
  - sardine population variability and, 46–8f
  - scale deposition rates and regional fish abundance variations, 55
  - variability of, SDR and chronostratigraphy estimates and, 51
- sensor arrays, SPF research using, 307
- SEPODYM model
  - geographic shifts, 104
  - tuna population dynamics, 100
- sequential t-test analysis to detect regime shifts (STARS) algorithm, NEMURO-3D model, 101–2
- shad
  - global harvesting, production, and trade practices, 257–61f
- site characteristics, historical records and, 57–8
- size-base partitioning. *See* resource partitioning
- Skeletonema*, trophic dynamics, Humboldt Current dietary studies, 118–23
- slumps, chronostratigraphy limitations and, 51
- Small Pelagic Fish and Climate Change (SPACC) project, 3–4
  - biophysical models, 88
  - overview, 89
  - climate change research, 36, 332–3
    - acidification and UV radiation effects, 328
    - future effects, research overview, 312–13
    - gaps in, 328–9
  - exploited fisheries research, 13
  - future research applications, 300–1, 308–9
  - global SPF production and economics
    - human dimensions of, 275–6
    - social–ecological research collaborations, 280–2
  - habitat comparisons, 12
  - scientific research and, 345–7
  - summary of habitat research, 38
  - synthesis and perspective, 344–9
  - trophic dynamics research summary, 146–7
- small pelagic fisheries, scientific history of, 6–10
- snapshot models, fishing collapse analysis, 164–5
  - flows to detritus data, 172–4f, 175t
  - time series data, 166
  - trophic pathways disruption, 182
- snipefish, Canary Current ecosystem populations, 79
  - snoek, sardine occurrence in, 77–8
  - social discount rates, climate-based fishery fluctuations and, 264–6, 272n.5
  - socially optimal harvest levels, climate-based fishery fluctuations, 269, 270f
  - social science research, global SPF production and economics
    - future social–ecological research collaborations, 280–2
    - human dimensions of, 275–6
    - scientific interaction with, 279–80
  - societal values, climate-based fishery fluctuations and, 264, 272n.4
- Soledad Basin, sedimentary pelagic fish records, 49
- Soutar box core, surface sediment recovery, 49
- South African anchovy fisheries, 7
  - Ecosystem Approach to, 238
- South African sardine fisheries, 205–7
  - upstream movement and offshore habitat utilization, 288
- South Atlantic Central Water (SACW), Canary Current ecosystem, climate change-based decadal-scale population dynamics, 81
- Southeast Asia, climate-related fisheries fluctuations in, 270–1
- Southern Africa Experiment (SAfE), Benguela Current upwelling system, sardine/anchovy models, 91
- Southern Extension of the Cromwell Current (SECC), Humboldt Current and, 18
- Southern Oscillation Index, future research applications, 306
- Southwestern Australia sardine
  - modeling and harvest strategies, 235
  - stock assessment and management, 208
- spatio-temporal scaling, climate change research, gaps and limitations, 330
- spawning habitat (SH)
  - Benguela Current, climate change and, 324
  - biophysical models, early life stages, small pelagic fish, 104
  - Humboldt Current ecosystem, biota changes and, 74–5
  - low-frequency population fluctuations, sardine upstream movement and offshore habitat utilization, 288–9
  - population expansion, contraction, and refugia, 35
- spawning patterns
  - Bay of Biscay anchovy habitat, 27–30
  - Benguela habitat, 23–5
  - biophysical models
    - Benguela Current upwelling system, sardine/anchovy models, 92–3
    - temporal and spatial patterns, 92
  - California Current, 18
  - Canary Current ecosystem biota, 78–9
  - climate change effects on circulation and, 326
  - Humboldt Current, 18–20
  - Japanese sardine, 33–4
  - NEMURO.FISH model, 101–2
  - Northern Benguela ecosystem, 77
  - Southern Benguela ecosystem, 77–8
  - stock assessment and management
    - Atlantic herring, North Sea autumn stock, 208–10
    - Atlantic herring, Norwegian spring stock, 210
    - Benguela anchovy, 199–200
    - Benguela sardine, 205–7
    - Black Sea, 202
    - California anchovy, 193–8
    - California sardine, 203
    - Chilean common sardine, 205
    - European anchovy, 200–1
    - European sardine, 207–8
    - Humboldt sardine, 203–5
    - Japanese anchovy, 192–3
    - Japanese sardine, 202–4
  - Spawning Stock Biomass (SSB)
    - modeling and harvest strategies
      - Arcto-Norwegian herring stock, 232
    - Benguela operational management, 219–20
    - Japanese stocks, 217–18
    - Southwestern Australia sardine, 235
    - stock assessment and management
      - egg production surveys, 214–15
      - hydroacoustic surveys, 215–17
      - Japanese sardine, 202–4
- species interrelationships
  - climate-based fishery fluctuations, California ecosystem model, 266–8
  - fishing collapse analysis and species proliferation, 178–9
  - low-frequency population fluctuations, competition mechanisms, 294
  - scale deposition rates and, 58–9
- species replacement theory, 7
- “spotlight and searchlight” technique, global SPF production and economics, social science/natural science collaborations, 279–80
- spotted seatrout, predator–prey models, 98
- sprats
  - drift index and recruitment patterns, 92
  - early life hydrodynamic model, 95

- global harvesting, production, and trade practices, 257–61f
- modeling and harvest strategies
  - Baltic Sea sprat, 232–3
  - North Sea sprat, 232
- prey quality of, 179
- stock assessment and management
  - Black Sea, 211
  - North Sea, 211–12
- trophic dynamics, Baltic Sea system, 135–7
  - climate change and, 145
- Sprattus sprattus*
  - drift index and recruitment patterns, 92
  - early life hydrodynamic model, 95
  - filter feeding patterns, 128–9
  - global harvesting, production, and trade practices, 257–61f
  - prey quality of, 179
  - stock assessment and management
    - Black Sea, 211
    - North Sea, 211–12
  - trophic dynamics, Baltic Sea system, 135–7, 145
- squid, Kuroshio Current ecosystem, biota changes, 71
- “stable ocean hypothesis,” fish recruitment mechanisms and, 1
- steady-state trophic strategy, Benguela Current sardines, 117
- stochastic equations
  - Baltic Sea herring models, 233–4
  - Baltic Sea sprat modeling, 232–3
  - biophysical models, fish egg and larval dynamics, 90
- stock assessment and management
  - anchovy stocks, 192–202
    - Benguela anchovy, 199–200
    - Black Sea anchovy, 202
    - California anchovy, 193–8
    - European anchovy–Bay of Biscay, 200–1
    - European anchovy–Mediterranean Sea, 201–2
    - Humboldt anchovy, 198–9
    - Japanese anchovy, 192–3
  - climate change and, 241, 263–4
  - current projects and limitations, 235–42
  - ecosystem approach, 238
  - fishery-independent surveys, 213–17
    - aerial fish spotting data, 217
    - bottom trawl surveys, 217
    - driftnet surveys, 217
    - egg production surveys, 213–15
    - hydroacoustic surveys, 215–17
    - surface trawl surveys, 217
  - herring, 208–11
    - Baltic Sea, 210–1
    - North Sea autumn spawning, 208–10
    - Norwegian spring spawning, 210
  - modeling and harvesting strategies, 217–42
    - Arcto-Norwegian herring spring spawning, 232
    - Baltic Sea herring, 233–4
    - Baltic Sea sprat, 232–3
    - Bay of Biscay anchovy, 221–4
    - Benguela operational management procedure, 219–20
    - California sardine and anchovy, 218–19
    - European sardine, 224
    - Humboldt fisheries, 219
    - ICES guidelines, 220–1
    - Japanese anchovy, 217–18
    - Mediterranean and Black Sea pelagic fisheries, 234–5
    - North Sea herring, autumn spawning stock, 224–32
    - North Sea sprat, 232
    - Southwestern Australia sardine, 235
  - recruitment and classical collapse theory, 6–7
  - research overview, 191–2
  - sardine stocks, 202–8
    - Benguela sardine, 205–7
    - Brazilian sardine, 205
    - California sardine, 203
    - Chilean common sardine, 205
    - Chilean sardine (South Peru/North Chile), 203–5
    - European sardine, 207–8
    - Japanese sardine, 202–4
    - Southwestern Australia sardine, 208
  - species stock structure, population trends, and fishery information, 192, 196t
    - Black Sea, 211
    - North Sea, 211–12
- storm frequency and intensity, climate change and, 327–8
- Strangomera bentincki*, stock assessment and management, 205
- subsurface processes, decadal-scale population variability, 64–5
  - climate-based regime shifts, 82
- subtropical surface water (SSW)
  - decadal-scale population variability, 64
  - synchronies and teleconnections, 82–3
- Humboldt Current ecosystem, climate-based decadal-scale population dynamics, 75–6
- Sufflogobius bibarbatu*s, Northern Benguela ecosystem, 77, 78
- super-individuals, biophysical models
  - climate change, 103
  - fish egg/larval mortality, 90–1
  - mackerel, 95
  - predator–prey models, 98
  - upwelling systems, 93
- surface flow fields, biophysical models, upwelling systems, 93–5
- surface layer stratification, oceanographic
  - climate change, future scenarios, 315–16
- swimming ability
  - biophysical models, upwelling systems, 93
  - low-frequency population fluctuations, anchovy nearshore habitat restriction, 289
- synchronous interdecadal mechanisms, low-frequency population fluctuations, 287–8
  - cyclic dominance model, 295
- teleconnections, decadal-scale variability
  - research
    - pelagic low-frequency fluctuations, 65
    - synchronies and, 82–3
- Temora longicornis*, herring and sprat diet, Baltic Sea system, 135–7
- temperature
  - acidification effects and, 38
  - biophysical models, fish egg/larval growth, 90
  - Canary Current ecosystem, climate change-based decadal-scale population dynamics, 80–1
  - disease rates and increases in, 328
  - low-frequency population fluctuations, 286–9
    - egg retention and developmental time thresholds, 289
    - temperature optima, 292t
  - niche replacement and, 305
  - ocean temperature, habitat change and, 36
  - pelagic fish population shifts and, 34–5
  - trophic dynamics and, 139–41
    - Baltic Sea herring/sprat recruitment, 145
- Tephra layers, chronostratigraphy interference, 49–51
- thermocline effects
  - California Current ecosystem, climate-based decadal-scale population dynamics, 70–1
  - decadal-scale population variability, climate-based regime shifts, 82
  - Humboldt Current ecosystem, climate-based decadal-scale population dynamics, 75–6f

- thermohaline circulation, climate change,  
 ocean surface layer stratification, 316
- three-dimensional modeling, NEMURO-3D  
 simulation, 101–2
- Thyrsites atun*, sardine occurrence  
 in, 77–8
- time scales of variability, scale deposition  
 rates, 53, 59–60t
- time series data  
 density dependence and, 304–5  
 fishing collapse analysis, 165–6  
 climate change effects, 181–3  
 stock assessment and management  
 Humboldt sardine, 204–5  
 larval species, 217
- Todarodes pacificus*, Kuroshio Current  
 ecosystem, biota changes, 71
- top-down effects of regime shifts, 159  
 fishing collapse, effects on small pelagics,  
 176–8
- future research application, 308  
 wasp-waist ecosystem research, 301
- Total Allowable Catch (TAC)  
 modeling and harvest strategies  
 Baltic Sea herring, 233–4  
 Baltic Sea sprat, 232–3  
 Benguela operational management,  
 219–20  
 environmental indices, 221–4  
 Humboldt stock, 219  
 Japanese stocks, 217–18  
 North Sea sprat, 232  
 Southwestern Australia sardine, 235  
 stock assessment and management  
 Baltic Sea herring and sprat catches,  
 212–13  
 limitations of, 235–6
- Trachurus capensis*, Northern Benguela  
 ecosystem, 77, 78
- Trachurus murphyi*, 22
- Trachurus trachurus*, filter feeding patterns,  
 128–9
- trade statistics, global production small  
 pelagic fish production and trade,  
 257–61f  
 commodity production, 259f  
 discounted present values, 265f  
 export values, 260f, 261f
- trade wind intensification. *See also*  
 winds  
 Canary Current ecosystem, climate  
 change-based decadal-scale  
 population dynamics, 80–1
- triggerfish, Canary Current ecosystem  
 populations, 79
- Tri-National Sardine Forum, 240
- trophic dynamics (trophic levels)  
 Baltic Sea system, 135–7, 136–42f, 140–4f,  
 146t  
 physical forcing, 142f
- Benguela Current  
 regime shifts, 78  
 summary of findings, 140–4t  
 upwelling system, 113–18f, 115–16f
- California Current  
 climate-based decadal-scale population  
 variability, 70–1  
 summary of findings, 140–4t  
 upwelling system, 118–23, 119–20ft
- cannibalism incidence and, 144t
- climate change and, 139–41  
 Baltic herring/sprat recruitment and, 145  
 food web considerations, 318–19f  
 menhaden distribution and, 145–6  
 phenology and match–mismatch  
 hypothesis, 318  
 plankton and fish communities, 319–20  
 research gaps and limitations, 329–30  
 summary of research, 147
- decadal-scale population dynamics, 81  
 defined, 184n.4
- fishing collapse analysis  
 catch and abundance data series, 164,  
 166–9f  
 collapse indicators, small pelagics as, 181  
 pathway disruption, 182  
 “static” (snapshot) models, 164–5  
 time series data models, 165–6  
 trophic flow models, 174–6
- Humboldt Current upwelling system,  
 123–6f, 125–30t, 140–4t
- Kuroshio Current, Northwestern Pacific,  
 126–8, 127–8f, 128–9f, 140–4t
- low-frequency population fluctuations, 294
- non-SPACC species results, 146–7
- Northeastern Atlantic and Mediterranean  
 systems, 128–33, 129–34f, 130–2t,  
 132–40t, 140–4t  
 comparison of indices, 143–6f
- Northwestern Atlantic system, 133–5f,  
 140–4t  
 research overview, 112–13t  
 summary of case studies, 137–46, 140–4t  
 wasp-waist pelagic fish models, 160f, 301
- Tropical North Atlantic (TNA) Index,  
 Canary Current ecosystem, climate  
 change-based decadal-scale population  
 dynamics, 81
- Tsushima (Warm) Current, 31  
 Japanese anchovy, 31–3  
 Japanese sardine, 33–4
- tuna population dynamics  
 forage model, 100  
 future research issues, 306
- SEPODYM model, 100
- Southern Oscillation Index and, 306
- turbulence, winds and changes in, 37
- turning points in regime shifts, 66, 67f
- Typical scales category, scale deposition rates  
 and scale shedding, 51–3, 54t
- US Fishery Conservation and Management  
 Act, California fisheries and, 268–9
- ultraviolet (UV) radiation, climate change  
 and, 328
- uncertainty  
 climate change research, 330  
 stock assessment and management and role  
 of, 236
- upwelling  
 Benguela Current  
 biophysical models, anchovies and  
 sardines, 91–3  
 fishing pressure in, 161–2  
 regime shifts, 78  
 biophysical models, 93–5  
 climate change and, 104–5
- California Current ecosystem,  
 climate-based decadal-scale  
 population dynamics, 70–1
- Canary Current ecosystem, climate  
 change-based decadal-scale  
 population dynamics, 80–1
- climate change and  
 circulation models, 326  
 fishing collapse and effects of, 183  
 wind intensification, 313–15
- coastal wind driven, 12, 13, 14, 18, 27, 34, 37
- decadal-scale population dynamics,  
 trophodynamic aspects, 81
- future research issues, 306
- Humboldt Current, fishing pressure, 162–3
- land–ocean exchanges, research gaps  
 concerning, 330–1
- low-frequency population fluctuations,  
 sardine-anchovy alternations and, 289
- trophic dynamics  
 Benguela Current, 113–18f, 115–16f  
 California Current, 118–23, 119–20ft  
 climate change and, 139–41  
 Humboldt Current, 123–6f, 125–30t,  
 140–4t  
 research overview, 112–13t  
 winds and changes in, 37  
 wind-stress, curl driven, 14, 15, 18, 34, 315,  
 332–3
- Uria aalge*, prey quality effects on, 179
- value-added production, global harvesting,  
 production, and trade practices, small  
 pelagic fisheries, 259f

- velocity modeling and measurement,
  - biophysical models, upwelling systems, 93
- vertical migration, biophysical models, fish egg/larval behavior, 91
- “Victoria” sea surface temperature pattern, California Current ecosystem regime shifts, 68
- Virtual Population Analysis (VPA)
  - Benguela Current population dynamics, 25
  - classical theories on pelagic collapse and, 6–7
  - limitations of, 236
  - modeling and harvest strategies
    - Baltic Sea sprat, 232–3
    - California stock, 218–19
    - Humboldt stock, 219
    - ICES stocks, 220–1
    - Japanese stocks, 217–18
    - Mediterranean and Black Sea pelagics, 234–5
    - North Sea autumn spawning herring stock, 224–32
  - Northern Benguela ecosystem biota, 77
  - stock assessment and management, 191
    - Benguela anchovy, 199
- von Bertalanffy equation
  - OSMOSE model, 100
  - tuna forage model, 100
- wasp-waist pelagic fish models
  - ecosystem analysis using, 158–60, 301
  - feedback instability, 301–2
  - fishing collapse analysis
    - climate change effects, 182–3
    - trophic flow control, 174–6
  - future research applications, 300–1
  - low-frequency population fluctuations, 294
  - niche replacement and, 305–6
  - time series data, 166
  - trophic dynamics and, 160f
  - tuna population dynamics, 306
- water quality model, bay anchovy population, 99–100
- wetfish species, climate-based fishery
  - fluctuations, ecosystem models, 266–70, 272n.7
- “what-if” climate scenario, fishing collapse analysis, modeling and testing with, 183
- white sharks, climate-based fishery
  - fluctuations, ecosystem models, 266–70
- winds. *See also* trade wind intensification
  - Aleutian Low pressure system, trophic dynamics and, 139–41
  - climate change and fishing collapse, 183
  - habitat changes and, 37
  - low-frequency population fluctuations, latitudinal source water shifts, 290–1
  - oceanographic climate change, future scenarios, 313–15
- Working Group 67, formation of 1
- World Climate Research Program (WCRP), decadal-scale variability research, 65
- zoogeography, distributional dynamics, future research in, 302–3
- zooplankton
  - biophysical models, upwelling systems, 93
  - California Current ecosystem, 68, 69f
  - climate-based decadal-scale population dynamics, 70–1
  - regime shifts, 70
- Canary Current ecosystem, climate change-based decadal-scale population dynamics, 80–1
- climate change effects
  - acidification and UV increases, 328
  - ocean surface layer stratification, 315–16
- decadal-scale population dynamics, trophodynamic aspects, 81
- density dependence and, 304–5
- fishing collapse analysis
  - gelatinous zooplankton simulations, 169–72f
  - species proliferation, 178–9
  - trophic flow control models, 174–6
- Humboldt Current ecosystem, 75, 162
- Kuroshio Current ecosystem
  - biota changes, 71
  - regime shifts, 71–2
- NEMURO.FISH model, 101–2
- Northern Benguela ecosystem biota, 77
- trophic dynamics
  - California Current, pelagic dietary patterns, 119–23
  - climate change effects on, 146
  - community structure and climate change, 319–20
  - dominance as dietary component, 137–40
  - food web models and climate change, 318–19
  - Humboldt Current dietary studies, 118–23
  - Northeast Atlantic/Mediterranean systems, 129–31
  - Northwestern Atlantic system, 133–5
  - research gaps and limitations, 329–30
  - research overview, 112–13
  - summary of research, 147