

INDEX

- Abnormal genes, 17, 20–21, 20f–21f
- Acetylcholine
 - anticholinergic agents and, 48f
 - dopamine receptor blockage and, 48f
 - relationship with dopamine, 47f
 - release, 46
- Affective flattening, 5
- Aggression
 - mania and, 99, 100f
 - symptoms in schizophrenia, 8f, 9
 - treatment for, 86, 86f
- Agitation, atypical antipsychotics in, 119f
- Agranulocytosis, 70
- AIDS therapy, 131
- Akathisia, 13
- Alcohol abuse, mania and, 99, 100f
- Allosteric modulators, 24, 27f
- Alogia, 5
- Alpha 1 adrenergic blocking, antipsychotic agents and, 46f, 49, 50f
- Alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid (AMPA), 23, 26f
- Alpha-7-nicotinic cholinergic agonists, 130
- Alzheimer's disease
 - psychosis and, 2, 3t
 - Aggressive symptoms and, 8f, 9
 - cholinergic deficiency and, 130, 131
 - cognitive symptoms and, 6, 7f
 - excitotoxicity and, 23, 28, 29f–33f
 - positive symptoms and, 4–5, 6f
- Amenorrhea, 16
- Amisulpride, 96, 97f
- Amyotrophic lateral sclerosis, 23
- Anhedonia, 5
- Anticholinergic agents, 46
 - acetylcholine and, 48f
- Anticonvulsants
 - bipolar combinations and, 119, 120, 122, 123f
 - in ion storms, 100f
 - as mood stabilizers, 102t, 108, 109, 109t, 110f–114f, 111–113, 115, 115f, 116f
- Antidepressants
 - combinations, 122, 123f
 - in bipolar disorder, 120, 122
- Antihistaminic properties, of conventional antipsychotic agents, 46f, 49, 49f
- Antipsychotic agents
 - atypical drugs (*See* Atypical antipsychotic drugs)
 - conventional drugs (*See* Conventional antipsychotic drugs)
 - future antipsychotics, 127–132
 - future combination chemotherapies, 131–132
 - novel neurotransmitter mechanisms, 130–131
 - novel serotonergic and dopaminergic mechanisms, 127–129
 - other antipsychotics, 127, 128f–129f
 - in treatment-resistant schizophrenia, 124–127, 124f–126f
- Anxious self-punishment, 4
- Anxious symptoms, in schizophrenia, 8f, 9–10
- Apathy, 3–4
- Apostasis, blocking, 29–30, 33f, 34f
- Aripiprazole, 96, 96f
- Attentional impairment, 5, 9f
- Attention deficit hyperactivity disorder, 8f, 9
- Atypical antipsychotic drugs, 51–66
 - binding to postsynaptic dopamine 2 receptors, 60f
 - bipolar combinations and, 120, 123f

- Atypical antipsychotic drugs (*continued*)
 in bipolar disorder, 97, 118–119, 119f–122f
 in clinical practice, 80–86
 clozapine, 68–70, 69f
 for cognitive symptoms, 84–85, 84f
 cytochrome P450 1A2, 76, 76f–77f
 cytochrome P450 3A4, 77–80, 79f–80f
 cytochrome P450 2D6, 76, 77f–78f
 four key dopamine pathways, 52–53
 for hostility, aggression, and poor impulse control, 86, 86f
 hypothetical action over time, 90f
 in ion flux reduction, 122f
 in manic symptoms of psychotic mania, 118–119, 119f–122f
 mesocortical pathway, 59–61, 61f–63f
 mesolimbic pathway, 62, 64
 for mood disorders, 82–84, 83f
 in mood stabilization, 118–119, 119f–122f
 for negative symptoms of schizophrenia, 85–86, 85f
 nigrostriatal pathway (*See* Nigrostriatal pathway)
 in nonpsychotic mania, 118–119, 121f, 122f
 olanzapine, 69f, 72, 73f, 120, 121f–122f
 pharmacokinetic considerations for, 75–80, 76f–80f
 for positive symptoms of schizophrenia, 81–82, 82f
 quetiapine, 69f, 72–73, 74f
 receptor binding properties of, 87–90, 88f, 90f
 risperidone, , 70–72, 71f
 serotonergic control of dopamine release, 52–66, 53f–58f, 61f, 63f–65f
 serotonin-dopamine antagonism, 52–53, 52f
 tuberoinfundibular pathway, 61–62
 ziprasidone, 73–75, 74f–75f
- Avalon, 5
- Basal ganglia, dopamine deficiency in, 13
- Benzamide antipsychotics, 96
- Benzodiazepines, 117, 123f
 bipolar combinations and, 119, 122, 123f
- Biological basis, of schizophrenia, 10–16
- Bipolar combination treatments, 119–122, 123f
- Bipolar disorder
 aggressive symptoms and, 8f, 9
 brainstorm of ions and, 96–99, 98f–106f
 depressive symptoms and, 8f, 9
 lithium in, 102, 102t, 107, 107f–108f, 109f
 positive symptoms and, 4, 6f
- Blame, 4
- BMY-14,802, 130
- Brain development, early, 16–21, 17f–21f
- Burnout, 22, 22f
- Cannabinoid antagonists, 130
- Carbamazepine, 109t, 112, 115f, 123f
- Carbonic anhydrase inhibitor, 116f
- Caspase enzymes, 30
- Caspase inhibitors, 29
- Cellular events, during excitotoxicity, 28, 30f–32f
- Chemotherapies, combination, 131–132
- Childhood psychosis, 5, 6f, 8f, 9–10
- Chlorpromazine, 39, 41t
- Cholecystokinin, 130
- Chorea, 13
- Clozapine, 68–70, 69f
- Cognitive symptoms
 in schizophrenia, 6, 7f
 treatment for, 84–85, 84f
- Conceptual disorganization, 3
- Conduct disorders, in children, 9
- Conventional antipsychotic drugs, 39–50, 41t, 51f
 alpha 1 adrenergic blocking properties, 46f, 49, 50f
 antihistaminic properties, 46f, 49, 49f
 blocking all four dopamine pathways, 44–45
 dopamine 2 receptor blockade, 39, 40f, 41–45, 42f–45f, 59, 59f
 haloperidol, 41t, 49, 50f–51f
 hypothetical action over time, 89f
 icon of, 46f
 “lead in” atypical antipsychotics, 124, 126f
 in mood stabilization, 117–118, 117f–118f, 120, 123f
- muscarinic cholinergic blocking properties of, 45–49, 46f–48f
- pharmacologic properties of, 45–46, 46f–50f, 49
- receptor binding properties of, 87f, 88–89, 89f
- risks and benefits of, 45
- side effects of, 45–46, 47f, 49, 49f, 50f
- “top up” atypical antipsychotics, 125, 126f
- Cross-titration, 124, 124f, 125f
- Cytochrome P450 1A2, 76, 76f–77f
- Cytochrome P450 3A4, 77–80, 79f–80f
- Cytochrome P450 2D6, 76, 77f–78f
- Dendrites, damage to, 28, 29f
- Depression. *See also* Antidepressants
 depressive psychosis, 3–4
 mania and, 97–99, 98f–99f
 symptoms in schizophrenia, 8f, 9–10
- Disorganized-excited psychosis, 3
- Disorientation, 3
- Divalproex, 108, 109t, 111–112, 111f–114f, 119–120, 123f
- Dopamine. *See also* Dopamine 2 receptors
 acetylcholine release and, 45, 46, 47f–48f
 hypothesis of positive psychosis symptoms, 10
 novel mechanisms, 127–128
 secondary deficiency, 61f
 stabilizers, 91–96, 91f–96f
 stimulant, 91f
- Dopamine 2 receptors
 binding in nigrostriatal pathway
 atypical antipsychotics, 60f
 conventional antipsychotics, 59f

- binding properties, 87–90, 87f–90f
- blockade of, 39–44, 40f–44f, 88–90, 89f–90f, 118f
 - dilemma of, 44–45
 - hyperprolactinemia and, 44f
 - long-term blockade, 43f
 - mesocortical pathway, 42f
 - negative symptoms and, 42f
 - in nigrostriatal pathway, 42f
 - pure D2 blocker, 40f
 - dopamine system stabilizers, 91–96, 91f–96f
 - hit and run actions, 87–90, 87f–90f
 - improved positive symptoms and, 62–64
 - extrapyramidal symptoms and, 41, 42f–43f, 43, 51, 55–59, 57f–58f
 - hyperprolactinemia and, 43–44, 44f, 61–62, 63f–65f
- Dorsolateral prefrontal cortex, 11, 13f, 14f
- D3 receptors, 128
- Drug abuse
 - mania and, 99, 100f
- Drug-induced psychoses, 5
- DU-127090, 96
- Dyskinesias, 13
- Dysphoria, 99, 99f
- Dystonia, 13

- Excitatory neurotransmission, 23, 28, 28f, 29f–32f
- Excitement, 3
- Excitotoxicity, 22–23, 29f–32f
 - cellular events during, 30f–32f
 - glutamate system and, 28, 28f–32f
- Executive functioning, 6
- Extrapyramidal symptoms, 41, 42f–43f, 43, 46, 48f, 118

- Fananserin, 128
- Free radicals
 - in excitotoxicity, 31f, 32f, 34f
 - scavengers, 29, 34f

- Gabapentin, 113, 116f, 123f
- Galactorrhea, 16
- Gamma-aminobutyric acid (GABA), 109, 110f, 111, 111f, 113, 115f, 116f
- Genetic basis in schizophrenia, 17–21, 20f–21f
- Glutamate
 - antagonists, 29, 33f
 - production, 24f
 - receptors, 23–25, 26f–27f
 - removal, 23, 25f
 - spectrum of excitation, 29f
 - systems, excitotoxic overactivity of, 11
- Glutamatergic neurotransmission, 23–28
 - excitotoxicity and glutamate system, 22, 28, 28f–32f

- Kainate, 23, 26f

- L-745,870, 128
- Lamotrigine, 109t, 112–113, 115f, 123f
- Lazaroids, 30, 34f
- Ligand-gated ion channels, 27f, 28f, 108, 110f
- Lithium, 102, 102t, 107, 107f–109f
 - bipolar combinations and, 117f, 119, 119f, 123f
- Long-term potentiation, 23
- Lou Gehrig's disease, 23
- Low extrapyramidal symptoms, pharmacology of, 55–59, 57f–60f
- Loxapine, 127, 128f
- LU-111,995, 128

- Magnesium, 24, 27f
- Mania, 97–101, 102f
atypical antipsychotics in, 118–119, 119f–122f
characterization of, 97, 98f
glutamate excitation and, 29f
anticonvulsants in, 102t, 108–111, 109t
mixed state, 99, 99f
mood stabilizing drugs and (*See* Mood stabilizing drugs)
other clinical features, 100f
psychosis associated with, 99, 100f, 102f, 117f, 117–119, 119f, 120f
rapid cycling, 98f
- Mazapertine, 128
- MDL-100,907, 128
- Mesocortical dopamine pathway, 10–13, 11f, 13f, 14f, 42f
improved negative symptoms and, 59–61, 61f–63f
- Mesolimbic dopamine pathway, 10, 11f, 12f, 40f
improved positive symptoms and, 62, 64
- Metabotropic glutamate receptor, 26f
- Mixed state, 99, 99f
- Mood disorders in schizophrenia, treatment for, 82–84, 83f
- Mood-stabilizing drugs, 102–122, 102t, 109t, 123f
anticonvulsants, 102t, 108–113, 109t, 110f–116f, 115
atypical antipsychotics, 118–119, 1119f–122f
benzodiazepines, 117, 119, 122, 123f
bipolar combinations, 119–122, 123f
carbamazepine, 109t, 112, 115f, 123f
conventional antipsychotics, 117–118, 117f–118f, 120, 123f
gabapentin, 113, 116f, 123f
lamotrigine, 109t, 112–113, 115f, 123f
lithium, 102, 102t, 107, 107f–109f
topiramate, 113, 115, 116f
valproic acid, 109t, 111–112, 111f–114f, 123f
- Muscarinic cholinergic receptors, blockade of, 45–49, 46f–48f
- Negative symptoms, 5f, 5–6, 7f, 9t, 41, 42f, 59–61, 61f–63f
- Nemonapride, 128
- Neurodegeneration
blocking, 29–30, 33f, 34f
causes of schizophrenia, 35f
hypothesis, 21–29
excitotoxicity, 22–23, 29f–33f
glutamatergic neurotransmission and, 23–28, 24f–32f
- Neurodevelopmental hypothesis, of schizophrenia, 16–21, 17f–21f
- Neurodevelopmental/neurodegenerative therapy hypothesis, 33, 35f
- Neurokinins, 130
- Neurolepsis, 39
- Neuroleptic antipsychotics, bipolar combinations and, 120, 123f
- Neuroleptic-induced deficit syndrome, 41
- Neuroleptic-induced tardive dyskinesia, 14
- Neuroleptic malignant syndrome, 45
- Neurons
migration of, 16, 17f, 18f
neuroprotection, 29f
- Neurotensin antagonists, 130
- Neurotransmission, glutamate, 23–28, 24f–32f
- NGD-94-4, 128
- Nigrostriatal dopamine pathway, 13–14, 15f
dopamine 2 receptor blockage in, 41, 42f, 43, 43f
pharmacology of low extrapyramidal symptoms, 55–59, 53f–60f
serotonin-dopamine interactions in, 53–54, 53f–56f
- N-methyl-D-aspartate (NMDA), 23–28, 26f, 27f, 28f, 29, 33f
- Olanzapine, 69f, 72, 73f, 120, 121f–122f
- OPC14523, 130
- OPC-14597, 96
- Organic dementias, 5, 9
- OSU-6162, 96
- Panic, 29f
- Paranoid projection, 3
- Paranoid psychosis, 3
- Parkinson's disease, 13, 23, 41
- Partial agonist action, 95
- PCP site, 25, 27f
- Perospirone, 127, 129f
- Phosphatidyl inositol, 105f, 107
- PNU-101,387G, 128
- Polyamines, 24, 27f
- Positive symptoms
pharmacology of, 62–64
in psychosis, 9t
in schizophrenia, 4–5, 5f, 6f, 9t, 10, 11f–12f, 40f
- Postsynaptic serotonin 2A receptors, 52–62, 52f–59f, 61f–65f, 64
- Precamol, 96
- Presymptomatic treatment, in schizophrenia, 30, 33
- Prolactin
conventional antipsychotic drugs and, 43–44, 44f, 64f
dopamine inhibition of, 63f
elevated level, 15f, 16
serotonin 2A antagonism and, 61–62, 65f
serotonin stimulation of, 64f
- Psychosis
associated disorders, 2, 3t
clinical description of, 2–4
defining symptom in, 2, 2t

- depressive psychosis, 3–4, 8f, 9–10
- related disorders, 2t
- disorganized-excited, 3
- mania and, 99, 100f, 102f, 117f, 117–119, 119f, 120f
- negative symptoms in, 5f, 5–6, 7f, 9t
- paranoid psychosis, 3
- positive symptoms in, 4–5, 5f–6f, 9t
- Psychotomimic agent phenylcyclidine (PCP)**, 25, 27f, 130

- Quetiapine**, 69f, 72–73, 74f

- Rapid cycling**, 98f, 99, 107, 111, 114f
- Rapid dissociation**, 87–90, 88f, 90f
- Receptor binding properties**
 - atypical antipsychotics, 87–90, 88f
 - conventional antipsychotics, 87f, 87–90
- Retardation**, 3–4
- Risperidone**, 70–72, 71f
- Ritanserin**, 128

- Schizoaffective disorders**, 4, 6f, 8f, 9
- Schizophrenia**
 - aggressive and hostile symptoms, 8f, 9
 - cognitive symptoms, 6, 7f
 - defined, 2–4
 - depressive and anxious symptoms, 8f, 9–10
 - dopamine pathways in, 10–16, 11f–15f
 - mesocortical dopamine pathway, 10–11, 11f, 13, 13f, 14f
 - mesolimbic dopamine pathway, 10, 11f, 12f
 - nigrostriatal dopamine pathway, 10, 13–14, 15f
 - tuberoinfundibular dopamine pathway, 14–16, 15f
 - excitotoxicity and, 22–23, 29f–33f
 - experimental therapeutic approaches, 29–33, 33f, 34f
 - five symptom dimensions, 4–10, 5f–8f, 9t
 - future combination chemotherapies for, 131–132
 - in general population, 4
 - glutamatergic neurotransmission and, 23–28, 24f–34f
 - negative symptoms, 5f, 5–6, 7f, 9t
 - neurodegenerative causes of, 21–30, 24f–35f
 - neurodegenerative hypotheses of, 21–29
 - neurodevelopmental hypothesis of, 16–21, 17f–21f
 - novel neurotransmitter mechanisms and, 130–131
 - novel serotonergic and dopaminergic mechanisms and, 127–128
 - positive symptoms, 4–5, 5f–6f, 9t
 - stages of, 21–22, 22f
 - treatment in clinical practice, 80–86 (*See also* Antipsychotic agents)
 - for cognitive symptoms, 84–85, 84f
 - for hostility, aggression, and poor impulse control, 86, 86f
 - for mood disorders, 82–84, 83f
 - for negative symptoms, 85–86, 85f
 - for positive symptoms, 81–82, 82f
 - treatment resistant, 124–127, 124f–126f
- Second-messenger system**, 102, 103f–108f, 107
- Serial learning**, 6
- Serotonergic mechanisms, novel**, 127–128
- Serotonin 2A receptors**
 - binding in cerebral cortex
 - atypical antipsychotics, 63f
 - conventional antipsychotics, 62f
 - improved positive symptoms and, 62–64
 - low EPS and, 51, 55–56, 57f–59f, 59
 - reduced hyperprolactinemia and, 61–62, 63f–65f
- Serotonin-dopamine antagonism**, 51–52, 52f, 55–56, 58f, 59–64, 65f
- serotonergic control of dopamine release**, 51–53
- Serotonin-dopamine interactions, in nigrostriatal pathway**, 53–54, 53f–58f
- Sexual dysfunction**, 16, 44
- Sigma antagonists**, 130
- SR 141716A**, 130
- SR-142948**, 130
- SR31742A**, 130
- Stevens-Johnson syndrome**, 113
- Stroke**, 23, 28, 29f
- Substance P**, 130
- Synapses**
 - formation, 16
 - inadequate selection, 17f
 - normal formation of, 18f
 - wrong connections, 19f

- Tardive dyskinesia**
 - conventional antipsychotics and, 46, 117, 118f
 - D2 receptor blockage and, 43, 43f
 - neuroleptic-induced, 14
- Therapeutic approaches in schizophrenia**, 29–33. *See also* Schizophrenia
 - blocking neurodegeneration and apostasis, 29–30, 33f, 34f
 - combined neurodevelopmental/neurodegenerative hypothesis, 33, 35f
 - presymptomatic treatment, 30, 33
- Therapeutics, genetic approach to**, 21, 21f
- 3-PPP**, 96
- Tics**, 13
- Topiramate**, 113, 115, 116f, 123f
- Toxic insult**, 17f
- Tuberoinfundibular dopamine pathway**, 14–16, 15f
- Tuberoinfundibular pathway, reduced**
 - hyperprolactinemia and, 43–44, 44f, 61–62, 63f–65f

142 Index

- Twin studies. *See* Genetics WAY135, 452, 96
Valproic acid, 108, 109t, 111–112, 111f–114f Weight gain, 44, 49, 49f
 bipolar combinations and, 119–120, 123f YM-43611, 128
Verbal fluency, impaired, 6 Ziprasidone, 73–75, 74f–75f
Vitamin E, 29 Zotepine, 127, 129f
Voltage-gated ion channels, 100f–101f, 102f–108g,
 112f–113f