

Index

-
- ABCDE approach to patient
 - assessment, 1
 - abdomen
 - CT scanning, 54–5
 - abdominal X-ray, 53
 - acid–base balance
 - bicarbonate buffer system, 150
 - Boston approach to diagnosing disorders, 151–2
 - buffering responses, 150–1
 - compensatory responses, 151–2
 - disorders in critical care patients, 148
 - factors that affect $[H^+]$ homeostasis, 149–50
 - haemoglobin as a blood buffer, 151
 - hydrogen ion concentration and pH, 148–9
 - normal controls, 148
 - phosphate buffer system, 150–1
 - physico-chemical (Stewart's) theory, 149–50
 - responses to acid–base changes, 150–2
 - role of carbonate and phosphate salts in bone, 151
 - standard base excess, 150
 - strong ion gap, 149–50
 - traditional theory, 149–50
 - acid–base disturbances
 - definition of acidaemia and acidosis, 152
 - definition of alkalaemia and alkalosis, 152
 - metabolic acidosis, 153–6
 - metabolic alkalosis, 156–7
 - physiological effects, 152–3
 - respiratory acidosis, 157
 - respiratory alkalosis, 157
 - underlying cause, 152
 - acute asthma
 - mechanical ventilation strategy, 224–5
 - acute coronary syndromes (ACS)
 - adjunctive therapy, 191–2
 - biomarkers, 189
 - classification, 185
 - complications, 192–3
 - creatinine kinase (CK-MB) biomarker, 189
 - definition of myocardial infarction (MI), 185
 - diagnosis, 185–90
 - diagnostic criteria, 185
 - ECG, 186–9
 - imaging, 190
 - laboratory tests, 189, 190
 - management of non-ST-elevation ACS, 191
 - management of ST-elevation ACS, 190–1
 - pathophysiology of MI, 185
 - patient history and examination, 185
 - risk stratification, 190
 - troponin biomarkers, 189
 - acute epiglottitis in children, 365
 - acute gastrointestinal bleeding
 - common presentations, 257
 - complications associated with, 257
 - haematemesis, 257
 - haematochezia, 257
 - lower GI bleeding, 262–3
 - management of a large GI bleed, 257–8
 - melaena, 257
 - non-variceal upper GI bleeding, 259–60
 - prevention of upper GI bleeding on ICU, 262
 - upper GI bleeding, 259–62
 - variceal bleeding, 260–2
 - acute heart failure
 - aetiology, 202
 - cardiogenic shock, 202
 - clinical evaluation, 204
 - clinical syndromes and classifications, 202
 - coronary angiography, 206
 - CT angiogram, 206
 - definition, 202
 - ECG, 204
 - endomyocardial biopsy, 206
 - imaging techniques, 204–5
 - invasive monitoring, 206
 - laboratory investigations, 205–6
 - pathophysiology, 202–3
 - presentation, 202–3
 - prognosis, 211
 - pulmonary oedema, 202
 - acute heart failure treatment
 - analgesia, 207
 - diuretics, 209
 - future agents, 211
 - haemodynamic support, 207–8
 - heart transplantation, 210
 - immediate resuscitation, 207
 - intra-aortic balloon pump (IABP), 208–9
 - mechanical assist devices, 208–9
 - natriuretic peptides, 209
 - prognosis, 211
 - respiratory support, 207
 - treatment of underlying disease and co-morbidities, 209–10
 - ultrafiltration, 209
 - vasodilators, 209
 - ventricular assist devices (VADs), 209
 - acute liver failure
 - acute-on-chronic liver disease, 290
 - adrenal dysfunction, 289
 - auxiliary or partial liver transplantation, 289
 - causes, 284–6
 - complications, 286–9
 - general supportive care, 286
 - hepatic encephalopathy (HE), 286–7
 - hepatitis infections, 286
 - intracranial hypertension (ICH), 287–8
 - liver transplantation, 289–90
 - living related lobe donation, 289
 - management, 284
 - paracetamol poisoning, 284–6
 - presentation, 284
 - prognostication, 289–90
 - renal failure complication, 288
 - acute lung injury (ALI), 236–8
 - mechanical ventilation strategy, 222–4
 - acute pancreatitis. *see* severe acute pancreatitis
 - acute renal failure
 - aetiological classification, 292–5
 - definitions, 292
 - diagnosis and assessment, 295–6

Index

- intrinsic causes, 294–5
 management, 296–8
 outcome, 298
 physiological considerations, 292
 postrenal causes, 295
 prerenal causes, 292–4
 susceptibility of kidneys to failure, 292
- acute respiratory distress syndrome (ARDS), 236–8
 mechanical ventilation strategy, 222–4
 prone positioning during ventilation, 251
 use of high-frequency oscillatory ventilation (HFOV), 252–3
 use of recruitment manoeuvres, 252
- adaptive support ventilation (ASV), 243–4
- adrenaline. *see* epinephrine
- advance decision (end of life care), 88–9
- advance directives, 19
- advanced airway management
 cricothyroidotomy, 14–15
 extubation/weaning protocols, 9–12
 intubation, 6–9
 mini tracheostomy, 15
 requirements of critically ill patients, 6
 tracheostomy, 12–15
 weaning. *see* extubation/weaning protocols; respiratory weaning
- Advanced Life Support™ courses, 1
- Advanced Trauma Life Support™ courses, 1
- airway assessment
 predicting a difficult intubation, 6
- airway support. *see* advanced airway management
- AIS (Abbreviated Injury Scoring), 27
- albumin, 131
- ALERT™ (Acute Life threatening Events – Recognition and Treatment) courses, 3
- alfentanil, 74
- Allen's test, 41
- alpha-1 (α_1) receptors, 59
- amikacin, 127
- aminoglycosides, 127
- amniotic fluid embolism, 355–6
- amphotericin B, 129
- analgesia. *see* pain control; sedation
- antibiotic therapy
 post-operative critical care, 162, 167
- antibiotics
 bacteriocidal mode of action, 125
 bacteriostatic mode of action, 125
 methicillin-resistant *Staphylococcus aureus* (MRSA), 125
 multi-resistant organisms, 125–6
 pharmacokinetics, 125
 post antibiotic effect (PAE), 125
 prescribing in renal failure, 125
 principles of rational antibiotic testing, 124
 prophylaxis, 126
 vancomycin-resistant Enterococci (VRE), 125, 127
- antibiotics used in the ICU, 126–9
 amikacin, 127
 aminoglycosides, 127
 antifungal drugs, 128–9
 beta-lactams, 126–7
 carbapenems, 127
 cephalosporins, 126–7
 ciprofloxacin, 127–8
 clarithromycin, 128
 erythromycin, 128
 gentamicin, 127
 glycopeptides, 127
 imipenem, 127
 levofloxacin, 127–8
 linezolid, 128
 macrolides, 128
 meropenem, 127
 metronidazole, 128
 nitroimidazoles, 128
 oxazolidinones, 128
 penicillins, 126–7
 quinolones, 127–8
 rifampicin, 128
 teicoplanin, 127
 vancomycin, 127
- antifungal drugs, 128–9
- APACHE (Acute Physiology and Chronic Health Evaluation), 27, 28
 versions, 32
- APACHE II score, 30
 use in severe acute pancreatitis, 270–1
- arterial blood gas sampling, 1
- arterial pulse contour analysis, 46–7
- arteriovenous CO₂ removal (AVCO₂R), 254
- ASCOT scoring system, 33
- asplenia
 infection risk, 122
- assessment
 ABCDE approach, 1
 initial assessment of the critically ill patient, 1
 physiology monitoring systems, 2–3
- atracurium, 8
- auditory evoked potentials, 78
- Awakening and Breathing Control (ABC) Trial, 83
- bacteraemia
 definition, 99
- balloon tamponade, 261
- bedside charting, 34–5
 electronic chart systems, 35
 ICU chart, 34
 paper chart, 34
 requirements of an electronic chart, 34–5
- Bentham, Jeremy, 85
- benzodiazepines, 81
- benzylpenicillin, 126
- best interests
 and consent, 88
- beta receptors, 59
- beta-1 (β_1) receptors, 59
- beta-2 (β_2) receptors, 59
- beta-lactams, 126–7
- bicarbonate buffer system, 150
- bispectral index (BIS), 78
- bleeding and clotting
 normal coagulation, 177–8
 normal haemostasis, 177–8
- bleeding disorders
 acquired, 180–3
 congenital, 179–80
 disseminated intravascular coagulation (DIC), 181–2
 drug induced, 136
 haemolytic uraemic syndrome (HUS), 182
 haemophilia A, 179–80
 haemophilia B, 179–80
 heparin-induced thrombocytopenia (HIT), 183
 major haemorrhage and massive transfusion, 182–3
 microangiopathic haemolytic anaemia, 182
 thrombocytopenia, 180–1
 thrombotic thrombocytopenia purpura (TTP), 182
 von Willebrand disease, 179–80
- blood gas analysis, 234
- blood pressure monitoring. *see* invasive blood pressure monitoring
- blood sugar control
 post-operative critical care, 167
- Bloomsbury scale, 78
- brain injury
 mechanical ventilation strategy, 225
see also traumatic brain injury

Index

- brainstem death, 92–3
 bronchiolitis
 in children, 365–6
 bronchiolitis obliterans organizing pneumonia (BOOP), 120
 burns, 338–41
- calcium levels, 144–7
 hypercalcaemia, 146–7
 hypocalcaemia, 145–6
 plasma calcium, 144–5
- cancer patients
 neutropenia and infection risk, 117–19
 ‘can’t intubate, can’t ventilate’ situation, 58, 60
- capacity
 and consent, 87–8
- capnography, 234
- carbapenems, 127
- cardiac arrest
 incidence, 170
 organ donation following brainstem death, 173
 outcomes, 170
 post resuscitation care, 170–3
 post resuscitation syndrome, 170
 prognostication, 173
- cardiac arrhythmias
 anti-arrhythmia drugs, 199
 arrhythmogenic mechanisms, 194–5
 atrial fibrillation, 194
 atrial flutter, 194
 automaticity, 194
 bradyarrhythmias, 195–6, 200
 classification, 195–8
 contributory arrhythmogenic factors, 195
 in the intensive care setting, 194
 management in the ICU, 199–201
 post-operative, 167
 re-entry, 194
 sinus node automaticity, 194
 tachyarrhythmias, 196–8
 tachyarrhythmias with adverse signs, 200
 tachyarrhythmias without adverse signs, 200–1
 triggered activity, 194–5
 ventricular tachyarrhythmia, 194
- cardiac output monitoring, 44–8
 arterial pulse contour analysis, 46–7
 clinical parameters, 44
 dye dilution technique, 45
 Fick’s principle, 46
 lithium dilution technique, 45
- methods of measuring cardiac output, 44
 pulse-induced contour cardiac output (PiCCO), 46–7
 thermodilution technique, 44–5
 transoesophageal Doppler, 45–6
 transoesophageal echocardiography, 47
- cardiac risk stratification, 164–5
- cardiogenic pulmonary oedema
 use of non-invasive ventilation, 246–7
- cardiogenic shock, 202
- cardiorespiratory arrest in children, 362–4
- cardiovascular system
 post-operative critical care, 164–7
- caspofungin, 129
- catheters and tubes
 X-ray assessment of positioning, 51–2
- CCRISP™ (Care of the Critically Ill Surgical Patient) courses, 3
- ceftazidime, 126
- cefuroxime, 126
- central venous catheters
 X-ray assessment of positioning, 52
- central venous pressure monitoring, 42–4
 advantages of central venous catheterization, 43
 cannulation sites, 43
 catheter-over-guidewire cannulation technique, 43–4
 changes in central venous pressure, 43
 complications, 44
 insertion of central venous catheter in IJV, 43–4
 Seldinger’s cannulation technique, 43–4
- cephalosporins, 126–7
- cervical spine
 CT scan, 54
- Chain of Survival, 170
- chest disease
 CT scanning, 54
- chest drains
 X-ray assessment of positioning, 52
- chest X-ray, 50–3
- Child–Pugh Score (liver failure), 27
- chlorpromazine, 83
- chronic liver disease
 acute on chronic liver disease, 290
- chronic obstructive pulmonary disease (COPD)
 mechanical ventilation strategy, 224
 use of non-invasive ventilation, 246
- cilastatin, 127
- ciprofloxacin, 127–8
- clarithromycin, 128
- clavulanic acid, 127
- clindamycin, 128
- clinical decision making
 applications for scoring systems, 29
 potential drawbacks of scoring systems, 29–30
 use of scoring systems, 29–30
- clinical record keeping
 electronic medical records, 35
- clinical scoring systems, 29–30
 assessment of sedation level, 78
- clonidine, 79–80, 81–2
- Clostridium difficile*, 124
- coagulation
 normal process and regulation, 177–8
- coagulation disorders. *see* bleeding disorders
- codeine, 74
- colloid fluids, 131
- colloid vs. crystalloid fluid, 132
- coma. *see* unconscious patient
- competency based training in intensive care medicine in Europe (CoBaTrICE) initiative, 22
- consciousness
 definition of normal state, 312
 spectrum of abnormal levels, 312
 see also unconscious patient
- consent, 87–8
 best interests, 88
 capacity, 87–8
 Mental Capacity Act 2005, 87–8
 necessity principle, 87
 organ donation, 95–6
- COPD. *see* chronic obstructive pulmonary disease
- coronary artery stent
 post-operative management of patients with, 165
- creatinine kinase (CK-MB)
 ACS biomarker, 189
- cricoid pressure, 7
- cricothyroidotomy, 14–15
- Critical Care Network, 21
- critical care outreach (CCO), 3–4, 17
- critical care team
 indications for patient referral, 4
 routes of referral, 4
- critical care unit
 cost of care, 16
 use of available resources, 16
- critical illness
 definition, 1

Index

- croup in children, 365
 crystalloid fluids, 130–1
 crystalloid vs. colloid fluid, 132
 CT pulmonary angiography (CTPA), 54
 CT scanning, 53–5
 basic principles, 49–50
 CT abdomen, 54–5
 CT cervical spine, 54
 CT chest, 54
 CT head, 54
 iodinated intravenous contrast, 53
 cytopathic hypoxia, 109
- data storage and archiving, 38
 dead space
 hypoxia caused by, 235
 decision support information
 bedside level, 37
 information systems, 37
 organizational level, 37
 ward level, 37
 deep vein thrombosis (DVT)
 obstetric emergencies, 351–2
 post-operative prophylaxis, 167
 deontology, 85
 dexmedetomidine, 82
 dextrose-containing solutions, 130–1
 diabetes insipidus, 136
 diamorphine, 74
 difficult airway situation, 9
 diffuse alveolar haemorrhage, 119
 diffusion impairment
 hypoxia caused by, 235–6
 disseminated intravascular coagulation (DIC), 181–2
 ‘Do not resuscitate’ order, 82
 dobutamine, 61
 dopamine, 61
 dopaminergic (D1 and D2) receptors, 59–60
 drug-induced bleeding disorders, 136
 drug overdose. *see* poisoning
 DVT. *see* deep vein thrombosis
 dye dilution technique
 cardiac output monitoring, 45
- early warning scoring systems, 31
 features, 31
 MEWS (modified early warning score), 31
 eclampsia. *see* pre-eclampsia and eclampsia
 electrocardiogram (ECG)
 acute coronary syndromes, 186–9
 acute heart failure, 204
 electroencephalogram (EEG)
 assessment of level of sedation, 78
 unconscious patient, 316
 electrolytes. *see* fluid and electrolyte disorders
 electromagnetic flow measurement, 153
 electronic chart systems, 35
 electronic medical records, 35
 electronic prescribing (physician order entry), 35–7
 encephalitis. *see* viral encephalitis
 end-of-life care, 88–9
 advance decision, 88–9
 Independent Mental Capacity Advocate (IMCA), 89
 refusal of life-prolonging care, 88–9
 welfare attorney, 89
 withholding and withdrawing life-prolonging care, 88
 endotracheal intubation, 9
 X-ray assessment of positioning, 51
 engraftment syndrome, 119
 enteral feeding (EF), 67–9, 71
 complications, 71
 types of feeds, 68–9
 see also nutritional support
 enteral tubes
 X-ray assessment of positioning, 52
 Entonox™, 75
 epidural analgesia, 75–6
 epinephrine (adrenaline), 60–1
 erythromycin, 128
 ethical decision making, 86–7
 ethical framework for critical care
 ethical theories, 85–6
 practical approach to decision making, 86–7
 ethical obligations of doctors, 85
 ethical theories, 85–6
 deontology, 85
 principlism, 85–6
 utilitarianism, 85
 ethics
 organ donation, 95–6
 etomidate, 8, 81
 examinations in intensive care medicine
 European Diploma in Intensive Care (EDIC), 370–3
 example multiple choice questions, 374–6
 example viva questions, 376–7
 Intercollegiate Diploma in Intensive Care Medicine (DICM), 369–70
 multiple choice answers, 377–8
 useful websites, 373
 viva answers, 378–9
- extracorporeal carbon dioxide removal (ECCO₂R), 254–5
 extracorporeal lung assist (ECLA), 254
 extracorporeal membrane oxygenation (ECMO), 253–4
 extravascular lung water (EVLW), 47
 extubation/weaning protocols, 9–12
 criteria for spontaneous breathing trials, 11
 extubation criteria, 11–12
 weaning assesment, 11
 weaning techniques, 11
 see also respiratory weaning
- failed airway situation, 9
 fentanyl, 74
 Fick’s principle, 46
 fluconazole, 129
 fluid and electrolyte disorders
 albumin, 131
 calcium levels, 144–7
 colloid fluids, 131
 colloid vs. crystalloid fluid, 132
 crystalloid fluids, 130–1
 dextrose-containing solutions, 130–1
 gelatins, 131
 Hartmann’s solution, 130
 hetastarch solutions, 131
 magnesium levels, 141–3
 measurement of fluid balance, 130
 normal (isotonic) saline, 130
 phosphorus in the body, 143–4
 potassium balance, 137–40
 sodium balance, 132–7
 starch solutions, 131
 types of fluid used in treatment, 130–1
 fluid balance
 post-operative critical care, 161–2, 165
 fungal infection
 antifungal drugs, 128–9
- gastrointestinal bleeding. *see* acute gastrointestinal bleeding;
 gastrointestinal perforation
 gastrointestinal perforation, 263–4
 bowel perforation, 263–4
 causes, 263
 lower GI perforation, 263–4
 upper GI perforation, 263
 gelatins, 131
 gentamicin, 127
 Glasgow Coma Scale (GCS), 27
 Glasgow Outcome Scale, 333
 Glasgow score (Imrie score), 270
 glycopeptides, 127
 gut hypothesis of multiple organ failure, 68

Index

- Haemacue™, 1
 haematological malignancies
 neutropenia and infection risk, 117–19
 haematopoietic stem cell transplant (HSCT) recipients
 post HSCT complications, 119–20
 haemodynamics monitoring
 cardiac output monitoring, 44–8
 central venous pressure monitoring, 42–4
 essential considerations in critically ill patients, 40
 invasive blood pressure monitoring, 40–2
 haemoglobin
 as a blood buffer, 151
 haemolytic uraemic syndrome (HUS), 182
 haemophilia A, 179–80
 haemophilia B, 179–80
 haemorrhage
 antepartum, 356–7
 postpartum, 356–7
 haemostasis
 normal processes, 177–8
 haloperidol, 79–80, 83
 Hartmann's solution, 130
 head
 CT head scan, 54
 head injury. *see* traumatic brain injury
 heart-beating donation (HBD), 91–3
 heart failure. *see* acute heart failure
 heart transplantation, 210
 HELLP syndrome, 344, 347
 Henderson–Hasselbach equation, 149
 heparin-induced thrombocytopenia (HIT), 183
 hetastarch solutions, 131
 high-resolution CT (HRCT), 54
 highly active antiretroviral therapy (HAART), 121–2
 HIV and AIDS patients
 complications of HIV and its therapy, 121–2
 HL7 (Health Level 7) standard for information exchange, 37
 Human Tissue Act 2004, 95
 hydrogen ion concentration. *see* acid–base balance
 hypercalcaemia, 146–7
 hyperkalaemia, 139–40
 hypermagnesaemia, 142–3
 hypernatraemia, 135–7
 hyperphosphataemia, 144
 hypocalcaemia, 145–6
 hypokalaemia, 138–9
 hypomagnesaemia, 141–2
 hyponatraemia, 133–5
 hyponatraemic encephalopathy, 134
 hypotension
 post-operative critical care, 165
 hypoventilation, 235
 hypoxia pathophysiology, 235–6
 dead space, 235
 diffusion impairment, 235–6
 hypoventilation, 235
 shunt, 235
 ventilation/perfusion (V/Q) mismatch, 235
 hypoxic hypoxia, 109
 ICU chart, 34
 idiopathic pneumonia syndrome, 120
 imaging
 acute coronary syndromes, 190
 acute heart failure, 204–5
 choice of technique, 49
 general considerations, 49
 patient and staff safety issues, 49
 picture archiving and communication system (PACS), 49
 risks to the critically ill patient, 49
 imaging modalities
 CT scanning, 53–5
 magnetic resonance imaging (MRI), 55–6
 nuclear medicine investigations, 56
 ultrasound imaging, 56–7
 X-ray imaging, 49–53
 imipenem, 127
 immunocompromised patient
 use of non-invasive ventilation, 247
 immunonutrients, 69–70
 immunosuppressed patients
 antiretroviral therapy in ICU, 121–2
 asplenia, 122
 cancer and haematological malignancies, 117–19
 causes of immunosuppression, 116
 highly active antiretroviral therapy (HAART), 121–2
 infections, 116–17
 patients with HIV and AIDS, 121–2
 post haematopoietic stem cell transplant (HSCT), 119–20
 respiratory failure, 117
 solid organ transplant recipients, 120–1
 Imrie score, 270
 Independent Mental Capacity Advocate (IMCA), 89
 induction agents, 7–8
 induction cardiography, 48
 infection control
 post-operative critical care, 162, 167
 infections
 immunosuppressed patients, 116–17
 see also sepsis
 information management
 challenges in critical care, 34
 implementation of a critical care information system, 38–9
 information system functions
 bedside charting, 34–5
 clinical record keeping, 35
 data access for audit and research, 38
 data access for quality and financial management, 38
 data mining, 38
 data storage and archiving, 38
 data warehousing, 38
 decision support, 37
 electronic medical records, 35
 electronic prescribing (physician order entry), 35–7
 HL7 (Health Level 7) standard for interoperability, 37
 integration with other hospital systems, 37
 multi-site communication, 37–8
 remote access, 37–8
 simulated environments, 38
 training and education, 38
 usability of the system, 35–7
 virtual ICU, 37–8
 inhalational induction, 8–9
 initial assessment and resuscitation, 1–2
 aim of intensive care, 1
 assessment of the critically ill patient, 1
 definition of critical illness, 1
 general considerations, 1
 importance of early intervention, 1, 2
 monitoring progress of resuscitation, 2
 organ failure, 1
 resuscitation, 1–2
 signs suggesting severe illness, 1
 inotropes
 complications and considerations
 intensive care
 aim of, 1
 Intensive Care National Audit and Research Centre (ICNARC) UK, 30
 intra-aortic balloon
 X-ray assessment of positioning, 52
 intra-aortic balloon pump (IABP), 208–9

Index

- intrathoracic blood volume (ITBV), 47
 intubation, 6–9
 airway assessment, 6
 ‘can’t intubate, can’t ventilate’ situation, 58, 60
 difficult intubation situation, 9
 endotracheal, 9
 failed intubation situation, 9
 indications for, 6
 induction agents, 7–8
 inhalational induction, 8–9
 muscle relaxants, 8
 predicting a difficult intubation, 6
 rapid sequence induction (RSI), 6–7
 invasive blood pressure monitoring, 40–2
 advantages of intra-arterial cannulation, 40
 Allen’s test, 41
 calibration of the monitoring system, 42
 cannulation sites, 40
 catheter-over-needle cannulation technique, 41
 catheter-over-guidewire cannulation technique, 41–2
 complications, 42
 components of the monitoring system, 42
 damping of the measuring system, 42
 Seldinger technique for cannulation, 41–2
 techniques for radial artery cannulation, 41–2
 isoflurane, 81
 ISS (Injury Severity Score), 27

 Kant, Immanuel, 258
 ketamine, 75, 80–1

 levofloxacin, 127–8
 levosimendan, 62
 linezolid, 128
 liquid ventilation (LV), 255–6
 lithium dilution technique
 cardiac output monitoring, 45
 liver failure. *see* acute liver failure; chronic liver disease
 lorazepam, 79–80, 81
 Lundberg waves, 329
 lungs
 appearance on chest X-rays, 52–3
 lusitropy, 60

 macrolides, 128
 magnesium levels, 141–3
 hypermagnesaemia, 142–3
 hypomagnesaemia, 141–2

 magnetic resonance imaging (MRI)
 basic principles, 55–6
 Mallampati score, 6
 malnutrition risk in critical care, 66
 mechanical ventilation
 epidemiology of patients, 212
 history and development, 212
 hypercarbic (Type 2) respiratory failure, 213–14
 hypoxaemic (Type 1) respiratory failure, 213
 indications for initiation, 212–13
 purpose, 212
 respiratory failure, 213–14
 strategy for acute asthma, 224–5
 strategy for acute lung injury (ALI), 222–4
 strategy for brain-injured patient, 225
 strategy for COPD, 224
 strategy for each individual patient, 222
 strategy for early ARDS, 222–4
 see also respiratory weaning
 mechanical ventilation adjuncts
 extracorporeal carbon dioxide removal (ECCO₂R), 254–5
 extracorporeal membrane oxygenation (ECMO), 253–4
 high-frequency oscillatory ventilation (HFOV), 252–3
 liquid ventilation (LV), 255–6
 prone positioning, 251
 recruitment manoeuvres, 252
 transtracheal gas insufflation (TGI), 256
 use of inhaled nitric oxide (NO), 251–2
 mechanical ventilation complications
 haemodynamic effects, 220–1
 ventilator-associated pneumonia (VAP), 221
 ventilator-induced lung injury, 221
 mechanical ventilation modes, 214–19
 airway pressure release ventilation (APRV), 219
 bilevel ventilation, 217
 classification and nomenclature, 214–15
 continuous mandatory ventilation (CMV), 216–17
 continuous positive airway pressure (CPAP), 218
 inverse ratio ventilation (IRV), 218–19
 mandatory breaths and spontaneous breaths, 216–17

 positive end expiratory pressure (PEEP), 218
 pressure control ventilation (PCV), 216
 pressure support ventilation (PSV), 218
 proportional assist ventilation (PAV), 219
 support during expiration, 218
 supporting spontaneous breaths and PSV, 218
 synchronized intermittent mandatory ventilation (SIMV), 216–17
 triggers, 217–18
 volume control ventilation (VCV), 215–16
 mechanical ventilation procedure
 endotracheal tubes, 219–20
 gastric protection, 220
 humidification, 220
 sedation, 220
 thromboprophylaxis, 220
 tracheal intubation, 219
 tracheostomy, 220
 mechanical ventilators
 airway pressure, 214
 design and function, 214
 flow and delivered volume, 214
 medical emergency teams (METs), 3–4
 meningitis
 aetiology, 319
 clinical features, 320
 definition, 319
 incidence, 319–20
 infective bacteria, 319–20
 investigations, 320
 management, 321–2
 mortality, 320
 pathophysiology, 319
 public health issues, 322
 Mental Capacity Act 2005, 87–8
 meropenem, 127
 metabolic acidosis, 153–6
 anion gap, 153–6
 osmolar gap, 153
 metabolic alkalosis, 156–7
 methicillin-resistant *Staphylococcus aureus* (MRSA), 125
 metronidazole, 128
 MEWS (modified early warning score), 31
 microangiopathic haemolytic anaemia, 182
 microcirculatory and mitochondrial distress syndrome (MMDS), 109, 112

Index

- midazolam, 79–80, 81
 Mill, John, 85
 mini tracheostomy, 15
 Modellflow, 47
 morphine, 74
 MPM (mortality probability model), 27, 32–3
 multiple organ dysfunction syndrome (MODS), 40
 definition, 108
 see also multiple organ failure
 multiple organ failure (MOF)
 clinical presentations of MODS, 111
 cytopathic hypoxia, 109
 definition, 108
 genetic disposition, 108
 gut hypothesis, 210
 hypoxic hypoxia, 109
 management of MOF, 111
 mechanisms and managements of MODS, 111–14
 microcirculatory and mitochondrial distress syndrome (MMDS), 109, 112
 multiple organ dysfunction syndrome (MODS), 108
 pathophysiology, 108–10
 possible predictors of poor outcome in MODS, 110–11
 prevention of MOF, 114–15
 secondary MODS, 108
 multiple systems organ failure (MSOF)
 definition, 108
 muscle relaxants, 8
 myocardial infarction (MI). *see* acute coronary syndromes

 necessity principle in consent, 87
 neck
 CT cervical spine, 54
 neuraxial blockade, 75–6
 neuromuscular blockade, 83
 nitric oxide (NO)
 inhalation during mechanical ventilation, 251–2
 nitroimidazoles, 128
 nitrous oxide
 analgesic properties, 75
 non-heart-beating donation (NHBD), 93
 non-invasive ventilation (NIV), 162–4
 complications and troubleshooting, 249–9
 conditions for NIV initiation, 247
 contraindications, 247–8
 definition, 246
 indications, 246–7
 instituting NIV, 248
 mechanism of action, 246
 patient selection, 247–8
 practical tips for successful NIV, 248
 role in respiratory weaning, 243
 use in acute exacerbation of COPD, 246
 use in cardiogenic pulmonary oedema, 246–7
 use in failure to wean from ventilation, 247
 use in the immunocompromised patient, 247
 non-ketotic hyperglycaemia, 136
 non-steroidal anti-inflammatory drugs (NSAIDs), 73
 norepinephrine (noradrenaline), 61
 normal (isotonic) saline, 130
 nuclear medicine investigations
 basic principles, 56
 potential risks for the critically ill, 56
 nutrition
 post-operative critical care, 167
 nutritional status assessment, 66
 anthropometric measurements, 66
 biochemical tests, 66
 body mass index (BMI), 66
 subjective global assessment, 66
 nutritional support
 enteral feeding (EF), 67–9
 estimating nutritional requirements, 66–7
 goal for critical care patients, 66
 immunonutrients, 69–70
 malnutrition risk in critical care, 66
 parenteral nutrition (PN), 69
 pharmaco/immunonutrition, 69–70
 routes of administration, 67–9
 nutritional support complications
 electrolyte imbalances, 71
 enteral feeding complications, 71
 hyperglycaemia, 70–1
 micronutrient deficiencies, 71
 overfeeding, 70
 parenteral nutrition complications, 71
 refeeding syndrome, 70

 obstetric emergencies
 amniotic fluid embolism, 355–6
 antepartum haemorrhage, 356–7
 deep vein thrombosis (DVT), 351–2
 incidence of ITU admissions, 349
 maternal physiology, 349–50
 mortality rates (maternal and neonatal), 349
 obstetric haemorrhage, 356–7
 ovarian hyperstimulation syndrome (OHSS), 357–8
 particular challenges for management, 349
 postpartum haemorrhage, 356–7
 pulmonary embolism, 350–5
 sepsis, 358
 thromboembolism, 350–5
 venous thromboembolism (VTE), 350–5
 see also pre-eclampsia and eclampsia
 oesophageal Doppler probes
 X-ray assessment of positioning, 52
 opioids
 analgesic properties, 74
 sedative agents, 81
 organ donation
 brainstem death, 92–3
 brainstem death after cardiac arrest, 173
 consent, 95–6
 contraindications and exclusion criteria, 93
 corneal donation, 94–5
 development of transplant services in England, 91
 early organ and tissue transplants, 91
 epidemiology, 91
 ethical issues, 95–6
 heart-beating donation (HBD), 91–3
 management of the potential donor, 95
 need for organs and tissues, 91
 non-heart-beating donation (NHBD), 93
 Organ Donor Register (NHS), 95
 organs and tissues that can be donated, 93–4
 transplantation success rates, 91
 types of donation, 91–3
 Organ Donation Taskforce (ODTF), 96–7
 organ failure in the critically ill patient, 1
 outcome probabilities
 development of outcome probability model, 28
 limitations of outcome probability model, 28
 outcome measures, 28
 use of scoring systems, 28
 ovarian hyperstimulation syndrome (OHSS), 357–8
 overdose. *see* poisoning
 oxazolidinones, 128
 oxygen cascade, 232
 oxygen delivery (DO₂) equation, 58

Index

- oxygen delivery and shock, 58
- oxygenation
 - blood gas analysis, 234
 - monitoring gas exchange, 232–4
 - pulse oximetry, 202
- oxygenation failure
 - acute lung injury (ALI), 236–8
 - acute respiratory distress syndrome (ARDS), 236–8
 - pathophysiology of hypoxia, 235–6
 - pulmonary embolism (PE), 238–9
- pacing wires
 - X-ray assessment of positioning, 31, 52
- paediatric emergencies
 - acute epiglottitis, 365
 - anatomy of children, 360–1
 - approach to paediatric emergencies, 361–2
 - bronchiolitis, 365–6
 - cardio-respiratory arrest in children, 362–4
 - common paediatric emergencies, 364–6
 - croup, 365
 - differences between children and adults, 360–1
 - non-accidental injury, 366
 - paediatric intensive care admission rate, 360
 - parental support, 362
 - physiology of children, 361
 - preparation for transfer to PICU, 366–8
 - recognizing the sick child, 361
 - shock, 364–5
 - stabilizing the child for retrieval, 366–8
 - status epilepticus, 366
 - weight determination for a child, 360
- pain control
 - assessment of pain severity, 72
 - health benefits of pain relief, 72
 - importance of pain relief, 72
 - occurrence of pain in critical care patients, 72
 - post-operative critical care, 159–61
 - severity of pain, 72
 - sources of pain, 72
 - types of pain, 72
 - see also* sedation
- pain relief modalities
 - adjuncts to acute pain management, 76
 - alfentanil, 74
 - anxiolysis, 76
 - chronic pain management, 76
 - codeine, 74
 - diamorphine, 74
 - epidural analgesia, 75–6
 - fentanyl, 74
 - ketamine, 75
 - morphine, 74
 - multimodality approach to analgesia, 72
 - neuraxial blockade, 75–6
 - nitrous oxide, 75
 - non-steroidal anti-inflammatory drugs (NSAIDs), 73
 - opioids, 74
 - paracetamol, 73
 - peripheral nerve blocks, 76
 - regional analgesia, 75–6
 - remifentanyl, 74
 - spinal analgesia, 75
 - systemic analgesia, 73–5
 - tramadol, 73
- pancreatitis *see* severe acute pancreatitis
- pancreatitis outcome prediction (POP) score, 271
- paracetamol, 73, 279–80
- parenteral nutrition (PN), 69, 71
 - complications, 71
 - see also* nutritional support
- Parsonnet score, 30
- partial liquid ventilation (PLV), 255
- patient admission
 - admission procedure, 19
 - classification of patient dependency, 16
 - criteria for provision of critical care, 16–17
 - decision not to admit, 19
 - decision to admit, 17
 - degree of dependency of patients, 16
 - indications for admission to critical care, 16
 - lack of available bed on the unit, 17–19
 - patient management when the unit is full, 17–19
 - use of critical care resources, 16
 - use of scoring systems to help decide, 16–17
- patient discharge
 - discharge criteria, 20
 - discharge procedure, 20
 - outcomes of premature discharge, 19–20
 - risks of delayed discharge, 19
 - use of critical care resources, 16
- patient transfer. *see* transfer of the critically ill
- penicillins, 126–7
- peripheral nerve blocks, 76
- pH. *see* acid–base balance
- pharmaco/immunonutrition, 69–70
- phenylephrine, 62
- phosphate buffer system, 150–1
- phosphodiesterase enzyme inhibitors (PDEI), 60, 62
- phosphorus levels in the body, 143–4
 - hyperphosphataemia, 144
 - hypophosphataemia, 143–4
- physician order entry (electronic prescribing), 35–7
- physiological scoring systems (PSS), 2–3
- physiology monitoring systems, 2–3
- piperacillin, 127
- poisoning
 - arterial blood gases investigation, 277
 - clinical examination, 276
 - deliberate self-poisoning, 275
 - diagnosis, 275
 - drug identification, 275
 - drug manipulation, 278
 - drug screening, 277–8
 - ECG investigation, 276
 - general supportive care, 278
 - ICU admissions, 275
 - incidence, 275
 - initial assessment and resuscitation, 275–6
 - investigations, 276–7
 - mortality rates, 275
 - patient history, 275
 - principles of care, 275
 - radiology investigations, 277
 - stimulant overdose, 282
 - toxicology search engines and databases, 719
- poisoning (specific management)
 - acetaminophen, 279–80
 - alcohols, 282
 - amphetamines, 282
 - aspirin, 279
 - barbiturates, 281
 - benzodiazepines, 281
 - beta-blockers, 283
 - calcium channel blockers, 283
 - carbon monoxide, 281
 - cocaine, 282
 - digitalis (digoxin), 282
 - lithium, 282–3
 - MDMA (ecstasy), 282
 - opioids, 281
 - organophosphates, 282
 - paracetamol, 279–80
 - phencyclidine, 282

Index

- poisoning (cont.)
 salicylates, 279
 sedatives, 281
 serotonin reuptake inhibitors (SSRIs), 282–3
 theophylline, 282
 tricyclic antidepressants (TCAs), 280–1
- post resuscitation care
 ABCDE assessment of the patient, 171
 control of seizures, 173
 correction of acidosis, 172–3
 glycaemic control, 172
 incidence of cardiac arrest, 170
 organ donation after brainstem death, 173
 outcomes of cardiac arrest, 170
 prognostication, 173
 role in the Chain of Survival, 170
 support for cardiac dysfunction, 172
 therapeutic hypothermia, 171–2
- post resuscitation syndrome, 170
- post-operative critical care
 analgesia, 159–61
 antibiotic therapy, 162, 167
 blood sugar control, 167
 cardiac arrhythmias, 167
 cardiovascular system, 164–7
 care of drains and wound, 167
 confused patient, 167
 deep vein thrombosis (DVT) prophylaxis, 167
 delirious patient, 167
 early mobilization, 168
 early post-operative period, 159–67
 fluid balance, 161–2, 165
 handover from the surgical team, 159
 hypotension, 165
 infection control, 162, 167
 late post-operative period, 167–8
 monitoring of vital parameters, 159
 non-invasive ventilation (NIV), 162–4
 nutrition, 167
 orientation of the patient, 167
 pain management, 159–61
 patients with coronary artery stents, 165
 prevention of pressure ulcers, 168
 pulmonary system, 162–4
 urine output, 162
 ventilatory support, 162–4
- potassium balance, 137–40
 hyperkalaemia, 139–40
 hypokalaemia, 138–9
 pseudohyperkalaemia, 139
- pre-eclampsia and eclampsia
 aetiology, 342
 assessment, 342–4
 control of blood pressure, 345
 definition of eclampsia, 342
 definition of pre-eclampsia, 342
 eclampsia, 346
 fluid balance, 346
 HELLP syndrome, 344, 347
 incidence, 342
 management, 345–7
 mortality, 342
 multi-disciplinary approach to care, 342
 pathogenesis, 342
 planning the delivery, 346
 postpartum care, 346–7
 prevention of convulsions, 345–6
- pressure support ventilation (PSV), 11
- pressure ulcers
 prevention, 168
- principlism, 85–6
- prognosis
 applications for scoring systems, 29
 potential drawbacks of scoring systems, 29–30
 use of scoring systems, 29–30
- prone positioning during mechanical ventilation, 251
- propofol, 8, 79–80
- proportional assist ventilation (PAV), 243
- proton pump inhibitors (PPI), 259–60
- pseudohyperkalaemia, 139
- pseudohyponatraemia, 133
- pulmonary artery (Swan–Ganz) catheter
 X-ray assessment of positioning, 52
- pulmonary embolism (PE), 238–9
 obstetric emergencies, 350–5
- pulmonary system
 post-operative critical care, 162–4
- pulse-induced contour cardiac output (PiCCO), 46–7
- pulse oximetry, 202
- PulseCO, 46
- pumpless extracorporeal lung assist (PECLA), 254
- quality and performance assessment
 Parsonnet score, 30
 SMR (standardized mortality ratio), 30
- quinolones, 127–8
- Ramsay Scale, 78
- Ransom Score (pancreatitis), 27
- Ranson scoring system, 270
- rapid sequence induction (RSI), 6–7
- recruitment manoeuvres
 use with mechanical ventilation, 252
- refeeding syndrome, 70
- referral to the critical care team
 indications for referral, 4
 routes of referral, 4
- remifentanyl, 74, 79–80
 analgesic-based sedation, 81
- renal failure. *see* acute renal failure
- renal replacement therapy (RRT)
 anticoagulation, 304
 associated therapeutic options, 303
 convection, 300
 CVVH (continuous veno-venous haemofiltration), 301
 CVVHD (continuous veno-venous haemodialysis), 301–2
 CVVHDF (continuous veno-venous haemodiafiltration), 302
 diffusion, 299–300
 fluid prescriptions, 303–4
 future possibilities in critical care, 304–5
 haemoperfusion, 303
 HVHF (high-volume haemofiltration), 303
 indications for RRT, 300–1
 intermittent vs. continuous, 302–3
 kidney functions replicated, 299–300
 modes of RRT, 301–3
 plasmapheresis, 303
 principles, 299–300
 role in critical care management, 299
 SCUF (slow continuous ultrafiltration), 301
 ultrafiltration, 299
- research
 use of scoring systems, 31–2
- resource allocation
 use of scoring systems, 30–1
- respiratory acidosis, 157
- respiratory alkalosis, 157
- respiratory failure
 immunosuppressed patients, 117
- respiratory support. *see* advanced airway management
- respiratory weaning
 adaptive support ventilation (ASV), 243–4
 assessment of readiness to wean, 241–2
 criteria for readiness to wean, 241–2
 definitions, 241
 difficult weaning, 241
 failure of attempts at weaning, 243
 goals of weaning, 241

Index

- knowledge-based expert system, 244
 methods of weaning, 243
 newer modalities for weaning, 243–4
 prolonged weaning, 241
 proportional assist ventilation (PAV), 243
 role of non-invasive ventilation (NIV), 243
 role of tracheostomy, 243
 simple weaning, 241
 specialized weaning units (SWU), 244
 spontaneous breathing trial (SBT), 242–3
 stages in the mechanical ventilation process, 241
 time required for, 241
 use of non-invasive ventilation, 247
see also extubation/weaning protocols
- resuscitation
 assessing adequacy of resuscitation, 2
 importance of early intervention, 2
 initial treatment of the critically ill patient, 1–2
 methods, 2
 monitoring patient progress, 2
 purpose, 1–2
 Revised Trauma Score (RTS), 27
 rifampicin, 128
 Rigler's sign, 53
 rocuronium, 8
- SAPS (Simplified Acute Physiology Score), 27, 32
- scoring systems
 anatomical types, 27
 functional types, 27
 generic types, 27
 physiological types, 27
 purpose and uses, 27
 specific types, 27
 systems used in critical care, 32–3
 types of, 27
- scoring systems applications, 27–32
 clinical decision making, 29–30
 early warning of 'at risk' and 'deteriorating' patients, 31
 outcome probabilities, 28
 pre-ICU patient screening, 31
 prognosis, 29–30
 quality and performance assessment, 30
 research, 31–2
 resource allocation, 30–1
 secondary multiple organ dysfunction syndrome (MODS), 108
- sedation
 aims in critical care, 77
 appropriate use in critical care, 77
 ideal properties of sedatives, 79
 importance in the critical care environment, 77
 loading dose, 79
 pharmacokinetic considerations, 80
 pharmacological management, 79
 pharmacology of commonly used sedatives, 79–80
 protocol-driven approach, 83
 required level of sedation, 77
 side effects of sedatives, 79
 use of daily sedation breaks, 83
 use of neuromuscular blockade, 83
- sedation level assessment, 77–8
 auditory evoked potentials, 78
 bispectral index (BIS), 78
 clinical scoring systems, 78
 EEG, 78
- sedative agents
 α 2-receptor agonists, 247–48
 benzodiazepines, 81
 chlorpromazine, 83
 clonidine, 79–80, 81–2
 dexmedetomidine, 82
 etomidate, 81
 haloperidol, 79–80, 83
 intravenous anaesthetic agents, 80–1
 ketamine, 80–1
 lorazepam, 79–80, 81
 midazolam, 79–80, 81
 neuroleptic agents, 83
 opioids, 81
 propofol, 79–80
 remifentanyl, 79–80, 81
 thiopentone, 80
 volatile anaesthetic agents, 81
- Seldinger technique for cannulation, 41–2, 45–4
- Sellick's manoeuvre, 7
- sepsis
 definition of sepsis, 99
 definition of septic shock, 99
 definition of severe sepsis, 99
 diagnostic criteria, 99
 goals for resuscitation, 2
 management, 103–6
 obstetric emergencies, 358
 pathophysiology, 99–103
 resuscitation, 103–6
 signs and symptoms of infection (SSI), 100
 Surviving Sepsis Campaign, 99
 systemic inflammatory response syndrome (SIRS), 99
- septic shock
 definition, 99
- severe acute pancreatitis (SAP)
 aetiology, 266–7
 APACHE II score, 270–1
 common complications, 266
 diagnosis, 268–9
 early and late phases of mortality, 269–70
 Glasgow Score (Imrie score), 270
 incidence, 266
 mortality rate, 266
 pancreatitis outcome prediction (POP) score, 271
 pathophysiology, 266
 prognostic factors, 269–71
 Ranson criteria, 270
 severity of illness scores, 270–1
 treatments, 271–3
- severe sepsis
 definition, 99
- shock
 and oxygen delivery, 58
 causes of, 58
 choice of vasoactive drug treatment, 62
 definition, 58
 in children, 364–5
 indications that resuscitative therapy is working, 64
 inotropes, complications and considerations, 65
 management of, 62–3
 monitoring the patient, 64–5
 role of vasoactive drugs, 62–3
 vasoactive drug treatment and responses, 63–4
 vasoactive drug treatment complications, 65
- shunt
 hypoxia caused by, 235
- simulated environments
 role in staff training, 38
- sodium balance, 132–7
 diabetes insipidus, 136
 hypernatraemia, 135–7
 hyponatraemia, 133–5
 hyponatraemic encephalopathy, 134
 non-ketotic hyperglycaemia, 136
 pseudohyponatraemia, 133
- SOFA (Sequential Organ Failure Assessment), 27, 33, 100
- solid organ transplant patients
 infection and complications, 120–1
- spinal analgesia, 75
- standard base excess, 150
- standardized mortality ratio (SMR), 30
- starch solutions, 131

Index

- status epilepticus
 aetiology, 306
 barbiturates, 310
 benzodiazepines, 309
 classification, 306
 definition, 306
 fosphenytoin, 310
 generalized convulsive status epilepticus (GCSE), 306
 incidence, 306
 long-term anti-epileptic drug (AED) therapy, 310–11
 mortality and morbidity, 306
 non-convulsive status epilepticus (NCSE), 306
 occurrence in critical care, 306
 outcome, 311
 paediatric, 366
 pathophysiology, 306–8
 phenytoin, 309–10
 propofol, 310
 refractory status epilepticus, 310
 terminating seizure activity, 308–10
 treatment of acute status epilepticus, 308–10
- stroke volume, 58
 strong ion gap, 149–50
 suxamethonium, 8
 synchronized intermittent mandatory ventilation (SIMV), 11
- tazobactam, 127
 teicoplanin, 127
 therapeutic hypothermia
 post resuscitation care, 171–2
 thermomodulation technique
 cardiac output monitoring, 44–5
 thiopentone, 8, 80
 thoracic electrical bioimpedance, 48
 thrombocytopenia, 180–1
 thromboembolism
 obstetric emergencies, 350–5
 thrombotic thrombocytopenia purpura (TTP), 182
 ticarcillin, 126
 TISS (Therapeutic Intervention Scoring System), 27, 30–1
 total liquid ventilation (TLV), 255
 total parenteral nutrition (TPN), 69
 toxicology. *see* poisoning
 tracheostomy, 12–15
 complications, 12–13
 contraindications, 12
 cricothyroidotomy, 14–15
 indications for, 12
 mini tracheostomy, 15
 percutaneous tracheostomy, 12
 surgical tracheostomy, 12
 techniques, 12
 timing, 12
 tracheostomy tubes, 13–14
 tracheostomy tube
 X-ray assessment of positioning, 52
 ‘track and trigger’ scores, 43–44
 training and education
 simulated environments, 38
 tramadol, 73
 transfer of the critically ill
 air transfer, 22
 considerations during patient transfer, 24–6
 Critical Care Network, 21
 equipment required, 23–4
 handover to the admitting team, 26
 indications for transfer, 21
 inter- and intra-hospital transfers, 21
 modes of transport, 22
 monitoring during transfer, 23, 24–6
 patient care during transfer, 24–6
 patient preparation for transfer, 23
 patient transfer record sheet, 24
 personnel, 22
 post transfer, 26
 preparation prior to transfer, 23
 primary transfers, 21
 risks associated with transfer, 21
 road transfer by ambulance, 22
 secondary transfers, 21
 specialist transfer team, 22
 training of the transfer team, 22
 transfer due to lack of capacity, 21
 transjugular intrahepatic portosystemic shunt (TIPS), 261
 transoesophageal Doppler, 45–6
 transoesophageal echocardiography, 47
 transtracheal gas insufflation (TGI), 256
- trauma
 ABCDE approach to primary survey, 334–8
 airway with cervical spine control, 334–5
 breathing, 335–6
 circulation with haemorrhage control, 336–8
 clearance of the cervical spine, 338
 definition of the ‘golden hour’, 334
 disability and exposure, 338
 pattern of trauma mortality, 334
 secondary survey, 334
 trauma scores, 33
 traumatic brain injury (TBI)
 cerebral contusions, 326
 classification, 325
 diffuse injury, 326
 extradural haematoma, 325
 focal injuries, 325–6
 incidence, 325
 initial assessment and resuscitation, 327
 intensive therapy management of TBI, 330–3
 intracerebral haematoma, 325
 monitoring the injured brain, 328–30
 mortality and morbidity, 325
 outcome, 333
 pathophysiology, 325–6
 pathophysiology of intracranial pressure, 328
 primary and secondary injury, 325–6
 subdural haematoma, 325
 TRISS (Trauma and Injury Severity Score), 27, 33
 troponins (I and T)
 ACS biomarkers, 189
 tubes and catheters
 X-ray assessment of positioning, 51–2
- ultrasound imaging
 basic principles, 56
 use in the critically ill, 57
- unconscious patient
 abnormal levels of consciousness, 312
 assessment, 314–15
 AVPU score, 315
 brainstem death, 312
 causes of unconsciousness, 312–13
 EEG, 316
 examination, 315
 Glasgow Coma Score (GCS), 314
 history, 315
 immediate assessment, 314
 investigations, 315–16
 locked-in syndrome, 312
 lumbar puncture, 316
 management, 316–18
 minimally conscious state, 312
 neurological conditions imitating coma, 312
 radiological tests, 315–16
 vegetative state, 312
- unconsciousness
 causes of, 312–13
 herniation of brain tissue, 313
- urine output
 post-operative critical care, 162
 utilitarianism, 85

Index

- vancomycin, 127
- vancomycin-resistant Enterococci (VRE), 125, 127
- vasoactive drug mechanisms, 59–60, 61–2
 - alpha-1 (α 1) receptors, 59
 - beta receptors
 - beta-1 (β 1) receptors, 59
 - beta-2 (β 2) receptors, 59
 - dopaminergic (D1 and D2) receptors, 59–60
 - phosphodiesterase enzyme enzyme inhibitors (PDEI), 60
 - vasopressin receptors, 61–2
- vasoactive drug treatment
 - deciding which drug to use, 62
 - indications that resuscitative therapy is working, 64
 - inotropes, complications and considerations
 - monitoring the patient, 64–5
 - patient responses, 63–4
 - role in management of shock, 62–3
 - use of vasoactive drugs, 63–4
- vasoactive drugs
 - definitions, 59
 - effects of inotropes, 58, 59
 - effects of vasopressors, 59
- vasoactive drugs commonly used
 - dobutamine, 61
 - dopamine, 61
 - epinephrine (adrenaline), 60–1
 - levosimendan, 62
 - norepinephrine (noradrenaline), 61
 - phenylephrine, 62
 - phosphodiesterase enzyme inhibitors (PDEI), 62
 - vasopressin, 61–2
- vasopressin, 61–2
- venous thromboembolism (VTE)
 - obstetric emergencies, 350–5
- ventilation
 - alveolar ventilation, 228
 - expiratory reserve volume (ERV), 226
 - flow volume loop, 227–8
 - functional residual capacity (FRC), 226
 - inspiratory reserve volume (IRV), 226
 - lung volumes and capacities, 226
 - monitoring pulmonary function, 227–8
 - residual volume (RV), 226
 - respiratory control system, 226–7
 - spirometry, 227
 - tidal volume (VT), 226
 - total lung capacity (TLC), 226
 - vital capacity (VC), 226
 - see also* advanced airway management, mechanical ventilation, non-invasive ventilation
- ventilation disorders
 - hyperventilation syndromes, 230–1
 - hypoventilation syndromes, 228–30
- ventilation/perfusion (V/Q)
 - mismatch, 235
- ventricular assist devices (VADs), 209
- viral encephalitis
 - aetiology, 322
 - clinical features, 322
 - definition and characteristics, 322
 - investigations, 322–3
 - management, 323–4
 - prognosis, 324
- virtual ICU, 37–8
- von Willebrand disease, 179–80
- voriconazole, 129
- weaning. *see* extubation/weaning protocols; respiratory weaning
- welfare attorney (end-of-life care), 89
- X-ray imaging, 49–53
 - abdominal X-ray, 53
 - appearance of lungs, 52–3
 - assessment of tubes and catheters, 51–2
 - basic principles, 49–50
 - chest X-ray, 50–3
 - detector, 49
 - interpretation of the antero-posterior chest X-ray, 51
 - radiation dose to the patient, 50
 - X-ray source, 49