

Index

- absorbing set, **117**
- almost bi-Lipschitz embeddings into
 Euclidean spaces, 97
 for metric spaces, 100
- analytic function, *see* real analytic function
- approximate inertial manifold, 143
- Assouad dimension, 81, 83, **85**
 almost bi-Lipschitz embeddings into Hilbert
 spaces, 94
 bi-Hölder embeddings into Euclidean
 spaces, 94
 bounds upper box-counting dimension, 85
 embedding into Euclidean spaces when
 $d_A(X - X) < \infty$, 97
 ill-behaved for difference sets, 89
 monotonicity, 85
 not sufficient for bi-Lipschitz embeddings
 into Euclidean spaces, 92
 of orthogonal sequences, 88–91
 of products, 86
 of \mathbb{R}^n , 85
 of unions, 85
- attracting set, **115**
- attractor, *see* global attractor
- Auerbach basis, 79, 127
 existence, 74
- $B_N = B_N(0, 1)$ (unit ball in \mathbb{R}^N), **42**
- $B_Z(0, r)$ (r -ball in the space Z), 123
- backwards uniqueness, 136, 144
- Baire Category Theorem, **13**, 17, 177
- ball, image under a linear map, 133, 146
- bi-Lipschitz embedding
 finite Assouad dimension not sufficient, 92
 finite box-counting dimension not sufficient,
 81
- bi-Lipschitz mapping, **84**
- Borel–Cantelli Lemma, **43**, 45, 77, 98
- bounded linear maps, 41
 decomposition in: Banach spaces, 78;
 Hilbert spaces, 57
- box-counting dimension, **31**
 lower, *see* lower box-counting dimension
 non-existence in general, 31
 of products, 35
 upper, *see* upper box-counting dimension
- Brouwer Fixed Point Theorem, **10**, 12
- Brunn–Minkowski inequality, 54, 56
- $C_c^\infty(\Omega)$ (C^∞ functions with compact support
 in Ω), 106
- Cantor set, 31, 58
- CAT(0) space, 182
- covering, **8**
 Lebesgue number, **13**, 18
 mesh size, **10**
 order, **8**
 refinement, **8**
- covering dimension, 7, **8**
 bounded by Hausdorff dimension, 27
 characterised in terms of: Assouad
 dimension, 85; box-counting dimension,
 39; Hausdorff dimension, 29
 embedding into Euclidean spaces, 17
 monotonicity, 8
 of compact sets, 13
 of products, 9, 40
 of unions, 9
- covering lemma, 30
- d_A , *see* Assouad dimension
 d_B , *see* upper box-counting dimension

- d_H , *see* Hausdorff dimension
 d_{LB} , *see* lower box-counting dimension
 decomposition of bounded linear maps
 Banach spaces, 78
 Hilbert spaces, 57
 degrees of freedom, 168
 density of prevalent sets, 48
 determining nodes, 167
 diameter of a set, **10**
 difference set, **41**, 161
 for metric spaces, 100
 ill-behaved with Assouad dimension, 89
 differential inequality
 a version of Henry's Lemma 7.1.1, 132
 Ghidaglia's Lemma, 143
 Uniform Gronwall Lemma, 121
 dim, *see* covering dimension
 dist (Hausdorff semi-distance), **10**
 doubling space, **84**
 if and only if homogeneous space, 84
 dual thickness, 65, **69**
 bounded by Lipschitz deviation in a Hilbert space, 72
 bounded by the upper box-counting dimension, 71
 is zero when thickness exponent is zero, 73

 ϵ -mapping, **13**
 eigenvalues, asymptotics, 137, 139, 143, 164
 embedding into Euclidean spaces
 for metric spaces, 100
 when $d_A(X - X)$ is finite, 97
 when $d_B(X)$ is finite, 43, 75
 when $d_H(X - X)$ is finite, 60
 when $\dim(X)$ is finite, 17
 Euclidean space, *see* \mathbb{R}^n

 fractal, 29
 fractional power spaces, 107
 and Sobolev spaces, 108
 compact embeddings, 108, 113
 interpolation inequality, 141, 143, 164
 Frostman's Lemma, **25**

 G_δ (countable intersection of open sets), **17**
 general position, **14**
 generic, **17**
 geometric independence, **15**
 global attractor, 105, **115**
 bound on upper box-counting dimension:
 for semilinear parabolic equations, 130;
 general method, 124; when Df has finite rank, 132; when $Df \in \mathcal{L}_1$, 130; when $Df \in \mathcal{L}_s$, $s < \frac{1}{4}$, 129
 contains unstable sets, 121
 dimension and degrees of freedom, 168
 existence for: Navier–Stokes equations, 119; semilinear parabolic equations, 118; necessary and sufficient condition, 117
 is connected, 121
 is the maximal compact invariant set, 115, 121
 is the minimal closed attracting set, 115, 121
 is union of all globally bounded solutions, 118
 of Navier–Stokes equations: has zero Lipschitz deviation when $f \in L^2$, 142; is bounded in $D(A)$ when $f \in H$, 122; is real analytic when f is real analytic, 168
 parametrised by point values, 160
 uniqueness, 115
 graph of a function, Hausdorff dimension, 29
 Gromov–Hausdorff metric, **92**

 $H^s(\Omega)$, *see* Sobolev spaces
 Haar null, 47
 Hahn–Banach Theorem, 60, 71, 79, 95
 Hausdorff dimension, **23**
 bounded by box-counting dimensions, 34
 bounds covering dimension, 27
 embedding into Euclidean spaces when $d_H(X - X) < \infty$, 60
 impossibility of general linear embedding, 58
 Mañé's embedding result, 57
 monotonicity, 24
 of a graph, 29
 of products, 25, 35, 40
 under Hölder continuous maps, 24
 Hausdorff measure, **21**
 and Lebesgue measure, 22
 spherical, **30**
 Hausdorff semi-distance, **10**
 Heisenberg group, 91
 Hölder continuity, 24, 34, 43, 149
 Hölder continuous inverse
 of linear embedding map, 75
 homogeneity, 83
 if and only if doubling, 84
 of subsets of Euclidean spaces, 83
 under bi-Lipschitz maps, 84

- I_n (unit cube in \mathbb{R}^d), **10**
 inductive dimensions, **7, 17, 18**
 equal to covering dimension in a separable
 metric space, **19**
 inertial manifold, **167**
 interpolation inequality in fractional power
 spaces, **141, 143, 164**
 invariant set, **115**
 isometric embedding of metric spaces
 ℓ^∞ , **101**
 $L^\infty(X)$, **100**
- Johnson–Lindenstrauss Lemma, **74**
- $\mathcal{K}(\mathcal{B})$ (compact linear maps from \mathcal{B} into \mathcal{B}),
125
- Kuratowski embedding, **100**
- $L^2_{\text{per}}(\Omega, \mathbb{R}^d)$ (periodic functions in
 $L^2(\Omega, \mathbb{R}^d)$), **160**
- \mathcal{L}_λ (sums of compact maps and contractions),
125
- \mathcal{L}^n (n -dimensional Lebesgue measure), **22**
 Lebesgue covering dimension, *see* covering
 dimension
 Lebesgue Covering Theorem, **10**
 Lebesgue number of a covering, **13, 18, 171**
 linear operator
 fractional powers, **107**
- Lipschitz deviation, **65, 68**
 bounded by the thickness exponent, **68**
 bounds the dual thickness in a Hilbert space,
 72
 is zero for the attractors of certain
 semilinear parabolic equations, **139**
- lower box-counting dimension, **32, 77**
 bounded by upper box-counting dimension,
 32
 bounds Hausdorff dimension, **34**
 in general no embeddings with Hölder
 inverse, **81**
 of orthogonal sequences, **36**
 of products, **35**
- $\mathcal{L}(X)$ (space of bounded linear maps from X
 into X), **41**
- $\mathcal{L}(X, Y)$ (space of bounded linear maps from
 X into Y), **41**
- measurable set, **21, 22**
 measure, **21**
 mesh size, **10**
- metric outer measure, **21**
 modified upper box-counting dimension, **40,**
63
 equal to packing dimension, **40**
 multi-index, **107, 161, 169**
- Navier–Stokes equations, **30, 109–113, 139,**
144
 as a semilinear parabolic equation, **113**
 attractor: existence, **119**; has zero Lipschitz
 deviation when $f \in L^2$, **142**; has zero
 thickness when f is smooth, **138**; is
 ‘smooth’ when f is smooth, **138**; is
 bounded in $D(A)$ when $f \in H$, **122**; is
 real analytic when f is real analytic, **168**
 determining nodes, **167**
 generate a semigroup, **110**
 heuristic theories of turbulence, **169**
 in functional form, **111**
 orthogonality properties of nonlinear term,
 111
 negatively invariant set, **123**
 nonlinear semigroup, *see* semigroup
- Ω_n (volume of unit ball in \mathbb{R}^n), **22**
 omega limit set, **116, 121**
 order of a covering, **8**
 order of vanishing, **162**
 and zero set, **163**
 orthogonal projection, **81**
 rank, **62**
 orthogonal sequence, **25, 63, 78, 82, 99**
 Assouad dimension, **88–91**
 box-counting dimensions, **36**
 dual thickness, **80**
 thickness exponent, **67**
- outer measure, **20**
- packing dimension, **40**
 equal to modified upper box-counting
 dimension, **40**
- periodic orbit
 minimal period: for ODEs, **153**; for
 semilinear parabolic equations, **156**
 obstruction to the Takens Time-Delay
 Theorem, **152**
- periodic point, **150**
 Poincaré inequality, **112, 114, 120**
- polyhedron, **15**
 prevalence, **46, 47, 48**
 implies density, **48**

- preserved under: countable unions, 49; finite unions, 49
 probability measure, **21**
 probe space, **48**
 products of sets
 Assouad dimension of, 86
 box-counting dimensions of, 35
 covering dimension of, 9, 40
 Hausdorff dimension of, 25, 35, 40
 projection, *see also* orthogonal projection
 rank in ℓ^p , 78
 quadrilateral inequality, 93, 100
 rank
 of orthogonal projection, 62
 of projection in ℓ^p , 78
 real analytic function, 160, **161**
 bounds on derivatives, 169
 has finite order of vanishing, 162
 zero set, 163
 refinement of a covering, **8**
 semigroup, **105**
 generated by: a semilinear parabolic equation, 109; the Navier–Stokes equations, 110
 semilinear parabolic equation, **108**
 existence of global attractor, 118
 finite-dimensional attractor, 130
 generates a semigroup, 109
 minimal period of periodic orbits, 156
 shyness, **47**
 for subsets of \mathbb{R}^n , 48
 preserved under finite unions, 48
 simplex, **15**
 singular values of a matrix, **146**, 158
 Sobolev spaces, 106
 and continuous functions, 107, 113
 and fractional power spaces, 108
 spherical Hausdorff measure, **30**
 squeezing property, 143
 Stokes operator, 111
 eigenvalues on $[0, 2\pi]^2$, 143
 Takens Time-Delay Embedding Theorem
 in \mathbb{R}^N , 149
 in a Banach space, 154
 obstruction from existence of periodic orbits, 152
 $\tau(X)$, *see* thickness exponent
 thickness exponent, 46, 64, **65**
 alternative definition, 73
 bounded by the upper box-counting dimension, 66
 bounds the Lipschitz deviation, 68
 in two spaces, 65
 of an orthogonal sequence, 67
 of set of real analytic functions, 163
 related to bounds in Sobolev spaces, 137
 zero for: ‘smooth’ sets, 67, 137; the Navier–Stokes attractor when f is smooth, 138
 zero implies zero dual thickness, **73**
 zero thickness implies zero dual thickness, 70
 turbulence, 169
 Uniform Gronwall Lemma, 121
 unit ball, volume, 22
 unit cube, hyperplane slices of, 52
 unstable set, **121**
 contained in global attractor, 121
 upper box-counting dimension, **32**
 alternative definitions, 39
 bounded by Assouad dimension, 85
 bounds: Hausdorff dimension, 34; the dual thickness, 71; the thickness exponent, 66
 can be calculated through a geometric sequence, 33
 embedding into: \mathbb{R}^k for subsets of \mathbb{R}^N , 43; Euclidean spaces, 75
 finite for attractors of semilinear parabolic equations, 130
 impossibility of bi-Lipschitz embeddings in general, 81
 in two spaces, 35
 modified, *see* modified upper box-counting dimension
 monotonicity, 33
 of an invariant set, 124
 of orthogonal sequences, 36
 of products, 35
 of set of real analytic functions, 163
 of unions, 33
 of unit cube in \mathbb{R}^n , 34
 under Hölder continuous maps, 34
 variation of constants formula, **108**, 113, 130, 156
 Vitali Covering Theorem, 23
 weak derivative, 106