

## Contents

---

<i>Preface</i>	<i>page xi</i>
<b>Introduction</b>	<b>1</b>
PART I: FINITE-DIMENSIONAL SETS	
<b>1 Lebesgue covering dimension</b>	<b>7</b>
1.1 Covering dimension	8
1.2 The covering dimension of $I_n$	10
1.3 Embedding sets with finite covering dimension	12
1.4 Large and small inductive dimensions	17
Exercises	18
<b>2 Hausdorff measure and Hausdorff dimension</b>	<b>20</b>
2.1 Hausdorff measure and Lebesgue measure	20
2.2 Hausdorff dimension	23
2.3 The Hausdorff dimension of products	25
2.4 Hausdorff dimension and covering dimension	26
Exercises	29
<b>3 Box-counting dimension</b>	<b>31</b>
3.1 The definition of the box-counting dimension	31
3.2 Basic properties of the box-counting dimension	33
3.3 Box-counting dimension of products	35
3.4 Orthogonal sequences	36
Exercises	39
<b>4 An embedding theorem for subsets of <math>\mathbb{R}^N</math> in terms of the upper box-counting dimension</b>	<b>41</b>

viii	<i>Contents</i>	
<b>5</b>	<b>Prevalence, probe spaces, and a crucial inequality</b>	<b>47</b>
5.1	Prevalence	47
5.2	Measures based on sequences of linear subspaces	49
	Exercises	56
<b>6</b>	<b>Embedding sets with <math>d_H(X - X)</math> finite</b>	<b>57</b>
6.1	No linear embedding is possible when $d_H(X)$ is finite	58
6.2	Embedding sets with $d_H(X - X)$ finite	60
6.3	No modulus of continuity is possible for $L^{-1}$	62
<b>7</b>	<b>Thickness exponents</b>	<b>64</b>
7.1	The thickness exponent	65
7.2	Lipschitz deviation	67
7.3	Dual thickness	69
	Exercises	73
<b>8</b>	<b>Embedding sets of finite box-counting dimension</b>	<b>75</b>
8.1	Embedding sets with Hölder continuous parametrisation	75
8.2	Sharpness of the Hölder exponent	77
	Exercises	81
<b>9</b>	<b>Assouad dimension</b>	<b>83</b>
9.1	Homogeneous spaces and the Assouad dimension	83
9.2	Assouad dimension and products	86
9.3	Orthogonal sequences	88
9.4	Homogeneity is not sufficient for a bi-Lipschitz embedding	91
9.5	Almost bi-Lipschitz embeddings	94
9.6	Sharpness of the logarithmic exponent	99
9.7	Consequences for embedding compact metric spaces	100
	Exercises	100
PART II: FINITE-DIMENSIONAL ATTRACTORS		
<b>10</b>	<b>Partial differential equations and nonlinear semigroups</b>	<b>105</b>
10.1	Nonlinear semigroups and attractors	105
10.2	Sobolev spaces and fractional power spaces	106
10.3	Abstract semilinear parabolic equations	108
10.4	The two-dimensional Navier–Stokes equations	109
	Exercises	113

<i>Contents</i>		ix
<b>11</b>	<b>Attracting sets in infinite-dimensional systems</b>	<b>115</b>
	11.1 Global attractors	115
	11.2 Existence of the global attractor	115
	11.3 Example 1: semilinear parabolic equations	118
	11.4 Example 2: the two-dimensional Navier–Stokes equations	119
	Exercises	121
<b>12</b>	<b>Bounding the box-counting dimension of attractors</b>	<b>123</b>
	12.1 Coverings of $T[B(0, 1)]$ via finite-dimensional approximations	125
	12.2 A dimension bound when $Df \in \mathcal{L}_{\lambda/2}(\mathcal{B})$ , $\lambda < \frac{1}{2}$	129
	12.3 Finite dimension when $Df \in \mathcal{L}_1(X)$	130
	12.4 Semilinear parabolic equations in Hilbert spaces	130
	Exercises	132
<b>13</b>	<b>Thickness exponents of attractors</b>	<b>136</b>
	13.1 Zero thickness	136
	13.2 Zero Lipschitz deviation	138
	Exercises	143
<b>14</b>	<b>The Takens Time-Delay Embedding Theorem</b>	<b>145</b>
	14.1 The finite-dimensional case	145
	14.2 Periodic orbits and the Lipschitz constant for ordinary differential equations	152
	14.3 The infinite-dimensional case	154
	14.4 Periodic orbits and the Lipschitz constant for semilinear parabolic equations	156
	Exercises	158
<b>15</b>	<b>Parametrisation of attractors via point values</b>	<b>160</b>
	15.1 Real analytic functions and the order of vanishing	161
	15.2 Dimension and thickness of $\mathcal{A}$ in $C^r(\Omega, \mathbb{R}^d)$	163
	15.3 Proof of Theorem 15.1	165
	15.4 Applications	167
	Exercises	169
	<i>Solutions to exercises</i>	170
	<i>References</i>	196
	<i>Index</i>	202