

# Index

- ‘A Cognitive Perspective on Social Phobia’ (Clark), 124
  - acceptance and commitment therapy (ACT)
    - for CYP, 321
    - efficacy for depression in dementia caregivers, 300
  - activity monitoring, 239, 321
    - brief BA for adolescents, 318
    - depression case study, 240
  - acute in-patients
    - concerns around feasibility of psychological therapies, 249
    - prevalence of adverse childhood experiences, 256
  - acute in-patients CBT, 248–261
    - adaptations to therapy, 255
    - and aims of crisis services, 248–249
    - assessing impact of past experiences, 257
    - average length of admissions, 249
    - BPS and ACP-UK guidelines, 252–253
    - case consultation and ‘indirect psychological interventions’, 258–259
    - components and features of in-patient & crisis services, 252–253
    - contracting of therapy, 253
    - formulating and conceptualising risk, 251–252
    - formulating maintenance cycles, 257
    - identification of exits and psychological interventions, 258
    - individual assessment, formulation and intervention, 253–258
    - managing disagreement with diagnosis or detention, 255
    - pacing and timing of assessment and formulation, 254–255
    - personal history assessments and predisposing factors, 256–257
    - and phased-based models of intervention, 249–250, 256
    - potential benefits of intensive psychological interventions, 249
    - reasons for admission, 248
    - role of the therapist, 258
    - team formulation, 259–261
    - transdiagnostic psychological therapy groups, 258
  - acute stress disorder (ASD), 335–336
    - diagnostic criteria, 335
    - adaptations for specific populations, 277–278
    - Arundell’s typology, 277
    - children, 277
    - cultural adaptations, 278
    - dementia caregivers, 278
    - Hinton and Patel’s contextually sensitive approach, 278
    - older people, 277
  - adverse childhood experiences (ACEs)
    - ACEs model of CBT for CYP, 313
    - prevalence in inpatient psychiatric settings, 256
  - agoraphobia
    - aetiology, 112
    - Clark et al.’s treatment model, 29
    - as common comorbidity with panic disorder, 105
  - Akkerman, R. L., 301
  - Alexopoulos, G. S., 282
  - Alzheimer’s disease and related dementias (ADRD)
    - prevalence, 295
    - see also* dementia; dementia caregivers.
  - antidepressant medication, comparison of CBT with, 5
  - anxiety, diagnoses where anxiety plays a key role, 170
  - anxiety disorders, 93–96
    - anxiety equation, 94–95
    - behavioural experiments, 56 for belief
    - disconfirmation, 95
    - behavioural theory of, 174
    - application, 93–94
  - Clark and Salkovskis’s treatment models, 4, 29, 95
  - cognitive change methods, 94–95
  - cognitive theory, 94–95
  - efficacy of CBT for, 28–30, 93
  - exposure for habituation, 93–94
  - factors in Beck’s cognitive model, 94–95
  - and graded exposure, 94
  - interventions for CYP, 319–320
  - key behavioural factors, 93
  - measures of anxiety, 29
  - NICE recommendation for CBT treatment, 25, 93
  - vicious flower model, 96
- APA guidelines, recommendation of CBT as treatment for a range of common mental health conditions, 22

- Aranda, M. P., 299
- Arundell, L. L., 277
- assertive defence of the self, 198
- assessment
- the process, 80–82
  - Structured Clinical Interview for DSM-5 (SCID-5), 81
- attachment theory
- and adaptations for working with children, 277
  - application in the PCMH context, 246
  - potential role in understanding a child's distress, 313
  - usefulness when applied to CBT, 265
- attention deficit hyperactivity disorder (ADHD), 157
- GAD case study, 157
  - potential impact on CYP, 324
  - recommendations for adapting CBT, 324–325
- attention training
- body dysmorphic disorder case study, 201
  - GAD case study, 162
  - in social anxiety disorder therapy, 128
- autism spectrum disorder (ASD), 314, 321
- adapting parent-led therapy for, 320
  - potential impact on CYP, 324
  - recommendations for adapting CBT, 324–325
- Avatar Therapy, evidence for the efficacy of for hearing voices, 270
- avoidance model of worry (AMW), 143–144
- avoidant personality disorder, similarity with social anxiety disorder, 124
- avoided situations, creating a hierarchy of, 113
- Baltes, P., 290
- Bandura, A., 298
- Barlow, D. H., 106, 153
- Barlow's Unified Protocol, 86
- 'Basics and Beyond' (Beck), 54
- Beck, Aaron T.
- cognitive model proposed by, 94
  - cognitive theory of GAD, 142
  - conceptualisation of GAD, 145
  - development of Cognitive Therapy, 1–4
  - first RCT of CBT conducted by, 5
  - influences, 3
  - main elements of Beck's approach to CBT, 2
  - original RCT evaluating efficacy of CBT for depression, 25
  - research on the efficacy of psychoanalytic psychotherapy, 2
  - stress-diathesis model of depression, 2
  - stress-diathesis model of psychopathology, 13–15
- Beck Anxiety Inventory (BAI), 29
- Beck Depression Inventory II (BDI-II), 231
- Beck's Psychopathology Model for CBT, 15
- Behar, E., 142
- behaviour therapy (BT), Beck's
- differentiation of cognitive therapy from, 4
- behavioural activation, 57–58
- adolescent model, 316–318
  - couples therapy, 361
  - creation of variety of activity, 58
  - dementia caregiving case study, 305
  - disconfirming of depressogenic beliefs with, 233, 237
  - disconfirming of depressogenic beliefs with, 57
  - efficacy, 6, 232
  - focus on bodily cycles, 58
  - GAD case study, 153
  - history of in treatment for depression, 232
  - Martell model, 6, 238–241
  - rationale for, 57
- behavioural experiments, 56, 61
- body dysmorphic disorder, 196–197
- depression, 235
- GAD case study, 158, 163
  - IAD, 206–207
  - OCD, 182–183, 187, 188
  - panic disorder, 107–110
  - personality disorder, 266–267
  - PTSD, 216, 222, 225
  - social anxiety disorder, 346
  - social anxiety disorder case study, 130–131
- behavioural therapy
- activity diaries and mood ratings, 239
  - depression, 238–241
  - case study, 240–241
  - phobias, case study, 99–100
  - problem-solving strategies, 239
  - rumination work, 239
- behaviourism, pioneering work of Mary Cover Jones, 28
- benchmarking therapy, 46
- Benjamin Rose Institute on Aging, 299
- Benne, K. D., 349
- Big I/Little i technique, 198
- body dysmorphic disorder case study, 202
- Bilbrey, A. C., 301
- bi-polar disorder
- characteristics of, 268
  - principles for effective engagement, 245
- bi-polar disorder CBT, 244, 268–270
- contents of therapy sessions
  - sessions 1–6, 270
  - sessions 6–24, 270
  - sessions 24–30, 270
- Blackburn, Ivy-Marie, 5
- body dysmorphic disorder (BDD), 193–202
- age of onset, 194, 202
  - comorbidities, 194
  - cosmetic surgery, cost benefit analysis, 198
  - definition, 193
  - diagnostic criteria, 193
  - evidence for CBT for, 320

- (BDD) (cont.)  
 impact on the patient, 194  
 prevalence, 194  
 safety behaviours, 195  
 screening measures, 194  
 treatment (Veale and Neziroglu's model), 194–202  
 case study, 198–202  
 cognitive change techniques, 198  
 complications, 198  
 of covert mental strategies, 196  
 efficacy data for CBT, 194  
 feedback, 197  
 guidelines for helpful use of mirrors in BDD, 197  
 key protocol, 194–196  
 managing overt behavioural strategies, 197
- borderline personality disorder, role of early life emotional neglect, 256
- Borkovec, T. D., 143, 151
- Bowlby, J., 313
- breathing skills, 113
- Breitholtz, E., 142
- Brodaty, H., 297
- Bryant, R. A., 335
- 'Building Better Caregivers', 302
- Butler, A.C., 27
- Butler, G., 145
- caregiver burden, cultural perspective, 299
- caregiver stress, sociocultural model, 299
- caregivers  
 evidence of benefits of CBT for, 30  
 mobile apps for, 372  
 potential impact of caregiving on mental health of, 295  
 psychoeducational programs for grandparents as, 304  
 technology-based CBT for, 371–372  
*see also* dementia caregivers.
- caregiving  
 Benjamin Rose Institute's best practice website, 299  
 stress process model of, 298
- caregiving burden  
 cultural perspective, 297  
 definition, 297  
 dementia caregivers, 297
- CareLoop, for relapse prevention in psychosis, 270
- caring for a person living with dementia (PLWD)  
 case study, 304–305  
 factors influencing the caregiver's experience, 296  
 as international concern, 296  
 mobile apps for, 372  
 positive and negative effects, 296  
 theoretical models, 298–300  
*see also* dementia caregivers.
- Chaplin, R., 23
- Child Outcomes Research Consortium (CORC), 314
- children, developmental adaptations, 277
- children and young people (CYP)  
 IAPT programme, 341  
 models and interventions  
 ACEs model, 313  
 anxiety disorders, 319–320  
 Brief Behavioural Activation for Adolescents, 316–318  
 building a therapeutic alliance, 315  
 CBT, 312–325  
 co-morbidities to consider, 314  
 comparison with adult models, 313  
 computerised/digital CBT, 322  
 developmental considerations, 312–313  
 assessing developmental needs, 323  
 attachment, 323
- environment, breaks and session length, 324  
 language, 323  
 neurodiversity, 324–325  
 play and creativity, 324  
 effectiveness, 316–321  
 effectiveness and long-term plan, 312  
 engagement, assessment and formulation, 314–316  
 exposure response prevention (ERP), 320  
 innovations and CYP-specific recommendations, 321–322  
 key CBT models, literature, and protocols, 317  
 low intensity interventions, 322  
 low mood, 316–318  
 obsessive compulsive and related disorders, 320  
 outcomes measures, 314–315  
 parent led CBT (PL-CBT), 319–320  
 parental involvement, 321–322  
 Piagetian theory and, 277, 313  
 PTSD, 320–321  
 research recommendations, 319  
 risk and safety planning, 314  
 Vygotsky's theories and, 277
- Clark, D. M., 4–5, 11, 29, 95, 106, 107, 112, 122, 124, 126, 129, 142, 145, 223, 227, 253
- classical conditioning, development of fear through, 93
- cognitive behavioural therapy (CBT), 37, 42  
 aims, 37, 43  
 definition, 37  
 features of good CBT, 38–41

- agenda, 39
- collaboration, 40
- conceptual integration, 41
- eliciting appropriate
  - emotional expression, 40
- eliciting key behaviours, 41
- eliciting key cognitions, 41
- feedback, 39
- guided discovery, 40
- homework, 41
- interpersonal
  - effectiveness, 40
- pacing and efficient use of time, 40
- strategy for change, 41
- techniques for change, 41
- understanding, 40
- key tenet, 60
- primary discussion style, 40
- processes, 38
- cognitive development,
  - Piaget's stages of, 313
- cognitive processing therapy (CPT), 215
- cognitive restructuring, 52–54
  - the basis of, 52–53
  - of core beliefs, 134
- definition, 15
- in dementia caregivers
  - treatment, 372
- devising an alternative belief, 53
- focus of the treatment, 156
- in GAD treatment, 144, 152, 154
  - case studies, 152, 155, 158
- graded exposure and, 100–101, 113
- importance of a clear focus to, 53
- importance of four symptom relief, 4
- level of belief and, 54
- in OCD treatment, 175
- older people and, 281
- in PD treatment, 106, 113
- positive problem orientation and, 59
- in PTSD treatment, 216, 221, 227
- questions to evaluate
  - negative thoughts, 53
- in SAD treatment, 131–132, 133
- the skill of, 113
  - in trauma-focussed therapy, 336
- cognitive therapy, depression
  - case study, 237
- Cognitive Therapy and Research* (journal), 5
- Cognitive Therapy of Depression* (Beck et al.), 5, 232, 344
- Cognitive Therapy Rating Scale (CTRS), 38
- Cognitive Therapy Scale, Revised (CTSr), 39
- Cohen, J. A., 24
- collaboration
  - in agenda setting, 67
  - CBT's focus on, 70
  - as feature of good CBT, 40
  - in formulating origins of psychotic experience, 262–263
  - in goal-setting, 45–47
- common cognitive distortions
  - in CBT, 17–19
- compensatory beliefs and strategies, 14
- competence, development of
  - for therapists, *see under* therapist competence.
- complex PTSD (CPTSD), 214
  - phase-based model of intervention for, 249
- Comprehend, Cope, Connect (CCC) model of psychological therapy, 253, 256
- identification of cycle-breaking interventions, 258
- computerised CBT self-help
  - approaches, 369–374
  - artificial intelligence and, 371
  - benefits of, 370
  - the concept, 369–370
  - costs and contraindications for, 370–371
  - efficacy data for, 370
  - interventions for CYP, 322
  - mobile apps for caregivers, 372
  - technology-based CBT for caregivers, 371–372
- conceptual integration, 41
- continuing professional development (CPD), 377, 378
  - see also* therapist competence,
- Coping Cat, 319
- 'Coping with Caregiving' (US programme), 302
- 'Coping with Depression' (booklet), 369
- Core Belief Worksheet, 235
- core beliefs, definition, 14
- Costello, E., 151
- couples therapy, 353–362
  - cognitive behavioural couples therapy (CBCT), 354
  - couples therapy for depression (CTfD), 354
- defining principles, 354
- development of CBT for, 353–354
- dyadic formulation
  - processes, 355–357
  - DEEP analysis, 356–357
  - Nine areas model, 355
  - polarisation focussing statements, 356
- efficacy data for, 354–355
- integrative behavioural couple therapy (IBCT), 354
- interventions, 358–362
  - behavioural activation, 361
  - building tolerance, 359
  - communication training, 361–362
  - coping methods, 359–360
  - empathic joining, 358
  - problem solving, 360–361
  - sexual problems, 360
  - unified detachment, 358
- traditional behavioural couple's therapy (TBCT), 353–354
- traditional vs integrative approaches, 355
- typical course of treatment, 355
- COVID-19 pandemic,
  - remote delivery CBT and, 364
- Covin, R., 30
- Craske, M. G., 106, 108, 112–117

- crisis services  
 aims of, 248–249  
*see also* acute in-patients CBT.
- Critical Incident Stress  
 Debriefing  
 (CISD), 333
- cue-controlled relaxation,  
 GAD case study, 150
- Cuijpers, P., 27, 29, 30, 146
- cultural adaptations, 277–278
- Curry, V., 315
- Dalglish, T., 13
- Davidson, K. M., 266
- Declarative-Procedural-  
 Reflective (DPR)  
 model of therapist  
 development, 377
- del-Pino-Casado, R., 297
- dementia  
 estimated number of  
 informal  
 caregivers, 295  
 neurological disorders  
 encompassed by the  
 term, 295  
 potential of CBT for  
 treatment of  
 depression and  
 anxiety in, 30  
 prevalence globally, 295
- dementia caregivers,  
 295–306  
 adaptations for, 278  
 assessment interview, 297  
 caregiving burden or  
 strain, 297  
 chronic stress reported  
 by, 297  
 culturally adapted  
 psychoeducational  
 programmes for, 303
- depression and anxiety  
 prevalence, 297  
 risk factors, 297
- effective forms of  
 psychotherapy  
 for, 300
- effectiveness of CBT by  
 delivery format/  
 modality/  
 intensity, 303
- efficacy of CBT for  
 individual  
 caregivers, 302
- evidence-based  
 psychoeducational  
 programmes for,  
 302–303
- generalisations about, 296
- individual and group  
 psychotherapy for  
 caregiver  
 depression, 301
- international support  
 initiatives, 296
- mobile apps for, 372
- popularity of  
 psychoeducational  
 programs for, 299
- psychoeducational programs  
 for caregiver  
 depression, 302–304
- psychological intervention  
 categories, 300
- reluctance to seek help, 300
- stress process model of  
 caregiving, 298
- Dementia Caring Response  
 App (DCRA), 372
- depression, 230–241  
 Beck's differentiation of CT  
 from BT, 4  
 common comorbidities, 231  
 comorbid with body  
 dysmorphic  
 disorder, 198
- definition, 241
- diagnostic and treatment  
 history, 230, 231–232
- diagnostic criteria, 230
- gender perspective, 231
- hopelessness theory of, 16
- Lewinsohn's behavioural  
 model of, 5–7
- and loss of motivation, 57
- measures used in research  
 and clinical  
 practice, 231
- NICE recommendation for  
 CBT treatment, 25
- prevalence, 231
- role of negative automatic  
 thoughts, 15
- case study, 237
- stress generation model of, 14
- stress-diathesis model of, 2
- study into the effective  
 treatment ingredients  
 of CBT for, 6
- treatment, 231–232
- Beck cognitive therapy of  
 depression protocol,  
 233–237
- behavioural therapy,  
 238–241
- case study, 240–241
- cognitive therapy case  
 study, 237
- core beliefs work, 235, 237
- efficacy data for CBT,  
 25–28, 232–233
- intermediate beliefs  
 work, 235
- Martell depression in  
 context protocol,  
 238–241
- most commonly used  
 approaches, 233
- protocol selection, 233
- TRAP and TRAC, 239
- typical age of onset, 231
- DeRubeis, R.J., 26
- Dialogue Guide, 362
- diaphragmatic breathing, 113
- disorder-specific CBT  
 protocols, 85–89
- APA recommendations, 89
- BABCP curriculum  
 guidance, 89
- choosing the right approach,  
 88–89
- efficacy data, 86
- evolution of, 85–86, 89
- NICE recommendations,  
 88–89
- Roth & Pilling at CORE  
 competency  
 framework, 89
- WHO recommendations, 88
- Dobson, K., 26
- Donkin, M., 297
- dual process theory, Power and  
 Dalglish's  
 proposal, 13
- Dugas, M. J., 47, 156–157
- Dunmuir, S., 315
- Early Intervention in Psychosis  
 services, 271
- Eden, J., 296
- effectiveness, difference  
 between efficacy  
 and, 23
- efficacy of CBT, 22
- clinical guidelines  
 recommendations, 22

- Cuijpers et al.'s evidencing of, 27–28  
 difference between efficacy and effectiveness, 23  
 evidence base, 25  
   anxiety disorders, 28–30  
   depression, 25, 28, 232–233  
   older people, 30–31  
 evidence-based practice, 23  
 Fordham et al.'s panoramic meta-analysis of systematic reviews, 22, 28  
   meta-analysis primer, 23–25  
 Ehlers, A., 216  
 Elkin, I., 26  
 Ellis, Albert, 2, 3  
 emotional processing theory, 215  
 empathy, importance for the therapeutic alliance, 70  
 Epictetus, 11  
 evidence-based practice (EBP), requirements, 23  
 exposure for habituation, 54–55  
   anxiety disorders, 93–94  
   exposure with response prevention (ERP), 55  
   hierarchical grading method, 55  
   imaginal exposure, 55  
   similarity to active behavioural experiments, 56  
   *see also* graded exposure.  
 exposure with response prevention (ERP), 55  
   aim of, 178, 179  
   evolution of the process, 174  
   example of, 170  
   in GAD treatment, 144, 153  
   NICE recommendations, 89  
   in OCD treatment, 175–176, 182, 190, 320  
   aims, 178  
   case study, 180  
   primary rationale, 175  
   in OCRD treatment, 170, 171  
   variation from standard exposure, 55  
 exposure work, two types of, 171
- Family Caregiver Distress* (Gallagher-Thompson et al.), 297, 301
- fear  
   development of through classical conditioning, 93  
   Mowrer's two-factor acquisition model, 93, 174  
   Pavlovian extinction of, 54, 56  
 features of good CBT, 38–41  
   agenda, 39  
   collaboration, 40  
   conceptual integration, 41  
   eliciting appropriate emotional expression, 40  
   eliciting key behaviours, 41  
   eliciting key cognitions, 41  
   feedback, 39  
   guided discovery, 40  
   homework, 41  
   interpersonal effectiveness, 40  
   pacing and efficient use of time, 40  
   strategy for change, 41  
   techniques for change, 41  
   understanding, 40  
 feedback  
   as feature of good CBT, 39  
   and therapist competence, 378–379  
 Foa, E. B., 172, 175, 176–177, 215, 217, 218, 219  
 Folkman, S., 298  
 Fordham, B., 22  
 formulation  
   overview, 72–80  
   complete generic formulation, 79  
   function of, 72, 84  
   importance of in CBT, 80  
   important roles served by formulations, 72  
   as key feature of socialisation, 65  
   key questions, 80  
   roles of  
     creating a problem list, 72  
     describing underlying psychological mechanism/key beliefs, 72–74  
     explaining how key beliefs cause and maintain the list of problems, 73–74  
     identifying origins of key beliefs in early life, 75, 77  
     identifying precipitants, 74  
     predicting obstacles to treatment, 77  
     socialisation techniques and, 49  
 Forrest plots in meta-analysis, 24  
 four components model of CBT, 3  
 Freeston, M. H., 142  
 Fuggle, P., 315
- Gallagher-Thompson, D., 302  
 Garber, J., 215  
 generalised anxiety disorder (GAD), 141–164  
   cardinal symptoms, 141  
   case studies  
     AR protocol, 149–150  
     Borkovec protocol, 152–153  
     CT protocol, 161–164  
     IUS protocol, 157–159  
     MAW protocol, 154–155  
   definition, 141–142  
   diagnostic criteria, 142  
   evidence base on the efficacy of CBT for, 29  
   prevalence, 142  
   recent developments in the cognitive theory of GAD, 160–161  
   review of theories and treatments, 142  
   treatment, 143–164  
     applied relaxation, 143  
     AR protocol, 149–150  
     avoidance model of worry, 143–144  
     Borkovec protocol, 150–153  
     CMW protocol, 142  
     cognitive therapy for GAD, 145–146  
     comorbidity and, 146–147

- (GAD) (cont.)  
 comparison of protocols, 156, 164  
 CT protocol, 159–164  
 efficacy data for CBT, 146  
 IUM protocol, 144–145  
 IUS protocol, 156–159  
 MAW protocol, 144, 153–155  
 NICE recommendations, 146, 164  
 protocol selection, 147–148  
 typical therapeutic content, 143  
 worry themes of people diagnosed with, 142  
 gerontology, *see also* older people, 279  
 Gloaguen, V., 26  
 goals of CBT, 43–47  
 aims of CBT, 43  
 benchmarking therapy, 46  
 collaborative goal-setting, 45–47  
 concept of positive goals, 45  
 evidence base on the effects of goal setting, 44  
 features of good goals, 44–45  
 purpose of goal setting, 44  
 SMART goal setting method, 44  
 therapist treatment targets, 47  
 good CBT, features of, *see* features of good CBT.  
 Gortner, E. T., 6  
 Gould, R. L., 30  
 graded exposure  
 anxiety disorders, 94  
 and cognitive elements of treatment, 100–101  
 emetophobia case study, 99–100  
 the process, 100  
 Grimmer, A. G., 356  
 group CBT, 344–350  
 benefits of, 346–347  
 considerations for provision of, 348–350  
 costs and contraindications, 347–348  
 current NICE guidelines, 345  
 development of, 344–345  
 efficacy data for, 345–346  
 patient suitability factors, 347  
 potential additional training requirements, 347  
 problematic member roles, 349  
 stages of group change, 350  
 guided discovery, 54  
 body dysmorphic disorder and, 196  
 as feature of good CBT, 40  
 Gustavsson, S. M., 146  
 Haan, A. de, 320  
 habitual thinking patterns, neuromodulation of, 51  
 habituation, exposure for, *see* exposure for habituation.  
 Haigh, E. A. P., 14  
 Hamilton Rating Scale for Anxiety (HAMA), 29  
 Hammen, C., 4  
 Harper, S. F., 246  
 Haye's Acceptance and Commitment Therapy, 86  
 Health Anxiety Inventory, 205  
 Hinton, L., 278  
 Hirsch, C. R., 142  
 history of CBT, 1  
 Beck's development of Cognitive Therapy, 1–4  
 Clark and Salkovskis's work, 4  
 first RCT conducted, 5  
 importance of behavioural activation for CBT, 5–7  
 Lewinsohn's behavioural model of depression, 5–7  
 recommended by NICE for common mental health conditions, 4  
 scientific evaluation of cognitive therapy, 4–5  
 Hoffman, D. L., 86  
 Hollon, S. D., 26, 215, 345  
 Holmes, E. A., 336  
 homework, as feature of good CBT, 41  
 Hopkinson, M. D., 30  
 hypochondriasis, 204–205, *see also* illness anxiety disorder (IAD)  
 illness anxiety disorder (IAD), 204–210  
 case study, 207–210  
 definition, 204–205  
 maintenance formulation, 206  
 measures for assessment, 205  
 treatment, 206–210  
 behavioural experiments, 207  
 challenges for, 206  
 efficacy data for CBT, 205  
 Salkovskis, Warwick & Deale model, 206  
 vicious flower model, 206, 208, 210  
 imagery, 61–62  
 aphantasia and, 61  
 tools for breaking links with distress, 61  
 imagery exposure, 155  
 imagery exposure, GAD case study, 155, 159  
 imagery rescripting  
 body dysmorphic disorder case study, 199–201  
 the technique, 196  
 imaginal exposure, 55  
 in combination with in-vivo exposure, 100  
 in GAD treatment, 144, 145, 154, 156, 159  
 case study, 153, 155  
 principles of, 100  
 in PTSD treatment, 219  
 reasons for use of, 55  
 in trauma-focussed therapy, 336  
 The Integrated Motivational-Volitional (IMV) Model of Suicide (O'Connor & Kirtley), 251  
 interacting cognitive systems (ICS) theory, application in the PCMH context, 246  
 International Trauma Questionnaire, 215  
 internet delivery of CBT, *see under* remote delivery CBT.

- interpersonal therapy (IPT),  
evaluation of CBT  
against, 26
- intervention, overview, 82–84
- intolerance of uncertainty  
model (IUM),  
144–145
- Jacobson, N. S., 6, 233
- Jacobsson, Edmund, 143
- Janoff-Bulman, R., 215
- Jones, Mary Cover, 28
- Kiddie-Schedule for Affective  
Disorders and  
Schizophrenia  
(K-SADS), 314
- Kishita, N., 30
- Klonsky, E. D., 251
- Knight, B. G., 299
- Kwon, O.Y., 30
- Laidlaw, K., 11, 30, 285
- Lang, P. J., 143
- Larson, E. B., 296
- Lazarus, R. S., 298
- Lejuez, C. W., 6
- Lewin, Kurt, 10
- Lewinsohn, P., 3, 301
- Lewinsohn's behavioural  
model of  
depression, 5–7
- Liebowitz Social Anxiety Scale  
(LSAS), 121
- Liu, Z., 297
- longitudinal model of  
depression, 10
- Losada, A., 302
- 'Managing Social Anxiety'  
(Hope, Heimberg &  
Turk), 124
- Manual of the Revised  
Cognitive Therapy  
Scale, 70
- Marks, I., 98
- Martell, C. R., 6, 239
- mastery of your anxiety and  
worry (MAW), 144,  
153–155
- Mavranzouli, I., 320
- Meichsner, F., 302
- meta-analysis  
Forrest plots, 24  
guidelines for producing  
robust MA, 24–25
- how it works, 23–25
- risk of bias, 24
- metaphor, 51
- example of, 51
- 'The Oxford Guide to  
Metaphors in  
CBT', 51
- Milne, D. L., 379
- mobile apps for caregivers,  
372
- Mobility Inventory, 105
- Monfils, M. H., 336
- monitoring, panic disorder  
therapy, 113
- Montoro-Rodriguez, J., 304
- Morrison, A. P., 262
- Mowrer's two-factor model of  
fear acquisition,  
93, 174
- National Institute for Health  
and Care Excellence  
(NICE)
- recommendation of CBT as  
treatment for a range  
of common mental  
health conditions, 22
- recommendations on  
disorder-specific CBT  
protocols, 88–89
- recommendations on  
treatment for anxiety  
disorders, 25, 93
- recommendations on  
treatment for  
depression, 25
- recommendations on  
treatment for GAD,  
146, 164
- recommendations on  
treatment for OCD,  
172, 176
- recommendations on  
treatment for  
PTSD, 335
- negative automatic thoughts  
(NATs), 11, 13, 54
- addressing through  
therapy, 233
- cognitive restructuring  
of, 235
- the concept of, 15
- depression case study, 237
- exploration of, 134, 235
- parent-led CBT and, 319
- role of in depression, 15
- teaching patients to identify  
and challenge, 133
- negative cognitive triad (NCT),  
16–20
- negative maintenance cycle,  
example of, 17–20
- negativity hypothesis, 16
- neuromodulation, of habitual  
thinking patterns, 51
- NHS Talking Therapies for  
Anxiety and  
Depression, *see also*  
NHSTT, 380
- NHSTT (National Health  
Service Talking  
Therapies), 338–342
- current picture and future  
hopes, 341–342
- data collection and  
performance  
monitoring, 341
- evolution of, 338–339
- international influence, 342
- manual for best practice  
across services, 380
- recovery rates of pilot  
schemes, 339
- service expectations, 341
- stepped care system,  
339–341
- success of pilot sites, 339
- NICE, *see* National Institute for  
Health and Care  
Excellence (NICE).
- Obsessive Compulsive  
Cognition Working  
Group, 175
- obsessive compulsive disorder  
(OCD), 172–190
- age of onset, 173
- behavioural theory of  
anxiety disorders  
applied to, 174
- case studies
- Foa and Kozak protocol,  
178–180
- Salkovskis protocol,  
183–185
- Steketee protocol,  
188–190
- characteristics, 172
- common themes, 173
- comorbidities, 173
- definition, 172–173
- diagnostic criteria, 172–173

- (OCD) (cont.)  
 and ERP, 55  
 evolution of treatment approaches, 172  
 imagery and, 61  
 key themes, 175  
 measures for assessment, monitoring and evaluation, 173  
 and theory A/B technique, 52  
 treatment  
   consensus on focus of, 175  
   efficacy data for CBT, 175–176  
   Foa and Kozak protocol, 176–180  
   focus of, 190  
   NICE recommendation, 172, 176  
   protocol selection, 176  
   Salkovskis's protocol, 176, 180–185  
   Steketee protocol, 176, 186–190  
   vicious flower model, 181, 184
- Obsessive Compulsive Inventory (OCI), 173
- obsessive compulsive or related disorders (OCRD), 170–171  
 behavioural theory applied to, 170–171  
 exposure with response prevention, 170–171  
 internal triggers of distress, 170  
 interventions for CYP, 320
- older people, 279  
 adaptations for, 277  
 age-appropriate approach to CBT, 282–283  
 dementia and pseudodementia, 282  
 and demographic change, 280  
 differences of CBT with, 281–282  
 efficacy of CBT treatment for, 30–31  
 the focus of CBT for, 279–280  
 late life emotional and psychological development, 281  
 late life mental health, 280–281
- life expectancy, 280  
 likelihood of benefiting from CBT, 23  
 positive aspects of ageing, 281  
 positive practice guide for psychological therapists, 280  
 reflection and review, 291–292  
 Sadavoy's 5 Cs of working with, 281–282  
 selective optimisation with compensation (SOC), 282, 290–291  
 suggested Socratic questions, 286  
 and the experience of ageing, 280  
 usefulness of gerontological theories, 280  
 vascular depression hypothesis, 282  
 wisdom enhancement in CBT with older people, 283  
 the concept, 291  
 lifeskills, 283–284  
 timelines, 284–286, 291  
   case studies, 285, 286–289
- operant conditioning, and the negative reinforcement of fear, 93
- Ost, L.-G., 100  
 Ostwald, S. K., 301  
 Oud, M., 316
- Padesky, C. A., 54, 266–267
- panic attacks  
 definition, 104–105  
 prevalence, 105
- panic disorder (PD), 104–117  
 aetiology, 112  
 agoraphobia as common comorbidity, 105  
 assessment measures, 105  
 best performing psychotherapies for, 29  
 Clark and Salkovskis's theory, 106  
 Clark et al.'s treatment model, 29  
 cognitive theory devised, 85
- Craske and Barlow's theory, 106  
 defining, 104–105  
 differentiating panic attacks from, 105, 117  
 early theories of, 105  
 efficacy data for CBT, 106  
 meta-analysis on the efficacy of CBT for, 29  
 prevalence, 105  
 treatment  
   case studies, 110–112, 114–117  
   Clark & Salkovskis protocol, 107–112  
   Craske & Barlow protocol, 112–117  
   protocol selection, 106–107  
   treatment for, 105–106
- Panic Disorder Severity Scale, 105
- Papola, D., 29, 106  
 paranoia, 262–263, 270  
 Patel, A., 278  
 patient feedback, and therapist competence, 378–379
- Patient Health Questionnaire – 9 (PHQ-9), 231
- Pavlovian extinction of fear, 54, 56
- persecutory delusions, 262, 270
- personality disorder CBT, 264–268  
 aims of formulation, 265–266  
 contents of therapy sessions  
   sessions 1–6, 266  
   sessions 6–24, 266–267  
   sessions 24–30, 267–268  
 schema therapy approach, 264
- personality disorders, 244  
 principles for effective engagement, 245
- Persons, J. B., 72
- persons living with dementia (PLWD), 295–297, 299
- phobias, 98  
 categories, 98–99  
 as commonest type of anxiety disorder, 98  
 diagnostic criteria, 99  
 emetophobia case study, 99–100

- graded hierarchy  
 case study, 101–103  
 co-construction, 101  
 history of treatment within CBT, 98, 103  
 principles of psychological treatment for, 100–101
- Piaget, J., 313
- Piagetian stages of cognitive development, 277
- Pilling, S., 89
- Pinquart, M., 299
- Post Traumatic Cognitions Inventory, 215
- post traumatic stress disorder (PTSD), 213–227  
 alternative approach to prevention or maintenance of, 335  
 case study, 225–227  
 clinical guidelines recommending CBT as treatment for, 22  
 comorbidities, 214  
 comparison with ASD, 335  
 complex PTSD, 214  
 defining, 213–215  
 diagnostic criteria, 214  
 explanatory theories, 57  
 growing recognition, 332  
 historical perspective, 213  
 measures for assessment, formulation, and monitoring of, 215
- NICE  
 recommendations, 335
- prevalence, 214
- re-experiencing as cardinal feature, 214
- symptoms, 213–214
- treatment, 215–216  
 consensus on, 216  
 CYP, 320–321  
 efficacy data for CBT, 216–217  
 Ehlers & Clark protocol, 223–227  
 Foa & Rothbaum protocol, 219–221  
 focus of, 227  
 protocol selection, 217–219  
 Resick, Monson & Chard protocol, 221–223
- unhelpful assumptions regarding ‘trauma’ and, 332–333
- post-stroke anxiety, 23
- Power, M. J., 13
- preoccupation  
 as key driver of distress in OCD, 171  
 as unifying feature of anxiety disorders, 170
- primary care, 338–342  
 rationale for a national mental health service, 338–339  
*see also* NHSTT (National Health Service Talking Therapies).
- problem solving, 58–60  
 couples therapy, 360–361  
 defining the problem and goals, 59  
 dementia caregiving case study, 305  
 developing a positive problem orientation, 58–59  
 evaluating and choosing solutions, 59–60  
 generating solutions, 59  
 key processes, 58  
 reviewing and refining solutions, 60
- in therapy for depression, 239
- progressive muscle relaxation, GAD case study, 155
- progressive relaxation training, GAD case study, 149
- prolonged exposure therapy (PET), 215
- psychoeducation, 49–50  
 GAD case study, 154–155, 157, 161
- panic disorder therapy, 112
- relevant topics, 49–50
- tailoring for the patient, 50
- in treatment for OCD, 178
- psychological disorders, DSM-4 definition, 43
- psychological first aid (PFA), trauma and, 334
- psychological wellbeing practitioners (PWP), 340
- psychopathology  
 Beck’s model, 10–12
- the cognitive model of, 11–12
- stress-diathesis model of, 13–15
- Psychopathology Model for CBT (Beck), 15
- psychosis, and therapist beliefs about aetiology, 245
- psychosis and complex mental health (PCMH), 244–271, 342  
 client difficulties in engaging with therapy, 270  
 digital CBT applications, 270  
 discussing diagnosis with clients, 246–247  
 duration of treatment indicated by evidence base, 247  
 importance of expert supervision, 247  
 key underlying principles for cognitive therapy, 245–246
- psychological theories to consider in context, 246
- role of third sector agencies, 244
- sessional planning, 247–248  
*see also* acute in-patients; bipolar disorder; personality disorder; psychosis.
- psychosis CBT, 261–264  
 collaborative formulation, 262–263  
 benefits of, 262  
 contents of therapy sessions  
 sessions 1–6, 263  
 sessions 6–24, 264  
 sessions 24–30, 264  
 enquiry regarding adversity recommended, 261  
 flexibility in formulation, 262  
 formulation, model framework, 262  
 identifying achievable goals, 262
- PTSD Symptom Checklist, 215
- quality assessment measures, comparison, 39

- Rachman, J., 28–29, 174
- randomised controlled trials (RCTs)
- function and process, 25
  - as model of good supervision, 380
  - original RCT evaluating the efficacy of CBT for depression, 25
  - panic disorder, 106
  - and the evidence base for CBT treatment, 25
- rape victims, emotions experienced by, 215
- rational-emotive behavior therapy (REBT), 2
- reciprocal inhibition, 94
- processes, 85
- relapse prevention
- GAD case study, 159
  - in panic disorder therapy, 114
- relaxation techniques, 94
- remote delivery CBT, 364
- benefits and risks, 367
  - considerations when offering remote delivery, 366–367
- telephone therapy, 364–365
- text-based therapy, 365–366
- and the impact of the COVID-19 pandemic, 364
- videoconferencing therapy, 365
- re-scripting, 128, 196
- Resick, P. A., 215, 217
- resilience, as most common response to trauma, 333
- Response to Intrusions Questionnaire, 215
- Responsibility Attitudes Scale (RAS), 173
- reverse role-play, for body dysmorphic disorder, 198
- The Revised Sociocultural Stress and Coping Model for Caregivers, 299
- Robichaud, M., 156–157
- Rogers, Carl, 4
- Rogers, J., 316
- Roth, A. D., 89
- Rubinstein, Arthur, 290
- Rush, A. J., 5
- Russell, Bertrand, 23
- Safety Behaviours Questionnaire, 215
- safety-seeking behaviours
- behavioural experiments designed to address, 147
- body dysmorphic disorder, 187, 195, 197, 199, 202
- examples of, 126
  - function, 95
  - GAD, 155, 160, 161, 162–163
  - IAD, 206–207, 209
  - identifying and resisting the use of, 101
  - and maintenance of exaggerated threat beliefs, 145–146, 160, 223
  - maintenance of social anxiety through, 122
  - OCD, 174, 185, 187, 321, 324
  - and prevention of disconfirming fears, 126
  - PTSD, 225
  - role of worry as, 146, 157
  - uncertainty as trigger for, 157
- Salkovskis, P.M., 4–5, 29, 95, 172, 175, 177
- Santayana, George, 1
- Saunders, R., 23, 30, 282
- ‘Savvy Caregivers’, 302
- schemas
- activation of, 13–14, 17
  - children’s development of their own schemata, 313
  - content formation, 14
  - maladaptive schemata, 14
  - role in processing meaning of events, 14
  - in the Beck model, 12–13
- schizophrenia spectrum disorders, 244
- Schulz, R., 296
- selective optimisation with compensation (SOC), 282, 290–291
- self-control desensitisation techniques, GAD case study, 152
- self-efficacy
- Bandura’s theory of, 298–299
  - dementia caregiving case study, 305
- self-harm, functions of, 251
- self-practice/self-reflection and therapist competence, 380–381
- Seligman, M. E. P., 3
- sexual problems, couples therapy, 360
- Shaw, B. F., 345
- Sheats, P., 349
- skills training, 60–61
- Slo-Mo therapy, efficacy for paranoia and persecutory delusions, 270
- slowed breathing, 113
- SMART (specific, measurable, assignable, realistic and time-related) goal-setting method, 44
- social anxiety disorder, 120–138
- assessment and monitoring measures, 121
  - case studies, 128–129, 134–137
  - central conundrum, 120
  - definition and diagnostic criteria, 120–121, 138
  - efficacy data for CBT, 123
  - efficacy data for CBT, Warnock-Parkes’ video-feedback study, 123
  - median age of onset, 121
  - meta-analysis on the efficacy of CBT for, 29
  - NICE guidance, 123
  - scientific interest in, 120
  - thinking errors, 133
  - treatment, 121–123
    - Clark & Wells protocol, 126–129
    - Clark & Wells’ theory, 122
    - co-morbidity and treatment, 124
    - Hope, Heimberg & Turk protocol, 131–137
    - key differences between theories of, 123
    - protocol selection, 124

- Rapee & Heimberg's theory, 122–123
- Social Attitudes Questionnaire, 121
- Social Behaviour Questionnaire, 121
- Social Cognitions Questionnaire, 121
- Social Phobia Inventory (SPIN), 121
- socialisation to therapy  
formulation as key feature of, 65  
techniques, 49  
theory A/B and, 52
- Socratic questioning, 54  
case study, 111  
in therapy for older people, 284, 286
- Sörensen, S., 299
- specific phobias, 98, *see also* phobias
- Stallard, P., 316, 324
- START (STrategies for RelaTives) program, 303
- State-Trait Anxiety Inventory (STAI), 29
- Steffen, A. M., 302
- Steketee, 172
- Steketee, G., 175, 177
- stepped care system  
benefits and risks, 340  
step 1: assessment and diagnosis, 339  
step 2: low intensity therapy options and medication, 340  
step 3: high-intensity therapy options and medication, 340  
step 4: combined treatment, multidisciplinary care and inpatient services, 340
- stress-diathesis model of depression, 2
- stress-diathesis model of psychopathology, 13–15
- Stroud, C., 296
- structure of CBT, 64–70  
assessment and formulation phases, 64  
intervention phase, 65  
phases of therapy, 64–65
- relapse prevention phase, 65
- sessional structure, 66–70  
commonly used ordering, 66  
mood check (1), 66  
feedback and summary of previous session (2), 67  
agenda setting (3), 67  
homework review (4), 67–68  
sessional work (5), 68–69  
summary and feedback (6), 69  
setting homework (7), 69–70
- socialisation phase, 65
- and the therapeutic alliance, 70
- treatment structure, 64–65
- Subjective Units of Distress Scale (SUDS), 132
- suicidal ideation  
correlation of body dysmorphic disorder with, 198  
role of defeat, entrapment and hopelessness, 251
- systemic desensitisation, 94, 98
- team formulation  
5P's approach recommended for, 260  
benefits of, 259–260  
focus of, 260  
importance of patient involvement, 261  
professional practice guidelines, 260  
recommended structure of meetings, 260–261  
required development time, 260
- Teasdale, J. D., 246
- techniques of CBT, 49–62  
behavioural activation, 57–58  
behavioural experiments, 56  
cognitive restructuring, 52–54  
examples of, 49  
exposure for habituation, 54–55  
guided discovery, 54  
imagery, 61  
intervention techniques, 52
- metaphor, 51
- problem solving, 58–60
- psychoeducation, 49–50
- skills training, 60–61
- socialisation techniques, 49
- theory A/B, 52, 65
- technology-based CBT for caregivers, 371–372
- telephone delivery of CBT, *see under* remote delivery CBT.
- text-based delivery of CBT, *see under* remote delivery CBT.
- 'The Oxford Guide to Metaphors in CBT' (Stott et al.), 51
- theory A/B technique, 52, 65, 182
- body dysmorphic disorder treatment, 196  
case study, 199
- GAD case study, 162, 163
- IAD case study, 208–209
- OCD treatment case study, 184–185
- PD case study, 110
- usefulness for OCRD, 171
- theory of CBT, 10–20  
application in practice, 11–13
- Becks model applied as formulation worksheet, 11
- Beck's Psychopathology Model for CBT, 15
- common cognitive distortions in CBT, 17–19
- content-specificity hypothesis, 17
- covert structure underlying negative thoughts, 16
- information processing errors, 15
- negative cognitive triad, 16–20
- stress-diathesis model, 13–15
- therapeutic alliance  
age and, 283  
behavioural activation and, 6  
building of in CBT for personality disorders, 265

- therapeutic alliance (cont.)  
 creation of in CBT for psychosis, 263  
 importance of empathy for, 70  
 as indicator of success in therapy, 313  
 interpersonal style of the therapist, 70  
 language used in therapy and, 323  
 role in effective CBT, 322  
 SRS as measure of, 315  
 support and maintenance with CYP, 315, 324  
 therapist competence, 376–381  
 the concept of, 376  
 continuing professional development (CPD), 377–378  
 developing competence, 377  
 DPR model of therapist development, 377  
 meta-competence, 376  
 patient feedback and, 378–379  
 self-practice/self-reflection and, 380–381  
 supervision and, 379–380  
 ‘Thinking Good – Feeling Better’ (Stallard), 324  
 thought suppression, OCD case study, 189  
 Tindall, L., 316  
 TRAC (triggers, responses, and alternative coping), 239  
 transdiagnostic CBT, 86  
 TRAP (triggers, responses, and avoidance patterns), 239  
 trauma, 332  
 and active monitoring/watchful waiting, 335  
 benefits and risks of debriefing, 333  
 best practice in early aftermath of, 336  
 definitions, 332  
 comparison of DSM-5 and ICD-11, 213  
 disasters and exposure to, 333  
 early intervention following, 335–336  
 ‘Funnel Approach’ to trauma assessment, 257  
 most common response to, 333  
 organisational support for those at risk of trauma exposure, 334  
 potential impact of traumatic events on a child’s development, 313  
 prevention *vs* early intervention, 333  
 and psychological first aid, 334  
 research areas, current and future, 336  
 TF-CBT for CYP, 320–321  
 unhelpful assumptions around PTSD and, 332–333  
*see also* post traumatic stress disorder (PTSD).  
 Trauma Risk Management (TRiM), 334  
 trauma-focussed cognitive interventions, 216, 333, 335, 336  
 treatment of depression  
 collaborative programme (TDCRP), .US-based evaluation of CBT efficacy for depression, 26  
 tricyclic antidepressant medication, comparison of CBT with, 5  
 trigger, response, avoidance pattern (TRAP), 6  
 Tuckman, B. W., 350  
 two-stage acquisition of fear, 85  
 underlying concepts of CBT, 85  
 vascular depression hypothesis, 282  
 Verduyn, C., 316  
 vicious flower model, 96  
 GAD case study, 161  
 IAD treatment, 206, 208, 210  
 OCD treatment, 181, 184  
 video delivery of CBT, *see under* remote delivery CBT.  
 in vivo exposure  
 in combination with imaginal exposure, 100  
 GAD case study, 314  
 for social anxiety, 147  
 Vygotsky, L. S., 277, 313  
 Werson, A., 22, 30–31  
 Whitfield, G., 347  
 Willhelm, S., 172, 175  
 Wolpe, Joseph, 28, 98, 143  
 Wood, A., 316  
 World Health Organization (WHO), recommendations relating to mental health, 88  
 Yale-Brown Obsessive Compulsive Scale (YBOCS), 173  
 Yalom, I. D., 344  
 Yerkes-Dodson law, 115  
 Zinbarg, R. E., 153  
 zone of proximal development (ZPD), 313