

Index

- alloimmunization, 64, 65
 amniotic fluid, 26
 anesthesia
 EXIT procedure, 181–182, 193–200
 and fetal brain development, 29–30
 fetal cardiac intervention, 111–113
 fetal endoscopic tracheal occlusion (FETO) procedure, 97–98
 for fetal intrauterine transfusion, 72
 lung masses, congenital, 159
 for maternal intrauterine transfusion, 70–71
 multidisciplinary approach to maternal-fetal surgery, 48–49
 myelomeningocele (MMC) surgical management, 147–150
 open fetal surgery, 159–161, 178–181
 selective fetoscopic laser photocoagulation (SFLP), 91, 92
 anesthesia, sacrococcygeal teratoma (SCT)
 background, 175–176
 early delivery with immediate debulking/resection, 182–186
 EXIT procedure, 181–182
 minimally invasive fetal therapy, 176–178
 open fetal surgery, 178–181
 antenatal fetal surveillance (AFS)
 biophysical profile (BPP), 127–128
 contraction stress test (CST), 126–127
 Doppler evaluation of umbilical artery, 129–130
 and fetal blood sampling (FBS), 119
 during fetal surgery, 130–132
 indications for, 119–120
 maternal-fetal movement assessment, 120
 modified biophysical profile (BPP), 128–129
 nonstress test (NST), 126
 objective of, 119
 biophysical profile (BPP)
 for fetal surveillance, 127–128
 modified, 128–129
 blood pressure during pregnancy, 3
 blood volume during pregnancy, 7
 bronchial atresia (BA), 155
 bronchogenic cysts (BC), 154–155
 bronchopulmonary sequestration (BPS), 154
 cardiac output during pregnancy, 2
 cardiovascular system during pregnancy
 anatomic changes, 1
 blood pressure, 3
 cardiac output, 2
 heart rate, 2
 hemodynamic changes
 during labor, 3–4
 pulmonary vascular resistance, 3
 stroke volume, 2–3
 systemic vascular resistance (SVR), 3
 cardiovascular system, fetal, 22–26
 central nervous system during pregnancy, 7
 cesarean section, 58–60
 Cincinnati staging system, 85
 circulation system, fetal, 24
 coagulation during pregnancy (blood), 8
 communication in maternal-fetal surgery, 60
 congenital cystic adenomatoid malformation (CCAM), 153–154
 congenital diaphragmatic hernia (CDH), 95–97
 congenital high airway obstruction syndrome (CHAOS), 155–156
 congenital lobar emphysema (CLE), 155
 consent in maternal-fetal ethics, 39
 contraction stress test (CST), 126–127
 critical titer, 67
 endocrine system during pregnancy, 10–11
 ethics in maternal-fetal surgery
 1982 conference on, 37
 consent, 39
 examples, 41–44
 fetal status paradigms, 39–40
 fetal therapy board, 37–39
 institutional review board standards, 60
 twins, 41
 work-up, 40–41
 ethnicity and prevalence of Rh (D)negative blood type, 65
 ex-utero intrapartum treatment. *see* EXIT procedure
 EXIT procedure
 anesthesia for, 22, 193–200
 congenital lung masses, 162, 193
 defined, 191
 history of, 191
 indications for, 192
 outcomes, 200–203
 procedure, 55–57
 sacrococcygeal teratoma (SCT) anesthesia, 181–182
 surgical steps, 200–203
 fetal access sites
 free loop of cord, 72–73

Index

- fetal access sites (cont.)
 intracardiac transfusion, 73
 intraperitoneal transfusion (IPT), 73
 placental cord insertion, 72
- fetal cardiac disease
 extracardiac anomalies, 105–106
 genetic conditions, 106
 prevalence of, 103
- fetal cardiac intervention
 anesthesia for, 111–113
 complications, 111
 fetal echocardiography, 103–105
 popularity of, 103
- fetal cardiac intervention categories
 aortic valvuloplasty, 107–108
 catheter-based fetal cardiac intervention, 106–107
 generalities, 106
- fetal cardiac intervention surgical approaches
 catheter-based, 108–109
 pulmonary atresia with intact intraventricular septum (PA/IVS), 111
 pulmonary valvuloplasty, 111
 restrictive septum, 108–110
 stent placement, 110
- fetal endoscopic tracheal occlusion (FETO)
 complications, 100
 history of, 97
- fetal endoscopic tracheal occlusion (FETO) indications
 congenital diaphragmatic hernia (CDH), 95–96
 severe pulmonary hypoplasia from ePPROM, 96
- fetal endoscopic tracheal occlusion (FETO) procedure
 maternal and fetal anesthesia, 97–98
 mother remaining close distance for, 60
 postoperative considerations, 98–99
 preoperative considerations, 97
 tracheal occlusion balloon insertion, 98
- tracheal occlusion reversal, 99–100
- fetal growth retardation (FGR), 129–130
- fetal intervention phases
 prenatal assessment, 53
 preoperative preparation, 53–54
 surgical intervention, 54–60
- fetal intervention types, 59
- fetal operating room nurse, 51
- fetal pain, 27–30
- fetal physiology
 effects of drugs on, 21–22
 environment, 17
 placenta, 17–20
 uteroplacental blood flow, 20–21
- fetal physiology by system
 amniotic fluid, 26
 blood pressure, 24
 blood volume, 24
 cardiac development, 22–24
 cardiovascular system, 22–26
 circulation, 24
 pulmonary, 24–26
 renal system, 26
- fetal therapy center, 45–46
- fetal therapy nurse
 coordinator, 50
- fetal therapy review board
 composition of, 38–39
 examples, 41–44
 necessity of, 37–38
- fetal thoracotomy, 161–162
- fetoscopic surgery, 60
- follow-up, maternal-fetal surgery, 61
- gallbladder changes during pregnancy, 10
- gastrointestinal system during pregnancy
 changes, 8–10
 gallbladder, 10
 liver, 10
- genetics, 48
- heart rate during pregnancy, 2
- hematologic system
 blood volume, 7
 coagulation, 8
 platelets, 8
 red blood cells, 7
 white blood cells, 8
- hemodynamic changes during labor, 3–4
- hemodynamic changes during pregnancy, 2
- hemolytic disease of the fetus and newborn (HDFN)
 epidemiology, 64
 etiology, 64
 history of, 64
 hypothyroidism, 10–11
- intraperitoneal transfusion (IPT), 73
- intrauterine transfusion in maternal-fetal surgery
 epidemiology, 64
 etiology, 64, 65
 history of, 64
 outcomes, 77
 patient selection criteria, 65–66
- intrauterine transfusion in maternal-fetal surgery anesthesia
 fetal, 72
 maternal, 70–71
- intrauterine transfusion in maternal-fetal surgery complications
 long-term, 76–77
 procedural, 76
- intrauterine transfusion in maternal-fetal surgery pre-procedure
 blood preparation, 70
 maternal preparation, 70
 setting and staff, 69–70
 timing, 69
- intrauterine transfusion in maternal-fetal surgery procedure
 fetal access site, 72–73
 subsequent procedure timing, 75–76
 technique, 74–75
 volume to transfuse, 73–74
- liver changes during pregnancy, 10
- lung embryology, 153
- lung masses, congenital
 anesthesia for, 159
 bronchial atresia (BA), 155
 bronchogenic cysts (BC), 154–155

Index

- bronchopulmonary sequestration (BPS), 154
- congenital cystic adenomatoid malformation (CCAM), 153–154
- congenital high airway obstruction syndrome (CHAOS), 155–156
- congenital lobar emphysema (CLE), 155
- defined, 152
- diagnosis, 156
- EXIT procedure, 162–163
- fetal thoracotomy, 161–162
- incidence of, 152–153
- open fetal surgery, 159–161
- postnatal treatment, 163
- prognosis, 156–157
- treatment, 158–159
- lung mechanics during pregnancy, 4–5
- maternal-fetal medicine specialists (MFM), 48
- maternal-fetal surgery
 - antenatal fetal surveillance (AFS), 119–132
 - congenital lung masses, 152–163
 - ethical considerations, 37–44
 - EXIT procedure, 191–204
 - fetal cardiac intervention, 103–113
 - fetal endoscopic tracheal occlusion (FETO) procedure, 95–100
 - intrauterine transfusion, 64–77
 - multidisciplinary approach to, 45–54
 - myelomeningocele (MMC), 137–150
 - phases of fetal intervention, 52–60
 - sacrococcygeal teratoma (SCT), 168–187
 - twin anemia polycythemia sequence (TAPS), 89–90
 - twin-twin transfusion syndrome (TTTS), 83–89
- maternal-fetal surgery, day of
 - communication, 60
 - ethical oversight, 60
 - follow-up, 61
 - operating room, 61
 - preparation for, 54–60
 - simulation, 60–61
- minimally invasive fetal therapy, 171–173, 176–178
- multidisciplinary approach to maternal-fetal surgery
 - anesthesia, 48–49
 - cardiology, 49–50
 - clinical team, 46–47
 - fetal therapy center, 45–46
 - genetics, 48
 - imaging, 47
 - maternal fetal medicine specialists (MFM), 48
 - neonatology, 50
 - nursing, 50–52
 - obstetrics, 48
 - psychosocial services, 52
 - surgeons, 47
- musculoskeletal system during pregnancy, 11
- myelomeningocele (MMC)
 - defined, 137
 - diagnosis, 138–140
 - etiology, 138
 - incidence of, 137
 - pathophysiology of, 140–141
 - in-utero repair of, 137–138
- myelomeningocele (MMC)
 - surgical management
 - anesthesia for, 147–150
 - indications for, 141
 - post-surgery, 150
 - procedure, 142–147
 - ultrasound to assess motor function, 142
- neonatal ICU nurse, 52
- neonatal surgical advance practice nurse, 51–52
- neurologic system during pregnancy
 - anatomic changes, 6
 - central nervous system, 7
- nonstress test (NST)
 - accelerations, 122–123
 - baseline, 121, 122
 - categories, 121
 - contraction stress test (CST), 126–127
 - decelerations, 123–124
 - intrapartum fetal monitoring, 124–126
 - prevalence of, 121
 - sinusoidal pattern, 124–125
- vibroacoustic stimulation, 121
- nursing for maternal-fetal surgery
 - fetal operating room nurse, 51
 - fetal therapy nurse coordinator, 50
 - neonatal ICU nurse, 52
 - neonatal surgical advance practice nurse, 51–52
 - obstetric nurse, 51
 - perinatal advance practice nurse, 50–51
- obstetric nurse, 51
- open fetal surgery
 - anesthesia for, 159–161, 178–181
 - background, 159–161
 - post-surgery, 58
 - for sacrococcygeal teratoma (SCT), 173–174
- operating room, maternal-fetal surgery, 61
- pain, fetal, 27–30
- perinatal advance practice nurse, 50–51
- pharmacology, effect on fetus, 21–22
- placenta
 - anatomy of, 18
 - at delivery, 19–20
 - functions of, 17–18
 - pathophysiology of during pregnancy, 18–19
- placenta accreta spectrum, 19–20
- platelets during pregnancy, 8
- pregnancy
 - cardiovascular system, 1–4
 - endocrine system, 10–11
 - gastrointestinal system, 8–10
 - hematologic system, 7
 - musculoskeletal system, 11
 - neurologic system, 6–7
 - renal system, 10
 - respiratory system, 4
- preterm premature rupture of membranes (ePPROM), 96
- psychosocial services, 52
- pulmonary system, fetal, 24–26
- pulmonary vascular resistance during pregnancy, 3

Index

- Quintero staging system, 84–85
- red blood cells during pregnancy, 7
- renal system during pregnancy, 10
- renal system, fetal, 26
- respiratory system during pregnancy
 anatomic changes, 4
 factors affecting obstetric airway, 4
 gas exchange, 6
 lung mechanics, 4–5
- sacrococcygeal teratoma (SCT)
 classification, 168
 defined, 168
 fetal and maternal complications, 169
 management options, 168–169
- sacrococcygeal teratoma (SCT)
 anesthesia
 background, 175–176
 early delivery with immediate debulking/resection, 182–186
 EXIT procedure, 181–182
 minimally invasive fetal therapy, 176–178
 open fetal surgery, 178–181
- sacrococcygeal teratoma (SCT)
 prenatal assessment
 echocardiography, 170–171
 magnetic resonance imaging (MRI), 171
 ultrasound, 170
- sacrococcygeal teratoma (SCT)
 treatment options
 early delivery, 174–175
 minimally invasive fetal therapy, 171–173
 open fetal surgery, 173–174
- selective fetoscopic laser photocoagulation (SFLP)
 anesthesia for, 91
 complications, 93
 local anesthesia versus sedation, 92
 most popular approach, 88
 selective coagulation, 88–89
 ultrasound guidance of, 88
- simulation, maternal-fetal surgery, 60–61
- stroke volume during pregnancy, 2–3
- surgical intervention steps
 cesarean section to immediate neonatal resection, 58–60
 clinical team requirements, 56
 day of, 54
 fetoscopic and ultrasound procedures, 58
 fetoscopic surgery, 60
 intraoperative phase, 55
 open and EXIT procedures, 55–57
- open fetal surgery, 58
 postoperative, 58–60
 preoperative preparation, 54–55
- systemic vascular resistance (SVR) during pregnancy, 3
- thrombocytopenia, 8
- tracheal occlusion balloon insertion, 98
 reversal, 99–100
- twin anemia polycythemia sequence (TAPS), 89–90.
see also twin-twin transfusion syndrome
- twin-twin transfusion syndrome (TTTS). *see also* twin anemia polycythemia sequence
 etiology of, 83–84
 management, 87–89
 pathophysiology of, 83–84
 screening for, 84
 staging systems for, 84–87
- twin-twin transfusion syndrome (TTTS)
 anesthesia
 intraoperative management, 92
 preoperative considerations, 90–92
- white blood cells during pregnancy, 8