

Index

- bracketing
 - Dirichlet–Neumann, 180
- calculus
 - functional, 18
- compactness
 - relative, 57
- conjecture
 - Glaser–Grosse–Martin, 470
 - Lieb–Thirring, 330, 467
 - Pólya, 197
- constant
 - best known
 - Cwikel–Lieb–Rozenblum summary, 466
 - Lieb–Thirring summary, 465, 466
 - comparison between $L_{\gamma,d}^{(1)}$ and $L_{\gamma,d}$, 330, 433, 434
 - comparison between $L_{\gamma,d}^{(1)}$ and $L_{\gamma,d}^{\text{cl}}$, 331
 - comparison between $L_{\gamma,d}^{\text{cl}}$ and $L_{\gamma,d}$, 323, 342, 433
 - one-particle $L_{\gamma,d}^{(1)}$, 325, 464
 - for $\gamma = 0$, 329
 - in dimension one, 326
 - in the dual formulation, 445
 - semiclassical $L_{\gamma,d}^{\text{cl}}$, 167, 169, 323, 464
 - in the dual formulation, 445
 - multiplication property, 219
 - sharp
 - Lieb–Thirring for $\gamma = \frac{1}{2}$ in dimension one for matrix potentials, 456
 - Lieb–Thirring for $\gamma = \frac{1}{2}$ in one dimension, 349
 - Lieb–Thirring for $\gamma = \frac{3}{2}$ in dimension one, 353
 - Lieb–Thirring for $\gamma = \frac{3}{2}$ in dimension one for matrix potentials, 395, 400
- Lieb–Thirring for $\gamma = \frac{3}{2}$ in higher dimension, 395
- Lieb–Thirring summary, 464
- continuity
 - absolute, 71
- convergence of operators
 - norm, 10
 - strong, 10
 - weak, 10
- convergence of vectors
 - norm, 9
 - strong, 9
 - weak, 9
- coordinates
 - spherical, 221
- counting function
 - spectral, 38
- covering, 283
 - multiplicity, 283
- derivative
 - in direction e , 67
 - weak
 - in direction e , 67
- differentiable
 - weakly, 69
- eigenvalue, 13
- eigenvector, 13
- embedding
 - compact, 34, 96
- equation
 - Riccati, 355
 - matrix, 402
- form
 - core, 29
 - quadratic, 29

- closed, 29
- compact, 57
- corresponding to an operator, 30
- lower semibounded, 29
- sesquilinear, 29
- densely defined, 29
- symmetric, 29
- formula
 - IMS localization, 193
 - trace, 363
 - in the matrix case, 412
 - Weyl (Weyl's law), 179, 212–214
 - for the negative eigenvalues of Schrödinger operators, 279, 281, 300
 - on the asymptotics of eigenvalues of the Dirichlet Laplacian, 181, 182, 184–186
 - on the asymptotics of eigenvalues of the Neumann Laplacian, 181, 182, 192, 196
- gradient, 69, 81, 82
- identity
 - polarization, 29
 - resolvent, 27, 56
- inequality
 - Bargmann, 309
 - Berezin, 199, 201, 214
 - for higher powers, 204
 - for lower powers, 205
 - reverse for the Neumann Laplacian, 209
 - Birman–Schwinger, 277
 - Cwikel–Lieb–Rozenblum (CLR), 282, 283
 - on domains, 289
 - optimality of conditions, 296
 - Faber–Krahn, 116, 176
 - Friedlander, 177
 - Friedrichs, 113, 115
 - Gagliardo–Nirenberg, 105, 325
 - sharp constant in one dimension, 106
 - Glaser–Grosse–Martin
 - lower bound on the number of negative eigenvalues in dimension two for radial potentials, 429
 - upper bound on the number of negative eigenvalues in dimension four for radial potentials, 429
 - Hardy
 - in dimension one and two, 138
 - on \mathbb{R}^d , 137, 138
 - on the half line, 128
 - on the half line with weights, 132
 - Hersch, 117, 176
 - Kröger
 - upper bound for eigenvalues of the Neumann Laplacian, 211
 - Kröger, 214
 - upper bound for sums of eigenvalues of the Neumann Laplacian, 212
 - Li–Yau, 201
 - for higher powers, 207
 - for lower powers, 207
 - on individual eigenvalues, 206
 - Lieb–Thirring (LT), 292, 323
 - counterexample of Helffer and Robert, 342
 - duality, 437
 - for $\gamma = 1$ in one dimension for matrix potentials, 457
 - for $\gamma = \frac{1}{2}$ in dimension one for matrix potentials, 456
 - for $\gamma = \frac{1}{2}$ in one dimension, 349
 - for $\gamma = \frac{3}{2}$ in dimension one, 353
 - for $\gamma = \frac{3}{2}$ in dimension one for matrix potentials, 395, 400
 - for $\gamma = \frac{3}{2}$ in higher dimension, 395
 - for antisymmetric functions, 440
 - for the Coulomb Hamiltonian, 337
 - for the harmonic oscillator in higher dimensions, 340
 - for the harmonic oscillator in one dimension, 338
 - for the Pöschl–Teller Hamiltonian, 335
 - optimality of conditions, 296
 - reversed, 303
 - Pólya, 197, 198, 208
 - Poincaré, 119, 123
 - Poincaré–Sobolev, 126
 - Rotfel'd, 51
 - Sobolev, 103
 - sharp constant, 109
 - Kato–Seiler–Simon inequality, 314
 - Laplacian
 - Dirichlet, 170
 - Laplace–Beltrami, 220
 - Neumann, 170
 - weak, 84
 - weak spherical, 84
 - lemma
 - Glazman on the spectral counting function, 38, 39
 - measure

- Jordan, 183
- projection-valued, 17
- spectral, 22
- measure space
 - separable, 16
- method
 - commutation, 251
- monotonicity
 - domain, 175
 - semiclassical, 418
- multiplicity
 - geometric, 13
- operator
 - absolute value, 36
 - adjoint, 9, 12
 - Birman–Schwinger, 63, 275, 276, 350
 - bounded, 9
 - closed, 12
 - compact, 9
 - continuous, 9
 - corresponding to a quadratic form, 31
 - densely defined, 12
 - embedding, 34
 - kernel $\ker T$, 12
 - Laplace
 - with Dirichlet boundary conditions, 170
 - with Neumann boundary conditions, 170
 - Laplace–Beltrami on \mathbb{S}^{d-1} , 220
 - lower semibounded, 24, 30
 - multiplication, 16
 - non-negative, 24
 - polar decomposition, 36
 - range $\text{ran } T$, 12
 - Schrödinger, 247
 - Coulomb potential, 258
 - definition, 247
 - harmonic oscillator, 251
 - Pöschl–Teller potentials, 256
 - with matrix potential, 388
 - self-adjoint, 15
 - symmetric, 15
 - unbounded, 12
 - with discrete spectrum, 34
- operators
 - T^*T and TT^* , 36
 - comparison of, 41
- orthogonal sum
 - of Hilbert spaces, 13
 - of operators, 13
 - of quadratic forms, 35
- perturbation
- of quadratic forms, 54
- formula for resolvent difference, 56
- polycubes, 182
- principle
 - Birman–Schwinger, 60, 62, 65
 - for Schrödinger operators, 275, 276
 - Courant–Fischer–Weyl
 - max-min, 41
 - min-max, 40
 - duality, 202
 - lifting in dimension, 396, 452
 - variational, 38–41
 - for sums of eigenvalues, 44, 45
- product
 - Blaschke, 370
 - inner, 8
- projection
 - orthogonal, 17
- property
 - extension, 99, 146, 192
 - above graph of a Lipschitz function, 149
 - half space, 147
 - uniform Lipschitz continuous boundary, 152
 - zero boundary measure, 155
- rearrangement
 - non-increasing, 132
 - symmetric decreasing, 109
- resolvent, 13
- Riesz mean, 49, 199, 204
- rule
 - chain, 72
 - product, 75
- sequence
 - singular, 26
- set
 - open
 - connected, 70
 - resolvent, 12
 - tiling, 197
- sharp lower bound
 - m_A of an operator, 24, 30
 - m_a of a quadratic form, 29, 33
- space
 - Hilbert, 9
 - phase, 212
 - separable, 9
 - Sobolev
 - $H^1(\Omega)$, 85
 - $H^1(\mathbb{R}^d)$, 92
 - $H^1(\mathbb{S}^{d-1})$, 219

- $H_0^1(\Omega)$, 90
 $\dot{H}^1(\mathbb{R}^d)$, 142
 $\tilde{H}^1(\mathbb{R}^d)$, 145
- spectrum, 13
discrete, 26, 34
essential, 26
of the Coulomb Hamiltonian, 258
of the Dirichlet Laplacian
on a ball, 232
of the Dirichlet Laplacian on a cube, 172
of the harmonic oscillator, 251
of the Laplace–Beltrami operator on \mathbb{S}^{d-1} , 221
of the Laplacian on \mathbb{R}^d , 172
of the Neumann Laplacian
on a ball, 230
of the Neumann Laplacian on a cube, 172
of the Pöschl–Teller Hamiltonian, 256
of the Schrödinger operator
basic properties, 265
discreteness of the negative spectrum, 266, 267
finiteness of the negative spectrum, 268–271, 273
point, 13
- theorem
Aizenman–Lieb on the monotonicity of $L_{\gamma,d}/L_{\gamma,d}^{\text{cl}}$ in γ , 323
- Besicovitch on coverings, 284
fundamental of calculus, 71
Glaser–Grosse–Martin
on counting negative eigenvalues in case of radial potentials, 427
Meyers–Serrin on $H = W$, 86
on the monotonicity of $L_{\gamma,d}^{(1)}/L_{\gamma,d}^{\text{cl}}$ in the dimension, 381
on the monotonicity of $L_{\gamma,d}^{(\text{mat})}/L_{\gamma,d}^{\text{cl}}$ in the dimension, 456
Rellich on compactness of
 $H^1(\Omega) \hookrightarrow L^2(\Omega)$, 100
 $H_0^1(\Omega) \hookrightarrow L^2(\Omega)$, 97
 $H_d^1(R) \hookrightarrow L_d^2(R)$, 101
Riesz on the representation of linear functionals in a Hilbert space, 9
spectral, 20
for compact operators, 28
spectral mapping, 24
Stubbe
on the monotonicity of $\hbar^d \operatorname{Tr} H(\hbar)^\gamma$ for $\gamma \geq 2$, 418
Weyl on the stability of the essential spectrum, 27
form version, 58
- trace, 45