

## TRANSCENDENTAL NUMBER THEORY

First published in 1975, this classic book gives a systematic account of transcendental number theory, that is, the theory of those numbers that cannot be expressed as the roots of algebraic equations having rational coefficients. Their study has developed into a fertile and extensive theory, which continues to see rapid progress today. Expositions are presented of theories relating to linear forms in the logarithms of algebraic numbers, of Schmidt's generalization of the Thue–Siegel–Roth theorem, of Shidlovsky's work on Siegel's *E*-functions and of Sprindžuk's solution to the Mahler conjecture.

This edition includes an introduction written by David Masser describing Baker's achievement, surveying the content of each chapter and explaining the main argument of Baker's method in broad strokes. A new afterword lists recent developments related to Baker's work.

**Alan Baker** was one of the leading British mathematicians of the past century. He took great strides in number theory by, among other achievements, obtaining a vast generalization of the Gelfond–Schneider Theorem and using it to give effective solutions to a large class of Diophantine problems. This work kicked off a new era in transcendental number theory and won Baker the Fields Medal in 1970.

**David Masser** is Professor Emeritus in the Department of Mathematics and Computer Science at the University of Basel. He is a leading researcher in transcendence methods and applications and helped correct the proofs of the original edition of *Transcendental Number Theory* as Baker's student.

CAMBRIDGE  
MATHEMATICAL LIBRARY

---

Cambridge University Press has a long and honourable history of publishing in mathematics and counts many classics of the mathematical literature within its list. Some of these titles have been out of print for many years now and yet the methods which they espouse are still of considerable relevance today.

The *Cambridge Mathematical Library* provides an inexpensive edition of these titles in a durable paperback format and at a price that will make the books attractive to individuals wishing to add them to their own personal libraries. Certain volumes in the series have a foreword, written by a leading expert in the subject, which places the title in its historical and mathematical context.

A complete list of books in the series can be found at [www.cambridge.org/mathematics](http://www.cambridge.org/mathematics). Recent titles include the following:

*Attractors for Semigroups and Evolution Equations*  
OLGA A. LADYZHENSKAYA

*Fourier Analysis*  
T. W. KÖRNER

*Transcendental Number Theory*  
ALAN BAKER

*An Introduction to Symbolic Dynamics and Coding (Second Edition)*  
DOUGLAS LIND & BRIAN MARCUS

*Reversibility and Stochastic Networks*  
F. P. KELLY

*The Geometry of Moduli Spaces of Sheaves (Second Edition)*  
DANIEL HUYBRECHTS & MANFRED LEHN

*Smooth Compactifications of Locally Symmetric Varieties (Second Edition)*  
AVNER ASH, DAVID MUMFORD, MICHAEL RAPOPORT &  
YUNG-SHENG TAI

*Markov Chains and Stochastic Stability (Second Edition)*  
SEAN MEYN & RICHARD L. TWEEDIE

TRANSCENDENTAL  
NUMBER THEORY

ALAN BAKER F.R.S.

*With an Introduction by*

DAVID MASSER  
*Universität Basel, Switzerland*



CAMBRIDGE  
UNIVERSITY PRESS

Cambridge University Press & Assessment  
978-1-009-22994-4 – Transcendental Number Theory  
Alan Baker, Foreword by David Masser  
Frontmatter  
[More Information](#)

CAMBRIDGE  
UNIVERSITY PRESS

University Printing House, Cambridge CB2 8BS, United Kingdom

One Liberty Plaza, 20th Floor, New York, NY 10006, USA

477 Williamstown Road, Port Melbourne, VIC 3207, Australia

314–321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre, New Delhi – 110025, India

103 Penang Road, #05–06/07, Visioncrest Commercial, Singapore 238467

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning, and research at the highest international levels of excellence.

[www.cambridge.org](http://www.cambridge.org)

Information on this title: [www.cambridge.org/9781009229944](http://www.cambridge.org/9781009229944)

DOI: 10.1017/9781009229937

© Cambridge University Press 1975

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 1975

Reprinted with additional material 1979

Reissued as a paperback with updated material in the Cambridge Mathematical Library series 1990

Reprinted with introduction 2022

*A catalogue record for this publication is available from the British Library.*

ISBN 978-1-009-22994-4 Paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

## Contents

---

	<i>Introduction by David Masser</i>	page ix
	<i>Preface</i>	xiii
<b>1</b>	<b>The origins</b>	
1	Liouville's theorem	1
2	Transcendence of $e$	3
3	Lindemann's theorem	6
<b>2</b>	<b>Linear forms in logarithms</b>	
1	Introduction	9
2	Corollaries	11
3	Notation	12
4	The auxiliary function	13
5	Proof of main theorem	20
<b>3</b>	<b>Lower bounds for linear forms</b>	
1	Introduction	22
2	Preliminaries	24
3	The auxiliary function	28
4	Proof of main theorem	34
<b>4</b>	<b>Diophantine equations</b>	
1	Introduction	36
2	The Thue equation	38
3	The hyperelliptic equation	40
4	Curves of genus 1	43
5	Quantitative bounds	44

<b>5</b>	<b>Class numbers of imaginary quadratic fields</b>	
1	Introduction	47
2	$L$ -functions	48
3	Limit formula	50
4	Class number 1	51
5	Class number 2	52
<b>6</b>	<b>Elliptic functions</b>	
1	Introduction	55
2	Corollaries	56
3	Linear equations	58
4	The auxiliary function	58
5	Proof of main theorem	60
6	Periods and quasi-periods	61
<b>7</b>	<b>Rational approximations to algebraic numbers</b>	
1	Introduction	66
2	Wronskians	69
3	The index	69
4	A combinatorial lemma	73
5	Grids	74
6	The auxiliary polynomial	75
7	Successive minima	76
8	Comparison of minima	79
9	Exterior algebra	81
10	Proof of main theorem	82
<b>8</b>	<b>Mahler's classification</b>	
1	Introduction	85
2	$A$ -numbers	87
3	Algebraic dependence	88
4	Heights of polynomials	89
5	$S$ -numbers	90
6	$U$ -numbers	90
7	$T$ -numbers	92
<b>9</b>	<b>Metrical theory</b>	
1	Introduction	95
2	Zeros of polynomials	96
3	Null sets	98

## CONTENTS

vii

4	Intersections of intervals	99
5	Proof of main theorem	100
<b>10</b>	<b>The exponential function</b>	
1	Introduction	103
2	Fundamental polynomials	104
3	Proof of main theorem	108
<b>11</b>	<b>The Siegel-Shidlovsky theorems</b>	
1	Introduction	109
2	Basic construction	111
3	Further lemmas	114
4	Proof of main theorem	115
<b>12</b>	<b>Algebraic independence</b>	
1	Introduction	118
2	Exponential polynomials	120
3	Heights	122
4	Algebraic criterion	124
5	Main arguments	125
	<i>Bibliography</i>	129
	<i>Original papers</i>	130
	<i>Further publications</i>	145
	<i>New developments</i>	155
	<i>Some developments since 1990 by David Masser</i>	162
	<i>Index</i>	166

Cambridge University Press & Assessment  
978-1-009-22994-4 – Transcendental Number Theory  
Alan Baker , Foreword by David Masser  
Frontmatter  
[More Information](#)

---



## Introduction

David Masser

On the first page of the Bibliography are listed earlier works about some of the topics treated in this monograph. In particular the books of Gelfond, Schneider and Siegel are universally regarded as milestones in the development of the theory of transcendental numbers. Each book was based largely on the author's own breakthroughs.

The present monograph represented a similar milestone. Chapters 2, 3, 4, 5, 9, 10, and to a lesser extent Chapters 6, 8, cover material due to the author Alan Baker. This material and Baker's own further developments of it earned him a Fields Medal in 1970.

Of course it is the material in Chapter 2 that constitutes the heart of his achievement. This is explained in the first two pages with a characteristic brevity and modesty. Here we wish to complement this with the following less brief and modest account.

The essential ideas can be conveyed through the special case of his Theorem 2.1 for  $n = 2, 3$ , even ignoring the extra 1 that appears there.

We start with  $n = 2$ . It amounts to the impossibility of

$$\beta \log \alpha = \log \alpha' \tag{1}$$

for  $\alpha, \alpha'$  non-zero algebraic numbers and  $\beta$  irrational algebraic. Of course this is the Gelfond–Schneider Theorem of 1934. It also follows from Theorem 6.1 of Chapter 6, and we proceed to sketch the argument.

We assume (1) and we will obtain a contradiction. Following Gelfond we construct a non-zero polynomial  $F$ , say in  $\mathbf{Z}[x, y]$ , such that

$$f(z) = F(e^z, e^{\beta z}) \tag{2}$$

has many zeroes. More precisely we need the derivatives

$$f^{(t)}(s \log \alpha) = 0 \tag{3}$$

for a certain range of integers

$$t < T, \quad 1 \leq s \leq S \tag{4}$$

with  $T, S$  integers to be suitably chosen later. Thus we have zeroes at  $\log \alpha, \dots, S \log \alpha$  and moreover of multiplicity at least  $T$ . The point is that the functions  $e^z, e^{\beta z}$  in (2) take the values

$$e^{\log \alpha} = \alpha, \quad e^{\beta \log \alpha} = e^{\log \alpha'} = \alpha'$$

at say  $z = \log \alpha$ ; and these are algebraic numbers. Similarly at  $s \log \alpha$  and with multiplicities. Thus the conditions (3) are homogeneous linear equations in the coefficients of  $F$ . Under appropriate assumptions relating  $T, S$  to the degree of  $F$ , these can be solved non-trivially; and using things like Lemma 1 of Chapter 2 or Lemma 1 of Chapter 6 one can make sure that the resulting coefficients are not too large.

Next, Gelfond used analytic techniques to show that the values  $f^{(t)}(s \log \alpha)$  are very small on a range larger than (3). Compare (8) of Chapter 2 and the use of Cauchy’s Theorem in section 5 of Chapter 6. These values are still algebraic numbers, and then arithmetic techniques show that they are in fact zero. Compare Lemma 3 of Chapter 2 or Lemma 3 of Chapter 6 (nowadays one tends to use heights, with a definition slightly different from that of Chapter 1).

Together these lead to (3) for the new range

$$t < 2T, \quad 1 \leq s \leq 2S \tag{5}$$

slightly larger than (4) (this is not quite consistent with Baker’s remark near the end of section 1 of Chapter 2 or the method of section 5 of Chapter 6, but it simplifies the proof a little).

And now the step from (4) to (5) can be iterated, and even indefinitely. This provides infinite multiplicities, and so this  $f$  must be identically zero. Taking into account the irrationality of  $\beta$ , we see that this implies that  $F$  is also identically zero; our required contradiction.

Next for  $n = 3$  we have to reach a similar contradiction from

$$\beta_1 \log \alpha_1 + \beta_2 \log \alpha_2 = \log \alpha' \tag{6}$$

instead of (1), where  $\alpha_1, \alpha_2, \alpha'$  are non-zero algebraic and  $\beta_1, \beta_2$  are algebraic, this time with  $1, \beta_1, \beta_2$  linearly independent over  $\mathbf{Q}$ . Even this was a new result.

Baker’s first step looks natural: to construct now

$$f(z_1, z_2) = F(e^{z_1}, e^{z_2}, e^{\beta_1 z_1 + \beta_2 z_2}) \tag{7}$$

INTRODUCTION

instead of (2) with many zeroes; but no-one had written this down before. Still less had anyone considered multiplicities, now defined by the partial derivatives

$$\left(\frac{\partial}{\partial z_1}\right)^{t_1} \left(\frac{\partial}{\partial z_2}\right)^{t_2} f(s \log \alpha_1, s \log \alpha_2) = 0 \tag{8}$$

instead of (3). Note that there is no  $(s_1 \log \alpha_1, s_2 \log \alpha_2)$  here, because we do not have the Cartesian product situation for  $\mathbf{C}^2$  mentioned in section 1 of Chapter 2. In fact our  $(s \log \alpha_1, s \log \alpha_2)$  lie on a complex line in  $\mathbf{C}^2$ .

Baker took a range

$$t_1 + t_2 < T, \quad 1 \leq s \leq S \tag{9}$$

instead of (4), and the problem is then to increase this as in (5).

Now the experts know that the world of two complex variables is very different from that of a single variable. Possibly Baker did not know this. Anyway, to this day no-one knows how to reach  $t_1 + t_2 < 2T$  as in (5).

He probably started by reducing to  $\mathbf{C}$  via

$$g(z) = f(z \log \alpha_1, z \log \alpha_2). \tag{10}$$

Then we deduce

$$g^{(t)}(s) = 0 \quad (t < T, \quad 1 \leq s \leq S).$$

The twin analytic-arithmetic argument then shows that  $g(s) = 0$  for a larger range of  $s$ , that is,  $f(s \log \alpha_1, s \log \alpha_2) = 0$ . However, as it stands we cannot deduce even  $g'(s) = 0$  because differentiation in (10) introduces transcendental numbers, so we cannot get at, say,

$$\left(\frac{\partial}{\partial z_1}\right) f(s \log \alpha_1, s \log \alpha_2) \tag{11}$$

in this way.

Now that we have set up the scene, Baker’s solution to this problem may seem in retrospect obvious: we use (10) with  $f$  replaced by  $(\partial/\partial z_1)f$ . For the new  $g$  we get almost (9), but now only for  $t < T - 1$ . However this tiny loss does not affect the argument, and we find indeed that (11) vanishes on the larger range of  $s$ .

And what about higher derivatives? To get at some

$$\left(\frac{\partial}{\partial z_1}\right)^{\tau_1} \left(\frac{\partial}{\partial z_2}\right)^{\tau_2} f(s \log \alpha_1, s \log \alpha_2)$$

we use (10) with  $f$  replaced by  $(\partial/\partial z_1)^{\tau_1} (\partial/\partial z_2)^{\tau_2} f$ . We then get (9) for  $t < T - \tau_1 - \tau_2$ . If we aim for all  $\tau_1, \tau_2$  just with  $\tau_1 + \tau_2 < T$ , then hardly

anything remains of the multiplicity; so it is wiser to restrict to say  $\tau_1 + \tau_2 < T/2$ , thus securing  $t < T/2$ . Now the loss is less tiny, but still acceptable.

We end up with (8) on the range, say,

$$t_1 + t_2 < T/2, \quad 1 \leq s \leq 8S. \quad (12)$$

As (9) is about  $T^2S/2$  conditions and (12) is about  $T^2S$  conditions, we do actually have more zeroes.

But now another problem arises: we cannot iterate indefinitely the step from (9) to (12).

In fact a related problem had turned up before Baker; for example when trying to show that the two sides of (1) cannot even be too near to each other. And indeed Baker was able to extend the classical methods; in this case the argument of section 5 of Chapter 2 amounts to the use of a non-vanishing Vandermonde determinant.

All this extends to  $n$  logarithms, and then to include 1 as in Theorem 2.1. In our notation assuming

$$\beta_0 + \beta_1 \log \alpha_1 + \cdots + \beta_{n-1} \log \alpha_{n-1} = \log \alpha', \quad (13)$$

we have to use

$$f(z_0, z_1, \dots, z_{n-1}) = F(z_0, e^{z_1}, \dots, e^{z_{n-1}}, e^{\beta_0 z_0 + \beta_1 z_1 + \cdots + \beta_{n-1} z_{n-1}})$$

in place of (7) at  $(s, s \log \alpha_1, \dots, s \log \alpha_{n-1})$  – compare Lemma 2 of Chapter 2.

This completes our account of Chapter 2. On the way, we have mentioned the problem of approximate versions of (1), and the corresponding generalizations to (6) and (13) are treated in Chapter 3. It is these that are needed for the applications in Chapters 4 and 5.

It is these applications, to Diophantine equations and class numbers, that were the most spectacular of his achievements. It is enough here to cite the first ever upper bounds for the solutions of Mordell's equation  $y^2 = x^3 + k$  with a history going back to 1621, and the verification of Gauss's conjectures from 1801 about imaginary quadratic fields with class numbers  $h = 1$  and  $h = 2$ .

But one should not overlook the less spectacular material in Chapter 6, whose subsequent developments (by others) will be described in the afterword.

## Preface

---

*Fermat, Euler, Lagrange, Legendre . . . introitum ad penetralia huius divinae scientiae aperuerunt, quantisque divitiis abundant patefecerunt*

*Gauss, Disquisitiones Arithmeticae*

The study of transcendental numbers, springing from such diverse sources as the ancient Greek question concerning the squaring of the circle, the rudimentary researches of Liouville and Cantor, Hermite's investigations on the exponential function and the seventh of Hilbert's famous list of 23 problems, has now developed into a fertile and extensive theory, enriching widespread branches of mathematics; and the time has seemed opportune to prepare a systematic treatise. My aim has been to provide a comprehensive account of the recent major discoveries in the field; the text includes, more especially, expositions of the latest theories relating to linear forms in the logarithms of algebraic numbers, of Schmidt's generalization of the Thue–Siegel–Roth theorem, of Shidlovsky's work on Siegel's  $E$ -functions and of Sprindžuk's solution to the Mahler conjecture. Classical aspects of the subject are discussed in the course of the narrative; in particular, to facilitate the acquisition of a true historical perspective, a survey of the theory as it existed at about the turn of the century is given at the beginning. Proofs in the subject tend, as will be appreciated, to be long and intricate, and thus it has been necessary to select for detailed treatment only the most fundamental results; moreover, generally speaking, emphasis has been placed on arguments which have led to the strongest propositions known to date or have yielded the widest application. Nevertheless, it is hoped that adequate references have been included to associated works.

Notwithstanding its long history, it will be apparent that the theory of transcendental numbers bears a youthful countenance. Many topics would

certainly benefit by deeper studies and several famous longstanding problems remain open. As examples, one need mention only the celebrated conjectures concerning the algebraic independence of  $e$  and  $\pi$  and the transcendence of Euler's constant  $\gamma$ , the solution to either of which would represent a major advance. If this book should play some small rôle in promoting future progress, the author will be well satisfied.

The text has arisen from numerous lectures delivered in Cambridge, America and elsewhere, and it has also formed the substance of an Adams Prize essay.

I am grateful to Dr D. W. Masser for his kind assistance in checking the proofs, and also to the Cambridge University Press for the care they have taken with the printing.

*Cambridge, 1974*

A.B.