

Index

- Abilify *see* aripiprazole
- abuse
 children and adolescents 211
 older adults 227
 substance *see* substance abuse
- accountable care organizations 295
- acculturation 277
- acute diversion units 16
- adolescents *see* children/adolescents
- AEIOU-TIPS assessment tool 309
- African Americans 274
- aftercare plan 12
- against medical advice, leaving
 see leaving against medical advice
- age
 agitated patients 158
 of onset 1
 and suicide risk 60
- aggressive/violent behavior
 170–175, 317
 armed patients 174
 assessment 206, 309
 children and adolescents 211
 proactive aggression 212
 reactive aggression 212
 de-escalation 171
 see also verbal de-escalation
 homeless persons 245
 interpersonal interventions 171
 legal issues 175
 medical illness as cause of 170
 pharmacologic interventions 172, 173
 physical restraint 173
 pregnant patients 260
 rural communities 283
 screening for 171
 show of force 173
 signs of 171
 treatment 206
 weapons screening 174
 see also behavioral disturbance
- agitation 89, 136, 155, 197, 317
 approach to patient 156, 157
 causes 155, 156
 definition 155
 differential diagnosis 198
 escalating 155
 length of hospital stay 168
 offering psychiatric medication 161
 older adults 167, 225
 pregnant patients 167, 260, 261
 restraint *see* chemical restraint;
 physical restraint
 safety of staff 157
 special populations 167
 children and adolescents 167
 older adults 167
 pregnant women 167
 treatment 164–168, 187, 201, 261
 antipsychotics 164, 165
 benzodiazepines 166
 goals of 155
 ketamine 166
 routes of administration 166
 rural communities 286
 specific interventions 168
 verbal de-escalation 155–162
- agoraphobia 81
- aguantarse* 274
- air conditioning 357
- akathisia 165
- Alaska Natives 277
- alcohol abuse 6, 37
 binge-drinking 37
 blood alcohol level (BAL) 37
 CAGE screen 227
 community impact 37
 emergency treatment 152
 homeless persons 245, 247
 laboratory findings 38
 leaving against medical advice 331
 management 37
 prevalence 37
 psychiatric symptoms 199
 and schizophrenia 150
 screening for 134
 suicide risk 36
- alcohol testing 33–35
- alcohol withdrawal 46
 delirium tremens 47, 91
 hallucinations 91
 patient disposition 48
 treatment 47
- alexithymia 30
- alopecia, eating disorders 145
- alprazolam
 agitation 166
 anxiety 78
 teratogenesis 264
- Alzheimer's disease 120
 age of onset 2
 depression in 221
 psychosis 91
 see also dementia
- amantadine, neuroleptic
 malignant syndrome 194
- American College of Emergency Physicians 34
- American Indians 277
- Americans with Disabilities Act 366
- amitriptyline
 metabolism 182
 side effects 135
 teratogenesis 262
- amnesia, post-traumatic 252
- amoxapine 183
- amphetamine
 psychosis 93
 toxicity 33
 urine testing 34
- amyl nitrite
 abuse 42
 sudden sniffing death syndrome 42
- angel's trumpet 92
- anhedonia 53
- anorexia nervosa 7, 140
- antibiotics, and mood disorders 57
- anticholinergics
 delirium 22, 90
 hallucinations 91
- anticonvulsants
 side effects 135
 mood disorders 57
 psychosis 92
 see also specific drugs
- antidepressants 182
 heterocyclic 183
 monoamine oxidase inhibitors
 see monoamine oxidase inhibitors
 somatoform disorders 73
 tricyclic *see* tricyclic antidepressants
 see also specific drugs
- antihypertensives, and mood disorders 57
- antipsychotics 58, 96, 98, 184
 agitation 164, 165
 atypical 185, 187
 potency 191
 rapidly acting 207
 delirium 120
 excited delirium syndrome 127, 128
 side effects 135, 185
 anticholinergic 165
 movement disorders 165, 172
 neuroleptic malignant syndrome 166, 183, 190–195, 207
 QT prolongation 187
 teratogenesis 95, 264
 typical 184
 potency 191
 rapidly acting 207
 see also specific drugs
- antisocial personality disorder 8
- anxiety disorders 3–5, 76–82, 197, 199
 agitated patients 161
 causes 77
 children and adolescents 215
 definition and diagnosis 76
 differential diagnosis 77, 80, 198
 DSM-IV criteria 77, 79
 evaluation 78, 81
 fight or flight response 76
 GAD *see* generalized anxiety disorder
 homeless persons 246
 ICD-10 criteria 77, 79
 panic disorder 80
 phobia 4, 80
 postconcussive 255
 pregnant patients 264
 prevalence 1, 4
 PTSD 1, 80
 social phobia 80
 symptoms 77
 treatment 78, 81
 see also specific disorders
- arguing with patients 160
- aripiprazole (Abilify) 58, 185, 186
 aggressive/violent behavior 173
 agitation 165, 201
 dose 201
 children and adolescents 213
 potency 191
 armed patients 174
 suicide risk 61
 see also weapons
- asenapine 186
- Asian Americans 274
- aspartate amino transferase (AST) 38
- assertive community treatment teams 292, 301
- assessment 308, 309
 aggressive/violent behavior 206, 309
 concussion 251

Index

- assessment (cont.)
 disaster victims 230
 factitious illness 114
 homeless persons 245, 247
 in-home 285
 malingering 114
 medical 314, 316
 psychiatric 315
 psychosis 206
 safety 314
 somatoform disorders 71
 suicidal ideation 309, 316
see also risk assessment; and specific conditions
 Atarax *see* hydroxyzine
 Ativan *see* lorazepam
 atomoxetine 93
 attention-deficit/hyperactivity disorder 1, 215
 attitude of clinician 157
 atypical antipsychotics 185, 187
 potency 191
 rapidly acting 207
see also specific drugs
 auditory hallucinations 28–30, 88, 96
 Australasian Triage Scale (ATS) 315
- baclofen withdrawal 92
 balance disorders,
 postconcussive 256
 Balance Error Scoring System (BESS) 253, 256
 Banana Cream Nuke 92
 barbiturates
 abuse 39
 intoxication 152
 withdrawal 48
Bartonella quintana 245
 “bath salts” 40, 92
 bathrooms 358
 battle fatigue 83
 behavior control
 restraint *see* restraint
 seclusion 180
 Behavioral Activity Rating Scale 314
 behavioral disturbance
 aggression *see* aggressive/
 violent behavior
 consequences of 159
 out of control children/
 adolescents 211, 212
 postconcussive 254
 SMI-SUD 151
see also aggressive/violent
 behavior
 behavioral health clients 350
 benzodiazepines
 abuse 38
 aggressive/violent behavior
 172, 173
 agitation 166
 anxiety disorders 78, 265
 excited delirium syndrome 127
 intoxication 152, 199
 neuroleptic malignant
 syndrome 194
 psychosis 96, 98, 207
 teratogenesis 95, 264
 urine testing 34
 withdrawal 48, 91, 199
see also specific drugs
 benztropine (Cogentin) 172
 bereavement 238
 complicated grief 239
 uncomplicated grief 239
 best practice 335–345
 emergency psychiatric
 evaluation 338, 340
 medical clearance 337
 medications 338
 protocols 335
 space 335, 340
 staff 335, 339
 qualifications 341
 training 341
 substance abuse 337
 throughput 339
 triage 336
 binge drinking 37
see also alcohol abuse
 binge eating disorder 7, 140, 141
 bipolar disorder 5, 55, 93, 199
 children 94
 depressive episode 55
 homeless persons 245
 manic episodes 55, 56
 pregnant patients 262, 264
see also mania
 Black Mamba 92
 blood alcohol level (BAL) 37
 Blueberry Posh 92
 boarding 13, 203
 limitations on 365
 body dysmorphic disorder 71
 body language 157
 Boerhaave’s syndrome 143
 borderline personality disorder
 (BPD) 7, 107
 brain abscess 91
 brain injury, traumatic *see*
 concussion
 brain lesions 91
 breaking bad news 238
 Breakthrough Collaborative 351
 formation and operation 351
 brief interventions 38
 initial assessment 314
 substance abuse 37
 triage 336
 Brief Mental Status Exam 22
 Brief Psychiatric Rating Scale 316
 broken heart syndrome 84
 bromocriptine, neuroleptic
 malignant syndrome 194
 Buchenwald syndrome 83
 Buddhism 278
 bulimia nervosa 140, 141
 bupropion 183
 side effects 183
 burnout 240
 buspirone 78
 butane, sudden sniffing death
 syndrome 42
 butyl nitrite, sudden sniffing
 death syndrome 42
- CAGE screen 227
 Canadian ED Triage & Acuity
 Scale (CTAS) 315
 cannabinoids 42
 intoxication 152
 side effects 43
 synthetic 92
 capacity 324–333
 lack of 329
 leaving against medical
 advice 329
 carbamazepine 186
 in breast milk 267
 overdose 186
 side effects 186
 teratogenesis 264
 carbohydrate-deficient
 transferrin (CDT) 38
 cardiac-related anxiety 77
 cardiomyopathy 134
 cardiovascular disease 137
 and eating disorders 142
 care coordination 204
 carotenoderma 145
 case managers 293
 catalepsy 89
 catatonia 89
 ceilings 357
 Center for Epidemiological
 Studies Depression
 (CES-D) 220
 Centers for Medicare and
 Medicaid Services
 (CMS) 322
 Change Package 352
 Charge Plus 92
 chemical restraint 96, 187, 206
 definition 177
 forms of 178
 indications 178
 legal issues 369
 medications for 178
 oversedation 187
see also physical restraint
 chest pain 4
 childhood abuse, and suicide
 risk 62
 children/adolescents 211–217
 abuse/neglect 62, 211
 agitation 167
 anxiety 211, 215
 bipolar disorder 94
 class and ethnicity issues 216
 deliberate self-injury
 211, 212
 depression 55, 211, 215
 eating disorders *see* eating
 disorders
 hallucinations 215
 law enforcement 376
 leaving against medical
 advice 330
 medications 213
 out of control 211, 212
 presentations 212
 psychosis 93, 211, 215
 functional 94
 organic 93
 research 379
 schizophrenia 94
 substance abuse 211, 214
 suicide attempts 61
 trauma 216
 ChillX 92
 Chinese spiritual beliefs 278
 chlordiazepoxide (Librium)
 agitation 166
 alcohol withdrawal 47
 teratogenesis 264
 chlorpromazine (Largactil) 185
 agitation 165
 potency 191
 Chronic Consumer Stabilization
 Initiative 374
 chronic illness and suicide
 risk 61
 chronic obstructive lung
 disease 137
 chronic traumatic
 encephalopathy 253
 cirrhosis 134
 citalopram 183
 Clinical Institute Withdrawal of
 Alcohol Scale 47, 48
 clinicians
 attitude to patients 157
 demeanor of 157
 inadvertent acceptance of
 patient’s projections 161
 judgmental behavior 160
 provocation of patients 160
 punitive/threatening
 behavior 160
 Clock Drawing 121
 clomipramine, teratogenesis
 262
 clonazepam (Klonopin)
 agitation 166
 children and adolescents 213
 psychosis 207
 teratogenesis 264
 clonidine, opiate withdrawal 50
 clorazepate, agitation 166
 clothing removal,
 mandatory 367
 Cloud Nine 92
 clozapine (Clozaril) 186
 and metabolic syndrome 135
 potency 191
 teratogenesis 264
 Clozaril *see* clozapine

- co-occurring disorder (COD) *see* SMI-SUD
- cocaine 39
intoxication 152
onset of effects 39
- COD *see* SMI-SUD
- coercion 12
- Cogentin *see* benztropine
- cognitive behavioral therapy (CBT)
anxiety disorders 79, 265
nonsuicidal self-injury 214
somatoform disorders 73
- cognitive impairment 117–123
approach to 117
depression 54
see also delirium; dementia
- “cold turkey” 49
- collateral information 21
- comfort 332, 340, 356
- Commission on Accreditation of Rehabilitation Facilities (CARF) 300
- communication
interpreters/translators 271
verbal de-escalation 158
- communication systems 360
- Community Mental Health Centers 335
- Community Mental Health Centers Act (1963) 298
- community psychiatric services 297–301
case management 300
crisis management 299
current structure 298
day treatment programs 300
history 298
as information providers 313
residential services 300
- comorbidity 132
- compassion fatigue 240
- competence 311, 324
lack of 329
- Comprehensive psychiatric Emergency Programs (CPEP) 14
- computers, siting of 356
- concussion 251–257
assessment 251
behavioral presentation 252
causes 251
discharge planning 256
epidemiology 253
evaluation 253
Glasgow Coma Scale (GCS) 252
neurobehavioral sequelae 252
pathophysiology 252
postconcussive symptoms
behavioral 254
cognitive 254
in nonconcussed patients 256
somatic 255
- Concussion Symptom Inventory (CSI) 257
- conduct disorder 1
- confianza* 274
- confidentiality 356, 370
- confrontation 159
- Confusion Assessment Method (CAM) 118
- Confusion Assessment Model for the Intensive Care Unit (CAM-ICU) 95, 118
- consent to medical research 379
- Consolidated Omnibus Budget Reconciliation Act (1985) 320
- Continuous Improvement 348
- contralase* 274
- conversion disorder 70
- cooling, excited delirium syndrome 129
- coordination of care
agencies involved 291
with community resources 297–301
design of 294
nonclinical collaboration 295
with psychiatric services 291–296
resource availability 293
strengths and limitations 294
teamwork 293
- cortical contusion 252
- Cotard’s syndrome 54
- counselors 292
- criminalization of mental illness 376
- crisis intervention 11, 12, 299
mobile crisis teams 16, 299
psychiatric crisis pyramid 285
voluntary crisis centers 16
- Crisis Intervention Response Teams 374
- crisis intervention teams 299
- Crisis Intervention Training 374
- crisis stabilization units (CSUs) 14, 288
- crisis state 28–30
- Crisis Triage Rating Scale 316
- Critical Incident Stress Debriefing (CISD) 231
- cultural competence 270
- cultural formulation 270
- cultural issues 158, 270–280
African Americans 274
agitated patients 158
Alaska Natives 277
American Indians 277
approach to treatment 279
Asian Americans 274
children and adolescents 216
explanatory models of illness 271
immigration and acculturation 277
language 271
minority populations 273
Pacific Islanders 274
psychopharmacology 279
religion 278
rural communities 283
SMI-SUD 150
- culture 270
organizational 349
- culture-bound syndromes 278
- Da Costa’s syndrome 83
- dantrolene sodium, neuroleptic malignant syndrome 194
- data collection 352
- day treatment programs 300
- de-escalation 200
verbal *see* verbal de-escalation
of violent behavior 171
- decision-making capacity 311
- decoration 359
- delirium 21, 118, 161, 198
approach to patient 117
causes 22, 90
clinical features 118
diagnostic evaluation 119
disposition 120
excited *see* excited delirium syndrome
management 119
older adults 95
and psychosis 89
screening instruments 119
vs. dementia 117
- delirium tremens 47, 91, 134
- delusions 88
inadvertent acceptance by clinician 161
trying to dissuade 160
- dementia 94, 120
and agitation 161
Alzheimer’s disease *see* Alzheimer’s disease
approach to patient 117
clinical features 121
diagnostic evaluation 122
disposition 122
law enforcement 376
Lewy body 122
management 122
and psychosis 91, 223
reversible 121
screening instruments 119
vs. delirium 117
- depression 5, 53–58, 136, 198, 199
children and adolescents 55, 211, 215
clinical features 53
bipolar disorder 5, 55
cognition 54
major depressive disorder 53, 54
mnemonic 54
mood 53
psychomotor activity 53
vegetative function 53
- diagnosis 58
- differential diagnosis 198
- drugs causing 57
- homeless persons 246
- management 58
rapid lysis 208
stabilization 58
suicide risk 58
- older adults 54, 219, 220
inpatient admission 222
- postconcussive 254
- postpartum 55, 266
- pregnant patients 261
- screening for 220
- suicidal ideation 36, 54
- systemic illness causing 56, 57
- detention *see* involuntary detention, limitations on
- dextromethorphan
abuse 41
hallucinations 91
- diabetic ketoacidosis, and eating disorders 145
- diabulimia 145
- Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV) *see* DSM-IV
- Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-V) *see* DSM-V
- dialectical behavioral therapy (DBT) 214
- diazepam (Valium)
agitation 166
pregnant patients 261
alcohol withdrawal 47
children and adolescents 213
excited delirium syndrome 127
teratogenesis 264
- diffusion tensor imaging 253
- Digit Symbol Substitution Test 254
- Dilaudid *see* hydromorphone
- Dios Quire* 274
- disability adjusted life years (DALYs) 1
alcohol abuse 6
- disasters/terrorism 230–233
acute stress reactions 231
disposition 232
impact of 230
medical assessment 230
pain relief 232
psychiatric risk assessment 231
psychological first aid 232
sequestration 232
staged assessments 230
toxicologic assessment 231
treatment and referral 232
see also PTSD; trauma
- discrimination 367

Index

- disposition 203
 alcohol withdrawal 48
 delirium 120
 dementia 122
 eating disorders 147
 neuroleptic malignant syndrome 195
 personality (axis II) disorders 110
 psychosis 99
 SMI-SUD 153
 dissociative anesthesia 91
 dizziness, postconcussive 255
 doctors *see* clinicians
 domestic violence 266
 doors 359
 dream interpretation 31
 droperidol (Inapsine) 96, 98, 185
 aggressive/violent behavior 172, 173
 agitation 165
 excited delirium syndrome 127
 long QT syndrome 310
 psychosis 207, 208
 drug abuse 7
 legal synthetic drugs 92
see also substance abuse
 Drug Abuse Screening Test Modified for ED (DAST-ED) 151
 drug testing 33–35
 false positives/negatives 34
 reasons against 34
 reasons for 33
 drug toxicity, and psychosis 91
 drug withdrawal syndromes 46–50
see also specific drugs
 drug-drug interactions 198
 drug-induced mood disorders 57
 DSM-IV 1, 53
 anxiety disorders 77, 79
 brief psychotic episode 206
 concussion 251
 delirium 118
 dementia 121
 depression 136
 eating disorders 141
 malingering 113, 114
 neuroleptic malignant syndrome 192
 personality disorders 104
 PTSD 83–86
 somatoform disorders 69
 DSM-V 77, 363
 personality disorders 104
 suicidal behavior 6
 duloxetine 138
 dumping 370
 duration of detention 364
 dysthymic disorder 5
 dystonia 166, 172
 Earth Impact 92
 eating disorders 7, 140–148, 211
 complications 141
 cardiovascular 142
 endocrine 144
 gastrointestinal 143
 metabolic and electrolyte abnormalities 144
 neurologic 145
 pulmonary 143
 disposition 147
 DSM-IV criteria 141
 hospitalization of patients 147
 impact of 140
 management 146
 medications used by patients 145
 not otherwise specified (EDNOS) 141
 pregnant patients 266
 prevalence 141
 recovery 145
 SCOFF questionnaire 147, 150–153
 screening 147
 signs and symptoms 141
 types of 141
 ecitalopram 183
 economic burden of mental illness 2
 ecstasy 40
 intoxication 152
 liquid *see* gammahydroxybutyrate
 trauma therapy 238
 Edinburgh scale 261, 267
el destino 274
 elder abuse 227
 elderly *see* older adults
 electrical outlets 357
 electrolyte disturbances in eating disorders 144
 electrolyte replacement in eating disorders 146
 electronic control devices 310
 elevation of patient rooms 356
 elopement 325
 emergency medical condition (EMC) 320
 Emergency Medical Treatment and Active Labor Act *see* EMTALA
 emergency psychiatric evaluation
 best practice 338, 340
 delivery models 11–17
 mental health wing of medical ED 14
 psychiatric consultant in medical ED 12–14
 psychiatric emergency services 14–16
 development of 11
 goals of 11–12
 aftercare plan 12
 exclusion of medical etiology 11
 stabilization of acute crisis 12
 therapeutic alliance 12
 emergency room visits 2
 epidemiology 3
 homeless persons 245
 Emergency Severity Index (ESI) 315
 Emergency Treatment Services (ETS) 14
 emetine 143
 empathy 26–28, 158
 disclaimer 320
 failure of 160
 EMTALA 3, 16, 283, 320–322, 324, 362
 violations 322
 endocrine disease, and eating disorders 144
 engagement with patients 26–28
 engineering controls, siting of 356
 environmental issues 356
 comfort 332, 340, 356
 physical plant 355–361
 escape 325
 esophagitis 143
 ethnicity *see* cultural issues
 Ethnomed 217
 excessive drinking 37
see also alcohol abuse
 excited delirium syndrome 89, 125–129
 diagnosis 126
 etiology 126
 history 125
 initial workup 126
 treatment 127
 benzodiazepines 127
 combination therapy 128
 cooling 129
 first-generation antipsychotics 127
 intravenous fluids 128
 ketamine 128
 second-generation antipsychotics 128
 sodium bicarbonate 129
 explanatory models of illness 271
 extended observation 15
 eye contact 157
 factitious illness 113–116, 200
 assessment 114
 case example 113
 clinical features 114
 definition 113
 diagnosis 114
 management 115
 patient awareness 113, 114
familismo 274
 family and friends 293
 as information providers 314
fatalismo 274
 fatigue, postconcussive 256
 fever 151
 fight or flight response 76
 Filipino Americans 276
 fire alarms 360
 firearms access 174
 and suicide risk 61
see also weapons
 flight of ideas 89
 floors 357
 flow of patients *see* improvements in ED flow
 fluoxetine 183
 in breast milk 267
 depression 55
 side effects 135
 teratogenesis 262
 fluphenazine
 agitation 165
 potency 191
 fluvoxamine 183
 focused investigation 30
 freon, sudden sniffing death syndrome 42
 functional illness 137
 functional MRI 253
 furniture 359
 gamma-glutamyl transferase (GGT) 38
 gammahydroxybutyrate 39
 withdrawal 49
 ganaxolone 238
 gastroesophageal reflux disease (GERD) 143
 gastrointestinal disorders 137
 and eating disorders 143
 gastroparesis 143
 gender
 agitated patient 158
 and suicide risk 60
 generalized anxiety disorder 4, 76, 80
 age of onset 1
 prevalence 4
 treatment 81
see also anxiety disorders
 genetics, and suicide risk 62
 Genie 92
 Geodon *see* ziprasidone
 geographic isolation 284
 geriatric depression scale 220
 Glasgow Coma Scale (GCS) 252
 global burden of mental illness 1
 grab bars 358
 grief *see* bereavement
 gross stress reaction 83
 guardians
 court-appointed 293
 patients leaving against medical advice 331
 guarding behavior 26–28, 30
 guidelines 22
 gustatory hallucinations 88
 Haldol *see* haloperidol
 hallucinations 88
 alcohol withdrawal 46

- auditory 28–30, 88, 96
- children and adolescents 215
- depression 54
- olfactory/gustatory 88
- hallucinogens
 - abuse 41
 - complications 135
 - psychosis 91
- haloperidol (Haldol) 96, 98, 185
- aggressive/violent behavior
 - 172, 173
- agitation 165, 201
 - older adults 167
 - pregnant patients 261
- alcohol intoxication 152
- chemical restraint 178, 310
- contraindications 179
- dementia 122
- dose 201
 - children and adolescents 213
- excited delirium syndrome 127
- potency 191
- psychosis 207, 208
- side effects 135
- hand-offs 317
- head injury *see* concussion
- headache, postconcussive 255
- health care access 270
- Health Care for the Homeless (HCH) 249
- Health Insurance Portability and Accountability Act (HIPAA) 362, 370
- heating 357
- heatstroke 194
- heavy drinking 37
- hepatitis, and eating disorders 143
- heterocyclic antidepressants 183
- Hinduism 278
- HIPAA *see* Health Insurance Portability and Accountability Act
- Hispanic Americans 273
 - culture-bound syndromes 278
- history and physical exam 20
 - risk stratification 21
- homeless persons 244–249
 - assessment 245, 247
 - medical stability 246
 - case studies 245, 246
 - disposition 247
 - emergency room visits 245
 - history
 - part medical 247
 - past psychiatric 247
 - of present illness 247
 - psychosocial 247
 - interventions 245
 - medical problems of 244
 - mental illness 245
 - mode of arrival 246
 - mortality 245
 - substance abuse 245
 - systems issues 248
 - treatment 248
- homelessness 244
- honesty 158
- hopelessness 64
- hormones, and mood disorders 57
- hostage situation 174
- housekeeping 356
- housing supervisors 292
- humiliation of patients 160
- Hurricane Charlie 92
- hydrocodone abuse 38
- hydromorphone (Dilaudid)
 - abuse 38
- hydroxyzine (Vistaril; Atarax) 213
- hyperstimulation 78
- hyperthermia, excited delirium
 - syndrome 129
- hypnotics
 - abuse 38
 - and mood disorders 57
 - withdrawal 48
- hypochloremia, eating
 - disorders 144
- hypochondriasis 70
- hypoglycemia, anxiety
 - symptoms 77
- hypokalemia, eating disorders
 - 144, 146
- hyponatremia, eating
 - disorders 144
- hypoparathyroidism, anxiety
 - symptoms 77
- hypotension, and eating
 - disorders 142
- hypothermia, eating disorders 144
- ICD-10
 - anxiety disorders 77, 79
 - concussion 251
 - somatoform disorders 69
- iloperidone 186
- immigration 277
- ImPACT 254
- improvements in ED flow
 - 347–353
 - achievements 352
 - background 347
 - Breakthrough Collaborative 351
 - Change Package 352
 - culture 349
 - data collection 352
 - patient satisfaction 350
 - quality improvement 349
 - system of care 350
 - systems change theories 348
- impulse control disorders 7
 - prevalence 1
- Inapsine *see* droperidol
- incompetence 329
 - determination of 330
- Indians 276
- infanticide 96
- infection control 360
- inhalants
 - abuse 42
- sudden sniffing death
 - syndrome 42
- inpatient psychiatric units 292
- insomnia, in depression 53
- intellectual developmental
 - delay 376
- International Statistical
 - Classification of Disease and Related Health Problems, 10th Revision *see* ICD-10
- interpreters 271
 - errors of interpretation 272
- interviewing techniques 25–31
 - occult danger to others 28–30
 - occult medical acuity 26–28
 - poor working conditions 30–31
- intravenous fluids, excited
 - delirium syndrome 128
- involuntary commitment 286
- involuntary detention,
 - limitations on 363
 - boarding 365
 - duration of detention 364
 - substantive criteria 363
- Islam 278
- isobutyl nitrite abuse 42
- isocarboxazid 184
- Ivory Wave 92
- jails
 - diversionary behavior in 373
 - mental health care 373
- Japanese 276
- jimson weed 92
- judgmental behavior 160
- ketamine 41
 - agitation 166
 - chemical restraint 178, 179
 - contraindications 179
 - excited delirium syndrome 128
 - hallucinations 91
 - intoxication 152
- Klein, Melanie 106
- Klonopin *see* clonazepam
- Koreans 276
- Korsakoff's dementia 134
- la belle indifférence* 70, 96
- laboratory testing, medical
 - mimics 20
- lamotrigine
 - in breast milk 267
 - teratogenesis 264
- language barriers 271
- lanugo hair, eating disorders
 - 145
- Largactil *see* chlorpromazine
- laryngospasm 172
- latch systems 359
- Latinos 273
- lavatories 358
- law enforcement 373–377
 - children 376
 - criminalization of mental illness 376
 - dementia 376
 - initiatives 374
 - jails 373
 - police-applied restraints 374
 - roles and realities 374
 - see also* legal issues
- law officers 293
 - Crisis Intervention Training 374
 - as information providers 313
 - interaction with 375
- Lean Production 348
- LEARN principle 279
- leaving against medical advice
 - 324–333
- hospital policy 327
- legal issues 328, 330
 - capacity standards 329
 - detention pursuant to involuntary commitment statutes 328
 - intoxication 331
 - lack of capacity 329
 - minors 330
 - persons under guardianship 331
 - psychiatric patients with medical complaints 331
- mitigation of potential harm 332
- liability for adverse outcomes 333
- to patients 332
- prevention 331
 - delay in mental health evaluation 332
 - environmental comfort 332
 - nicotine, alcohol and drug dependence issues 331
 - peer patients 332
 - rewarding of skilled staff members 332
- reasons for 325
- recommended staff practices 326
- leaving before being seen 325
- legal issues 362–371
 - aggressive/violent behavior 175
 - Americans with Disabilities Act 366
 - confidentiality 356, 370
 - discrimination 367
 - dumping 370
 - HIPAA 370
 - leaving against medical advice 328
 - detention pursuant to involuntary commitment statutes 328
 - intoxication 331

Index

- legal issues (cont.)
 lack of capacity 329
 minors 330
 persons under guardianship 331
 psychiatric patients with medical complaints 331
 limitations on ED detention 363
 boarding 365
 duration of detention 364
 substantive criteria 363
 malpractice 363, 369
 mandatory clothing removal 367
 Rehabilitation Act 366
 restraint/seclusion 368
 chemical restraint 369
 physical/mechanical restraint 368
 systemic cases 365
see also law enforcement; law officers
 legal system 362
 lesbians, gays and bisexuals, suicide risk 62
 Lewy body dementia 122
 Librium *see* chlorthalidopoxide
 Licensed Clinical Social Workers 335
 light fixtures 360
 lithium 186, 199
 in pregnancy 263, 264
 side effects 186
 teratogenesis 264
 long QT syndrome 187, 310
 lorazepam (Ativan)
 aggressive/violent behavior 172, 173
 agitation 166, 201
 older adults 167
 pregnant patients 261
 alcohol withdrawal 47
 anxiety 78
 chemical restraint 178, 310
 dose 201
 children and adolescents 213
 excited delirium syndrome 127
 ketamine intoxication 41
 psychosis 96, 98, 207, 208
 teratogenesis 264
 loss of consciousness 252
 loxapine
 agitation 165
 potency 191
 lurasidone 186
 lysergic acid (LSD) 41
 hallucinations 91
 intoxication 152
 lysis of psychosis 206–209
 M-Tab *see* risperidone
 macular degeneration 88
 major depressive disorder 5, 53
 DSM-IV-TR criteria 54
see also depression
 malignant hyperthermia 193
 malingering 113–116, 200
 assessment 114
 case example 113
 clinical features 114
 definition 113
 diagnosis 113
 management 115
 patient awareness 113, 114
 Mallory-Weiss tears 143
 malpractice 363, 369
 mania 55, 198
 differential diagnosis 198
 drugs causing 57
 DSM-IV-TR criteria 56
 symptoms 56
see also bipolar disorder
 maprotiline 183
 marijuana 42
 hallucinations 91
 side effects 43
 toxicity 33
 marital status, and suicide risk 61
 MDMA *see* ecstasy
 mechanical restraint *see* physical restraint
 medical access, limited 132
 medical causes of psychiatric symptoms 197, 198
 medical clearance 19–23, 294
 areas of consensus 19
 best practice 337
 guidelines and protocols 22
 history and physical exam 20
 screening 19, 21
 Medical Clearance Exam 343
 medical evaluation 314, 316
 medical illness 132–138
 comorbidity incidence/prevalence 132
 duality of approach 133
 leaving against medical advice 331
 noncompliance 133
 risk factor assessment 134
 medical mimics 19, 20
 laboratory testing 20
 recognition of 20
 urine drug screens 21
 medical screening exam (MSE) 320
 medications 182–187
 best practice 338
 children and adolescents 213
 cultural issues 279
 rapidly acting 206–209
 rural communities 286
 safety 356
 side effects 198
see also specific drugs/drug groups
 Memory Impairment Screen 121
 mental health clinics 291
 mental health professionals 291
see also clinicians
 mental health services 297–301
 organization of 297
 Mental Health Study Act (1955) 298
 Mental Health Triage Scale 315, 316, 336
 mental healthcare infrastructure 2
 mental illness
 economic burden 2
 emergency room visits 2, 3
 global burden 1
 homeless persons 245
 life cycle extent 1
 perception in rural communities 282
 prevalence 1
 social/physical health impact 2
 mental status 21
 mescaline 41
 hallucinations 91
 intoxication 152
 metabolic alkalosis, eating disorders 144
 metabolic disease
 and eating disorders 144
 and psychosis 94
 metabolic syndrome 135
 metal detectors 356
 methadone
 abuse 38
 in opiate withdrawal 49
 urine testing 34
 methamphetamine 40
 intoxication 152
 N-methyl d-aspartate receptor antagonists *see* NMDAR antagonists
 3,4-methylenedioxymethamphetamine *see* ecstasy
 methylphenidate 41
 psychosis 93
 midazolam (Versed)
 aggressive/violent behavior 172, 173
 agitation 166
 chemical restraint 310
 excited delirium syndrome 127
 psychosis 96, 98
 Mini-Cog 118
 Mini-Mental State Examination 22, 72, 118, 121
 minority populations 273
 Hispanic Americans 273
 minors *see* children/adolescents
 mirrors
 bathroom 358
 observation 360
 mirtazepine 183
 side effects 183
 mitral valve prolapse, and eating disorders 143
 mobile crisis teams 16, 299
 monoamine oxidase inhibitors 184
 anxiety 78
 overdose 184
 mood disorders 5–6
 age of onset 2, 5
 depression *see* depression
 drugs causing 57
 pregnant patients 261
see also specific disorders
 mood stabilizers 186
 morale 295
 morning glory seeds 92
 motivational interviewing 28
 movement disorders, drug-induced 165
 Munchausen syndrome 114
 National Clearinghouse of the Agency for Healthcare Research and Quality 118
 National Hospital Ambulatory Medical Care Survey (NHAMCS) 3
 National Mental Health Act (1946) 298
 National Triage Scale 336
 nausea, postconcussive 255
 neuroleptic malignant syndrome 166, 183, 190–195, 198, 207
 diagnosis 192
 history 192
 laboratory tests 192, 193
 physical examination 192
 vital signs 192
 differential diagnosis 193
 disposition 195
 epidemiology 190
 pathophysiology 190, 191
 treatment 194
 neurological disease, and eating disorders 145
 neuropsychologic testing 254
 nicotine 134
 nitrites, abuse 42
 NMDAR antagonists 91
 noise reduction 360
 nonclinical collaboration 295
 noncompliance 133
 nonpsychiatric physicians 291
 nonsuicidal self-injury 214
 nortriptyline, teratogenesis 262
 nurse call systems 356
 nursing homes, dumping 370
 nutmeg 92
 observation
 extended 15
 mirrors 360
 status 322
 obsessive-compulsive disorder 80, 81
 occult danger to others 28–30
 occult medical acuity 26–28
 Ocean Snow 92

- oculogyric crisis 172
 OG potpourri 92
 olanzapine (Zydis; Zyprexa) 40, 58, 185
 aggressive/violent behavior 173
 agitation 165, 201
 pregnant patients 261
 dementia 122
 dose 201
 children and adolescents 213
 excited delirium syndrome 128
 potency 191
 psychosis 98, 207, 208
 side effects 135
 metabolic syndrome 135
 older adults 219–227
 agitation 167, 225
 delirium 95
 depression 54, 219, 220
 inpatient admission 222
 elder abuse 227
 presenting characteristics 224
 psychosis 94, 223
 substance abuse 226
 suicide attempts 222
 olfactory hallucinations 88
 Opana *see* oxymorphone
 opiates
 abuse 38
 intoxication 152
 overdose 38
 urine testing 34
 withdrawal 49, 91
 oppositional-defiant disorder 1
 organic illness 22
 organizational culture 349
 outreach programs 292
 oversedation 187
 Overt Aggression Scale 317
 Overt Agitation Severity Scale 317
 oxazepam
 agitation 166
 teratogenesis 264
 oxcarbazepine, teratogenesis 264
 oxycodone abuse 38
 oxymorphone (Opana) abuse 38

 Pacific Islanders 274
 pain 138
 agitated patients 161
 post-disaster 232
 pain disorder 70
 paliperidone 186
 pancytopenia 134
 panic alarms 356
 panic attacks 77
 homeless persons 246
 panic disorder 4, 80
 age of onset 1
 prevalence 4
 treatment 81
 underdiagnosis 4
 Panic Screen Score (PSS) 77
 paper towel dispensers 359
 paranoia 161

 paroxetine 183
 PTSD 86
 Patient Health Questionnaire 72
 patients
 arguing with 160
 armed 174
 belongings 357
 clinician's attitude to 157
 cultural issues *see* cultural issues
 engagement with 26–28
 leaving against medical advice 324–333
 pregnant *see* pregnant patients
 satisfaction 350
 see also specific conditions
 peer patients 332
 perphenazine, agitation 165
personalismo 274
 Personality Assessment Inventory 72
 personality (axis II) disorders 7, 103–111
 alliance building 107
 borderline 7, 107
 comorbidities
 addictive illness 104
 medical illness 105
 mental illness 105
 diagnosis 103
 disposition 110
 documentation 111
 etiology 103
 interpersonal issues 105
 interview 106
 medication 110
 mobilization of social support 109
 postconcussive 254
 prevalence 103
 psychodynamic perspective 106
 referral and aftercare 111
 risk assessment 108
 life events 108
 risk management 111
 phencyclidine (PCP)
 abuse 41
 hallucinations 91
 intoxication 152
 toxicity 33
 urine testing 34
 phenelzine 184
 pheochromocytoma 194
 anxiety symptoms 77
 phobic disorders 4, 80
 children and adolescents 215
 social phobia 80
 physical examination 20, 21
 see also assessment
 physical plant 355–361
 see also specific items
 physical restraint 98, 157, 310
 aggressive/violent behavior 173
 agitated patients 167
 definition 177
 forms of 179
 legal issues 368
 staff training 179
 see also chemical restraint
 physicians *see* clinicians
 police *see* law enforcement; law officers
 polypharmacy 135
 Positive and Negative Syndrome Scale (PANSS) 314
 Positive and Negative Syndrome Scale-Excited Component (PANSS-EC) 197
 positron emission tomography (PET) 253
 post-traumatic stress disorder *see* PTSD
 postpartum depression 55, 266
 postpartum psychosis 95, 267
 prednisone, hallucinations 92
 pregnant patients 260–268
 agitation 167, 260
 anxiety disorders 264
 bipolar disorder 262
 depression 261
 postpartum 55, 266
 domestic violence 266
 eating disorders 266
 effects of medications on infant *see* teratogenesis
 false pregnancy 71
 psychosis 95, 265
 postpartum 95, 267
 self-injurious behavior 260
 substance abuse 265
 suicide risk 260
 violent behavior 260
 prescribers 291
 prevalence of mental illness 1
 primary therapist care model 16
 private logic 89
 Programs of Assertive Community Treatment (PACT) 301
 propane, sudden sniffing death syndrome 42
 protocols 22
 Provisional Diagnostic Criteria for Depression in Alzheimer's Disease (PDC-dAD) 221
 provocation of patients 160
 pseudo-Bartter's syndrome 146
 pseudocyesis 71
 psilocybin 41
 hallucinations 91
 intoxication 152
 psychiatric clearance 294
 psychiatric consultants 12–14
 types of 13
 see also clinicians
 psychiatric crisis pyramid 285
 psychiatric emergency services 14–16
 EMTALA 16
 extended observation 15
 structure and design 15
 treatment model 14
 treatment models 16
 triage model 14
 psychiatric evaluation 315
 psychiatric illness, and suicide risk 60
 psychiatric issues 308–311
 electronic control devices 310
 patient assessment *see* assessment
 refusal of care 311
 scene safety 308
 treatment and transport 309
 Psychiatric Medical Clearance Checklist 342
 psychiatric symptoms
 benzodiazepine intoxication/withdrawal 199
 medical causes 197, 198
 medication side effects/interactions 198
 psychiatric causes 199
 recreational drug use 199
 see also specific conditions
 psychiatric urgent care 16
 psychiatrists 291
 psychological first aid 232
 psychologists 292
 psychomotor activity 53
 psychosis 7, 28–30, 88–100, 161, 197, 199
 assessment 206
 causes 90
 functional 92, 94
 organic 89, 93
 children and adolescents 93, 211, 215
 functional 94
 organic 93
 conditions presenting as 89
 and dementia 91, 223
 differential diagnosis 198
 disposition 99
 drug-induced 92
 features of 88
 management 96, 97, 98, 99
 rapid lysis 206–209
 see also antipsychotics
 older adults 94, 223
 postpartum 95, 267
 pregnant patients 95, 265
 PTSD 80, 83–86, 255
 acute reactions 231
 age of onset 1
 children/adolescents 216
 delayed onset 85
 diagnosis 83
 differential diagnosis 84
 diseases associated with 84
 epidemiology 236

Index

- PTSD (cont.)
 history 83
 management 85
 presentation and recognition 84
 risk assessment 231
 screening for 85
 subthreshold presentation 85
 treatment 81
 pulmonary disease, and eating disorders 143
 punitive/threatening behavior by clinicians 160
 pupil diameter 151
- QT prolongation *see* long QT syndrome
 qualified medical personnel (QMP) 320
 Quality Chasm's Ten Rules to Guide Redesign of Health Care 344
 quality improvement 349
 quality improvement organization (QIO) 322
 quetiapine (Seroquel) 58, 186
 agitation 165
 children and adolescents 213
 dementia 122
 and metabolic syndrome 135
 potency 191
 psychosis 98, 207, 208
 Quick Confusion Scale 22
- race *see* ethnicity issues
 racing thoughts 89
 RAND screening instrument 261
 rapidly acting drugs 206–209
 rasagiline 184
 red cell volume (MCV) 38
 Red Dove 92
 refeeding syndrome 146
 referral 204
 personality (axis II) disorders 111
 refusal of care 311, 321
 Rehabilitation Act 366
 religion 278
 research 295, 378–381
 barriers and obstacles 378
 children/adolescents 379
 consent 379
 facilitation of 379
 funding 380
 residential services 300
 resilience 237
 Respect for People 348
respeto 274
 restraint 177–181, 206
 alternatives to 179
 case history 178
 chemical *see* chemical restraint
 complications 180
 definition 177
 documentation 180
 indications 177, 314
 legal issues 368
 physical *see* physical restraint
 police-applied 374
 policy 180
 rhabdomyolysis 310
 risk assessment 21, 285
 disasters/terrorism 231
 personality disorders 108
 PTSD 231
 suicidal patients 65
 risk levels 357
 Risk of Suicide Questionnaire 316
 risperidone (M-Tab) 40, 58, 185
 aggressive/violent behavior 173
 agitation 165, 201
 older adults 167
 pregnant patients 261
 children and adolescents 213
 dementia 122
 potency 191
 psychosis 98, 207, 208
 side effects 135
 rule of 8s 202
 “rum fits” 46
 rural communities 282–288
 agitation management 286
 crisis stabilization units 14, 288
 cross-cultural issues 283
 definition 282
 enhanced EMS care 287
 geographic isolation 284
 in-home assessment/triage 285
 involuntary commitment 286
 lack of treatment centers 285
 medical stabilization 283
 medication management 286
 patient privacy 283
 patient transport 284, 288
 perception of behavioral disorders 282
 providers
 expectations 286
 recruitment and retention 287
 shortages of 285
 short-term treatment units 288
 substance abuse 282
 suicide and violence 283
 telepsychiatry 287
 Russell's sign 145
- SADHART study 133, 136, 137
 SADPERSONS scale 62
 safety 308–311, 340
 assessment 314
 evaluation 202
 medications 356
 physical plant 355–361
 staff 157
 suicidal patients 65
 salvia leaves 92
 SBIRT 37, 38
 Scarface 92
- scene safety 308–311
see also safety
 schizophrenia 1, 7, 93
 and alcohol abuse 150
 children 94
 hallucinations 88
 homeless persons 246
 and substance abuse disorder 150
 suicide risk 61
 schizophreniform disorder 93
 SCOFF questionnaire 147, 150–153
 Screener and Opioid Assessment for Patients with Pain (SOAPP) 73
 screening 19
 accuracy of 21
 algorithms 22
 Screening, Brief Intervention, Referral to Treatment *see* SBIRT
 sealants 357
 seclusion 180
 contraindications 180
 documentation 180
 legal issues 368
 requirements for 180
see also restraint
 security 356
 sedation
 agitated patients 178
 oversedation 187
see also specific drugs
 seizures
 alcohol withdrawal 46
 postconcussive 256
 selective serotonin reuptake inhibitors *see* SSRIs
 selegiline 184
 self-injury, deliberate
 211, 212
 behavioral chain 214
 means restriction 214
 nonsuicidal 214
 pregnant patients 260
 seeking information 212
see also suicidal ideation; suicidal patients
 Sence 92
 separation anxiety disorder 1
 serious mental illness with substance abuse disorder *see* SMI-SUD
 Seroquel *see* quetiapine
 serotonin discontinuation syndrome 198
 serotonin syndrome 183, 191, 198
see also neuroleptic malignant syndrome
 sertraline 183
 PTSD 86
 side effects 135
 Sextacy 92
- sexual orientation, and suicide risk 62
 “shakes” 46
 shell shock 83
 short-term treatment units 288
 sialadenitis 146
 signage 360
 single photon emission tomography (SPECT) 253
 Six Aims of Quality Healthcare 344
 Six Sigma system 348
 Six-Item Screener (SIS) 118
 Skunk 92
 sleep disturbance, postconcussive 256
 small bowel obstruction 143
 SMI-SUD 147, 150–153
 acute phase 152
 behavioral disturbance 151
 disposition 153
 emergency room assessment 150
 epidemiology 150
 ethnic variation 150
 treatment
 emergency room 152
 outpatient 153
 smoking 134
 soap dispensers 359
sobreponerse 274
 social phobia 80, 81
 Social Readjustment Rating Scale 108
 social workers 292
 sodium bicarbonate, excited delirium syndrome 129
 soldier's heart 83
 somatization disorder 69
 somatoform disorders 69–74, 200
 assessment 71
 consultation 72
 emergency room evaluation 71
 body dysmorphic disorder 71
 clinical characteristics 69
 conversion disorder 70
 diagnosis 72
 differential diagnosis 73
 hypochondriasis 70
 management 72
 not otherwise specified 71
 pain disorder 70
 somatization disorder 69
 treatment 72
 cognitive therapy 73
 pharmacotherapy 73
 undifferentiated 70
 space 335, 340
 speech 157
 Spice Gold 92
 Spice Smoke Blend 92
 Sport Concussion Assessment Tool 2 (SCAT2) 253
 sprinklers 360

- SSRIs 137, 183
 anxiety 78
 in breast milk 267
 postpartum depression 267
 PTSD 86
 somatoform disorders 73
 teratogenesis 262
 autism spectrum disorder 262
see also specific drugs
 stabilization 200, 320
 of acute crisis 12
 depression 58
 homeless persons 246
 rural communities 283
see also de-escalation
 staff
 best practice 335, 339
 qualifications 341
 training 341
 patients wanting to leave 326
 restraint training 179
 safety of 157
 skill-related awards 332
 verbal de-escalation 157
 Standardized Assessment of
 Concussion (SAC) 253
 START model 300
 State Mental Health Agencies
 (SMHAs) 298
 status epilepticus 194
 stereotyping 367
 stimulant abuse 39
 stress disorder
 acute 80
 post-traumatic *see* post-
 traumatic stress disorder
 stun guns 310
 substance abuse 6, 36–43
 best practice 337
 brief interventions 37
 CAGE screen 227
 children and adolescents
 211, 214
 homeless persons 245, 247
 leaving against medical
 advice 331
 medical comorbidity 37
 older adults 226
 postconcussive 255
 pregnant patients 265
 prevalence 1, 36
 psychiatric comorbidity 36
 psychiatric symptoms 199
 rural communities 282
 screening for 134
 with serious mental illness 147,
 150–153
 service utilization 37
 suicide risk 36
 urine drug screens 21, 33–35
see also specific drugs
 Substance Abuse and Mental
 Health Services
 Administration
 (SAMHSA) 36, 150, 298
 sudden sniffing death
 syndrome 42
 suicidal ideation 30, 63
 assessment 316
 depression 36, 54, 58
 rapid lysis 208
 eating disorders 140
 substance abuse 36
 suicidal patients 6, 60–66
 assessment 309
 children and adolescents 61
 documentation 66
 epidemiology 60
 management 65
 patient evaluation 63
 protective factors 62
 risk assessment 65
 key indicators 65
 risk factors 60
 age 60
 alcohol abuse 36
 childhood abuse 62
 chronic illness 61
 family history and
 genetics 62
 firearms access 61
 gender 60
 marital status 61
 previous suicide attempt 61
 psychiatric illness 60,
 197–204
 sexual orientation 62
 rural communities 283
 SADPERSONS scale 62
 safety planning 65
 suicide attempts 64
 older adults 222
 pregnant patients 260
 risk of suicide 61, 222
 superior mesenteric artery
 syndrome (SMA) 143
 sympathomimetics
 psychosis 91
see also specific drugs
 sympathy 158
 synesthesia 41
 Syrup of Ipecac 143
 system of care 350
 systemic lupus erythematosus
 (SLE), psychosis 93
 systems change theories 348
 tasers 310
 tauopathy 253
 telephones 360
 telepsychiatry 303–306
 advantages of 303
 emergency guidelines 305
 implementation 305
 literature review 304
 patient identification 306
 rural communities 287
 teratogenesis 262
 antipsychotics 95
see also specific drugs
 terrorism *see* disasters/terrorism
 tetrahydrocannabinol 33
 therapeutic alliance 12
 thioridazine
 agitation 165
 potency 191
 thiothixene 165
 thought disorganization 89
 throughput 339
 thyrotoxicosis 194
 toilet paper dispensers 358
 toilets 358
 torsades de pointes 187
 Trail Making Test 254
 training 295
 best practice 341
 transfer 321
 appropriate 321
 physician refusal of 322
 translators 271
 transport 309
 rural communities 284, 288
 tranlycypromine 184
 trash cans 360
 trauma 160, 161, 235–241
 biochemical response 237
 brain injury 251–257
 breaking bad news 238
 children and adolescents
 216
 definition 235
 disasters *see* disasters/
 terrorism
 management 238
 neurophysiology 237
 neuropsychology 237
 resilience 237
 response to 237
 vicarious 240
see also bereavement; PTSD
 traumatic events 235, 236
 consequences 236
 traumatic grief 240
 traumatic neurosis 83
 trazodone 183
 treatment 201, 309
 acute 197
 cultural issues 279
 homeless persons 248
see also medications; and
 specific conditions
 triage 14, 313–318
 best practice 336
 definition 313
 direct admissions 317
 factors affecting 336
 flowsheet 318
 hand-offs 317
 in-home 285
 initial screening 314
 medical evaluation 314
 model 14
 pre-arrival information 313
 community providers and
 crisis hotlines 313
 family and friends 314
 law enforcement agents 313
 psychiatric evaluation 315
 restraint 314
 safety assessment 314
 transfers 317
 waiting intervals 316
 triazolam 166
 trichloroethylene, sudden sniffing
 death syndrome 42
 tricyclic antidepressants 182
 anxiety 78
 cardiovascular toxicity 183
 overdose 182
 side effects 135, 182
 somatoform disorders 73
 teratogenesis 262
 urine testing 34
see also specific drugs
 trifluoperazine 165
 typical antipsychotics 184
 potency 191
 rapidly acting 207
see also specific drugs
 urine drug screens 21,
 33–35
 US Preventative Services Task
 Force rapid screen 261
 Valium *see* diazepam
 valproic acid 187
 in breast milk 267
 side effects 135, 187
 teratogenesis 263, 264
 vegetative function 53
 venlafaxine 183
 side effects 183
 ventilation 357
 verbal de-escalation 155–162
 agreeing with patient 159
 capturing patient's attention
 159
 clinician's demeanor 157
 communication techniques
 158
 cultural, ethnic, age and
 gender issues 158
 giving instructions 159
 interview errors 160
 offering choices 159
 rapport with patient 157
 rationale for 156
 setting limits 159
 special presentations 161
 staffing 157
 techniques 156
 wants/needs of patient 158
 Versed *see* midazolam
 vicarious traumatization 240
 violence *see* aggressive/violent
 behavior
 visiting nurses 292
 Vistaril *see* hydroxyzine
 vital signs 151

Index

vital signs (cont.) neuroleptic malignant syndrome 192	Wernicke’s encephalopathy 38, 134	World Mental Health Surveys 1	agitation 165, 201 pregnant patients 261
voluntary crisis centers 16	White Dove 92	Yale Delirium Prevention Trial 119	alcohol intoxication 152
waiting room, patient management 316	White Lightning 92	years of potential life lost (YPLL) 1	dose 201 children and adolescents 213
walls 357	White Rush 92	young people <i>see</i> children/ adolescents	excited delirium syndrome 128
water 357	wild children 211, 212	Yucatan Fire 92	potency 191
weapons 308 screening for 174 <i>see also</i> firearms access	Wilson’s disease, psychosis 94	ziprasidone (Geodon) 58, 185 aggressive/violent behavior 173	psychosis 98, 207, 208
well-being 295	windows 358		Zoom 92
	World Health Organization (WHO) Global Burden of Disease Study 2		Zydis <i>see</i> olanzapine
	World Mental Health Surveys 1, 2		Zyprexa <i>see</i> olanzapine