

CAMBRIDGE TRACTS IN MATHEMATICS

General Editors

B. BOLLOBÁS, W. FULTON, A. KATOK, F. KIRWAN,  
P. SARNAK, B. SIMON, B. TOTARO

---

**194 Mathematics of Two-Dimensional Turbulence**

## CAMBRIDGE TRACTS IN MATHEMATICS

General Editors:

B. BOLLOBÁS, W. FULTON, A. KATOK, F. KIRWAN, P. SARNAK, B. SIMON, B. TOTARO

A complete list of books in the series can be found at  
[www.cambridge.org/mathematics](http://www.cambridge.org/mathematics).

Recent titles include the following:

165. Projective Differential Geometry Old and New. By V. OVSIENKO and S. TABACHNIKOV
166. The Lévy Laplacian. By M. N. FELLER
167. Poincaré Duality Algebras, Macaulay's Dual Systems, and Steenrod Operations. By D. MEYER and L. SMITH
168. The Cube-A Window to Convex and Discrete Geometry. By C. ZONG
169. Quantum Stochastic Processes and Noncommutative Geometry. By K. B. SINHA and D. GOSWAMI
170. Polynomials and Vanishing Cycles. By M. TIBĂR
171. Orbifolds and Stringy Topology. By A. ADEM, J. LEIDA, and Y. RUAN
172. Rigid Cohomology. By B. LE STUM
173. Enumeration of Finite Groups. By S. R. BLACKBURN, P. M. NEUMANN, and G. VENKATARAMAN
174. Forcing Idealized. By J. ZAPLETAL
175. The Large Sieve and its Applications. By E. KOWALSKI
176. The Monster Group and Majorana Involutions. By A. A. IVANOV
177. A Higher-Dimensional Sieve Method. By H. G. DIAMOND, H. HALBERSTAM, and W. F. GALWAY
178. Analysis in Positive Characteristic. By A. N. KOCHUBEI
179. Dynamics of Linear Operators. By F. BAYART and É. MATHERON
180. Synthetic Geometry of Manifolds. By A. KOCK
181. Totally Positive Matrices. By A. PINKUS
182. Nonlinear Markov Processes and Kinetic Equations. By V. N. KOLOKOLTSOV
183. Period Domains over Finite and  $p$ -adic Fields. By J.-F. DAT, S. ORLIK, and M. RAPOPORT
184. Algebraic Theories. By J. ADÁMEK, J. ROSICKÝ, and E. M. VITALE
185. Rigidity in Higher Rank Abelian Group Actions I: Introduction and Cocycle Problem. By A. KATOK and V. NIȚICĂ
186. Dimensions, Embeddings, and Attractors. By J. C. ROBINSON
187. Convexity: An Analytic Viewpoint. By B. SIMON
188. Modern Approaches to the Invariant Subspace Problem. By I. CHALENDAR and J. R. PARTINGTON
189. Nonlinear Perron Frobenius Theory. By B. LEMMENS and R. NUSSBAUM
190. Jordan Structures in Geometry and Analysis. By C.-H. CHU
191. Malliavin Calculus for Lévy Processes and Infinite-Dimensional Brownian Motion. By H. OSSWALD
192. Normal Approximations with Malliavin Calculus. By I. NOURDIN and G. PECCATI
193. Distribution Modulo One and Diophantine Approximation. By Y. BUGEAUD
194. Mathematics of Two-Dimensional Turbulence. By S. KUKSIN and A. SHIRIKYAN
195. A Universal Construction for R-free Groups. By I. CHISWELL and T. MÜLLER
196. The Theory of Hardy's Z-Function. By A. IVIĆ
197. Induced Representations of Locally Compact Groups. By E. KANIUTH and K. F. TAYLOR
198. Topics in Critical Point Theory. By K. PERERA and M. SCHECHTER
199. Combinatorics of Minuscule Representations. By R. M. GREEN

# Mathematics of Two-Dimensional Turbulence

SERGEI KUKSIN

*Ecole Polytechnique, Palaiseau*

ARMEN SHIRIKYAN

*Université de Cergy-Pontoise*



**CAMBRIDGE**  
UNIVERSITY PRESS

Cambridge University Press & Assessment  
 978-1-107-02282-9 — Mathematics of Two-Dimensional Turbulence  
 Sergei Kuksin, Armen Shirikyan  
 Frontmatter  
[More Information](#)



CAMBRIDGE  
 UNIVERSITY PRESS

Shaftesbury Road, Cambridge CB2 8EA, United Kingdom  
 One Liberty Plaza, 20th Floor, New York, NY 10006, USA  
 477 Williamstown Road, Port Melbourne, VIC 3207, Australia  
 314–321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre, New Delhi – 110025, India  
 103 Penang Road, #05–06/07, Visioncrest Commercial, Singapore 238467

Cambridge University Press is part of Cambridge University Press & Assessment,  
 a department of the University of Cambridge.

We share the University's mission to contribute to society through the pursuit of  
 education, learning and research at the highest international levels of excellence.

[www.cambridge.org](http://www.cambridge.org)

Information on this title: [www.cambridge.org/9781107022829](http://www.cambridge.org/9781107022829)

© Sergei Kuksin and Armen Shirikyan 2012

This publication is in copyright. Subject to statutory exception and to the provisions  
 of relevant collective licensing agreements, no reproduction of any part may take  
 place without the written permission of Cambridge University Press & Assessment.

First published 2012

*A catalogue record for this publication is available from the British Library*

*Library of Congress Cataloging-in-Publication data*

Kuksin, Sergej B., 1955–

Mathematics of two-dimensional turbulence / Sergei Kuksin, Armen Shirikyan.

p. cm. – (Cambridge tracts in mathematics ; 194)

ISBN 978-1-107-02282-9 (hardback)

1. Hydrodynamics – Statistical methods. 2. Turbulence – Mathematics.

I. Shirikyan, Armen. II. Title.

QA911.K85 2012

532'.052701519 – dc23 2012024345

ISBN 978-1-107-02282-9 Hardback

Cambridge University Press & Assessment has no responsibility for the persistence  
 or accuracy of URLs for external or third-party internet websites referred to in this  
 publication and does not guarantee that any content on such websites is, or will  
 remain, accurate or appropriate.

To  
*Yulia, Nikita, Masha*  
and  
*Anna, Rafael, Gabriel*

## Contents

---

	<i>Preface</i>	<i>page ix</i>
<b>1</b>	<b>Preliminaries</b>	<b>1</b>
	1.1 Function spaces	1
	1.2 Basic facts from measure theory	5
	1.3 Markov processes and random dynamical systems	22
	Notes and comments	35
<b>2</b>	<b>Two-dimensional Navier–Stokes equations</b>	<b>36</b>
	2.1 Cauchy problem for the deterministic system	36
	2.2 Stochastic Navier–Stokes equations	58
	2.3 Navier–Stokes equations perturbed by a random kick force	62
	2.4 Navier–Stokes equations perturbed by spatially regular white noise	69
	2.5 Existence of a stationary distribution	87
	2.6 Appendix: some technical proofs	93
	Notes and comments	99
<b>3</b>	<b>Uniqueness of stationary measure and mixing</b>	<b>101</b>
	3.1 Three results on uniqueness and mixing	104
	3.2 Dissipative RDS with bounded kicks	115
	3.3 Navier–Stokes system perturbed by white noise	126
	3.4 Navier–Stokes system with unbounded kicks	141
	3.5 Further results and generalisations	145
	3.6 Appendix: some technical proofs	159
	3.7 Relevance of the results for physics	168
	Notes and comments	169

<b>4</b>	<b>Ergodicity and limiting theorems</b>	<b>173</b>
4.1	Ergodic theorems	173
4.2	Random attractors and stationary distributions	182
4.3	Dependence of a stationary measure on the random force	202
4.4	Relevance of the results for physics	209
	Notes and comments	210
<b>5</b>	<b>Inviscid limit</b>	<b>211</b>
5.1	Balance relations	211
5.2	Limiting measures	218
5.3	Relevance of the results for physics	241
	Notes and comments	244
<b>6</b>	<b>Miscellanies</b>	<b>245</b>
6.1	3D Navier–Stokes system in thin domains	245
6.2	Ergodicity and Markov selection	251
6.3	Navier–Stokes equations with very degenerate noise	264
	<b>Appendix</b>	<b>269</b>
A.1	Monotone class theorem	269
A.2	Standard measurable spaces	270
A.3	Projection theorem	271
A.4	Gaussian random variables	271
A.5	Weak convergence of random measures	274
A.6	The Gelfand triple and Yosida approximation	275
A.7	Itô formula in Hilbert spaces	277
A.8	Local time for continuous Itô processes	282
A.9	Krylov’s estimate	283
A.10	Girsanov’s theorem	285
A.11	Martingales, submartingales, and supermartingales	286
A.12	Limit theorems for discrete-time martingales	288
A.13	Martingale approximation for Markov processes	289
A.14	Generalised Poincaré inequality	291
A.15	Functions in Sobolev spaces with a discrete essential range	292
	<b>Solutions to selected exercises</b>	<b>293</b>
	<i>Notation and conventions</i>	304
	<i>References</i>	307
	<i>Index</i>	319

## Preface

---

### Equations and forces

Two-dimensional (2D) statistical hydrodynamics studies statistical properties of the velocity field  $u(t, x)$  of a (imaginary) two-dimensional fluid satisfying the stochastic 2D Navier–Stokes equations

$$\begin{aligned} \dot{u}(t, x) + \langle u, \nabla \rangle u - \nu \Delta u + \nabla p &= f(t, x), \quad \operatorname{div} u = 0, \\ u &= (u^1, u^2), \quad x = (x_1, x_2). \end{aligned} \quad (1)$$

Here  $\nu > 0$  is the kinematic viscosity,  $u = u(t, x)$  is the velocity of the fluid,  $p = p(t, x)$  is the pressure, and  $f$  is the density of an external force applied to the fluid. The space variable  $x$  belongs to a two-dimensional domain, which in this book is supposed to be bounded. Suitable boundary conditions are assumed. For example, one may consider the case when the domain is a rectangle  $(0, a) \times (0, b)$ , where  $a$  and  $b$  are positive numbers, and the equations are supplemented with periodic boundary conditions; that is to say, the space variable  $x$  belongs to the torus  $\mathbb{R}^2 / (a\mathbb{Z} \oplus b\mathbb{Z})$  (in the case of periodic boundary conditions we will assume that space-mean values of the force  $f$  and the solution  $u$  vanish). Equations (1) are *stochastic* in the sense that the initial condition  $u_0 = u(0, x)$ , or the force  $f$ , or both of them, are random, i.e., depend on a random parameter. So the solutions  $u$  are random vector fields. The task is to study various characteristics of  $u$  averaged in ensemble, or to study their properties which hold for most values of the random parameter. In this book, we assume that the force is random and refer the reader to [FMRT01] for a mathematical treatment of the Navier–Stokes equations with zero (or deterministic) force and random initial data.

The Reynolds number  $R$  of a random velocity field  $u(t, x)$  is defined as

$$R = \frac{\langle \text{characteristic scale for } x \rangle \cdot (\mathbb{E}E(u))^{1/2}}{\nu},$$



where  $E(u) = \frac{1}{2} \int |u(x)|^{1/2} dx$  is the kinetic energy of the fluid and  $\mathbb{E}$  denotes the average in ensemble. Since the forces we consider are smooth, then the solutions  $u$  of (1) are regular in  $x$  and their space-scale is of order one. So  $R \sim \nu^{-1}(\mathbb{E}E(u))^{1/2}$ . A velocity field  $u$  is called *turbulent* if  $R \gg 1$ . Turbulent solutions for (1) are of prime interest.

If the motion of a “physical” three-dimensional fluid is parallel to the  $(x_1, x_2)$ -plane and its velocity depends only on  $(x_1, x_2)$ , i.e.,  $u = u(t, x_1, x_2)$  and  $u^3 = 0$ , then  $(u^1, u^2)(t, x_1, x_2)$  satisfy (1). Such flows are called *two-dimensional*. Turbulent flows of real fluids are never two-dimensional (i.e., two-dimensional flows are never observed in experiments with high Reynolds number). Still, the 2D equations (1) and the 2D turbulence which they describe are now intensively studied by mathematicians, physicists, and engineers since, firstly, they appear in physics outside the realm of hydrodynamics (e.g., they describe flows of 2D films, see Figure 1 on p. xv), secondly, they provide a model<sup>1</sup> for the 3D Navier–Stokes equations and 3D turbulence, and, thirdly, the 3D statistical hydrodynamics in thin domains is approximately two-dimensional; see Section 6.1 of this book. Accordingly, two-dimensional statistical hydrodynamics is important for meteorology to model intermediate-scale flows in atmosphere (see Figures 6.1 and 6.2 in Chapter 6).

Statistical properties of the random force  $f$  are very important. It is natural and traditional to assume that

- (a) the random field  $f(t, x)$  is smooth in  $x$ , and
- (b) it is stationary in  $t$  with rapidly decaying correlations.

If the space domain is unbounded, we should also assume that

- (c) the space correlations of  $f$  decay rapidly.

However, (c) is not relevant for this book since we only consider flows in bounded domains.

In mathematics, the point of view<sup>2</sup> that turbulence in dimensions 2 and 3 should be described by the Navier–Stokes equations with a random force satisfying (a)–(c) goes back to A. N. Kolmogorov; see in [VF88]. Also see that book for some results on stochastic Navier–Stokes equations in the whole space  $\mathbb{R}^d$ ,  $d = 2$  or  $3$ , with a random force satisfying (a)–(c).

We consider three classes of random forces:

**Kick forces.** These are random fields of the form

$$f(t, x) = h(x) + \sum_{k \in \mathbb{Z}} \delta(t - \tau_k) \eta_k(x), \quad (2)$$

<sup>1</sup> This model is not perfect since it is well known that the Navier–Stokes equations in dimensions 2 and 3 are very different. Still, it may be the best available now. Another popular model for the 3D Navier–Stokes system is the Burgers equation; see the review [BK07] by Bec and Khanin. For the stochastic 1D Burgers equation, see [Bor12].

<sup>2</sup> which is not at all a unique insight on turbulence!

where  $h$  is a smooth deterministic function,  $\tau_k = k\tau$  with some  $\tau > 0$ , and  $\{\eta_k\}$  are independent identically distributed random vector functions, which we assume to be divergence-free. For  $t \in (\tau_{k-1}, \tau_k)$  (i.e., between two consecutive kicks) a solution  $u(t, x)$  for (1), (2) satisfies the deterministic equations  $(1)_{f=h}$ , and at the time  $\tau_k$ , when the  $k^{\text{th}}$  kick  $\eta_k(x)$  comes, it has an instant increment equal to that kick; see Section 2.3. The kick forces are singular in  $t$  and are not stationary in  $t$ , but statistically periodic (the difference between the two notions is not great if the time  $t$  is much longer than the period  $\tau$  between the kicks). An advantage of this class of random forces is that the kicks  $\eta_k$  may have any statistics.

**White in time forces.** These are random fields of the form

$$f(t, x) = h(x) + \frac{d}{dt}\zeta(t, x), \quad (3)$$

where  $h$  is as above and  $\zeta(t) = \zeta(t, \cdot)$  is a Wiener process in the space of smooth divergence-free vector functions. Such random fields are stationary and singular in  $t$ . A disadvantage is that they must be Gaussian; see Section 2.4.

**Compound Poisson processes.** These are kick forces (2) for which the periods  $\tau_k - \tau_{k-1}$  between kicks are independent exponentially distributed random variables.

A big technical advantage of these three classes of random forces is that the corresponding solution  $u(t, x)$ , regarded as a random process  $u(t, \cdot) =: u(t)$  in the space of vector fields, is a Markov process. At the time of writing it is not clear how to extend the results of this book to arbitrary smooth random forces  $f$  satisfying (a) and (b).

### What is in this book?

We are concerned with basic problems and questions, interesting for physicists and engineers working in the theory of turbulence. Accordingly Chapters 3–5 (which form the main part of this book) end with sections where we explain the physical relevance of the obtained results. These sections also provide brief summaries of the corresponding chapters.

In Chapters 3 and 4, our main goal is to justify, for the 2D case, the statistical properties of a fluid's velocity field  $u(t, x)$  which physicists assume in their work. We refer the reader to the books [Bat82; Fri95; Gal02], written in a sufficiently rigorous way and where the underlying assumptions are formulated in a clear manner.<sup>3</sup> The first postulate in the physical theory of turbulence is that

<sup>3</sup> Apart from a few pages at the end, the book [Bat82] is about 3D flows. But all discussions and most of the results may be literally translated to the 2D case.

*statistical properties of a turbulent flow  $u(t, x)$  converge, as time goes to infinity, to a statistical equilibrium independent of the initial data.* Mathematically speaking, this means that a process  $u(t, \cdot)$ , defined by Eq. (1) in the space of vector fields, has a unique stationary measure, and every solution converges to this measure in distribution. That is, the law of the random field  $x \mapsto u(t, x)$  (which is a time-dependent measure in a function space) converges, when  $t \rightarrow \infty$ , to the measure in question. Random processes possessing this property of “short-range memory” are said to be *mixing*.

In Chapter 3, we study the problem of convergence to a statistical equilibrium for Markov processes corresponding to equations with the three classes of random forces as above. We prove abstract theorems which establish the exponential mixing for certain classes of Markov processes. Next we show that these theorems apply to Eq. (1) if a random force  $f$  satisfies certain mild non-degeneracy assumptions. This establishes the convergence to a unique statistical equilibrium and proves that it is exponentially fast.

If the viscosity  $\nu$  and the force  $f$  continuously depend on a parameter in such a way that the former stays positive and the latter stays non-degenerate, then the stationary measure continuously depends on this parameter. For any fixed initial data  $u(0)$  the law of the corresponding solution  $u(t)$  continuously depends on the parameter as well. In Section 4.3, we show that this continuity is uniform in time  $t \geq 0$ . That is, in two space dimensions the statistical hydrodynamics is stable, no matter how big the Reynolds number, whereas the “usual” hydrodynamics of large Reynolds numbers is very unstable.

The mixing has a number of important consequences, well-known in physics, but taken for granted there. Namely, consider any observable quantity  $F(u)$ , such as the first or second component of the velocity field  $u = (u^1, u^2)$ , or the energy  $E = \frac{1}{2} \int |u|^2 dx$ , or the enstrophy  $\frac{1}{2} \int (\text{curl } u)^2 dx$ . Then  $F(t) = F(u(t, \cdot))$  is an ergodic process. That is, *its time average converges to the ensemble average with respect to the stationary measure*. We show that the difference between the two mean values (in time and in ensemble) decays as  $T^{-\gamma}$ , where  $\gamma < 1/2$  and  $T$  is the time of averaging; see Section 4.1.1. Next, if the ensemble average for an observable  $F(u)$  vanishes, then the process  $F(t)$  satisfies the central limit theorem: the law of the random variable

$$\frac{1}{\sqrt{T}} \int_0^T F(t) dt$$

converges, as  $T \rightarrow \infty$ , to a normal distribution  $N(0, \sigma)$ . For non-trivial observables  $F$ , the dispersion  $\sigma$  is strictly positive. In particular, for large  $T$  the random variables  $T^{-1/2} \int_0^T u^j(t, x) dt$ ,  $j = 1, 2$ , are almost Gaussian. Physicists say that *on large time-scales a turbulent velocity field is approximately Gaussian*. These and some other related results are proved in Chapter 4.

In Chapter 5 we study velocity fields  $u(t, x)$ , corresponding to solutions of (1) with a force (2) or (3) where  $h = 0$ , when the viscosity  $\nu$  is small and the Reynolds number is large. There we only discuss stationary measures and stationary-in-time solutions  $u_\nu$  (i.e., solutions  $u_\nu(t, x)$  such that the law  $\mathcal{D}(u_\nu(t))$  for each  $t$  equals the stationary measure). First we observe that for a limit of order one to exist as  $\nu \rightarrow 0$ , the force  $f$  should be proportional to  $\sqrt{\nu}$ ; see Section 5.2.4. So the equations read as

$$\dot{u}(t, x) + \langle u, \nabla \rangle u - \nu \Delta u + \nabla p = \sqrt{\nu} f(t, x), \quad \operatorname{div} u = 0,$$

where  $f$  is the force (2) or (3) with  $h \equiv 0$ . This is in sharp contrast with the 3D theory, where it is believed that a limit of order one exists for the original scaling (1), without the additional factor  $\sqrt{\nu}$  on the right-hand side.<sup>4</sup> In that chapter we restrict ourselves to the case when the space domain is the square torus  $\mathbb{T}^2 = \mathbb{R}^2/2\pi\mathbb{Z}^2$ . The results remain true for the non-square tori  $\mathbb{R}^2/(a\mathbb{Z} \oplus b\mathbb{Z})$ , but the argument does not apply to the equations in a bounded domain with the Dirichlet boundary condition.

Denote by  $\mu_\nu$  the unique stationary measure. We show that the set of measures  $\{\mu_\nu, 0 < \nu \leq 1\}$  is tight (i.e., relatively compact) and that any limit point  $\mu_0 = \lim_{\nu_j \rightarrow 0} \mu_{\nu_j}$  is a non-trivial invariant measure for the Euler system

$$\dot{u}(t, x) + \langle u, \nabla \rangle u + \nabla p = 0, \quad \operatorname{div} u = 0.$$

It is supported by the set of divergence-free vector fields from the Sobolev space  $H^2$  of order two. This result agrees well with the popular belief that *the Euler equation is “responsible” for 2D turbulence*. We do not know if a limiting measure  $\mu_0$  is unique, i.e., if  $\mu_0 = \lim_{\nu \rightarrow 0} \mu_\nu$ . But we know that the measures  $\mu_\nu$  satisfy, uniformly in  $\nu > 0$ , infinitely many algebraical relations, called the *balance relations*. These relations depend only on two scalar characteristics of the force  $f$ . This indicates some *universality features of 2D turbulence*. Such universality is another physical belief. In Section 5.1.3, we use the balance relations to prove that for any  $t$  and  $x$  the random variables  $u_\nu(t, x)$  and  $\operatorname{curl} u_\nu(t, x)$  have finite exponential moments uniformly in  $\nu \geq 0$ . In Section 5.2, we study further properties of the limiting measures  $\mu_0$ . In particular, we establish that any  $\mu_0$  has no atoms and that its support is an infinite-dimensional set.

The results of Chapter 5 provide a foundation of the mathematical theory of space-periodic 2D turbulence. In Section 5.3, we discuss the relation of these results with the existing heuristic theory of 2D turbulence, originated by Batchelor and Kraichnan.

<sup>4</sup> Note that for the small-viscosity Burgers equation the right scaling of the force is also trivial, i.e., without any additional factor; see [BK07; Bor12].

The difference between 2D turbulence and real physical 3D turbulence is very great. In Chapter 6, we discuss a few rigorous results on 3D turbulence, related to the 2D theory presented in the preceding sections. Namely, in Section 6.1 we discuss (without proof) the convergence of the statistical characteristics of a flow in a thin 3D layer, corresponding to the 3D Navier–Stokes system with a random kick force, to those of a 2D flow in the limiting 2D surface. In contrast with similar deterministic results, the convergence holds uniformly in time. So a class of *anisotropic 3D turbulent flows may be approximated by 2D flows* like those which we consider in our book. Section 6.2 contains a discussion of results due to Da Prato, Debussche, and Odasso, and Flandoli and Romito, showing that weak solutions of the stochastic 3D Navier–Stokes system perturbed by a white-in-time random force (which a priori are non-unique) may be arranged as a Markov process. This process is mixing if the force is rough as a function of the space variable. Finally, in Section 6.3, we invoke the methods of control theory to study further properties of stationary measures for Eqs. (1), (3).

### Other equations

The abstract theorems from Chapters 3 and 4 and the methods developed there to study the solutions of Eq. (1) apply to many other stochastic equations. For instance, one can consider the stochastic complex Ginzburg–Landau equation with a conservative nonlinearity,

$$\dot{u} + i\Delta u - i|u|^{2m}u = \Delta u - u + f(t, x), \quad (4)$$

where  $x \in \mathbb{T}^d$ ,  $d \leq 3$ . If  $d = 1$  or  $2$ , then  $m \geq 0$ , while if  $d = 3$ , then one can take, say,  $m \in [0, 1]$ . Such equations describe *optical turbulence*. If  $f$  is a bounded kick force, then direct analogues of the theorems in Chapters 3 and 4 remain true for (4) with the same proof.

However, if the force  $f$  is white in time, then the methods of Chapters 3 and 4 apply only to Eq. (4) with  $m = 1$  if  $d = 1$  and  $m < 1$  if  $d \geq 2$  (while the equation defines a good Markov process for any  $m$  as above). That is, for some deep reason, the arguments developed to treat the stochastic Navier–Stokes equations (1) with white-in-time forces apply only to PDEs with conservative nonlinearities of degree  $\leq 3$ ;<sup>5</sup> see Section 3.5.5.

### Readers of this book

The book is aimed at mathematicians and physicists with some background in PDEs and in stochastic methods. Standard university courses on these subjects are sufficient since the book is provided with preliminaries on function

<sup>5</sup> But the method applies to Eq. (4) with  $m > 1$  if we add a strong nonlinear damping  $-|u|^{2m'}u$ ,  $m' \geq m$ , on the right-hand side.




Figure 1 The onset of 2D turbulence. Panels 1–4 represent the down-motion of a soap film, punctured by a comb at the top. The Reynolds number is increasing from panel to panel. This is 2D turbulent motion described by the 2D Navier–Stokes system (1). Reprinted with permission from [Rut96]. Copyright 1996, American Institute of Physics.

spaces (Section 1.1), on the 2D Navier–Stokes equations (Chapter 2), and on stochastics (Sections 1.2 and 1.3). There the reader will find all the needed non-standard results.

### Acknowledgements

Our interest in the stochastic 2D Navier–Stokes equations (1) and the problem of 2D turbulence appeared during research on the qualitative theory of randomly forced nonlinear PDEs which we undertook at Heriot–Watt University in Edinburgh during the years 1999–2002. The second author was a post-doc, supported by two EPSRC grants and by the university’s funds. We are grateful

to the two institutions for financial support. This book is a much extended version of the lecture notes [Kuk06a] for a course which the first author taught at the Mathematical Department of ETH-Zürich during the winter term of the year 2004/05. He thanks the Forschungsinstitut at ETH for their hospitality and for help in the preparation of the lecture notes.