
Index

- (−1)-curve, 335
 - contraction of a , 433
 - extended, 367
- 1-noetherian, 186
- (−2)-curve, 335, 431
- 2-elementary lattice, 360, 434
- 2-reflective lattice, 358

- $\alpha(X)$, 466, 481
- $\mathfrak{A}(\Delta)$, 105
- $A_{>0}$, 23
- A_2 -maximal, 146
- A_2 -property, 122, 134
- A_2 -variety, 134
- $A_k(X)$, 81
- A_k -variety, 209
- A_σ , 76
- $A(w)$, 130
- Ample(X), 115, 174, 177, 183

- $\beta(X)$, 466
- $\mathfrak{B}(U)$, 108
- $B|D|$, 173
- $\mathbf{B}(w)$, 173
- $b_{i,w}$, 375
- B -lifting, weak, 222
- $B_k(X)$, 81
- $\text{Br}(\mathbb{K})$, 457
- $\text{Br}(X)$, 457
- $\text{Br}_0(X)$, 458
- $\text{Br}_1(X)$, 459
- $\text{Bs}|D|$, 173
- $\text{Bs}(w)$, 173

- $c(X)$, 258
- $c_{X,H}$, 466
- $\text{CAut}(\bar{X}, H_X)$, 252

- $\text{CDiv}(X)$, 61
- $\text{Cl}(X)$, 3, 25, 30, 32, 35, 39, 55, 81, 144, 298
- $\text{Cl}(X, x)$, 61, 114, 169
- $\text{cov}(\Phi)$, 144
- $\text{cov}(\Theta)$, 111

- δ -collection, 97
- $\Delta(\mathfrak{B})$, 108
- $\Delta(r)$, 203
- $\Delta(\Theta)$, 108
- $|D|$, 27, 337
- $D_1 \equiv D_2$, 335
- $D_1 \cdot D_2$, 335
- D -divisor, 27, 32
- $[D]$ -divisor, 50, 54, 55
- D_X^{ij} , 200
- D_X^k , 200
- D -localization, 27
- $[D]$ -localization, 51, 54, 55
- $\text{div}_{[D]}(f)$, 50
- $\text{div}_D(f)$, 27, 32
- $\text{div}(f)$, 25

- (e-e), 405
- (e-p), 405
- $\text{Eff}(X)$, 115, 172, 177, 183, 184, 268, 278, 315, 338, 339, 345, 356, 368, 431
- $\text{Env}(\Phi)$, 162
- $\text{Env}(G)$, 324

- \mathfrak{F} -bunch, 143
 - maximal, 147
 - true, 143
- \mathfrak{F} -face, 126, 143
- \mathbb{F}_a , 326

- γ -collection, 97

- $\Gamma(X, \mathcal{O}_X)$, 17
 $\mathbb{G}_{m, \mathbb{K}}$, 439
 $(G, 2)$ -maximal subset, 138
 $G(2, 5)$, 234
 g_X , 439
 G/F -embedding
 small, 304
 G_x , 18
 $G \cdot x$, 18
 G -action, 20
 continuous, 439
 strongly stable, 65
 G -algebra, 452
 G -ample cone, 327
 G -closedness, 22
 G -equivariant morphism, 16
 G -invariant morphism, 20
 G -linearization, 244
 of a sheaf of graded algebras, 248
 G -maximal, 108
 G -saturated, 108, 137
 G -separation, 22
 G -set
 good, 107, 137
 G -maximal subset, 108
 G -saturated subset, 108
 G -sheaf structure
 graded, 249
 G -variety, 16

 $H' \times^H \mathcal{X}$, 443
 $H^1(\mathbb{K}, H)$, 441
 $H^i(X, D)$, 336
 $h^i(X, D)$, 336
 $H^i(X, \mathcal{F})$, 336
 $h^i(X, \mathcal{F})$, 336
 H_L , 237
 H^L , 237
 H_X , 48, 57, 453
 $\mathcal{H}\mathcal{X}$, 443
 H -factorial variety, 53, 66, 137, 210, 284,
 285
 H -homogeneous rational function, 53
 H -prime divisor, 53

 $I_p(\eta_1, \dots, \eta_r)$, 481
 inv_v , 457

 $\mathcal{J}_{\text{irr}}(X)$, 64, 85, 118

 $\kappa(D)$, 339
 \mathcal{K}_X , 178
 \mathbb{K}_v , 454
 \mathbb{K}^* -surface (rational) 4, 213, 405

 A_2 -singular del Pezzo, 145, 165, 171, 177,
 180, 419
 E_6 -singular del Pezzo, 198, 201, 203, 207,
 283, 421, 489
 anticanonical divisor, 423
 defining data, 331, 405
 Gorenstein log del Pezzo, 422
 intersection numbers, 411
 rational elliptic, 354
 resolution of singularities, 417
 with ample canonical divisor, 350, 429
 \mathbb{K} -torsor, 441
 \mathbb{K} -variety, 437
 morphism of, 438
 k -boundaries, 81
 $(-k)$ -curve, 335, 366
 k -cycle, 81
 prime, 81
 K -factorial graded algebra, 53, 143, 186, 191
 K -graded R -algebra, 7
 automorphism of, 254
 moving cone of a, 182
 K -graded homomorphism, 7
 K -height of a prime ideal, 53
 K -homogeneous
 component, 7
 element, 7
 ideal, 9
 K -prime
 element, 52, 143
 ideal, 53
 K -principal ideal, 53

 $\lambda(w)$, 130
 $\Lambda(\mathcal{W})$, 93
 $\Lambda(X, H)$, 131
 $\Lambda(X)$, 131, 321
 \mathbb{L} -rational point, 438

 $\mu(k)$, 468
 $\mu(w_{i_1}, \dots, w_{i_{n+d}})$, 179
 \tilde{M} , 240
 $\text{Mov}(\mathcal{W})$, 94
 $\text{Mov}(R)$, 182
 $\text{Mov}(X)$, 115, 173, 177, 183, 184, 268, 271,
 338
 $\text{MW}(\pi)$, 352, 353

 \mathcal{N}_X , 178
 $N^1(X)$, 335
 $\text{Nef}(X)$, 272, 336, 338, 340, 341, 344, 361

 $\omega(\eta)$, 480
 ω_X , 124

- $\Omega_{\mathbb{K}}$, 454
 Ω_X , 124
 $\text{ord}_D(f)$, 25
 $\varphi^*(q)$, 493
 π -admissible, 220
 P_{\downarrow} , 99
 P^{\uparrow} , 99
 $p_a(X)$, 336
 (p-e) 405
 (p-p) 405
 $p_X: \widehat{X} \rightarrow X$, 56, 57, 65, 453
 P -drop, 99
 P -lift, 99
 $\text{PDiv}(X, x)$, 61
 $\text{Pic}(X)$, 59, 61, 114, 169, 240, 298, 328
 $\text{Proj}(A)$, 24
 \mathbb{Q}_{∞} , 454
 \mathbb{Q} -divisor, 184
 \mathbb{Q} -factorial (pre-)variety, 28
 \mathbb{Q} -factorial point, 62, 115, 170, 171
 \mathbb{Q} -Gorenstein, 179
 Q_{\downarrow} , 98
 Q^{\uparrow} , 98, 111
 $q(X)$, 337
 Q -drop, 98
 Q -lift, 98
 $\varrho(X)$, 335
 \mathcal{R} , 35, 40
 $\mathcal{R}_{[D]}$, 35, 40
 $\mathcal{R}_{[D]}(X)$, 35, 40
 $\mathcal{R}(L)$, 237
 $\mathcal{R}(X)$, 35, 40
 \mathcal{R}^x , 40
 \mathcal{R}_X , 451
 $\mathcal{R}_X(X)$, 451
 $R(A, P_0)$, 192
 $R(A, P)$, 197, 406
 $R[K]$, 8
 R -algebra
 K -graded, 7
 $\text{rlv}(\Phi)$, 144
 σ° , 79
 Σ , 111
 $\widehat{\Sigma}$, 83, 111
 $\widehat{\Sigma}_1$, 164
 Σ^{\sharp} , 92
 Σ_1 , 164
 $\Sigma_1 \leq \Sigma_2$, 91
 $\Sigma(\lambda)$, 93
 $\Sigma(P)$, 407
 S_X , 169
 $\text{SAmple}(X)$, 115, 174, 177, 182, 183, 268,
 271, 272, 338, 340, 341, 344, 431
 SL_2 -embedding, affine, 332
 SL_3 , 305
 SL_n , 151
 SO_n , 299
 $\text{Spec}_X \mathcal{S}$, 27, 28
 $\text{Spec}(\Phi)$, 248
 $\text{Supp}(D)$, 26
 $\tau_H(X)$, 466
 $\tau(w_{i_1}, \dots, w_{i_{n+d}})$, 179
 Θ^{\sharp} , 92
 $\Theta(\lambda)$, 93
 $\Theta'_r(C)$, 481
 $\Theta(U)$, 108
 T_N , 76
 T -variety of complexity 1, 197, 202, 281, 282,
 330, 405
 $\text{trop}(Y)$, 204
 $\text{type}(\mathcal{X})$, 443
 $U(\Phi)$, 135
 $U(x)$, 135
 \mathcal{V} -collection, 98
 \mathcal{V} -cone, 91
 \mathcal{V} -(quasi-)fan, 91
 $w_{i_1} \cdots w_{i_{n+d}}$, 180
 \mathcal{W} -bunch, 92
 \mathcal{W} -collection, 98
 $\text{WDiv}_{\mathbb{Q}}(X)$, 184
 $\chi(\mathcal{O}_X)$, 336
 $\mathbb{X}(H)$, 439
 \overline{X} , 63, 66, 144, 453
 $\overline{X}(\gamma_0)$, 167
 \widehat{X} , 48, 57, 453
 $\widehat{X}(A, P, \Phi)$, 200
 $\widehat{X}(\mathfrak{F})$, 147
 $\widehat{X}(R, \mathfrak{F}, \Phi)$, 144
 \widetilde{X} , 48, 58, 167
 $\widetilde{X}(L)$, 237
 $X_+(G)$, 152
 $X(A, P)$, 410
 $X(A, P, \Phi)$, 200
 $X_{D,f}$, 27
 X_{γ_0} , 144
 $X(\gamma_0)$, 167
 $X_{\mathbb{L}}$, 438
 $X(R, \mathfrak{F}, \Phi)$, 144
 X_{σ} , 76
 $X^{ss}(w)$, 129

- X -torsor, 442
- $Y(D_1, \dots, D_r)$, 386
- $Z_k(X)$, 81
- abelian covering, 386
- accumulating subvariety, 465
 - weakly, 465
- acting torus, 76
- action
 - algebraic group, 16, 440
 - complexity of, 197, 258
 - continuous G -, 439
 - diagonal quasitorus, 16
 - effective, 20
 - lifting of an, 244
 - quasitorus, 16, 18, 57, 70, 124
 - strongly stable G -, 65
 - torus, 16, 75, 124, 186, 279, 405
 - twisted, 441
- ADE singularity, 418, 471
- adjunction formula, 336
- admissible operation, 330
- admissible
 - π -, 220
- affine SL_2 -embedding, 332
- affine G -algebra, 452
- affine algebra, 16
- affine algebraic group, 14, 151, 438
- affine algebraic monoid, 324
- affine algebraic semigroup, 324
- affine intersection, 33, 71
- affine morphism, 20
- affine semigroup, 117
- affine variety, 14
- algebra
 - affine, 16
 - affine G -, 452
 - factorially graded, 53, 143, 186, 191
 - graded, 7, 69
 - graded affine G -, 452
 - graded factor, 9
 - integral, 8
 - monoid, 8
 - Rees, 230, 275
 - sheaf of divisorial, 25, 28, 248
- algebraic Brauer group, 459
- algebraic Brauer-Manin obstruction, 459
- algebraic group, 14, 438
 - action of, 16, 440
 - affine, 14, 438
 - character group of, 14
 - character of, 14
 - homomorphism of, 14, 438
 - reductive, 20
- algebraic modification, 220
- algebraic monoid
 - affine, 324
- algebraic semigroup
 - affine, 324
- almost Fano variety, 465
- almost free grading, 143, 157, 199
- almost general position, 367, 470
- almost homogeneous variety, 75, 117, 297
- ambient modification
 - toric, 226
- ample cone, 4, 115, 174, 177, 183
 - G -, 327
- ample divisor, 65, 174
 - very, 29
- ample group, 239
- anticanonical embedding, 464
- anticanonical height, 464
- anticanonical model, 368
- arithmetic genus, 336
- Artin's criterion, 335
- automorphism
 - of K -graded algebra, 254
 - of K3 surfaces, 356
 - of complete toric variety, 329
 - of Mori dream space, 251
 - weak, 252
- base locus, 173, 174, 176, 337
 - stable, 173, 174, 176, 337
- base point free theorem, 338
- Betti number
 - Picard graded, 375
- big cell, 320
- big divisor, 339
- big open subset, 146
- Birch's singular locus, 462
- blow-up, 226, 232, 273, 274, 277, 278, 279, 282, 315, 344, 354, 366, 367, 368, 385, 436
- Borel subgroup, 151, 258, 314
- boundary divisor, 272, 314
- branch divisor, 386
- Brauer group, 457, 458
- Brauer-Manin obstruction, 458, 459, 463
- Brauer-Manin set, 458
 - étale, 463
- Brauer-Severi surface, 470
- bunch
 - \mathfrak{F} -, 143
 - \mathcal{W} -, 92
 - lattice, 111

- maximal \mathcal{F} -, 147
- of orbit cones, 134
- true \mathcal{F} -, 143
- bunched polynomial ring, 144
- bunched ring, 143
- bundle
 - étale principal, 48
 - line, 210, 241, 243, 244, 464
- Calabi-Yau type, 274
- canonical Cox sheaf, 40
- canonical embedding, 302
- canonical linearization, 260
- canonical resolution of singularities, 417
- canonical section, 54
- canonical toric embedding, 162
- Cartier divisor, 28
- Castelnuovo criterion, 335
- categorical quotient, 23, 121, 297
- chamber
 - GIT, 130, 184
 - GKZ, 93
 - Mori, 4, 184
 - positive Weyl, 152
- character
 - of algebraic group, 14
 - identifying, 39
 - Galois-equivariant identifying, 450
- character group, 14
- characteristic quasitorus, 4, 48, 7, 57, 453
- characteristic space, 4, 6, 48, 57, 65, 82, 453
- Chevalley's theorem, 297
- Chow group, 81
- circle method
 - Hardy-Littlewood, 462
- class group
 - divisor, 3, 25, 30, 32, 35, 39, 55, 81, 144, 298
 - local, 61, 114, 169
- coarsened grading, 9
- coarsest common refinement, 94
- Coble surface, 363
- cocycle, 261, 441
- cohomology
 - Dolbeault, 358
 - Galois, 441
 - sheaf, 336
- collection
 - δ -, 97
 - γ -, 97
 - \mathcal{V} -, 98
 - \mathcal{W} -, 98
 - of relevant faces, 144
 - covering, 112, 144
- true lattice, 110
- color, 314
- column set, 158
- complete intersection, 178, 193
- complete linear system, 27, 337
- complete (quasi-)fan, 80
- complexity, 258
- complexity 1
 - T -variety, 197, 279, 435
 - grading of, 192
 - torus action, 197, 279, 435
- cone
 - G -ample, 327
 - \mathcal{V} -, 91
 - ample, 174
 - effective, 172
 - lattice, 76
 - moving, 94, 173, 182
 - nef, 336
 - orbit, 124
 - pointed, 76
 - polyhedral, 8
 - projected, 111, 126, 143
 - regular, 80
 - relative interior of a, 79
 - semiample, 174
 - simplicial, 80
 - weight, 8, 124
- cones of divisors, 115, 172
- conic, 366, 374
- connected γ -collection, 98
- constant part of the Brauer group, 458
- continuous G -action, 439
- continuous 1-cocycle, 441
- contraction
 - of a divisor, 226, 431
 - of a (-1) -curve, 433
- convergent one-parameter subgroup, 77
- coordinate space
 - total, 1, 63, 66, 144, 453
- coprimality condition, 476
- covering collection, 112, 144
- covering
 - cyclic, 386
 - abelian, 386
- Cox coordinates, 163, 232, 430
- Cox ring, 1, 35, 40, 451
 - of a cubic surface, 473, 474
 - of a del Pezzo surface, 234, 369, 374, 375, 422
 - of an embedded variety, 216
 - of a homogeneous space, 300
 - of a rational elliptic surface, 353
 - of a spherical variety, 316

- Cox ring (*cont.*)
- of a toric variety, 82
 - of a variety with torus action, 200, 279, 407
 - of a wonderful variety, 321, 322
 - of blow-up, 226, 229, 231
 - of Gorenstein log terminal del Pezzo surfaces, 234, 422
 - of K3 surface, 400
 - algebraic properties, 47
 - divisibility properties, 52
 - finite generation, 230, 257
 - well-definedness, 41
- Cox sheaf, 35, 39, 57, 237, 451
- canonical, 40
 - Galois-equivariant, 451
 - geometric realization, 56
 - not locally of finite type, 330
- Cox-Nagata ring, 274
- Cox's construction, 86, 162
- cubic surface
- E_6 -singular, 198, 201, 203, 207, 283, 421, 489
 - Cox ring of, 473
- curve
- (-1)-, 335
 - (-2)-, 431
 - (- k)-, 335, 366
 - cubic, 273, 352, 431, 435
 - extended (-1)-, 367
 - on a surface, 334
- cyclic covering, 386
- Daniilov-Gizatullin surface, 331
- degree
- in a graded algebra, 7
 - of a del Pezzo surface, 368, 471
- degree vector, 178
- del Pezzo surface, 366, 423, 469
- Cox ring of a, 234, 369, 374, 375, 423
 - degree of a, 368, 471
 - family, 436
 - Gorenstein log terminal, 234, 423
 - singular, 145, 198, 234, 366, 423
 - smooth, 146, 366, 433
 - type of a, 471
 - weak, 367, 469
- Demazure root, 329
- descent, 459
- diagonal action, 16
- diagonalizable group, 13
- discriminant group, 360
- distinguished point, 79
- distinguished polynomial, 370
- distinguished section, 369
- divisor class group, 3, 25, 30, 32, 35, 39, 55, 81, 144
- divisor
- D -, 27
 - $[D]$ -, 50
 - H -prime, 53
 - \mathbb{Q} -, 184
 - ample, 65, 174
 - big, 339
 - boundary, 314
 - branch, 386
 - Cartier, 28
 - cones of, 172
 - effective, 25
 - littaka dimension of, 339
 - locally principal, 28
 - movable, 115, 173
 - of a function, 25
 - prime, 24
 - principal, 25
 - rational, 184
 - semiample, 115, 174
 - support of a, 26
 - very ample, 29
 - Weil, 24
- divisorial (pre-)variety, 240
- divisorial algebra
- sheaf of, 25, 28, 248
- Dolbeault cohomology group, 358
- dominant weight, 152, 321
- du Val singularity, 418
- effective action, 20
- effective cone, 115, 172, 177, 183, 184, 268, 278, 315, 338, 339, 345, 356, 368, 431
- effective divisor, 25
- numerically, 335
- effective grading, 192
- elementary obstruction, 449
- elliptic fixed point, 409
- elliptic surface
- extremal rational, 350
- embedding
- SL_2 -, 332
 - anticanonical, 464
 - canonical, 302
 - canonical toric, 162
 - equivariant, 304
 - neat, 161
 - small G/F -, 304
- Enriques surface, 361
- envelope, 162
- enveloping monoid, 324
- epimorphic subgroup, 307

- equivalent
 - linearly, 25
 - Mori, 183
 - numerically, 335
- equivariant embedding, 304
 - morphism of, 304
- equivariant morphism, 16, 441
- étale H_X -principal bundle, 48
- étale Brauer-Manin obstruction, 463
- étale Brauer-Manin set, 463
- Euler characteristic, 336
- Euler function, 493
- even lattice, 356
- everywhere locally soluble, 454
- exceptional weight, 229
- exponential sequence, 432
- extended (-1) -curve, 367
- extended Dynkin diagram, 490
- extended multi-Rees algebra, 275
- extremal elliptic fibration, 348
- extremal rational elliptic surface, 350, 351

- face ring, 118
- face subalgebra, 73
- factor algebra
 - graded, 9
- factorial point, 62, 170, 171
 - \mathbb{Q} -, 62, 115, 170, 171
- factorial (pre-)variety
 - H -, 53
 - \mathbb{Q} -, 28
 - locally, 28
- factorially K -graded algebra, 53, 143, 186, 191
- fake weighted projective plane, 122
- fan, 77
 - $L_{\mathbb{Q}}$ -projectable, 104
 - \mathcal{V} -, 91
 - complete, 80
 - generating, 262
 - GIT, 131
 - lattice, 77
 - nondegenerate, 80, 82
 - normal, 80
 - polytopal, 80
 - quasi-, 77
 - support of a, 80
- Fano type, 273
- Fano variety, 179, 273, 366, 465
 - almost, 465
- fiber formula, 79
- fibration, 347
 - extremal, 348
 - jacobian, 351
 - relatively minimal, 347
- finite generation of Cox rings, 230, 257
- fixed point
 - elliptic, 409
 - hyperbolic, 409
 - parabolic, 409
- flag variety, 151, 381
- full intrinsic quadric, 159, 212
- function
 - divisor of a, 25
 - homogeneous, 16
 - order along a prime divisor, 25
- fundamental exact sequence, 444
- fundamental representation, 382
- fundamental weight, 152, 324

- Gale dual, 3, 89, 54, 145
- Gale transform, 90, 119
- Galois cohomology group, 441
- Galois-equivariant Cox sheaf, 451
- Galois-equivariant identifying character, 450
- Galois group, 437
- Gelfand-Kapranov-Zelevinsky decomposition, 93
- general position, 366
 - almost, 367, 470
 - very, 277
- generating fan, 262
- generating pair, 262
- generic isotropy, 279
- generic K3 surface, 397
- genus
 - arithmetic, 336
- genus formula, 336
- geometric invariant theory, 65
 - for affine quasitorus action, 110, 124
- geometric line bundle, 243
- geometric Picard group, 443
- geometric quotient, 21, 48, 63, 89, 106, 240, 279, 411
- geometrically connected, 438
- geometrically irreducible, 438
- geometrically rational, 438
- GIT, 65
 - for affine quasitorus action, 124
- GIT chamber, 130, 184
- GIT cone, 130, 184
- GIT (quasi-)fan, 131, 183
- GKZ decomposition, 93
- global Torelli theorem, 358
- good G -set, 107, 137
- good quotient, 21, 28, 48, 57, 66, 72, 103, 124, 144, 209, 237, 260
 - separated, 104
 - toric, 103
- good toric ambient modification, 230

- Gordan's lemma, 10
 Gorenstein, 179
 Gorenstein del Pezzo surface, 234, 423
 graded G -sheaf structure, 249
 graded affine G -algebra, 452
 graded algebra, 7, 69
 K -factorial, 53, 143, 186, 191
 K -homogeneous component of a , 7
 K -homogeneous element of a , 7
 automorphism of K -, 254
 coarsened grading of a , 9
 degree in a , 7
 factor algebra of, 9
 factorially, 53, 143, 186, 191
 homomorphism of, 7
 irrelevant ideal of a , 23
 lifting of, 9
 morphism of, 7
 moving cone of a K -, 182
 trivial extension of a , 9
 weight cone of a , 8
 graded homomorphism, 7
 gradiator, 156
 grading, 7
 almost free, 143
 coarsened, 9
 effective, 192
 maximal diagonal, 156
 of complexity 1, 192
 pointed, 192
 Grosshans extension, 300
 Grosshans subgroup, 300
 group action, 16, 440
 lifting of a , 244
 group
 affine algebraic, 14, 438
 algebraic, 14, 438
 ample, 239
 Chow, 81
 diagonalizable, 13
 discriminant, 360
 divisor class, 25
 geometric Picard, 443
 isotropy, 18, 62
 local class, 61
 Mordell-Weil, 352
 multiplicative, 439
 Néron-Severi, 335
 of multiplicative type, 439
 orbit, 19, 124
 Picard, 61
 reductive algebraic, 20
 weight, 8
 Weyl, 329, 358, 432, 470, 481
 Halphen transform, 354
 Hamilton's quaternions, 457
 Hardy-Littlewood circle method, 462
 Hasse norm principle, 455
 Hasse principle, 454
 Hasse-Minkowski theorem, 455
 height
 anticanonical, 464
 Weil, 464
 height condition, 476
 Hesse surface, 354
 highest weight, 152, 321, 382
 Hilbert's fourteenth problem, 277, 308
 Hirzebruch surface, 86, 326, 345, 367
 Hodge index theorem, 336
 homogeneous component of a graded algebra, 7
 homogeneous element in a graded algebra, 7
 homogeneous function, 16
 weight of, 16
 homogeneous ideal, 9
 homogeneous multiplicative monoid, 186
 homogeneous rational function, 53
 homogeneous space, 151, 297
 canonical embedding of, 302
 divisor class group of, 298
 Cox ring of, 300
 Picard group of, 298
 principal, 441
 homogeneous toric variety, 329
 homogeneous variety
 almost, 297
 homomorphism
 K -graded, 7
 of algebraic groups, 14, 438
 of graded algebras, 7
 hyperbolic fixed point, 409
 hyperbolic lattice, 356
 ideal
 K -height of a prime, 53
 K -homogeneous, 9
 K -prime, 53
 K -principal, 53
 irrelevant, 23, 64, 85, 118
 Stanley-Reisner, 118
 identifying character, 39
 Galois-equivariant, 450
 Iitaka dimension, 339
 integral algebra, 8
 intersection multiplicity, 335
 intersection number, 82, 334, 411
 self-, 335, 411
 intersection product, 81, 335

- intrinsic quadric, 500
 - full, 159
- invariant morphism, 20
- invariant separation lemma, 96
- involution, 45, 362, 434
 - hyperelliptic, 432
 - nonsymplectic, 397
- irreducible
 - geometrically, 438
- irregularity, 337
- irrelevant ideal, 1, 23, 64, 85, 118, 230, 303
- isotropy group, 18, 34, 62, 279

- jacobian elliptic fibration, 351

- K3 surface, 356, 390
 - automorphism group, 356, 358
 - Cox ring, 400
 - curves on a, 434
 - generic, 397
 - Mori dream, 358, 359
 - Picard group, 434
 - Torelli theorem, 358
- Kawamata log terminal pair, 273
- Kawamata-Viehweg vanishing theorem, 337
- Kempf-Ness set, 211
 - toric, 118
- King's theorem, 208
- Kleiman-Chevalley criterion, 212
 - toric, 122
- Kleinschmidt's classification, 116
- klt, 273
- Koszul complex, 375
- Kummer surface, 434

- lattice, 76, 356
 - 2-elementary, 360, 434
 - 2-reflective, 358
 - even, 356
 - hyperbolic, 356
 - nondegenerate, 356
 - transcendental, 358
- lattice bunch, 111
 - map of, 112
 - maximal, 111
 - true, 111
- lattice cone, 76
 - map of, 76
- lattice fan, 77
 - map of, 77
- Laurent monomial, 14
- lifting
 - of a group action, 244
 - of graded algebras, 9
- weak B -, 222
- line bundle, 243
 - G -linearization of, 244
 - geometric, 243
 - metrized, 464
 - pullback of a, 244
 - total space of a, 210
- linear form
 - separating, 77
 - separating $L_{\mathbb{Q}}$ -invariant, 96
- linear system, 27
- linearization
 - G -, 244, 248
 - canonical, 260
- linearization problem, 70
- linearly equivalent, 25
- local class group, 61, 114, 169
- local factor, 481
- local invariant map, 457
- localization
 - D -, 27
 - $[D]$ -, 51
- locally factorial variety, 28
- locally principal divisor, 28
- log canonical pair, 273
- log canonical singularity, 273
- log del Pezzo surface, 423
- log terminal pair
 - Kawamata, 273
 - purely, 273
- log terminal singularity, 273, 423

- Manin's conjecture, 466, 489
- maximal
 - A_2 -, 146
 - G -, 108
 - $(G, 2)$ -, 138
 - qp-, 138
- maximal δ -collection, 97
- maximal \mathfrak{F} -bunch, 147
- maximal γ -collection, 97
- maximal γ -collection, 98
- maximal \mathcal{V} -(quasi-)fan, 91
- maximal \mathcal{V} -collection, 98
- maximal \mathcal{W} -bunch, 92
- maximal bunch of orbit cones, 134
- maximal lattice bunch, 111
- maximal toric variety, 112
- metrized line bundle, 464
- Möbius function, 468
- Möbius inversion, 468
- model
 - anticanonical, 368
 - relatively minimal, 347

- modification
 - algebraic, 220
 - toric ambient, 226
- monoid
 - affine algebraic, 324
 - enveloping, 324
 - homogeneous multiplicative, 186
 - orbit, 19, 124
 - weight, 8
- monoid algebra, 8
- Mordell-Weil group, 352, 353
- Mori dream space, 3, 181, 251
 - G -ample cone, 327
 - automorphism of, 251
 - Calabi-Yau type, 274
 - chamber structure of the effective cone, 183, 268
 - characterization by Hu and Keel, 268
 - Fano type, 273
 - GIT quotient of, 260
 - modification of, 230
 - Zariski decomposition in, 184, 185
- Mori dream surface, 4, 334
 - of positive anticanonical lita dimension, 341
 - characterization of, 272
- Coble, 364
- Enriques, 361, 362
- $K3$, 356
- rational elliptic, 353, 354
- Mori equivalence, 183
- morphism
 - G -equivariant, 16
 - G -invariant, 20
 - affine, 20
 - equivariant, 441
 - of G -varieties, 16
 - of \mathbb{K} -varieties, 438
 - of equivariant embeddings, 304
 - of graded affine G -algebras, 452
 - of graded algebras, 7
 - of quotient presentations, 237
 - of small G/F -embeddings, 304
 - of varieties with action of algebraic groups, 16, 440
 - toric, 76
- movable divisor, 115, 173
- moving cone, 94, 115, 173, 177, 183, 184, 268, 271, 338
 - of a K -graded algebra, 182
- multi-Rees algebra, 274
- multiplicative group, 439
- multiplicative monoid, 186
- multiplicative type, 439
- Néron-Severi group, 335
- Néron-Severi torus, 444
- neat embedding, 161, 163
- nef cone, 272, 336, 338, 340, 341, 344, 361
- nef divisor, 335
- Newton polytope, 128
- Nishimura's lemma, 455
- nondegenerate fan, 80
- nondegenerate lattice, 356
- nondegenerate Laurent polynomial, 219
- nonsymplectic involution, 397
- norm 1 torus, 499
- normal (pre-)variety, 25
- normal (quasi-)fan, 80
- normal ring, 47
- normal sheaf, 178
- number field, 454
- numerically effective, 335
- numerically equivalent, 335
- observable subgroup, 300
- obstruction
 - Brauer-Manin, 458
 - descent, 459
 - elementary, 449
- one-parameter subgroup, 76
 - convergent, 77
- orbit, 18
- orbit cone, 124
 - bunch of, 134
- orbit decomposition, 79
- orbit group, 19, 124
- orbit monoid, 19, 124
- orbit structure, 79
- order of a function along a prime divisor, 25
- Orlik-Wagreich graph, 416
- parabolic fixed point, 409
- parabolic subgroup, 151
- Picard group, 59, 61, 114, 169, 240, 298, 328, 434
 - geometric, 443
- Picard number, 335
- Picard quasitorus, 444
- Picard quotient presentation, 240, 327, 500
- Picard-graded Betti number, 375
- Picard-Lefschetz reflection, 360
- place, 454
- plt, 273
- point
 - \mathbb{Q} -factorial, 62, 115, 170, 171
 - distinguished, 79
 - factorial, 62, 170, 171
 - normal, 25

- rational, 438
- semistable, 129
- pointed cone, 76
- pointed grading, 192
- polyhedral cone, 8
- polytopal (quasi-)fan, 80
- position
 - almost general, 367, 470
 - general, 366
 - very general, 277
- positive part, 338
- positive root, 323
- positive Weyl chamber, 152
- prevariety, 14
 - \mathbb{Q} -factorial, 28
 - divisorial, 240
 - irrelevant ideal of a, 64
 - normal, 25
 - separated, 14, 71
- prime k -cycle, 81
- prime divisor, 24
 - H -, 53
- prime element, 52, 143
- principal bundle, étale, 48
- principal divisor, 25
 - locally, 28
- principal homogeneous space, 441
- principal ideal, 53
- Proj construction, 24, 72
- projected cone, 111, 126, 143
- pullback
 - of characters, 14
 - of a Cartier divisor, 30
 - of a line bundle, 244
 - of an invariant Weil divisor, 161
 - of a section of a Cox ring, 230
 - of a Weil divisor, 32
 - of a Weil divisor class, 32
- purely log terminal pair, 273
- pushforward
 - of a Weil divisor, 66
 - of one-parameter subgroup, 77
 - of a section of a Cox ring, 226
 - of a torsor, 443
- qp-maximal subset, 138
- quadric, 374, 455
 - full intrinsic, 159
 - intrinsic, 500
- quotient of a, 155
- quasifan, 77
 - \mathcal{V} -, 91
 - complete, 80
 - GIT, 131
 - normal, 80
 - polytopal, 80
 - support of a, 80
- quasimodification, 183
- quasitorus, 14, 70, 438
 - characteristic, 48, 57, 453
 - Picard, 444
- quasitorus action, 16, 18, 57, 70, 124
- quiver, 208
- quotient
 - A_2 -, 133, 144
 - of H -factorial affine varieties, 137
 - of quadrics, 155, 212
 - categorical, 23, 121, 297
 - geometric, 21, 48, 63, 89, 106, 240, 279, 411
 - good, 21, 28, 48, 57, 66, 72, 103, 124, 144, 209, 237, 260
 - semistable, 128, 182
 - separated good, 104
- quotient presentation, 237
 - morphism of, 237
 - Picard, 240, 327, 500
 - toric, 120
- rational divisor, 184
- rational double point, 357, 418, 469
- rational elliptic surface, 350
- rational point, 438
- rational representation, 16, 17
- rational surface, 345
- rational variety
 - geometrically, 438
- reducing pair, 186
- reductive algebraic group, 20
- Rees algebra
 - extended multi-, 275
 - saturated, 230
- refinement
 - of \mathcal{V} -(quasi-)fan, 91
 - of \mathcal{W} -bunch, 92
 - of bunch of orbit cones, 134
 - coarsest common, 94
- reflection
 - Picard-Lefschetz, 360
- regular cone, 80
- relative interior of a cone, 79
- relative spectrum, 27, 28, 30, 32, 57, 237
- relatively minimal elliptic fibration, 347
- relatively minimal model, 347
- relevant face, 144
- representation
 - fundamental, 382
 - of a quiver, 208
 - rational, 16, 17

Cambridge University Press

978-1-107-02462-5 - Cox Rings

Ivan Arzhantsev, Ulrich Derenthal, Jürgen Hausen and Antonio Laface

Index

[More information](#)

528

Index

- resolution of singularities, 204, 417
- Riemann-Roch formula, 336
- root, 323
 - Demazure, 329
 - positive, 323
 - simple, 323, 356
 - spherical, 321
- Rosenlicht's lemma, 31, 241
- saturated
 - with respect to a map, 135
 - G -, 108, 137
- saturated δ -collection, 97
- saturated γ -collection, 97
- saturated γ -collection, 98
- saturated \mathcal{V} -collection, 98
- saturated Rees algebra, 230
- section
 - canonical, 54
 - distinguished, 369
- self-intersection number, 335, 411
- semiample cone, 115, 174, 177, 182, 183, 268, 271, 272, 338, 340, 341, 344, 431
- semiample divisor, 115, 174
- semistable point, 129
- semistable quotient, 128, 182
- separated δ -collection, 97
- separated γ -collection, 97
- separated \mathcal{V} -collection, 98
- separated good quotient, 104
- separated prevariety, 14, 71
- separating linear form, 77
 - $L_{\mathbb{Q}}$ -invariant, 96
- separation, 280
 - G -, 22
- Serre's duality, 336
- sheaf
 - Cox, 35, 39, 57, 237, 451
 - normal, 178
 - of divisorial algebras, 25, 28, 248
 - of Veronese subalgebras, 237
- sheaf structure
 - graded G -, 249
- Shioda-Tate-Wazir exact sequence, 352
- simple root, 323, 356
- simplicial complex, 118
- simplicial cone, 80
- singular del Pezzo surface, 145
- singular locus
 - Birch's, 462
- singularity
 - \mathbb{Q} -factorial, 62, 115, 170, 171
 - ADE, 418, 469
 - canonical resolution of, 417
 - du Val, 418
 - factorial, 62, 170, 171
 - log canonical, 273
 - log terminal, 273, 423
 - normal, 25
 - singularity type of a del Pezzo surface, 471
 - sink, 409
 - small G/F -embedding, 304
 - morphism of, 304
 - small quasimodification, 183
 - smooth del Pezzo surface, 366
 - of degree 5, 146
 - smooth point, 62, 115, 170, 171
 - source, 409
 - spectrum
 - relative, 27
 - spherical root, 321
 - spherical variety, 258, 314
 - of semisimple rank 1, 332
 - split torus, 499
 - split weak del Pezzo surface, 470
 - stable base locus, 173, 337
 - standard 1-torus, 439
 - standard n -torus, 14
 - Stanley-Reisner ideal, 118
 - Stanley-Reisner ring, 118
 - stratification, 167
 - strongly stable G -action, 65
 - strongly stable torus quotient, 212
 - subalgebra
 - face, 73
 - Veronese, 10, 12, 22, 69, 73, 130, 154, 189, 237
 - subrepresentation of a quiver, 208
 - support
 - of a (quasi-)fan, 80
 - of a Weil divisor, 26
 - surface, 334
 - Brauer-Severi, 470
 - Coble, 363, 364
 - cubic, 283, 473
 - del Pezzo, 145, 146, 366, 423, 433, 469
 - Enriques, 341, 361, 362
 - Hesse, 354
 - Hirzebruch, 86, 326, 345, 367
 - $K3$, 356, 390
 - Kummer, 434
 - Mori dream, 334
 - rational, 345, 431
 - rational \mathbb{K}^* -, 405
 - rational elliptic, 350
 - Zariski decomposition, 338

- Tamagawa number, 466
- Torelli theorem, 358
- toric ambient modification, 226
 - good, 230
- toric embedding, 162
- toric geometry, 78
- toric Kempf-Ness set, 118
- toric Kleiman-Chevalley criterion, 122
- toric morphism, 76
- toric quotient presentation, 120
- toric quotient, 103
- toric variety, 75
 - A_2 -property, 122
 - associated with (K, \mathcal{W}, Θ) , 112
 - associated with \mathfrak{B} , 111
 - characteristic quasitorus of a , 82
 - characteristic space of a , 82
 - Cox ring of a , 82
 - homogeneous, 329
 - maximal, 112
 - total coordinate space of a , 82
- torsor
 - \mathbb{K} -, 441
 - X -, 442
 - universal, 444, 453, 467
- torus, 14, 438
 - acting, 76
 - Néron-Severi, 444
 - norm 1, 499
 - split, 499
 - standard 1-, 439
 - standard n -, 14
- torus action
 - complexity 1, 197, 202, 281, 282, 330, 405
- total coordinate space, 1, 63, 66, 82, 144, 453
- total space of a line bundle, 210
- transcendental Brauer-Manin obstruction, 463
- transcendental lattice, 358
- trivial X -torsor, 442
- trivial extension of a graded algebra, 9
- tropical variety, 204
- true δ -collection, 97
- true \mathfrak{F} -bunch, 143
- true γ -collection, 97
- true \mathcal{V} -(quasi)-fan, 91
- true \mathcal{V} -collection, 98
- true \mathcal{W} -bunch, 92
- true \mathcal{W} -collection, 98
- true lattice bunch, 111
- true lattice collection, 110
- twist, 443
- twisted action, 441
- type
 - Calabi-Yau, 274
 - Fano, 273
 - of a del Pezzo surface, 471
 - of a torsor, 444
- unique factorization, 32, 55, 72, 73, 151, 156, 192, 193
- unirational variety, 258
- universal torsor, 444, 453, 467
- vanishing theorem
 - Kawamata-Viehweg, 337
- variety
 - A_k -, 209
 - G -, 16
 - \mathbb{Q} -factorial, 28
 - affine, 14
 - almost Fano, 465
 - almost homogeneous, 297
 - Brauer group of a , 457
 - divisorial, 240
 - factorial H -, 53
 - Fano, 179
 - flag, 151
 - geometrically connected, 438
 - geometrically irreducible, 438
 - geometrically rational, 438
 - irrelevant ideal of a , 64
 - normal, 25
 - over \mathbb{K} , 437
 - pre-, 14
 - spherical, 314
 - toric, 75
 - unirational, 258
 - weight cone of a , 124
 - wonderful, 320
- Veronese subalgebra, 10, 12, 22, 69, 73, 130, 154, 189, 237
- very ample divisor, 29
- very general position, 277
- weak B -lifting, 222
- weak approximation, 454
- weak automorphism, 252
- weak del Pezzo surface, 367, 469
- weak tropical resolution, 204
- weakly accumulating subvariety, 465
- weakly tropical, 204
- weight
 - dominant, 152, 321
 - exceptional, 229
 - fundamental, 152, 324
 - highest, 152, 321, 382
 - of a homogeneous function, 16

Cambridge University Press

978-1-107-02462-5 - Cox Rings

Ivan Arzhantsev, Ulrich Derenthal, Jürgen Hausen and Antonio Laface

Index

[More information](#)

530

Index

weight cone

of a graded algebra, 8

of a variety, 124

weight group, 8

weight monoid, 8

Weil divisor, 24

ample, 174

movable, 173

semiample, 174

support of a, 26

Weil height, 464

Weil restriction, 440

Weyl chamber, 152

Weyl group, 329, 358, 432, 470, 481

wonderful compactification, 323

wonderful variety, 320

Zariski decomposition, 184, 338