

Index

- accountability of the physician 7
 - truing required
 - by patients 129–30
- addiction *see* case studies, Mary
- adjustment disorder *see* case studies, Neal
- affordability of MPCP care 161–2
- agendas, influence on clinical
 - interactions 69
- algorithms to guide treatment 25, 36–7
 - limitations in clinical practice 51
 - limitations in complex cases 63
 - manualized protocols 49–50
- American Academy of Family Physicians 4
- American Diabetes Association 5–6
- anxiety *see* case studies, Lucy; case studies, Nathan
- asymptote concept 13
- attention deficit disorder *see* case studies, Victor
- attention deficit/hyperactivity disorder (ADHD)
 - see* case studies, Larry; case studies, Neal
- Beck Depression Inventory-II (BSI-II) 42
- best practice guidelines 25
- bipolar disorder 10–12
 - see also* case studies, Stanley; case studies, Victor
- care delivery
 - appropriate cases for MPCP care 150–3
 - comprehensive treatment protocol 145–7
 - implementation of the MPCP model 167–8
 - influence of the MPCP 134–9
 - models of care 3–4
 - MPCP model 3–4
 - MPCP role in complex cases 148
- case studies
 - Alan 51–4, 59, 63–4, 69, 71–2, 75, 81, 85, 110–11, 131
 - Alison 106–7, 109, 111–12
 - Diane 38
 - Dorothy 38–9, 127–8
 - examples of appropriate MPCP cases 150–3
 - Flora 39–40
 - Hilary 141–3
 - John 51
 - Joshua 40–1
 - Keith 27–32, 34, 85–6, 88–9, 92, 108–9, 127, 130
 - Larry 134–9, 143–4
 - Lucy 75–6
 - Margie 45–7
 - Marty 111–12, 116, 129–30
 - Mary 44–5, 47
 - Michael 119–22
 - Natalie 112–17, 122, 127
 - Nathan 107–8, 123–5, 127, 130, 134–9, 143, 146–7
 - Neal 54–64, 70–2, 77–9, 83–4, 87–9, 98–102, 126, 139–40
 - Nel 30–4
 - Randy 22–3
 - Solomon 65–6, 72, 77, 83–5, 91–2, 94–8, 102–6, 110, 139–40
 - Stanley 9–10, 14–16, 151–3
 - Victor 42–3
- childhood sexual abuse *see* case studies, Natalie
- chronic fatigue syndrome *see* case studies, Joshua
- clinical accuracy
 - contribution of truing devices 127–9, 142
 - use of truing measures 139–40
- clinical assessment, false positives and false negatives 38–41
- clinical complexity 1–3, 148
 - and clinical judgment 76
 - and the MPCP role 133
 - and treatment strategy 66–8
- dimensions of 19–23
- sources of 19–20
- use of the SO framework 76–9
- see also* case studies, Alan; case studies, Neal; case studies, Solomon

Cambridge University Press

978-1-107-02515-8 - Comprehensive Care for Complex Patients: The Medical–Psychiatric Coordinating Physician Model

Steven A. Frankel, James A. Bourgeois and Philip Erdberg

Index

[More information](#)

- clinical decision making 49–51, 76
 - clinical judgment 8
 - incremental validity 48
 - practical dilemma for physicians 82
 - see also* treatment decisions
- clinical experience, and the MPCP role 79–80
- clinical intuition
 - and experience-based clinical judgment (EBCJ) 78
 - and physician choices 79–80
 - versus strategy 71–2
- clinical judgment 49
 - and clinical complexity 76
 - and clinical strategy 9–13, 90
 - and intuition 78–80
 - and physician choices 79–80
 - and the MPCP role 85–6
 - effects of failure 85–6
 - interaction with truing measures 130–1
 - linking with truing measures 123–6
 - objective and subjective components 8
 - role of clinical judgment 85
 - selection of truing mechanisms 88–9
 - successful case 87–8
 - support from truing measures 79–80, 84–5
- clinical measurement, successive approximations 13
- clinical practice
 - contributions from the patient 73
 - contributions from the physician 73
 - factors affecting precision 130–1
 - influence of subjectivity 74
 - inherent complexity 65
 - limitations of algorithms 51
 - list of factors at play 72–4
 - non-verbal aspects 73
 - outcome measures 89
 - person-specific variables 14–16
 - realities of 146–7
 - role of subjectivity 14–16
 - use of algorithms (manualized protocols) 49–50
- clinical progress
 - and the conjunctive sequence 126–7
 - continuities and discontinuities 126–7
 - interpersonal experiences 126–7
 - monitoring 143–5
- clinical standards 16–17
- clinical strategy
 - and clinical complexity 66–8
 - and clinical intuition 71–2
 - and clinical judgment 9–13, 90
 - and interpersonal collaboration 70–1
 - and the physician–patient relationship 91–102
 - complex interpersonal aspects 91–3
 - definitions 91
 - emerging complications 93–4
 - in psychiatry (Neal) 98–102
 - leadership role 70–1
 - MPCP perspective 91–3, 102–4
 - primary care interface with psychiatry (Solomon) 94–8
 - subjective complications 93–4
 - support from truing methods 102
- clinical testing
 - ambiguity in 37–8
 - as a truing device 47
 - truing methods 37
- clinical tools, range of tools available 48–9
- clinical truth, and truing devices 13
- CNS-based illness and psychiatry 5
- cognitive behavioral therapy (CBT) 50–1, 55
- collaborative care xi–xii
 - and the MPCP 63–4
 - coordination by the MPCP 102–4, 156–7
 - Level 2 truing by the MPCP 140–1
 - see also* physician–patient collaboration; treatment team
- communication, role of the MPCP 53–4, 85
- co-morbid systemic medical and psychiatric illnesses 1–2
- complex cases
 - and the MPCP model of care 3–4, 148
 - definition 17–18
 - examples of appropriate MPCP cases 150–3
 - limitations of evidence-based treatment protocols 63
 - models of care 3–4
 - need for the MPCP 1–2
 - steps in evolution of 54–64
 - truing sequence in a complex case 54–64
 - utilization of healthcare resources 1–2
- complex partial seizures *see* case studies, Larry
- complex patients, definition 148
- complexity
 - dimensions in clinical work 2–3
 - see also* clinical complexity; diagnostic complexity; management complexity; operational complexity
 - comprehensive personality and neuropsychological assessment components of 42
 - unlocking complex cases 42–7

Cambridge University Press

978-1-107-02515-8 - Comprehensive Care for Complex Patients: The Medical–Psychiatric Coordinating Physician Model

Steven A. Frankel, James A. Bourgeois and Philip Erdberg

Index

[More information](#)**180** Index

- comprehensive treatment protocol, and the MPCP 145–6
- conjunctive sequence
 - and clinical progress 126–7
 - at MPCP level 135
 - physician–patient alliance 120–2
- consensus *see* physician–patient consensus
- continuous quality improvement 26
- coordinating physician model 4, 133, 149, 155
- costs, impact of MPCP management 161–2
- countertransferences, influence on clinical interactions 69
- cyclothymic disorder *see* case studies, Stanley
- decision making *see* clinical decision making; treatment decisions
- decision trees 25 *see also* algorithms
- delivery of care *see* care delivery
- dependent variable in medicine 7–8
- depression *see* case studies, Mary; case studies, Neal; case studies, Victor
- depressive disorder *see* case studies, Larry
- diabetes mellitus
 - diagnostic criteria 5–6
 - integrated approach to treatment 6–8
 - see also* case studies, Michael
- diagnosis and treatment
 - approaches to 29–30
 - patient's perspective on 127–9
- diagnostic ambiguity *see* case studies, Alison
- Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV-TR) 49
- schizoaffective disorder 50–1
- schizophrenia 6
- diagnostic complexity 2–3, 38–41
- diagnostic testing *see* clinical testing
- difficult-to-manage patients 17–18
- dual-trained psychiatrists, appeal of the MPCP role 3–4
- eating disorders *see* case studies, Randy
- empirically-supported treatments (ESTs)
 - 36–7, 49–51
 - limitations 81, 84–5
- enactments, influence on clinical interactions 69
- Erdberg, Philip 58
- evidence-based guidelines 25
- evidence-based medicine (EBM) 36–7
- evidence-based treatments (EBTs) 36–7, 49–51
 - limitations 81, 84–5, 63
- executive function derangement *see* case studies, Larry
- experience-based clinical judgment (EBCJ)
 - 36–7, 49, 82, 88
 - and intuition 78
- false negatives in clinical assessment 38–41
- false positives in clinical assessment 38–41
- family issues *see* case studies, Larry; case studies, Nathan
- gambling addiction *see* case studies, Larry
- Hamilton Rating Scale for Depression 42
- healthcare resources, utilization by complex cases 1–2
- imatinib 51
- implementation of the MPCP model 167–8
- incremental validity of clinical decisions 48
- independent variables
 - accounting for 7–8
 - influence on clinical judgment 8
- individual healthcare professional functions (Level 1) 2
- inherent complexity 19–20
- innovation and the MPCP 134
- integrated treatment model of care 1–2
- interferon 51
- International Classification of Diseases (ICD-10) (WHO) 49
- schizoaffective disorder 50
- inter-subjectivity between physician and patient 68
- intuition *see* clinical intuition
- laboratory testing *see* clinical testing
- leadership in the clinical situation 70–1
- Level 1 (microscopic) functions of the MPCP 2
- Level 2 (treatment team/macroscopic) functions of the MPCP 2, 26–9, 130–1, 140–1, 145–6
- management challenge *see* case studies, Alan
- management complexity 2–3, 148
- management-intensive, complex patients
 - 3, 17–18
- medical model 29–30
- medical–psychiatric coordinating physician (MPCP)
 - affordability 161–2
 - and patient-centered treatment 71
 - application of the conjunctive sequence 135
 - as a medical subspecialization 145, 148–9, 158–9

Cambridge University Press

978-1-107-02515-8 - Comprehensive Care for Complex Patients: The Medical–Psychiatric Coordinating Physician Model

Steven A. Frankel, James A. Bourgeois and Philip Erdberg

Index

[More information](#)

- becoming an MPCP 157–8
- benefits in complex cases 3–4, 146–8
- career paths 148–9
- clinical experience required 79–80
- clinical judgment 85, 138–9
- clinical needs targeted 148
- communication role 85
- complexities of clinical work 133
- complexities of the role 33–4
- comprehensive treatment protocol 145–6
- conflict management within the treatment team 156–7
- contribution to the treatment process 34–5
- coordination of care delivery 17–18
- coordination of clinical strategy 91–3
- coordination of complex cases 63–4
- coordination of the multidisciplinary treatment team 148, 156–7
- desirability of psychiatric training 153–4
- evidence for need for 1–2
- examples of appropriate cases 148, 150–3
- features of the MPCP model of care 148
- functioning of the treatment team 85–6
- future development of the MPCP model 168
- implementation of the MPCP model 167–8
- in outpatient psychosomatic medicine 160–1
- influence of patient collaboration 84
- influence on the direction of treatment 134–8
- innovation 134
- integration of psychiatry and systemic medicine 4–5
- involvement in ongoing patient care 153–4
- job description 153–4
- knowledge required 32
- leadership role 34–5
- Level 1 (microscopic) functions 2
- Level 2 (treatment team/macrosopic) responsibilities 2, 26–9, 130–1, 140–1, 145–6
- limits of certainty 141–3
- medical training required 32
- modification of the treatment plan 134–9
- need for a communication facilitator 53–4
- need for formalization of the role 85–6
- perspective on clinical strategy 102–4
- pilot study of MPCP outpatient cases 162–7
- psychiatrist in the role of 3–4, 133
- realities of clinical work 146–7
- response to crisis situations 134–8
- responsibilities 2
- role in complex cases 23
- role in operationally complex cases 91–3
- skills required 32, 148–9, 154–6
- specific training for the role 142–3
- strategic role 102–4
- target clinical subpopulation 3–4
- therapeutic impact 134
- training and experience required 153–6
- training models 157–8
- treatment coordinator 130–1
- treatment monitoring role 79
- truing at Level 2 (treatment team) 26–9, 140–1
- use of outcome measures 89
- use of SO analysis 78–9
- medical specializations 158–9
- mental dysfunctions, scope of psychiatry 5
- mind-body divide, fallacy of 4–5
- Minnesota Multiphasic Personality Inventory (MMPI) 42, 45–6, 124
- models of care for complex cases 3–4
- monitoring progress 143–5
- MPCP *see* medical–psychiatric coordinating physician
- multidisciplinary treatment team *see* treatment team
- multiple sclerosis *see* case studies, Diane
- myelodysplasia *see* case studies, Alan
- myelofibrosis *see* case studies, Alan
- Myers-Briggs Type Indicator Career Report 46
- neuropsychological assessment 42, 88–9
- non-verbal aspects of the clinical process 73
- non-verbal contributions to consensus 116–17
- one-subject scientific project *see* single subject research model
- operational complexity 2–3, 20, 38–41, 148
- organ transplantation 153
- outcome criteria 89
 - collaboratively defining 123–6
 - measuring results 143–5
- outpatient psychosomatic medicine, potential for MPCP practice 160–1
- pain management *see* case studies, Stanley
- Parkinson's disease *see* case studies, Nathan
- patient care
 - integrated treatment model (MPCP) 1–2
 - place for the MPCP 1–2
 - principles of 7
- Patient-Centered Medical Home (PCMH) model 167–8
- patient-centered treatment 36–7, 70–1

Cambridge University Press

978-1-107-02515-8 - Comprehensive Care for Complex Patients: The Medical–Psychiatric Coordinating Physician Model

Steven A. Frankel, James A. Bourgeois and Philip Erdberg

Index

[More information](#)**182** Index

- patient–physician relationship
 - see* physician–patient relationship
- patients
 - accountability of physicians 129–30
 - clinical subpopulation targeted by MPCP care 3–4
 - contributions to the clinical process 73
 - personal input about treatment 107–9
 - reporting of their medical condition 127–9
 - response to diagnosis 127–9
 - suitability for MPCP care 148
- personality assessment 42
- Personality Assessment Inventory (PAI) 45, 124
- personality disorder *see* case studies, Neal
- physician
 - accountability 7
 - contributions to the clinical process 73
 - decision-making dilemma 82
 - limits of certainty 141–3
 - personal accountability to patients 129–30
 - role of intuition in clinical work 71–2
 - subjectivity 14–16
- physician–patient alliance, conjunctive sequence 120–2
- physician–patient collaboration 88–9
 - as a truing device 82, 119–20
 - consequences of not collaborating 107–9
 - defining outcome measures 123–6
 - impact on physician judgment 82–4
 - patient’s input about treatment 107–9
 - see also* case studies, Keith; case studies, Nathan
- physician–patient consensus
 - components of 117–18
 - cooperation between team members 110–11
 - how it works 109–12
 - mixed systemic medical–psychiatric case 112–17
 - non-verbal contributions 116–17
 - physician’s distortions 111–12
 - rapport in treatment 109
 - role of truing 107
 - truing measures 118
 - working consensus 117–18
- physician–patient reciprocity 70–1, 82–4
- physician–patient relationship
 - agendas 69
 - and the clinical strategy 91–102
 - contaminating factors 69
 - countertransferences 69
 - developmental stages 120–2
 - dyadic versus team-based treatment 26–9
 - effects of uncertainty and subjectivity 64
 - enactments 69
 - interpersonal collaboration 68, 70–1
 - inter-subjectivity 68
 - leadership role 70–1
 - responsibility for results 70–1
 - responsibility for treatment 68
 - shared influence and responsibility 70–1
 - transferences 69
 - treatment alliance approach 67–8
- physician self-discipline, Self-Other Rapid Assessment Method 73–5
- physician self-monitoring 59, 79–80
- pilot study of MPCP outpatient cases 162–7
- polycythemia vera *see* case studies, Keith
- posttraumatic stress disorder (PTSD) 115–16
- precision in clinical work
 - ambiguity in clinical testing 37–8
 - and choice of truing device 139–40
 - incremental validity of clinical decisions 48
 - interpersonal precision 119–27, 129–31
 - limits of certainty 141–3
 - practical dilemma for physicians 82
 - range of truing devices 13–14
 - technical precision 127–31
- primary care
 - and systemic illnesses 5
 - definition 5
 - interface with psychiatry 94–8
 - interpersonal factors 127
- primary care physician (PCP)
 - examples of complex cases 150–3
 - relevance of the MPCP model 4
- projective identification mechanism 73
- projective testing 58
- psychiatric co-morbidity, and the MPCP
 - role 133
- psychiatrically co-morbid, management-intensive, complex patients 3, 17–18, 133
- psychiatrist in the MPCP role 3–4
- psychiatry
 - and CNS-based illnesses 5
 - and mental dysfunctions 5
 - integration with systemic medicine 4–8
 - interface with primary care 94–8
- psychological assessment 42
 - as a truing device 47
 - components of 42
 - factors influencing use of 49
 - unlocking complex cases 42–7
- psychometric testing *see* clinical testing

Cambridge University Press

978-1-107-02515-8 - Comprehensive Care for Complex Patients: The Medical–Psychiatric Coordinating Physician Model

Steven A. Frankel, James A. Bourgeois and Philip Erdberg

Index

[More information](#)

- psychosomatic medicine, potential for MPCP practice 160–1
- psychosomatic psychiatrists as MPCPs 3–4
- randomized controlled trials 25
- reciprocity between physician and patient 70–1, 82–4
- risk-taking behavior *see* case studies, Michael Rorschach inkblot test 42, 45–6, 124
- schizoaffective disorder
formal diagnostic criteria 50
see also case studies, John
- schizophrenia
diagnostic criteria 6
integrated approach to treatment 6–8
scientific approach to treatment 7–8
SCL-90 (Symptom Check List-90) 42
scleroderma *see* case studies, Solomon
- Self-Other Rapid Assessment Analysis (SO Analysis) 59
use by the MPCP 78–9
- Self-Other Rapid Assessment Framework (SO framework) 73–5
use in clinically complex contexts 76–9
- Self-Other Rapid Assessment Method (SO Method) 59, 74–5, 82, 88–9
application to systemic medicine 75–6
- Self-Other Relational Configuration (SO configuration) 74–5
- self-supervision method (Casement) 76
- sexual abuse in childhood *see* case studies, Natalie
- single-subject research model 7–8, 144
- Sjögren's syndrome *see* case studies, Natalie
- SO analysis *see* Self-Other Rapid Assessment Analysis
- SO configuration *see* Self-Other Relational Configuration
- SO framework *see* Self-Other Rapid Assessment Framework
- SO method *see* Self-Other Rapid Assessment Method
- Stevens–Johnson syndrome 51
- strategy *see* clinical strategy
- Strong Interest Inventory 46
- study of MPCP outpatient cases 162–7
- subjectivity in clinical work 14–16
complicating the clinical strategy 93–4
in the physician–patient relationship 64
influence on the clinical process 74
role in treatment decisions 82
- substance abuse *see* case studies, Randy
- systemic illness and primary care 5
- systemic medicine, integration with psychiatry 4–8
- technical complexity 20
- Thematic Apperception Test (TAT) 42, 46
- transferences, influence on clinical interactions 69
- treatment algorithms *see* algorithms to guide treatment
- treatment alliance 67–8, 109
and clinical strategy 91–3
deconstruction of the situation 109–12
mixed systemic medical–psychiatric case 112–16
- treatment approach
integration of systemic medicine with psychiatry 6–8
scientific argument 7–8
- treatment decisions
impact of physician–patient collaboration 82–4
limitations of objective approaches 81
practical dilemma in complex cases 82
role of subjectivity 82
use of truing devices 82
see also clinical decision making
- treatment monitoring responsibility 143–5
- treatment plan
influence of the MPCP 134–8
modification by the MPCP 138–9
- treatment strategy *see* clinical strategy
- treatment team
complexities of 27–9
conflict management by the MPCP 156–7
cooperation between team members 110–11
coordination by the MPCP 17, 145–6, 148, 156–7
Level 2 functions of the MPCP 2, 26–9, 130–1, 140–1, 145–6
operational complexity 2–3
truing measures 26–9, 118, 140–1
- truing, definition 13
- truing devices/methods/measures 1–2, 48
and clinical truth 13
clinical testing 37, 47
conjunctive sequence 120–2
consequences of omission 30–2
contribution to clinical accuracy 88–9, 127–9, 139–40, 142

Cambridge University Press

978-1-107-02515-8 - Comprehensive Care for Complex Patients: The Medical–Psychiatric

Coordinating Physician Model

Steven A. Frankel, James A. Bourgeois and Philip Erdberg

Index

[More information](#)**184** Index

- truing devices/methods/measures (cont.)
 - contribution to physician–patient consensus 107
 - definition 25–6
 - factors influencing choice of 49
 - incremental validity 88–9
 - Level 1 (direct service providers) 26–9
 - Level 2 (treatment team) 26–9, 118, 140–1
 - linking with clinical judgment 79–80, 123–6, 130–1
 - list of devices 13–14
 - patients’ need for 129–30
 - physician self-monitoring 59
 - principle of successive approximations 25–6, 48
 - psychometric assessment 47, 49
 - range of methods 119–20
 - selection of 88–9
 - support for clinical judgment 84–5, 88–9
 - support for clinical strategy 102
 - treatment team level 26–9, 118, 140–1
 - use in treatment decision making 82
 - truing sequence in a complex case (Neal) 54–64
 - appreciation of patient’s assets 58, 63
 - approach to treatment plan formulation 58–9
 - descriptive distillation of patient’s status 59–60
 - diagnosis 60–1
 - early impressions and clinical strategy 57
 - history of present illness 56
 - initial contact and review of records 56
 - initial impression of the patient 55
 - initial intervention and treatment planning 57
 - initial interview and clinical evaluation 56
 - limitations of evidence-based treatment protocols 63
 - medication 57
 - neurological workup 57
 - neuropsychological assessment 57
 - physician decisions and actions 61–2
 - physician self-monitoring 59
 - psychometric assessments 56–8
 - role of the MPCP 63–4
- Wechsler Adult Intelligence Scale (WAIS-III) 46, 124
- working consensus 117–18