

Author Index

- Abbiss, J. B., 648
 Abraham, B., 645
 Acheson, D., 228, 642
 Adams, J. W., 197, 642
 Ahmed, N., 217, 642
 Akaike, H., 493–4, 642
 Alavi, A. S., 151, 201, 642
 al Wasel, I., 215, 645
 Amrein, M., 262–3, 642
 Ander, M. E., 644
 Andersen, N. O., 509, 642
 Anderson, T. W., 35, 37, 166, 215, 447, 503,
 540, 552, 642
 Arun, K. S., 16, 650
- Babadi, B., 359, 642
 Baggeroer, A. B., 485, 642
 Balogh, A., 645
 Barbé, K., 424, 642
 Barnes, J. A., 162, 327, 329, 642
 Barrett, J. F., 635, 642
 Bartlett, M. S., 215, 269–70, 275, 302, 344, 414,
 642–3
 Beauchamp, K. G., 138–40, 643
 Bell, B. M., 560, 643
 Beltrão, K. I., 307, 643, 648
 Beran, J., 407, 643
 Berk, K. N., 452, 485, 643
 Bishop, T. N., 35, 494, 656
 Blackman, R. B., 185, 201, 264, 343, 643
 Bloomfield, P., 92, 154, 160, 232, 237, 252, 280,
 307, 513, 526, 550, 592, 643
 Bluestein, L. I., 94
 Bogert, B. P., 227, 301, 473, 643
 Bohman, H., 278, 643
 Bollenbacher, A. F., 649
 Bolt, B. A., 204, 643
 Bondon, P., 31, 232, 653
- Boorstyn, R. R., 551, 654
 Bøvikken, E., 541, 643
 Box, G. E. P., 16, 34–5, 40, 264, 333, 431, 446,
 557, 643
 Bracewell, R. N., 55, 61, 69, 72, 137, 643
 Branch, J. L., 94, 651
 Briggs, W. L., 92, 643
 Brillinger, D. R., 175–6, 195, 203–4, 206, 215,
 262, 643, 647, 653, 656
 Brockwell, P. J., 21, 28, 40, 43, 158, 203, 333,
 404, 448, 465, 471, 473, 482–3, 485, 488,
 599–601, 615, 621, 643
 Broersen, P. M. T., 174, 492, 643
 Bronez, T. P., 355, 359, 374, 382, 424, 643–4
 Brown, E. N., 359, 642
 Bruce, A. G., 486, 644
 Bunch, J. R., 462, 644
 Burg, J. P., 457, 466–7, 471–5, 577–8, 644
 Burrus, C. S., 647
 Burshtein, D., 485, 488, 491, 644
 Businger, P. A., 564, 644
 Butler, R. P., 645
- Canavero, F. G., 578, 653
 Carter, G. C., 257, 414, 420, 644, 652
 Cartwright, D. E., 512, 522, 652
 Chaika, M., 605, 616, 655
 Chambers, J. M., 621, 644
 Chambers, M. J., 605, 644
 Champeney, D. C., 49, 53, 55, 62, 69, 84, 137,
 165, 170, 644
 Chan, G., 217, 644
 Chan, K.-S., 228, 615, 644
 Chandna, S., 30, 644
 Chatfield, C., 151, 644
 Chave, A. D., 196, 356, 419, 644, 656
 Chen, W. W., 197, 648
 Chen, W. Y., 553, 558–60, 644

- Chi, A. R., 642
 Childers, D. G., 642, 644, 648–50, 656, 658
 Chilès, J.-P., 165, 340, 644
 Choi, B. S., 492, 644
 Chonavel, T., 110, 644
 Chung, K. L., 120, 588, 644
 Clarkson, I. V. L., 653
 Clayton, R. W., 494–5, 554, 656
 Cleveland, W. S., 181, 280, 644
 Coble, P. A., 655
 Colosi, J. A., 651
 Conover, W. J., 215–6, 645
 Conradsen, K., 40, 645
 Constantine, W. L. B., 228, 230, 611, 622, 631, 645, 653
 Cooper, G. R., 485, 649
 Courant, R., 63, 645
 Craigmile, P. F., 440, 548, 601, 645, 657
 Cramér, H., 108, 645
 Crowell, D. H., 648
 Cumming, A., 529, 645
 Cutler, L. S., 642
 Czerny, B., 407, 651
- Daniell, P. J., 271, 645
 Danioux, E., 655
 Davies, N., 631–2, 645
 Davies, R. B., 219, 601, 645
 Davis, H. T., 404, 645
 Davis, R. A., 21, 28, 40, 43, 158, 203, 333, 404, 448, 465, 471, 473, 482–3, 485, 488, 599–601, 615, 621, 643
 Davison, A. C., 593, 605, 618, 631, 645
 Day, N. L., 655
 de Gooijer, J. G., 492, 494, 645
 Delfiner, P., 165, 340, 644
 Denison, D. G. T., 548, 645
 de Paula, E. R., 553, 649
 Dietrich, C. R., 601, 605, 645
 Diggle, P. J., 215, 645
 Doetsch, G., 55, 645
 Doodson, A. T., 512, 645
 Durbin, E. J., 653
 Durrani, T. S., 648
 Dzhaparidze, K. O., 307, 645
 Dzieciuch, M. A., 651
 Dziuba, S., 40
- Efron, B., 356, 593, 618, 645
 Einstein, A., 219, 645
 Elliott, A. C., 658
 Emery, W. J., 337, 645
 Enck, P., 650
 Eom, K.-B., 408, 649
 Epanechnikov, V. A., 275, 646
 Eubank, S., 656
- Fairfield Smith, H., 264, 646
 Falk, R., 3, 646
 Fan, J., 307, 646
 Farebrother, R. W., 543, 646
- Farmer, J. D., 656
 Fay, G., 215, 646
 Ferguson, T. S., 45, 646
 Figueiras-Vidal, A. R., 650
 Filliben, J., 39–40
 Findley, D. F., 651
 Fisher, R. A., 1, 539–40, 646
 Flannery, B. P., 653
 Fodor, I. K., 355, 646
 Folland, G. B., 61, 646
 Forsyth, R. J., 645
 Fougere, P. F., 181, 183, 194, 558–60, 646
 Fox, W., 321–2
 Franke, J., 474, 646
 Friedlander, B., 554–5, 646
 Frijling, D. J., 221, 315, 657
 Fuller, W. A., 164, 166, 203, 214–5, 646
- Galdrikian, B., 656
 Gao, H.-Y., 307, 412, 646
 Geçkinli, N. C., 197, 646
 Geweke, J., 408, 646
 Gneiting, T., 601, 646
 Godolphin, E. J., 484, 646
 Gold, B., 80, 145, 156, 653
 Golub, G. H., 89, 375, 464, 505, 564, 599, 644, 646
 Gong, G., 356, 593, 618, 645
 Gonzalez, R. C., 217, 647
 Gould, A., 645
 Gradshteyn, I. S., 50, 345, 647
 Grandy, W. T., Jr., 644, 646
 Granger, C. W. J., 151, 407, 633, 647
 Gray, H. L., 658
 Graybill, F. A., 375, 647
 Greene, D. H., 25, 647
 Greenhall, C. A., 355, 369, 402, 437, 441, 647
 Grenander, U., 145, 166, 185, 251, 257, 647
 Grünbaum, F. A., 94–5, 647
 Gujar, U. G., 633, 647
 Gunn, D. E., 123, 658
- Hall, P., 644
 Hamming, R. W., 145, 647
 Han, J., 655
 Hannan, E. J., 262, 404, 523, 527, 540, 647, 653
 Hannig, J., 307, 312, 647
 Hanssen, A., 355, 647
 Hansson, M., 382, 647
 Hansson-Sandsten, M., 424–5, 647
 Hardin, J. C., 123, 647
 Harish, M., 219, 652
 Harris, B., 656
 Harris, F. J., 196, 277, 535, 647
 Harris, P. E., 407, 649
 Harte, D. S., 601, 645
 Hasan, T., 160, 232, 592, 647
 Hastie, T. J., 309, 647
 Hayashi, Y., 30, 647
 Haykin, S., 652, 656–7
 He, H., 655
 Healey, D. J., 642

- Healy, M. J., 643
 Heideman, M. T., 92, 647
 Heilig, B., 655
 Henson, V. E., 92, 643
 Herrera, R. H., 655
 Hilbert, D., 63, 645
 Hill, I. D., 633, 648
 Hill, R., 648
 Hinkley, D. V., 593, 605, 618, 631, 645
 Hipel, K. W., 599–600, 639, 651
 Hogan, J. A., 359, 648
 Holder, R. L., 648
 Hosken, J. W. J., 407, 657
 Hosking, J. R. M., 407, 648
 Howe, B. M., 651
 Hua, Y., 566, 648
 Hudson, J. E., 648
 Hurvich, C. M., 197, 307, 495, 648
- Ibragimov, I. A., 165, 648
 Ihara, S., 474, 648
 Ishida, M., 298, 300, 649
 Isserlis, L., 30, 648
- Jackson, D. D., 564–6, 648
 Jackson, P. L., 559–60, 648
 Jacovitti, G., 131, 648
 Jaynes, E. T., 473, 648
 Jenkins, G. M., 151, 160, 169, 201, 251, 264,
 290, 642–3, 648
 Jensen, O. G., 407, 656
 Jessup, A., 225
 Jiménez, C., 169–70, 185, 651
 Johnson, D. H., 647
 Johnson, D. M., 140, 650
 Johnson, G. E., 636, 648
 Johnson, N. L., 633, 648
 Jones, N. B., 16, 648
 Jones, R. H., 16, 210, 404, 482, 485, 494, 510,
 555, 645, 648–9
 Joughin, I., 443, 649
 Joyeux, R., 407, 647
- Kahan, W., 564, 646
 Kaiser, J. F., 196, 649
 Kalveram, K.-T., 650
 Kane, R. P., 553, 649
 Kantz, H., 615, 649
 Kapuniai, L., 648
 Kasyap, R. L., 408, 649
 Katz, R. W., 648
 Kavanagh, R. J., 633, 647
 Kaveh, M., 485, 649
 Kay, S. M., 123, 478, 480, 485, 493–5, 502,
 506, 530, 554, 558, 560–2, 595, 598–9, 624,
 636–7, 639, 649
 Keeling, R. F., 332, 649
 Kendall, D. G., 302, 643
 Kerner, C., 407, 649
 Kesler, S. B., 647, 649, 650–1, 653–4
 Kikkawa, S., 298, 300, 649
- King, M. E., 366, 649
 Kleiner, B., 644
 Knuth, D. E., 25, 647
 Kokoszka, P., 215, 649
 Kooperberg, C., 307, 649
 Koopmans, L. H., 108, 134, 210, 649
 Koslov, J. W., 485, 649
 Kowalski, A., 140, 650
 Kreiss, J.-P., 631, 650
 Kreuzberger, E., 307, 646
 Krishnaiah, P. R., 647
 Kromer, R. E., 485, 490, 650
 Kubrick, S., 215, 219, 228, 297, 307–9, 311–12,
 337, 540, 615, 644–6, 650, 652–3
 Kullback, S., 297, 345, 650
 Kumaresan, R., 564, 566–7, 656
 Kung, S. Y., 16, 650
 Künsch, H. R., 262–3, 642
 Kuo, C., 366, 650
 Kuo, F. F., 649
 Kürster, M., 529, 544, 658
- Lacoss, R. T., 494–5, 650
 Lago, P. J., 648
 Lagunas-Hernández, M. A., 474, 650
 Lahiri, S. N., 215, 631, 650
 Lakey, J. D., 359, 648
 Lampard, D. G., 635, 642
 Landers, T. E., 494–5, 650
 Lang, S. W., 32, 650
 Lanning, E. N., 140, 650
 Lawrance, A. J., 485, 650
 Lee, T. C. M., 296, 307, 312, 647, 650
 Leeson, D. B., 642
 Leibler, R. A., 297, 345, 650
 Li, J., 655
 Li, T.-H., 529–30, 650
 Lii, K. S., 370, 650
 Lilly, J. M., 655
 Lindberg, C. R., 366, 650, 652
 Linnik, Yu. V., 165, 648
 Liu, B., 633–4, 636, 650
 Liu, T.-C., 355, 650
 Ljung, G. M., 643
 Lomb, N. R., 528, 650
 Longbottom, J., 632, 650
 Longtin, A., 656
 Lopes, A., 230, 650
 Lopes, S., 650
 Lotz, S., 655
 Loupas, T., 30, 650
 Lynch, D., 76–8, 178–9, 192, 200–1, 218–9, 294,
 300–1, 306, 337, 382, 458–9, 489
 Lysne, D., 466, 484, 650
- MacDonald, G. J. F., 407, 652
 MacNeill, I. B., 643
 Madan, R. N., 648
 Maitani, T., 30, 651
 Makhoul, J., 28, 156, 473–4, 493, 555–6, 649,
 651
 Mallows, C. L., 633, 651

- Mann, H. B., 486, 651
 Mann, W. R., 25, 656
 Manolakis, D. G., 241, 653
 Marcy, G. W., 645
 Marple, S. L., Jr., 478, 480, 494–5, 506, 530,
 554, 558, 560–2, 649, 651
 Martin, R. D., 196, 227, 486, 492, 644, 651
 Matsakis, D., 326, 593, 651
 McClellan, J. H., 32, 650
 McCloud, M. L., 423, 651
 McCoy, E. J., 408–9, 412, 651, 657
 McCullagh, P., 308, 651
 McDicken, W. N., 30, 650
 McGunigal, T. E., 642
 McHardy, I., 407, 651
 McKilliam, R. G., 653
 McLeod, A. I., 169–70, 185, 599–600, 639, 651
 McPherson, D. A., 605, 651
 McQuarrie, A. D. R., 477, 651
 McWhirter, J. G., 648
 Mercer, J. A., 330, 651
 Mikosch, T., 215, 649
 Miller, K. S., 30, 518, 546, 651
 Miller, R. G., 356, 651
 Mitchell, R. L., 605, 651
 Mix, A. C., 124, 653
 Mohr, D. L., 408, 651
 Mombeni, H., 307, 651
 Monro, D. M., 94, 651
 Moon, F. C., 228, 651
 Moore, T. A., 648
 Morettin, P. A., 140, 462, 651–2
 Moses, R., 160, 232, 492, 592, 655
 Moulin, P., 307, 412, 652
 Mullen, J. A., Jr., 642
 Mullis, C. T., 359, 444, 651–2, 654
 Munk, W. H., 407, 512, 522, 652
 Munson, D. C., Jr., 633–4, 636, 650
 Murphy, A. H., 648
 Murray, M. T., 520, 652
 Musial, F., 650
- Nadarajah, S., 651
 Nakagawa, J. K., 648
 Narasimhan, S. V., 219, 652
 Natarajan, T., 642
 Neave, H. R., 277, 652
 Nelder, J. A., 308, 651
 Nelsen, R. B., 3, 652
 Newsam, G. N., 601, 605, 645
 Newton, H. J., 35, 170, 451, 463–4, 470, 480,
 483–5, 528, 652
 Nicewander, W. A., 3, 654
 Nicholls, D. F., 404, 647
 Nitzberg, R., 474, 652
 Nowroozi, A. A., 540, 652
 Nuttall, A. H., 197, 257, 414, 420, 652
 Nyquist, H., 405, 440, 653
- O’Sullivan, F., 307, 653
 Olhede, S. C., 655
 Ombao, H. C., 297, 307–9, 311–12, 652
- Ooe, M., 494, 656
 Oppenheim, A. V., 84, 92, 94, 579, 652
- Pagano, M., 484–5, 528, 652
 Paparoditis, E., 631, 650
 Papoulis, A., 21, 114, 278, 452, 457, 463, 562,
 579, 631, 634, 652
 Parekh, A., 648
 Park, J., 359, 366, 650, 652
 Parzen, E., 251, 275, 280, 644, 652
 Pawitan, Y., 307, 653
 Percival, D. B., 307, 408, 412, 560, 605, 611,
 622, 631, 643, 651, 653–4, 657
 Picinbono, B., 31, 232, 636, 653
 Pignari, S., 578, 653
 Pillai, S. U., 21, 114, 562, 579, 631, 634, 652
 Pintelon, R., 642
 Piper, S. C., 649
 Pisarenko, V. F., 554, 653
 Pisasias, N. G., 124, 653
 Politis, D. N., 631, 653
 Pollak, H. O., 66, 655
 Poor, H. V., 650
 Porter-Hudak, S., 408, 646
 Poskitt, D. S., 644
 Prescott, P., 38, 616, 657
 Press, H., 197, 653
 Press, W. H., 506, 528, 542, 653
 Priestley, M. B., 21, 37, 108, 110, 113–5, 117–8,
 213, 240, 250–2, 256, 269, 275, 292–3, 404,
 412, 446, 448–9, 463, 477, 516, 519, 522,
 534, 543–4, 588, 653
 Proakis, J. G., 241, 653
 Pukkila, T., 405, 440, 653
- Quinn, B. G., 526, 528, 540, 653
- Rabiner, L. R., 80, 145, 156, 447, 653
 Rabinowitz, P., 599, 653
 Radoski, H. R., 646
 Ralston, A., 599, 653
 Ramsey, F. L., 462, 653
 Rao, C. R., 514, 588, 653
 Rao, K. R., 642
 Ray, B. K., 407, 654
 Raz, J. A., 652
 Reid, J. S., 485, 654
 Reinsel, G. C., 643
 Rezaei, S., 651
 Rice, J. A., 527, 654
 Rice, S. O., 298, 654
 Richardson, G. A., 655
 Riedel, K. S., 355, 391–2, 400, 412, 434, 654
 Rife, D. C., 535, 551, 654
 Roberts, R. A., 444, 654
 Robinson, L., 645
 Rogers, J. L., 3, 654
 Rosenblatt, M., 145, 166, 257, 370, 463, 494,
 527, 643, 647, 650, 654
 Rovine, M. J., 3, 654
 Rowe, D. B., 30, 654
 Rutman, J., 431, 654
 Ryzhik, I. M., 50, 345, 647

- Sakai, H., 451, 485, 654
 Salomonsson, G., 382, 647
 Samarov, A., 166, 654
 Sandgren, N., 307, 655
 Santamaría-Perez, M. E., 650
 Sarkar, T. K., 566, 648, 656
 Satorius, E. H., 553, 654
 Satterthwaite, F. E., 264, 654
 Scarano, G., 131, 648
 Scargle, J. D., 528–9, 654
 Schafer, R. W., 84, 92, 94, 447, 579, 652–3
 Scharf, L. L., 30–1, 232, 359, 366, 651–2, 654, 657
 Scheffé, H., 486, 654
 Scher, M. S., 655
 Schmidt, L., 326
 Schmitz, A., 615, 654
 Schoukens, J., 642
 Schreiber, T., 615, 649, 654
 Schreier, P. J., 30–1, 232, 654
 Schuster, A., 539, 654
 Selén, Y., 492, 655
 Serroukh, A., 636, 654
 Shimshoni, M., 571, 654
 Shumway, R. H., 21, 203, 404, 655
 Sidorenko, A., 355, 391–2, 400, 412, 434, 654
 Siegel, A. F., 541–2, 655
 Singleton, R. C., 94, 655
 Sitaram, A., 61, 646
 Sjöholm, P. F., 301, 304, 655
 Slepian, D., 57–8, 61–3, 66, 85, 87–8, 105, 384, 655
 Sloane, E. A., 195, 371, 655
 Smith, C. R., 644, 646
 Smith, W. L., 642
 Soeda, T., 654
 Sondhi, M. M., 633–4, 636, 655
 Soulier, P., 215, 646
 Souza, R. R., 650
 Spann, M., 95, 658
 Spedding, T., 645
 Spencer-Smith, J. L., 33, 503, 655
 Spliid, H., 40, 645
 Stark, P. B., 355, 646
 Stegen, G. R., 553, 558–60, 644
 Stein, M. L., 340, 655
 Stephen, R., 651
 Stephens, M. A., 216, 655
 Stevens, W. L., 541, 655
 Stoffer, D. S., 21, 140, 203, 404, 655
 Stoica, P., 160, 232, 307, 492, 528, 592, 655
 Stone, C. J., 649
 Strang, G., 217, 655
 Strawderman, R. L., 652
 Subba Rao, S., 653
 Subba Rao, T., 653
 Sun, T. C., 605, 616, 655
 Sutcliffe, P. R., 553, 655
 Swanepoel, J. W. H., 618, 624–5, 631, 638, 655
 Sydnor, R. L., 642
 Sykulski, A. M., 30, 655
 Szegő, G., 185, 647
 Taqqu, M. S., 166, 654
 Tary, J. B., 553, 655
 Tavella, P., 593, 651
 Taylor, A. E., 25, 656
 Teukolsky, S. A., 653
 Theiler, J., 615, 656
 Therrien, C. W., 464, 509, 656
 Thomas, J. B., 658
 Thompson, R., 605, 656
 Thomson, D. J., 176, 196, 211–12, 227, 351–2, 355–7, 359, 366, 374, 381–3, 386, 390–1, 415–6, 424, 434, 492, 512, 544, 547–9, 574, 582, 644, 650–2, 656
 Thomson, R. E., 337, 645
 Tiao, G. C., 656
 Tibshirani, R. J., 309, 647
 Titchmarsh, E. C., 78, 120, 167, 656
 Tjøstheim, D., 466, 484, 650
 Todd, H. A. C., 33, 503, 655
 Toodeschuck, J. P., 407, 656
 Tokumaru, H., 654
 Toman, K., 558, 656
 Tong, H., 228, 554–5, 615, 644, 656
 Traganitis, A. P., 658
 Truong, Y. K., 649
 Tsai, C.-L., 477, 495, 648, 651
 Tsakiroglou, E., 201, 656
 Tsay, R. S., 407, 654
 Tseng, F. I., 535, 656
 Tufts, D. W., 564, 566–7, 656
 Tugnait, J. K., 555, 656
 Tukey, J. W., 160, 174, 185, 197–8, 201, 264, 343, 592, 643, 653, 656
 Tukey, P. A., 644
 Ulrych, T. J., 35, 494–5, 553–4, 559–60, 656
 Umphrey, G. J., 643
 Unwin, J. M., 484, 646
 van der Baan, M., 655
 Van Loan, C. F., 89, 375, 464, 506, 564, 599, 646
 Van Schooneveld, C., 221, 315, 657
 Van Veen, B. D., 355, 366, 650, 657
 Van Wyk, J. W. J., 618, 624–5, 631, 638, 655
 Vernon, F. L., III., 652
 Vessot, R. F. C., 642
 Vetterling, W. T., 653
 Vincent, G. A., 535, 654
 von Eye, A., 3, 654
 von Sachs, R., 652
 von Storch, H., 215, 335, 657
 Wahba, G., 301, 306, 657
 Wald, A., 486, 651
 Walden, A. T., 30, 38, 194, 196, 201, 210, 213, 232, 262–3, 280, 290, 297, 300, 307, 353, 359, 366, 400, 402, 405, 407, 410, 412, 440, 512, 522, 542, 548, 616, 633, 644–5, 650–1, 653–4, 656–7
 Walker, A. M., 526, 554, 657
 Walker, J., 16, 657

666

Author Index

- Walker, J. S., 649
 Wang, X. L., 658
 Warburg, H. D., 512, 645
 Watson, W., 645
 Watts, D. G., 160, 169, 290, 648
 Wei, L., 548, 657
 Weiner, D. D., 656
 Weinstein, E., 485, 488, 491, 644
 Weisberg, S., 408, 440, 513, 516, 543, 549, 552,
 588, 621, 657
 Weiss, G., 484–5, 657
 Welch, B. L., 264, 657
 Welch, P. D., 351, 413, 415, 658
 Well, A. D., 3, 646
 Wen, Q. H., 544, 658
 White, R. E., 262, 290, 297, 300, 650, 657
 Whittle, P., 307, 526, 543, 588, 658
 Wiener, N., 78, 658
 Wilson, R., 95, 658
 Winkler, G. M. R., 642
 Wise, G. L., 635, 658
 Wong, A., 658
- Wood, S., 39
 Woods, J. W., 601, 658
 Woods, R. E., 217, 647
 Woodward, W. A., 610, 658
 Worcester, P. F., 651
 Wright, J. H., 407, 658
 Wunsch, C., 123–4, 658
- Yaglom, A. M., 21, 165, 214, 219, 307, 340,
 645, 658
 Yajima, Y., 408, 658
 Yavuz, D., 197, 646
 Yuen, C. K., 194–5, 658
- Zawalick, E. J., 646
 Zechmeister, M., 529, 544, 658
 Zeidler, J. R., 553, 654
 Zhang, H.-C., 461, 658
 Zwiers, F. W., 215, 335, 657

Subject Index

- 100% cosine data taper, 189
- absolute deviations sinusoidal amplitude estimator
 (Laplace method), 529
- absolutely
 continuous, 120
 summable, 165
- acausal
 autoregressive (AR) process, 446, 506
 filter, 147–8, 152, 155
 purely
 autoregressive (AR) process, 447–8
- ACF (*see* autocorrelation function)
- acoustic properties of human speech
 AR model, 447
- ACLS (*see* approximate conditional least squares)
- ACS (*see* autocorrelation sequence)
- ACVF (*see* autocovariance function)
- ACVS (*see* autocovariance sequence)
- adaptive multitaper spectral estimator, 389
 ACVS issue, 615
 distribution of, 390
 EDOFs, 390
 ocean wave data, 427–8
 Parseval's theorem in expectation, 390, 439
- aggregated processes, 42
- AIC (*see* Akaike's information criterion)
- AICC (*see* Akaike's information criterion corrected for bias)
- Akaike's information criterion, 494
 corrected for bias (AICC), 495
 order selection, 494
 centered time series, 495
- Akaike's information criterion corrected for bias
 (AICC), 495
 order selection, 495
 centered time series, 495
 Willamette River data, 575–6
- aliases, 82
- aliasing, 83, 108, 113, 122–3, 569
 and averaging, 161
 antialiasing filter, 57, 224
 different sampling intervals example, 131
 discrete time/continuous frequency, 81
 sea level residuals, 522
- amplitude spectrum, 54
- analytic series, 114, 561, 578–9
 imaginary part of, 579
 Willamette River data, 578, 592
- angular frequency, 8
 antialiasing filter, 57, 224
- applications of spectral analysis, 16–7
- approximate conditional least squares (ACLS)
 conditions needed for ACLS to become ECLS
 complex-valued case, 517
 real-valued case, 515
 sinusoidal amplitude estimator, 515, 525, 549,
 569–70, 572–3, 585, 589
 white noise variance estimator, 549, 570, 589
- approximate least squares, 516, 585
- approximate least squares sinusoidal amplitude
 estimator
 expected value of, 516, 585
- AR (autoregressive) process (*see* autoregressive
 (AR) process)
- AR(1) process
 asymptotic stationarity and causality, 448–9
 autocovariance sequence, 44
 backward least squares autoregressive parameter
 estimator, 510
 Burg autoregressive spectral estimator, 510
 forward least squares autoregressive parameter
 estimator, 510
 from sampling a continuous parameter/time
 process, 507
 infinite-order moving average form, 44
 maximum likelihood estimator, 482

- model for colored noise, 335
- parameter estimation from estimated ACVS, 335–6
- simulation of Gaussian
 - via circulant embedding, 639–40
 - via Gaussian spectral synthesis method (GSSM), 639–40
 - via Kay's method, 639
- Yule–Walker autoregressive parameter estimator, 510
- AR(2) process, 34, 168, 172–3
 - alternative models, 440
 - asymptotic time series bandwidth, 347
 - bias in periodogram, 177
 - Burg autoregressive spectral estimator, 509
 - cumulative periodogram test, 216
 - direct spectral estimator, 186
 - and variability, 208
 - discretely smoothed periodogram, 309–10, 348–9
 - form of FBLS parameter estimator, 510
 - innovation variance estimators, 406–7
 - lag window spectral estimator, 294, 348–9
 - log SDF estimator, 304–5
 - maximum likelihood autoregressive spectral estimator, 489
 - numerical examples
 - Burg's algorithm, 509
 - periodogram, 172–3, 510
 - Yule–Walker estimator, 507–8
 - simulated examples, 34
 - direct spectral estimator properties, 242
 - periodogram properties, 242
 - simulation of Gaussian
 - via circulant embedding, 602–3
 - via Gaussian spectral synthesis method (GSSM), 608–9
 - via Kay's method, 597
 - tapering, 195
 - unimodality of spectrum and bandwidth of time series, 300
 - WOSA estimator, 416–7
 - Yule–Walker autoregressive spectral estimator, 509
- AR(4) process, 35, 168, 172–3
 - and end-point matching, 239
 - asymptotic time series bandwidth, 347
 - bias in periodogram, 178
 - Burg autoregressive spectral estimator, 468–9, 509
 - DCT-based periodograms, 218
 - direct spectral estimator, 186
 - discretely smoothed direct spectral estimator, 310–11, 348–9
 - eigenspectra, 361, 363, 368, 395, 397, 437
 - FBLS autoregressive spectral estimator, 478, 509
 - innovation variance estimators, 406–7
 - lag window spectral estimator, 295, 301, 349
 - log SDF estimator, 305–6
 - low-rank weighted multitaper spectral estimator, 422–3
 - maximum likelihood autoregressive spectral estimator, 482–3, 490
 - multitaper spectral estimator, 358–9, 364–5, 398–9, 400, 419
 - multitaper versus WOSA spectral estimation, 418–9
 - nonunimodality of spectrum and bandwidth of time series, 300–1
 - numerical examples
 - Burg's algorithm, 468–9, 509
 - periodogram, 172–3
 - Yule–Walker estimator, 458–9, 461–2, 509
 - prewhitening, 198–201
 - SDF and expectation of direct spectral estimator, 193
 - simulated examples, 34
 - direct spectral estimator properties, 242
 - periodogram properties, 242
 - simulation of Gaussian
 - via circulant embedding, 640–1
 - via Gaussian spectral synthesis method (GSSM), 606, 641
 - via Kay's method, 638–9
 - smoothing window
 - leakage, 289
 - parameter choice, 292
 - tapering, 185
 - WOSA estimator, 417–8
 - Yule–Walker autoregressive spectral estimator, 458–9, 461–2, 509
 - Yule–Walker versus periodogram spectral estimators, 508
- ARIMA (autoregressive, integrated, moving average) process, 16
- ARMA (*see* autoregressive moving average (ARMA) process)
- ARMA(1,1) process, 44
 - autocovariance sequence (ACVS), 44
 - infinite-order moving average form, 44
 - kurtosis, 632
 - skewness, 632
 - inequality, 633, 641
 - variance, 632
- asymptotic equivalence of autoregressive estimators, 484
- asymptotic relative efficiency, 165
- asymptotic stationarity and causality, 448–9
- asymptotically efficient estimator, 166, 522–3
- atmospheric CO₂ data, 332–3
- atomic clock data, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
- atomic clock fractional frequency deviates, 326, 430
 - associated time differences, 325–6
 - direct spectral estimator, 327
 - periodogram, 327
 - Burg autoregressive spectral estimator, 501
 - innovation variance estimation, 431–2, 499–500

- simulation, 628–9
- lag window spectral estimator, 328, 501
- multitaper spectral estimator, 501
- periodogram 328
- atomic clock timekeeping, 162, 593
- augmented Yule–Walker equations, 452
- autocorrelated cosine lag window, 284, 345
- autocorrelation, 72, 96–8, 100, 102, 112
- autocorrelation function (ACF), 27
- autocorrelation method, 480
- autocorrelation sequence (ACS), 4, 9, 27
 - conjugate symmetry, 30
 - estimation, 5, 19–20
 - MA(2) process, 159
 - sample, 3
 - white noise, 19
- autocorrelation width, 73, 97–8, 100, 102
 - smoothing window, 251
 - spectral window, 191–2, 194
- autocovariance function (ACVF), 27, 38
 - and the integrated spectrum, 114
 - and the SDF, 114
 - conjugate symmetry, 30
 - Gaussian-shaped, 131
 - positive semidefinite, 30
 - symmetry, 27
- autocovariance sequence (ACVS), 27, 29
 - and integrated spectrum, 111
 - ARMA process, 44, 600
 - AR process, 44, 460, 508
 - band-limited white noise, 379, 604
 - biased estimator of, 2–7, 166, 168–9, 232, 295, 505, 561, 591
 - for complex-valued process, 231, 562
 - positive definite, 451
 - classification examples, 121
 - complex-valued process, 29, 42
 - corresponding to
 - direct spectral estimator, 188, 611–12
 - lag window spectral estimator, 248, 612
 - multitaper estimator, 433
 - periodogram, 170–1
 - WOSA estimator, 420, 435
 - Yule–Walker autoregressive spectral estimator, 504, 508
 - decay to zero, 120
 - estimation of, 166, 231
 - for process squared, 43
 - fractionally differenced (FD) process, 440
 - GSSM-based approximation, 606–7, 640
 - harmonic process, 36–7, 117, 591
 - complex-valued, 519, 562
 - inverse Fourier transform of SDF, 111, 126
 - moving average process, 32
 - MA(2), 159
 - piecewise-constant SDF, 348
 - positive semidefinite, 28, 30, 43
 - power-law process, 391, 605
 - randomly phased sinusoid, 35, 511, 553, 598
 - subsampling, 130
 - symmetry and conjugate symmetry, 27, 29
 - unbiased estimator of, 166, 168–9, 236–7, 271
 - for complex-valued process, 562
 - versus spectral density function, 124
 - white noise, 31, 117
 - Wold’s theorem, 117, 126
- autoregressive (AR) process, 33, 445–6
 - acausal, 446, 506
 - acoustic properties of human speech, 447
 - ACVS, 460, 508
 - asymptotic stationarity and causality, 448–9
 - causality, 446, 448, 506, 618, 638
 - and parameter estimators, 448, 617
 - assessment, 463
 - conditions for, 34, 446
 - characterization, 509
 - order, 446
 - order selection (*see* order selection for autoregressive process)
 - plus white noise is an ARMA process, 554, 591
 - pseudo-, 553
 - purely acausal, 447
 - simulation, 595
 - zero initial conditions, 595
 - simulation of Gaussian
 - stationary initial conditions, 595, 599, 624, 637
 - via Kay’s method, 595–7
 - via McLeod and Hipel’s method, 599
 - sinusoidal variations, 447
 - spectral density function (SDF), 144–5, 446, 485, 609, 617–8
 - stationarity condition, 34, 145, 446, 448, 557
- autoregressive estimator
 - and prewhitening, 491, 506
 - asymptotic equivalences, 484
 - Burg’s algorithm, 466, 505
 - complex-valued process, 502
 - least squares estimators, 475, 478, 505
 - maximum entropy method, 471
 - maximum likelihood estimators, 480, 506
 - pure parametric approach, 506
 - variability assessment
 - via large-sample theory, 485
 - via simulation, 626
 - Yule–Walker method, 449–51, 505
 - complex-valued process, 502
- autoregressive moving average (ARMA) process, 35, 492, 597–8
 - ACVS, 600
 - AR process plus white noise, 159, 554, 591
 - causal, 600, 631
 - maximum entropy, 473–4
 - pseudo-AR process plus white noise, 554
 - SDF, 145, 159
 - simulation of Gaussian, 593–4, 597–8
 - stationary initial conditions, 599–600, 637, 639
 - via Kay’s method, 598, 639
 - via McLeod and Hipel’s method, 600, 639
 - simulation of non-Gaussian, 631, 638
 - simulation via step-down scheme, 461
 - with distinct AR and MA coefficients, 554, 591
 - with identical AR and MA coefficients, 554

- autoregressive moving average (ARMA) spectral estimator, 492, 504
- autorelation sequence, 29, 31
- average value of a function, 71

- background continuum, 511, 513, 523, 544, 547, 549, 551, 553, 583, 588
- backward least squares (BLS)
 - autoregressive parameter estimator, 477, 484
 - AR(1) example, 510
- backward linear prediction, 455
 - complex-valued process, 502, 564
- backward linear prediction error, 455
- band-limited function, 57–8, 93, 139
 - extrapolation of, 66
 - recovering the continuous from the sampled, 84
- band-limited white noise, 128, 131, 379, 604
- band-pass filter, 146
- band-pass process, 299, 347
- bandwidth
 - of AR(2) and AR(4) time series, 347
 - of spectrum, 292, 296, 346
 - estimation of, 297
 - of time series, 300–1, 321–2, 347
 - estimation of, 300
 - regularization, 357, 377, 425, 428, 434
 - smoothing window, 245, 251, 268, 281, 292, 296, 300–1, 318, 322, 341, 346–7
 - Bartlett–Priestley design, 344
 - choice of, 252
 - finer frequency grid, 255
 - for sea level residuals, 522
 - Grenander, 251, 268, 279, 281
 - lag window spectral estimator, 279
 - Parzen, 251
 - repeated lag window estimator, 349–50
 - spectral window, 206, 226–7, 229, 235, 268, 294, 296, 317–8, 323, 325, 328, 328, 331, 335, 341, 347, 568, 623–4, 628
 - approximation for sinusoidal multitaper spectral estimator, 400, 409
 - basic multitaper spectral estimator, 437, 575
 - definition for basic multitaper spectral estimator, 353
 - definition for direct spectral estimator, 194
 - definition for eigenspectrum, 353
 - definition for lag window estimator, 256
 - definition for weighted multitaper spectral estimator, 353
 - definition for WOSA, 414–5
 - for sea level residuals, 521–2
 - frequency range for statistical results, 213, 244
 - versus equivalent width, 214, 355, 369, 402
 - versus regularization bandwidth, 402
 - WOSA, 429
 - spectral window versus smoothing window, 257
- Bartlett
 - lag window, 268–9, 281, 344, 412
 - smoothing window, 269–70
 - spectral window, 269–70
- Bartlett–Priestley
 - design window, 273–4, 344
 - lag window, 268, 274, 281, 320, 344
 - smoothing window, 274–5
 - spectral window, 274–5
- basic multitaper spectral estimator, 352–5, 432–4, 611, 613
 - ACVS estimator for, 433
 - and Gaussian white noise, 370–4, 402–3
 - and improved estimation of innovation variance, 403–7
 - exponent of power-law process, 407–12, 440
 - log SDF, 412
 - and periodicity testing, 544–8
 - approximated by WOSA spectral estimator, 424
 - AR(4) data, 358–65, 368–70, 393–400, 405–7, 419
 - atomic clock fractional frequency deviates, 430–2, 501, 628–30
 - bias of, 359, 392
 - distribution of, 354, 356–7, 381–2, 409, 434
 - energy preservation of, 366–7
 - equivalent degrees of freedom (EDOFs) of, 354–5, 434
 - equivalent width, 355
 - expected value of, 353, 433
 - in circulant embedding simulation method, 611
 - jackknifing of, 356–7, 369–70
 - ocean wave data, 425–9
 - spectral window for, 353, 433
 - bandwidth of, 353, 359, 433
 - approximation (sinusoidal), 400–1, 434
 - sinusoidal, 398–400
 - Slepian, 358–9, 364–5
 - variance of, 353–4, 359
 - Gaussian white noise, 372–4, 402–3
 - via quadratic spectral estimator, 374
- basis for the class of band-limited functions, 66
- Bessel function, 196, 234
- best (backward) linear predictor, 455
 - complex-valued process, 502, 564
- best (forward) linear predictor, 452–3
 - complex-valued process, 502, 564
- best linear predictor, 404, 455, 460–1, 463, 472
- best predictor, 463
- bias
 - and dynamic range, 177
 - broad-band, 359, 380–2, 387–8, 390
 - defined, 378
 - indicator, 379–81, 439
 - minimizing indicator of, 380
 - due to leakage, 185, 192, 351
 - due to smoothing window, 256, 290
 - indicator for minimum-bias multitaper spectral estimator, 392, 441
 - local, 290, 359, 381–2
 - defined, 378
 - magnitude indicator, 378–9, 381
 - of direct spectral estimator, 192
 - of multitaper spectral estimator, 359
 - of periodogram, 163

- of weighted multitaper spectral estimator, 391, 441
- bivariate standard Gaussian PDF, 634
 - Mehler's expansion, 635
- Blackman–Tukey estimator, 252
- block averaging, 270–1, 412–14
- Bluestein's substitution, 94
- blurring function, 93
- bootstrapping, 593, 611, 618, 631, 638
 - confidence intervals for innovation variance, 630
 - second approach in AR case, 630
 - confidence intervals for log of SDF, 623–5
 - second approach in AR case, 626
- boundary conditions (*see* circular or reflecting)
- Brent's method, 528, 558
- broad-band bias, 359, 380–2, 387–8, 390
 - defined, 378
 - indicator, 379–81, 439
 - minimizing indicator of, 380
- Burg autoregressive estimator, 466–8, 505
 - AR(1) process, 510
 - AR(2) data, 509–10
 - AR(4) data, 468–9, 509
 - assessing variability (SDF), 485, 624–6
 - atomic clock fractional frequency deviates, 501
 - causality, 466, 505
 - complex-valued process, 502, 562
 - FPE criterion, 494
 - ocean wave data, 496–9
 - order selection via PACS, 498–9
 - of innovation variance, 629
 - peak in SDF
 - height, 576
 - shifting and splitting, 560, 562, 576, 591
 - width, 576
 - Willamette River data, 575–7
- Burg's algorithm, 466, 468, 470, 473, 478, 482, 491, 493, 505, 560, 575–6
 - analytic series, 562
 - and Yule–Walker method, 471
 - AR(2) numerical examples, 509–10, 556
 - observed prediction error, 466
 - two-step, 560
- carbon dioxide and global temperature, 366
- cascaded filter, 145
- Cauchy inequality, 257, 262, 371, 442
- causal process
 - autoregressive (AR), 34, 446, 448, 506, 618, 638
 - autoregressive moving average (ARMA), 600, 631
- causal filter, 147, 161
- causality, 446, 448, 506, 618, 638
 - and parameter estimators, 448, 617
 - assessment, 463
 - conditions for, 34, 446
- centering
 - AIC and AICC, 494–5
 - and direct spectral estimator, 195, 241, 244, 548
 - and periodogram, 184, 213, 240
 - and simulations, 619, 627
 - and tapering of a time series, 195
 - and WOSA spectral estimator, 419–20
 - of data for harmonic analysis, 583
- central peak (lobe), 175
- cepstrum, 301, 473, 503
 - estimator, 302–3
- Cesàro summability and sums, 78, 165
- chaotic beam data, 228
 - direct spectral estimator, 228–30
 - periodogram, 228–9
- Chebyshev's inequality, 164
- chi-square distribution, 37, 204, 209–10, 539
 - noncentral, 542, 552
- chi-square RV, 37, 203, 209–10, 235, 408, 539, 546
 - and confidence intervals, 438, 486, 490
 - logarithm of, 209, 412
 - expected value and variance, 302
 - scaled, 264, 298, 343
 - sum of chi-square RVs, 299
 - weighted sum of chi-square RVs, 264
- chirp transform algorithm, 94, 216
- Cholesky decomposition, 599, 604, 637, 639
 - modified, 464, 480, 599
- circular embedding simulation method, 440, 601–4, 611, 620, 627, 629, 631
 - ACVS properties, 602, 640
 - and direct SDF estimator, 611
 - and lag window SDF estimator, 611–12
 - and multitaper SDF estimator, 611, 613
 - and periodogram, 615, 641
 - and WOSA SDF estimator, 611, 613
 - compared to method of surrogate time series (MSTS), 616
 - computational efficiency, 603
 - issues with ACVS, 615
 - overcoming negative weights, 604, 637, 640
 - weights, 601–2, 605, 611, 629, 637, 640
 - AR(1) process, 640–1
 - direct SDF estimator, 612
 - lag window SDF estimator, 612
 - multitaper SDF estimator, 613
 - WOSA SDF estimator, 613
- circular boundary conditions, 424
- circularized sequence, 601–2, 640
- classification of spectra, 120–2
- CO₂ data, 332
 - lag window spectral estimator for filtered, 335
 - periodogram for filtered, 335
- coefficient of determination, 570, 572, 621
- colored noise, 120, 126, 219, 241, 377–8, 381, 389, 511, 513, 518, 520, 523, 539, 544, 550, 574, 584
 - proper complex-valued, 519, 585
- complementary covariance, 31
- complete stationarity, 215
- complex demodulation, 160, 232, 592
- complex exponentials, 17, 48, 134
 - linear combinations of, 132, 137
- complex-valued colored noise (proper), 519, 585
- complex-valued stochastic process
 - continuous parameter/time, 23

- discrete parameter/time, 23
- complex-valued Gaussian or normal RV (proper), 30, 546–7
- complex-valued process
 - by complex demodulation, 160, 374
 - resulting ACVS, 160
 - resulting SDF, 160, 374
- example of, 127
- complex-valued roots, 566–7, 580, 592
- complex-valued stationary process, 29, 231, 374, 501–2, 519, 585
 - time reverse of conjugate, 502
- complex-valued white noise (proper), 517–8, 564
- compound periodicity, 540–1
- computational pathways, 221, 224, 315
- concentration measure
 - continuous time/continuous frequency, 62, 65
 - discrete time/continuous frequency, 85, 105, 155, 386
 - discrete time/discrete frequency, 94
 - for eigenspectrum spectral window, 368
- condition number of covariance matrix, 185
 - and dynamic range, 185
- confidence interval for
 - innovation variance, 629
 - bootstrapping, 630
 - via simulation, 630
 - jump in the integrated spectrum, 590
 - log of SDF, 205, 227, 229, 266–7, 323, 328, 331, 335, 342, 354, 359, 425, 521
 - bootstrapping, 623–5
 - SDF, 205, 265, 267, 342, 357, 485, 488–90
 - chi-square versus jackknifing, 438
- conjugate symmetry, 29–30, 48, 54, 75, 92
- consistent estimator of
 - mean, 164
 - SDF
 - discretely smoothed direct spectral estimator, 246
 - lag window spectral estimator, 260, 342
- constructed multitaper spectral estimator, 355, 376
- continuous parameter deterministic sinusoid, 558
- continuous parameter white noise, 38
- continuous parameter/time stochastic process, 22, 38, 113–4, 131, 507
 - sampling and aliasing, 122–3
 - sampling and averaging, 161
- convergence
 - factors, 80, 152–3, 157, 579, 592
 - in mean square, 49, 53
 - pointwise, 50, 53, 84
- convolution, 67, 96, 98–9, 101
 - as an LTI filter, 137, 157
 - continuous parameter, 142, 247
 - discrete parameter, 143, 247
 - as smoothing operation, 69–70, 247, 385
 - cyclic, 101, 106
 - Fourier transform of, 67, 96, 98–9, 101
 - illustration of, 68–9, 104
 - in frequency domain, 96, 98–9, 101, 105
 - of periodic functions, 96, 105
- copper wires, 57
- correlation distortion due to zero-memory nonlinearity (ZMNL), 635–6
- correlation length, 131
- correlation of direct spectral estimator, 212, 243
- correlation time, 214
- correlogram method power spectral density estimator, 252
- cosine data taper, 189–90, 260
- cosine lag window, 290
- covariance
 - complex-valued case, 25, 41
 - real-valued case, 24
- covariance matrix, 28, 42, 44, 184, 376, 379, 382, 391, 440, 450, 457, 465, 485
 - Cholesky decomposition, 599, 604, 637, 639
 - modified, 464, 480, 599
 - condition number, 185
- covariance method, 480
- covariances of
 - direct spectral estimator, 205, 209–13, 243
 - in WOSA blocks, 415, 442
 - eigenspectra, 354, 372, 436, 438
 - lag window spectral estimator, 260–1, 342
 - periodogram, 203–4, 235
 - subtle issues, 214–5
 - weighted multitaper spectral estimator, 437
 - weighted sums of time series values, 242
- CPDF (*see* cumulative probability distribution function)
- cross-correlation (complex), 72, 96, 98–9, 101–2
- cumulant, 215, 632
 - inequality for fourth, 632–3
- cumulative periodogram test, 215–6
 - examples
 - AR(2) data, 216
 - ocean noise, 230–1
 - Willamette River data, 550, 573–4
- cumulative probability distribution function (CPDF), 23
 - N -dimensional, 23
- cyclic
 - autocorrelation, 220
 - convolution, 220, 273
 - convolution theorem, 101, 106
 - convolutions via FFTs, 106
- cyclical frequency, 8
- Daniell
 - design window, 272
 - lag window, 268, 273, 281, 290
 - smoothing window, 250, 272–3, 320
 - spectral window, 272–3
- data taper (*see* taper)
- data window, 186
- DC (direct current), 518
- DCT-based periodogram, 217, 244
 - AR(4) process, 218
 - asymptotic unbiasedness of, 218
 - bias of, 218–9
 - statistical properties, 219
- decay
 - of envelope of sidelobes, 269
 - rate of sidelobes

- Bartlett smoothing window, 269, 276
- Bartlett–Priestley smoothing window, 275
- Daniell smoothing window, 273
- Gaussian smoothing window, 277
- Papoulis smoothing window, 278
- Parzen smoothing window, 276
- spectral windows, 281, 328
- decibel (dB) scale, 13, 209, 225–7, 266
- deconstructed multitaper spectral estimator, 355, 376, 420–3
- decorrelation time, 214
- default data taper, 188, 190
- degenerate kernel, 86
- degrees of freedom
 - for chi-square RV, 37, 202
 - one, 13, 203
 - two, 13, 203
 - zero, 542
- equivalent (EDOFs), 264, 266
 - discretely smoothed direct spectral estimator, 343
 - discretely smoothed periodogram, 347
 - lag window spectral estimator, 264, 279, 342
 - multitaper spectral estimator, 354–5, 390, 434
 - WOSA spectral estimator, 415–6, 442–3
 - in a time series, 298, 507
 - of order 2, 298
- derivative as an LTI filter, 157
- design window, 247
 - for Bartlett–Priestley, 273–4, 319, 344
 - for Daniell, 272, 319
- determinant of a matrix, 480
- deterministic
 - function, 48
 - sequence, 48
 - sinusoid, 553
- detrending (*see* trend removal and quadratic trend removal)
- deviance, 308
 - generalized cross-validated (GCV), 309, 313, 324, 339
- DFT (*see* discrete Fourier transform)
- diagonal matrix, 464, 565–6, 599
- differencing
 - for linear trend removal, 40
 - for seasonality/periodicity removal, 40, 42, 333
 - higher-order, 340
 - to achieve stationarity, 340, 431
- digamma function, 210, 296, 302, 345, 405, 409
- Dirac delta function, 74, 113, 120, 157, 250, 524
- direct current (DC), 518
- direct spectral estimator, 164, 186, 201, 233, 340, 375, 535–6, 545, 611
 - and centering a time series, 195–6, 241, 244, 548
 - atomic clock data, 327–9
 - average value of, 241
 - averaging of uncorrelated, 443
 - bias, 192
 - chaotic beam data, 228–9
 - computation of, 219–24
 - correlations, 212, 243
 - corresponding ACVS estimator, 188, 452, 611
 - covariances, 205, 209–13, 243
 - discretely smoothed, 246–7
 - equivalent degrees of freedom (EDOFs), 347
 - spectral window bandwidth, 347
 - distribution of, 204, 235
 - for complex-valued process, 232
 - eigenspectrum, 352
 - expected value of, 186, 234
 - for real-valued harmonic process, 535, 585
 - exponential averaging, 444
 - for blocks in WOSA, 414
 - harmonic process examples, 536, 588
 - in circulant embedding simulation method, 611–12, 627–8
 - inconsistent estimator of SDF, 207, 235
 - linear combination of independent, 441–2
 - ocean wave data, 225–6, 496
 - relationship to quadratic spectral estimator, 376
 - ship altitude data, 330–1
 - simulating properties of, 242
 - smoothing of, 245
 - spectrum of, 213
 - use with AR order selection, 493
 - use with Yule–Walker autoregressive estimator, 452, 461–2, 505
 - variability, 208
 - variance of, 204, 235, 624
 - for complex-valued process, 232
 - white noise, 189, 371
- Dirichlet’s kernel, 17–8, 76–7, 80, 85, 93, 106, 127, 233, 383, 537,
- discrete cosine transform, 184, 217
- discrete Fourier transform (DFT), 74–5, 92, 219, 594, 602, 608
 - as a convolution, 93–4
 - chirp algorithm, 94
 - conjugate symmetry, 75, 92
- discrete parameter/time stochastic process, 22
- discrete power spectrum, 50, 139–40
- discrete prolate spheroidal sequence (DPSS), 87, 155, 189, 357
- discrete prolate spheroidal wave function (DP-SWF), 87, 105–6, 383–5, 439
- discrete spectrum, 120, 511–13, 518, 538, 584–6
- discrete time sequence, 74
 - autocorrelation width, 100
 - convolution theorem, 99
 - equivalent width, 100
 - Fourier representation, 75, 99
 - Parseval’s theorem, 75, 99
 - segment of, 91
 - autocorrelation width, 102
 - convolution theorem, 101
 - equivalent width, 102
 - Fourier representation, 92, 100
 - Parseval’s theorem, 92, 101
 - spectral properties, 101
 - spectral properties, 99
- discretely smoothed direct spectral estimator, 246–7
 - computation of, 316
 - consistent estimator of SDF, 246

- equivalence to lag window spectral estimator, 249–50
- finer frequency grid, 253
- discretely smoothed periodogram, 307
 - bias/variance trade-off, 309
 - computational scheme, 347
 - corresponding ACVS estimator, 312–13, 348, 615
 - equivalent degrees of freedom (EDOFs), 347
 - evaluating SDF estimates, 349
 - ice profile data, 324
 - parameter choice, 307, 309–10, 312, 324, 339, 348
 - spectral window bandwidth, 347
- distribution of
 - adaptive multitaper spectral estimator, 390
 - direct spectral estimator, 204, 235
 - complex-valued process, 232
 - lag window spectral estimator, 264, 342, 490
 - maximum likelihood AR parameter estimator, 483
 - other AR parameter estimators, 484
 - multitaper spectral estimator, 354, 356, 381, 409, 434
 - periodogram, 203, 235, 408
 - WOSA spectral estimator, 436
- Doppler measurements of blood flow, 131
- double orthogonality
 - of eigenfunctions, 65, 87
 - of Slepian sequences (DPSSs), 106
- double-sided versus single-sided SDF, 118
- DPSS (*see* discrete prolate spheroidal sequence)
- DPSWF (*see* discrete prolate spheroidal wave function)
- duration–bandwidth product, 66, 357
- dynamic range, 177, 184, 197, 492, 641
 - and condition number, 185
- echoes, 227
- ECLS (*see* exact conditional least squares)
- EDOFs (*see* equivalent degrees of freedom)
- effective bandwidth (*see* spectral window bandwidth)
- eigenfunction, 64
 - double orthogonality, 65, 87
 - for a linear filter, 135
- eigenspectra, 352, 385, 432, 545
 - bandwidth of spectral windows, 353, 358
 - covariances of, 354, 372, 436, 438
 - plots of for AR(4) data, 361, 363, 368, 395, 397
 - regularization bandwidth, 357
 - weights associated with, 352, 388–9
- eigenvalue, 64, 86–7, 95, 185, 375, 380, 392, 565, 567, 580, 591
 - for a linear filter, 135
- eigenvector, 87, 95, 380, 392, 591
 - left and right, 565, 567
- end-point mismatch, 181, 183
- energy
 - of a nonperiodic function, 54–5
 - of a periodic function, 49
- energy spectral density function, 55, 97, 99, 132
- ensemble, 21–2
- entire function, 57
- entropy, 471–3
- envelope of sidelobes, 269, 273
- Epanechnikov lag window (*see* Bartlett–Priestley lag window)
- equal
 - by definition, 8, 23
 - in distribution, 203, 485, 599
 - in the mean square sense, 49, 53
- equivalence of time and frequency domains, 55
- equivalent degrees of freedom (EDOFs), 264, 266
 - adaptive multitaper spectral estimator, 390
 - example (ocean wave data), 427–8
 - basic multitaper spectral estimator, 354–5, 434
 - example (ocean wave data), 427
 - discretely smoothed direct spectral estimator, 343
 - discretely smoothed periodogram, 347
 - lag window spectral estimator, 264, 279, 342
 - example (ocean wave data), 318–20
 - via eigenvalues, 423–4
 - weighted multitaper spectral estimator, 354
 - WOSA spectral estimator, 415–6, 442–3
 - example (ocean wave data), 429–30
- equivalent width, 58–60, 73, 97–8, 100, 102, 213–4
 - multitaper spectral estimator, 355, 437
 - sinusoidal multitaper spectral estimator, 401, 441
- smoothing window, 251
- versus spectral window bandwidth, 214, 355, 369
 - sinusoidal multitapers, 402
 - Slepian multitapers, 369
- ergodic theorems, 165
- Euler relationship, 17
- Euler’s constant, 210, 302, 404, 408
- exact conditional least squares (ECLS)
 - including mean, 552
 - sinusoidal amplitude estimator, 549, 569–70, 572, 589–90
 - white noise variance estimator, 549, 552, 570, 589–90
- exact unconditional least squares (EULS)
 - including mean, 552
 - sinusoidal amplitude estimator, 549, 569–70, 572
 - sinusoidal frequency estimator, 549
 - white noise variance estimator, 550, 552, 570
- expected value (expectation), 4
- expected value of
 - direct spectral estimator, 186, 234
 - real-valued harmonic process, 535, 585
 - lag window spectral estimator, 255, 341, 343
 - multitaper spectral estimator
 - adaptive, 389–90, 439
 - basic, 353, 433
 - weighted, 352
 - periodogram, 174, 233, 240
 - complex-valued harmonic process, 524, 586
 - real-valued harmonic process, 524, 585
 - conditional on phase value, 537–8
 - quadratic spectral estimator, 376–7, 439

- sinusoidal frequency estimator using periodogram maximum, 526
 WOSA spectral estimator, 414, 435
 exponential averaging, 444
 exponential distribution, 202, 209, 215, 598
 exponential RV, 209
 extrapolation of a band-limited function, 66
- fader, 186
 fast Fourier transform (FFT), 92, 94, 237, 602, 608
 FBLS (*see* forward and backward least squares)
F-distribution, 547, 574, 582, 587
 non-central, 552
 percentage points (special case), 544
 Fejér's kernel, 80, 174–6, 180, 233, 240, 524, 532, 534, 537
 different expressions for, 236
 sampling of, 524
 sidelobes, 178–80, 328, 527, 532, 535–6
 FFT (*see* fast Fourier transform)
 filter
 acausal, 147–8, 152, 155
 analog, 133
 antialias, 57, 224
 band-pass, 146
 cascaded, 145
 causal, 147
 demodulation, 160
 digital, 140
 effect on integrated spectrum (filtering theorem)
 analog, 136, 157
 digital, 141
 example of an LTI digital, 148
 finite impulse response (FIR), 147–8, 152, 155, 161, 503
 first difference, 333
 for pilot analysis, 160
 high-pass, 146
 as trend remover, 151
 formed by subtracting output of low-pass filter from its input, 149
 infinite impulse response (IIR), 147, 503
 Kalman, 482
 kurtosis norm of, 632
 linear time-invariant (LTI), 132, 631, 633
 analog, 133
 and convolution, 137
 digital, 140
 low-pass, 146, 592
 cutoff frequency, 151
 design by least squares, 152
 design using Slepian sequences, 155
 prediction error, 198
 prewhitening, 197, 234, 241, 447, 482, 491–3, 496, 510, 615
 seasonal differencing, 333
 transfer function, 136, 141
 transformation or operator, 133
 final prediction error (FPE), 493
 finite impulse response (FIR) filter, 147–8, 152, 155, 161, 503
 FIR (*see* finite impulse response)
 first difference filter, 40, 333
 effect of filter on SDF, 158, 326
 Fisher's test for simple periodicity, 539–40, 550, 586
 critical value, 581
 approximation, 540, 589
 ocean noise data, 581
 Willamette River data residuals, 571, 573
 flicker frequency noise, 431
 folding frequency (Nyquist frequency), 82
 forward and backward least squares (FBLS) and SVD approach, 565
 autoregressive parameter estimator, 477, 484
 AR(2) examples, 510, 555–6
 autoregressive spectral estimator
 AR(4) example, 478, 482
 Willamette River data, 592
 same as modified covariance method, 480
 forward least squares (FLS)
 autoregressive parameter estimator, 477, 484
 AR(1) example, 510
 same as covariance method, 480
 forward linear prediction, 452–3
 complex-valued process, 502, 564
 forward linear prediction error, 453, 596
 Fourier coefficient, 49
 Fourier frequencies, 8, 91, 171, 515, 523–4
 fundamental, 357
 grid of, 171
 alternative grid for uncorrelated SDF estimators, 206, 208, 212
 Fourier integral representation, 54
 Fourier representation of a periodic function, 49, 95
 Fourier synthesis, 54
 Fourier theory
 continuous time/continuous frequency, 53, 97
 continuous time/discrete frequency, 48, 95
 discrete time/continuous frequency, 74, 99
 discrete time/discrete frequency, 91, 100
 Fourier transform pair, 54, 96–7, 99–100
 examples of, 171, 181, 188, 212, 220–2, 237, 247–8, 301–2
 for convolution of two functions, 67
 for derivative of a function, 61
 FPE (*see* final prediction error)
 fractionally differenced (FD) process, 166, 407–8, 440
 autocovariance sequence (ACVS), 440
 SDF, 407–8
 simulation by circulant embedding, 440
 Fredholm integral equation
 of the first kind, 383
 of the second kind, 63, 86
 frequency grid, 171, 527, 531, 558, 578, 605
 for circulant embedding, 601
 for uncorrelated SDF estimators, 206, 208, 212
 frequency response function, 136
 Hilbert transform, 579
 function, special type of
 blurring, 93
 digamma, 210

- Dirac delta, 74
- gamma, 440
- modified Bessel, 196
- periodic, 48
- positive-part, 541
- trigamma, 296
- function width
 - autocorrelation, 73, 97–8, 105, 191, 194, 288
 - equivalent, 58, 73, 97–8, 105, 213
 - half-power, 191–2, 288
 - variance, 60, 73, 105, 191–2, 194, 288
- fundamental Fourier frequency, 357
- fundamental frequency, 569

- gain function, 136
 - squared
 - for differencing, 334
 - for Hilbert transform approximation, 592
 - for ideal low-pass, high-pass and band-pass filters, 146
- gamma function, 297, 345, 440, 542, 633
- Gaussian distribution, 265
 - generalized, 633
- Gaussian kernel, 69
- Gaussian lag window, 268, 276, 278, 281, 320, 330, 430
- Gaussian or normal process, 29, 445, 463, 480, 490, 544
 - autoregressive, 34, 445, 595
 - complex-valued, 30
 - entropy, 472
 - moving average, 33, 595
 - white noise, 31, 201, 210, 539, 594
- Gaussian or normal RV
 - complex-valued, 30, 210
 - PDF and its Fourier transform, 55
- Gaussian smoothing window, 276–8
- Gaussian spectral synthesis method (GSSM), 605–8, 637, 640–1
 - ACVS approximation, 607
 - adjustment for power-law process, 610
 - and adaptive multitaper spectral estimator, 615
 - and prewhitening, 615
 - role of sampling interval, 609
 - weights for AR(1) process, 640–1
- Gaussian spectral window, 278
- Gaussian white noise, 31, 201, 210, 539, 594
 - multitaper spectral estimator, 370
- Gaussian-shaped ACVF and SDF, 131
- Gegenbauer process, 610
- generalized cross-validated (GCV) deviance, 309, 313, 324, 339
- generalized Gaussian (GG) distribution, 633
- generalized inverse, 566
- generalized least squares, 412, 522
- Gibbs phenomenon, 77, 79, 81, 152
- goodness-of-fit test
 - Kolmogorov, 215
- grid size, 179, 527, 531, 558, 568, 578, 605
- group delay, 136
- GSSM (*see* Gaussian spectral synthesis method)
 - half-power width of function, 191–2, 288
 - Hamming lag window, 343
 - Hanning
 - data taper, 189–90, 222, 230, 234, 260, 305, 327, 331, 345, 415–8, 535
 - alternative form, 240, 285
 - lag window, 343, 579
 - harmonic analysis, 511
 - ocean noise data, 581
 - ocean wave data, 582
 - parametric approach, 553
 - AR examples, 555–6
 - problems with, 558
 - singular value decomposition (SVD) approach, 563
 - summary, 584
 - Willamette River data, 567
 - centering of time series, 583
 - harmonic process, 35, 108, 511
 - ACVS for, 36–7, 117
 - expected value of, 36
 - periodicity, 603–4
 - random phase, 37, 45
 - simulation, 593–4, 598–9, 616, 637, 641
 - examples, 36
 - variance of, 36, 108, 110
 - variance spectrum of, 108
 - harmonic process with additive noise
 - complex-valued, 519, 585–6
 - ACVS for, 519, 562, 585–7
 - expected value of, 519, 585–6
 - known frequencies, 517, 586
 - unknown frequencies, 586
 - variance of, 519, 585–6
 - real-valued, 512, 544, 584–5
 - ACVS for, 519, 584, 591
 - expected value of, 516, 584
 - known frequencies, 513, 515, 585
 - random phase, 516
 - signal-to-noise ratio, 526, 532
 - tapered, 535–6, 588
 - unknown frequencies, 523, 585
 - variance of, 519, 526, 532, 584
 - harmonic tidal prediction, 512
 - at Southampton, England, 513
 - harmonics
 - in tidal prediction, 522
 - of fundamental frequency, 569
 - Heisenberg's uncertainty principle, 61–2, 104
 - Hermite polynomials, 634–5
 - Hermitian Toeplitz matrix, 30
 - high-cut frequency, 57
 - high-pass
 - filter, 146
 - process, 299, 347
 - creation of from a low-pass filter, 160
 - Hilbert transform, 114, 562, 579
 - frequency response function of, 579, 592
 - impulse response sequence of, 579, 592
 - transfer function of, 579, 592
 - ice profile data, 321
 - lag window spectral estimator, 322–3
 - periodogram, 321, 323

- plot of, 322
- IID (independent and identically distributed)
 - noise, 32, 631
 - process, 32
- IIR (*see* infinite impulse response)
- imaginary part of
 - analytic series, 579
 - complex-valued number, 551
 - complex-valued process, 602, 606, 640
- impulse response
 - function, 142
 - sequence, 143
 - for Hilbert transform, 579, 592
 - normalization of, 150
- inconsistency of
 - direct spectral estimator, 235
 - periodogram, 163–4, 207–8
- index-limited sequence, 85
- indirect nonparametric estimator, 252
- infinite impulse response (IIR) filter, 147, 503
- initial conditions
 - AR process simulation, 595
 - Gaussian AR process simulation, 595, 599
 - Gaussian ARMA process simulation, 600, 639
- inner product, 470
- innovation process, 446
 - estimation of, 17
- innovation variance, 403–4, 446, 449, 455, 468, 471–2, 510
 - Burg autoregressive estimator, 499–500, 629
 - confidence interval for, 629
 - different estimators compared, 406, 440
 - direct spectral estimator, 405
 - mean square linear prediction error for AR(p), 455
 - multitaper spectral estimator, 405, 432, 629
 - variability assessment via simulation, 630
 - periodogram-based estimator, 404
 - simulation, 628–9
 - variability assessment via, 629
 - smoothed periodogram estimator, 404
 - Yule–Walker autoregressive estimator, 499–500
- integrable (in square sense), 49, 53–4
- integral equation, 63, 86
- integral time scale, 131, 214
- integrated magnitude square error, 52, 76
- integrated spectrum (spectral distribution function), 110, 117, 163, 511
 - absolutely continuous, 120
 - and AR SDF (parametric sinusoidal power estimation), 578
 - and LTI filters, 135–6, 141
 - cascade of filters, 146
 - basic properties, 116
 - classification examples, 121
 - continuous parameter stationary process, 114
 - continuous singular function, 120
 - differentiability, 120
 - discrete parameter stationary process, 110
 - jumps (steps), 120, 511, 549, 551, 589–90
 - confidence interval for, 590
 - distribution of estimator, 552, 590
 - Willamette River example, 570, 572, 575
- intrinsically stationary process
 - order d , 340
 - order unity, 431
- inverse Fourier transform (*see* Fourier transform pair)
- invertibility of moving average process, 43
- irregularly sampled observations, 355, 528, 543
- Isserlis theorem, 30, 43, 210, 236, 241, 298, 371
- jackknifing
 - and multitaper spectral estimator, 356, 369–70
 - for confidence interval calculation, 438
- Johnson system of distributions, 633
- joint moment, 27
- joint stationarity, 29
- jump process, 109
- jumps in the integrated spectrum, 120, 511, 549, 551, 589–90
 - confidence interval for, 590
 - distribution of estimator, 552, 590
 - Willamette River example, 570, 572, 575
- Kalman filter, 482
- kernel
 - degenerate, 86
 - Dirichlet's, 17–8, 76, 93
 - Fejér's, 80, 174–5
 - Gaussian, 69, 290
 - of an integral equation, 63, 86
 - rectangular, 71, 290
- Kolmogorov goodness-of-fit test, 215
- Kronecker's delta function, 44
- k th-order Slepian data taper, 357, 382, 384
 - plots of, 360, 362
- Kullback–Leibler (KL) discrepancy, 297, 312, 346, 349
 - expected value, 297
 - for scaled chi-square RVs, 345
 - symmetrized form, 345–6, 349
 - with swapped arguments, 345–6, 349
- kurtosis, 631
 - ARMA(1,1) process, 632
 - norm of a filter, 632
- lag, 2, 27
 - lag 1 scatter plot, 2
 - lag p differencing, 40
 - lag τ scatter plot, 2
- lag window, 248, 341
 - choice of, 287
 - compared to data taper, 282–3, 285
 - constructed from data taper, 283, 344–5
 - parameter m (controls smoothing), 247, 287
 - choice of, 291, 304, 313–4
 - examples of, 318, 322, 328, 330, 339
 - properties of, 250
 - reshaped, 257, 281–2, 375
 - type
 - autocorrelated 100% cosine, 283–5, 344–5
 - Bartlett, 268–9, 281, 344, 412
 - Bartlett–Priestley, 268, 273, 281, 320, 344
 - cosine (100%), 283–4, 290

- Daniell, 268, 271, 281, 290, 320
 Gaussian, 268, 276, 278, 281, 320, 331, 430
 Hamming, 343
 Hanning, 343, 579
 Papoulis, 268, 278–9, 281, 345
 Parzen, 248, 268, 275, 281, 301, 320–1, 323, 331, 344, 346, 496, 521–2
 Slepian, 285, 345
 lag window spectral estimator, 245, 248, 340–1, 492, 612
 alternative names for, 252
 as a quadratic spectral estimator, 375
 as deconstructed multitaper estimator, 420–2
 asymptotic distribution of, 264–5, 297, 342, 490
 asymptotic unbiasedness of, 256, 341
 asymptotic variance of, 258–9, 262, 265, 279, 341
 verification of validity by simulation, 622–3
 atomic clock data, 328–9, 501
 computation of, 314–7
 consistent estimator of SDF, 260, 342
 corresponding ACVS estimator, 248, 341, 612
 covariances, 260, 342
 equivalence to discretely smoothed direct spectral estimator, 249–50
 equivalent degrees of freedom (EDOFs), 264, 279, 342
 expected value of, 255, 341, 343
 filtered CO₂ data, 335
 gamma PDF, 297
 ice profile data, 321–4
 in circulant embedding simulation method, 611–12, 619–20, 627
 ocean wave data, 316–9, 349–50, 430, 496
 parameter m (controls smoothing), 247, 287
 choice of, 291, 304, 313–4
 examples of, 318, 322, 328, 330, 339
 repeated, 349–50
 sea level residuals, 521–2
 ship altitude data, 330–2
 smoothing window bandwidth B_W , 251, 279
 alternative measure β_W , 251, 279
 use in prewhitening, 491
 Laplace periodogram, 529–30
 law of total expectation, 588
 leakage, 77–8, 178, 180, 196, 233, 287, 358, 532, 592
 and end-point mismatch, 181, 239
 bias in spectra due to, 225, 227, 233, 351
 reduction of, 78, 185
 smoothing window, 268, 287–9, 294–5, 316, 318–9, 498
 least squares (LS), 40, 50, 332, 409, 430–1, 475–8, 513–6
 approximate, 516, 585
 approximate conditional (*see* approximate conditional least squares)
 complex-valued, 517, 546, 565
 exact conditional (*see* exact conditional least squares)
 exact unconditional (*see* exact unconditional least squares)
 filter design, 152, 155
 generalized (GLS), 412, 522
 ordinary (OLS), 409, 430, 440, 522
 least squares (LS) autoregressive estimators, 475–9, 505–6
 AR(1) process, 510
 assessing variability (SDF), 485
 backward, 477, 484, 510
 causality not guaranteed, 479, 506, 510, 617
 forward, 475–7, 484, 510
 forward and backward, 477–8, 484, 509, 555–6, 565
 relation to Yule–Walker scheme, 479–80
 Lebesgue decomposition theorem, 120, 518
 Levinson–Durbin recursions, 452, 456–7, 461–2, 464–6, 509, 556, 604, 639
 for complex-valued time series, 502
 linear algebra derivation, 463–4
 step-down scheme, 460–1, 507–8, 597
 causality assessment, 461, 463, 509
 generation of ACVS for an AR process, 460, 508
 simulation of ARMA process, 461, 595–8
 likelihood function, 308, 480
 and order selection criteria, 494–5
 log of, 480–1
 reduced (profile), 482–4
 Whittle, 307
 line spectrum, 120, 511, 522
 linear and log scales for SDFs, 336–9
 linear model, 2, 39, 46, 621
 linear regression, 151, 408–10, 412, 440, 446, 501
 complex-valued, 517, 545
 real-valued, 513–5, 529
 robust, 530
 linear taper, 186
 linear time-invariant (LTI) filter, 132
 analog, 133
 digital, 140
 linearity of, 133, 140
 linear trend removal
 by differencing, 40
 by linear least squares estimation, 40
 linear window, 186
 local bias, 290, 359, 381–2
 defined, 378
 indicator, 378–9, 381
 minimizing indicator of, 381
 log and linear scales for SDFs, 336–9
 log spectral density function, 301
 estimation of
 log direct spectral estimator, 301–2
 with smoothing, 303–6, 324
 Lomb–Scargle periodogram, 528–9, 543–4, 588
 long memory process, 407
 long-range dependence, 407–8, 411
 Lorenzian SDF, 131, 507
 loss of resolution, 76, 78, 271, 290, 357, 359, 400, 434, 535
 low-cut frequency, 57
 low-pass
 filter, 146, 592
 process, 299, 347
 use of to create a high-pass filter, 160

- low-rank approximation to SDF estimator, 420–3
 lower triangular matrix, 464, 484, 509, 599, 639
 LTI (*see* linear time-invariant)
- MA (moving average) process (*see* moving average (MA) process)
 male speech, 57
 Mann–Wald theorem, 486
 marginal distribution, 45, 631, 634
 matching condition, 135–6, 141–3
 matrix
 Cholesky decomposition of, 599, 604, 637, 639
 modified, 464, 480, 599
 covariance, 28, 42, 44, 184, 376, 379, 382, 391, 440, 450, 457, 465, 485
 determinant of, 480
 diagonal, 464, 565–6, 599
 generalized inverse of, 566
 lower triangular, 464, 484, 509, 599, 639
 rank of, 375–6, 380–1, 420, 422–3, 490, 565–6
 Schur, 484
 singular value decomposition (SVD) of, 425, 564–5
 and short time series, 567
 approach to harmonic analysis, 563
 Willamette River data, 578–80, 592
 singular value, 565
 singular vector, 565
 Toeplitz, 29, 156, 457, 462, 480, 484
 Hermitian, 30
 trace of, 376, 378, 439
 tridiagonal, 88, 95
 maximum confusion spectral analysis, 473
 maximum entropy
 principle of, 471, 473
 for ARMA process, 473–4
 spectral analysis (MESA), 194, 447, 466, 471–4, 503
 example for MA processes, 475–6
 maximum likelihood autoregressive estimator (MLE), 480–4, 506
 approximations to, 484, 510
 AR(1) process, 482
 AR(2) data, 489
 AR(4) data, 482–3, 490
 assessing variability (SDF), 485
 distribution of, 483
 recursive, 485
 mean, 4, 24
 complex-valued case, 25
 estimation of, 3, 164–6, 184, 195–6, 213, 548
 mean correction (*see* centering)
 mean integrated square error (MISE), 303
 mean square convergence, 49, 52–3
 mean square error (MSE), 167, 236, 293, 346, 349, 386–8, 404, 463, 526, 537
 normalized (NMSE), 296, 314, 346, 349
 mean square linear prediction error, 453–5, 457, 460, 468, 493, 509
 same as AR(p) innovation variance, 455
 mean square log error (MSLE), 296, 346, 349
 mean square prediction error, 463
 Mehler's expansion, 635
 method of surrogate time series (MSTS), 615–6, 641
 and harmonic processes, 641
 compared to circulant embedding, 616
 minimum-bias multitapers, 392–3, 441
 bias indicator of, 392, 441
 plots of, 393
 relationship to sinusoidal multitapers, 392–3
 mixed spectrum, 120–1, 359, 511, 513, 518–9, 521–2, 539, 584–5
 modified
 Cholesky decomposition, 464, 480, 599
 covariance method, 480
 Bessel function, 196
 Daniell spectral estimator and windows, 280
 periodogram, 186
 moment
 coefficient of kurtosis, 631
 coefficient of skewness, 631
 first and second, 27
 joint, 27
 moving average (MA) process, 32, 43, 503–4, 594–5
 autocovariance sequence (ACVS), 32, 128
 invertibility, 43
 noninvertibility, 474
 MA(1)
 ACVS, 504
 as infinite-order AR process, 43
 simulated examples, 33
 MA(2), 159–60
 simulation of, 594–5, 637, 639
 spectral density function (SDF), 130, 144, 503
 textile slivers, 503
 moving average (MA) spectral estimator, 503
 and truncated periodogram, 504
 MSTS (*see* method of surrogate time series)
 multitaper spectral estimator, 351–5, 432–4
 adaptive, 389–90
 ACVS (issue of being ill-defined), 615
 distribution of, 390
 versus weighted multitaper estimator, 390
 and improved estimation of
 innovation variance, 403–7
 exponent of power-law process, 407–12, 440
 log SDF, 412
 and periodicity testing, 544–8
 approximated by WOSA spectral estimator, 424
 AR(4) data, 358–65, 368–70, 393–400, 405–7, 419
 atomic clock fractional frequency deviates, 430–2, 501, 628–30
 basic, 352–5, 432–4, 611, 613
 ACVS estimator for, 433
 and Gaussian white noise, 370–4, 402–3
 bias of, 359, 392
 distribution of, 354, 356–7, 381–2, 409, 434
 energy preservation of, 366–7
 EDOFs of, 354–5, 434
 equivalent width, 355

- expected value of, 353, 433
 - jackknifing of, 356–7, 369–70
 - spectral window for, 353, 433
 - bandwidth of, 353, 359, 433
 - approximation (sinusoidal), 400–1, 434
 - sinusoidal, 398–400
 - Slepian, 358–9, 364–5
 - variance of, 353–4, 359
 - Gaussian white noise, 372–4, 402–3
 - constructed, 355, 376
 - deconstructed, 355, 376
 - eigenspectra (*see basic entry for eigenspectra*)
 - in circulant embedding simulation method, 611–13
 - information recovery, 195, 366, 372
 - minimum-bias, 391–3
 - ocean wave data, 425–9
 - peak-matched multitapers, 382
 - sinusoidal multitapers, 355, 391–4, 396, 434
 - Slepian multitapers, 355, 357–8, 360, 362, 368, 434
 - versus prewhitening, 424
 - versus WOSA spectral estimator, 414, 417–9, 424
 - via quadratic spectral estimator, 374
 - via regularization, 382
 - weighted, 352–4, 388, 433, 611–13
 - and quadratic spectral estimator, 376
 - bias of, 391
 - covariances of, 437
 - distribution of, 354–5
 - EDOFs of, 354–5
 - equivalent width, 355, 437
 - expected value of, 352, 389
 - low-rank approximation, 422–3
 - Parseval's theorem in expectation, 390
 - spectral window for, 352–3, 391
 - bandwidth of, 353
 - variance of, 354
 - multitaper-based test for periodicity, 544–8
 - sinusoidal amplitude estimator (complex-valued), 546, 551, 590
 - Willamette River data, 574–5
 - white noise variance estimator, 546, 551, 590
 - multivariate Gaussian (normal), 4, 29–30, 483, 485, 510
 - mutually (pairwise) uncorrelated random variables, 8, 13, 108, 203–4, 206, 235, 246, 258, 302, 306, 381, 410, 443, 465, 546, 595–6
- Newton–Raphson method, 528, 558
- noise
 - colored, (*see entry for colored noise*)
 - flicker frequency, 431
 - IID, 32, 631
 - in signal-to-noise ratio, 526
 - random walk frequency, 431
 - white, (*see entry for white noise*)
 - with heavy-tailed distribution, 530
- noncentral
 - F -distribution, 552
 - chi-square distribution, 542, 552
- nonergodic stationary process, 128
- non-Gaussian (non-normal) process, 36, 210, 412, 463, 530, 631–6
 - plots with examples of, 31
- nonindependent but uncorrelated random variables, 45–6, 108
- nonlinear lag relationship, 5–6
- nonnegative definite (*see positive semi-definite*)
- nonperiodic function, 53–4, 107, 143
- nonstationary process
 - complex-valued, 42
 - differencing for stationarity, 333, 340, 431
 - examples of, 39–41, 44–6, 332
- nonwhite noise (*see colored noise*)
- normal equations, 514, 517
- normal process (*see Gaussian process*)
- normalized mean square error (NMSE), 296, 314, 346, 349, 607
- normalized observed prediction error
 - backward, 618
 - forward, 618, 624, 626, 630, 638
- normalized root mean-square error, 406, 440
- Nyquist
 - frequency, 82, 84, 108, 122–3, 142
 - rate, 84, 123
- observed prediction error
 - backward, 466, 492, 618
 - normalized, 618
 - forward, 466, 491, 496, 617–8
 - normalized, 618, 624, 626, 630, 638
- ocean noise data, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - periodogram for, 231
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
 - testing for white noise, 11, 230–1, 581–2
- ocean tide prediction, 512
- ocean wave data, 224–8, 316–21, 425–30, 496–9, 582–3, 619–26
 - adaptive multitaper spectral estimator, 428
 - equivalent degrees of freedom (EDOFs), 427–8
 - Burg autoregressive spectral estimator, 496, 500, 625
 - direct spectral estimator, 226
 - estimated PACS, 498–9
 - harmonic analysis of, 582–3
 - lag window spectral estimator, 318–9, 349–50, 430, 496, 500, 623
 - multitaper spectral estimator, 426, 582
 - periodogram, 226–7
 - plot of, 225, 620
 - prewhitening, 496–7
 - reshaped spectrum, 582
 - simulated series, 620
 - uses for, 626
 - Thomson's F -test for periodicity, 582

- WOSA spectral estimator, 430
 Yule–Walker autoregressive spectral estimator, 496
- octave, 269, 273, 276–7, 282, 288–9
- one-sided versus two-sided SDF, 118
- orchestral music, 57
- order
 of autoregressive process, 33, 446
 of autoregressive moving average process, 35
 of autoregressive spectral estimator, 458
 of linear predictor, 452–3, 455
 of moving average process, 32, 503
 of taper in multitapering scheme, 357, 392–3
- order selection for autoregressive process, 492–5
 AIC method, 494
 centered time series, 495
 AICC method, 495
 centered time series, 495, 575–6
 direct spectral estimator criterion, 493
 examples of use of, 496, 498–500
 FPE method, 493
 PACS method, 493, 498
- ordinary least squares (OLS), 409, 430, 440, 522
- orthogonal
 data tapers, 354
 increments, 109
 process, 109
- orthogonality
 of Hermite polynomials, 635
 principle, 454
 corollary to, 508
 relationships (trigonometric), 19, 202, 516
- orthonormality
 basis exhibiting, 138, 384
 nonuniqueness of, 138
 of data tapers, 432
 of eigenvectors, 565
- oscilloscope, 21–2
- outliers, 196, 227
- oversmoothing, 71
- oxygen isotope ratios, 366
- PACS (*see* partial autocorrelation sequence)
- pairwise uncorrelated random variables (*see* mutually (pairwise) uncorrelated random variables)
- Papoulis
 lag window, 268, 278–9, 281, 290, 345
 smoothing window, 278–9
 spectral window, 279
- parametric approach to harmonic analysis, 553–8
 AR(2) example, 555–6
 AR(16) example, 556–7
 problems with, 558–62
 Willamette River data, 575–8
- parametric spectral estimation, 14, 445, 482, 491–2, 503, 506
 AR versus MA, 503
 summary, 505
- Parseval’s theorem, 49, 55, 75, 92
 in expectation
 adaptive multitaper spectral estimator, 390, 439
 weighted multitaper spectral estimator, 390
- one discrete time sequence, 99
 example of usage, 85, 111, 115, 188, 194, 251, 260, 303, 385, 390
- one nonperiodic function, 97
 example of usage, 61
- one periodic function, 96
 example of usage, 49, 52, 137–8
- one segment of discrete time sequence, 101
- two discrete time sequences, 99
- two nonperiodic functions, 97
- two periodic functions, 96
- two segments of discrete time sequences, 101
- partial autocorrelation
 coefficient, 457, 466, 469, 509
 sequence (PACS), 462–3
 complex-valued process, 502
 estimated for ocean wave data, 498–9
 for AR order selection, 493, 498
- Parzen
 lag window, 248, 268, 275–6, 281–2, 292, 316, 320–1, 331, 344, 346, 496, 521–2
 smoothing window, 275, 280–1, 344
 spectral window, 275
- PDF (*see* probability density function)
- peak-matched multitapering, 382
- peak shifting, 559–60, 576
 and splitting, 560
 analytic series, 561
 multiple sinusoids, 562
 with Burg’s algorithm, 560, 562, 591
 with Yule–Walker approach, 560–2, 591
- peak width, 576
- Pearson product moment correlation coefficient, 3
- percentage of block overlap in WOSA, 415
- periodic extension, 54, 92
- periodic function, 48–52, 74, 95–7
 convolution theorems, 96–7
 Fourier representation, 49, 95–6
 Parseval’s theorem, 49, 96
 spectral properties, 50, 96
- periodicity
 compound, 540–1, 587
 elimination by differencing, 40
 simple, 540–1, 586
 testing with white noise, 538–44, 586–7
 Fisher’s test for (simple periodicity), 539, 586
 example of use, 571, 573, 581
 irregular sampling (Lomb–Scargle), 543–4
 Schuster’s test for (simple periodicity), 539
 Siegel’s test for (compound periodicity), 541, 587
 example of use, 581
- Thomson’s *F*-test for (colored noise), 544–8, 587
 example of use, 574–5, 582–3, 592
- periodogram, 170–185, 232–3
 and DFT of time series, 116, 171
 and real-valued harmonic process plus white noise, 523

- average value of, 241
- averaging of independent, 441
- bias, 163, 176–7, 218
 - AR(2) process (low dynamic range), 177
 - AR(4) process (high dynamic range), 178–80
- asymptotic (unbiasedness), 175
- average value of, 239
 - due to use of log periodogram, 304–5, 307
- bootstrapping and, 615–6, 631
- computation of, 219–24
- covariances, 203–4, 235
 - verification via simulation, 242
 - subtle issues, 214
- DCT-based, 217–9, 244
- deficiency of power after removal of lines, 550–1
- definition, 170, 232
 - for complex-valued time series, 231
- discretely smoothed (*see* discretely smoothed periodogram)
- distribution of, 203, 235, 307–8, 408
 - verification via simulation, 242
- effect on periodogram of
 - centering of time series, 184, 213, 240
 - at nonzero Fourier frequencies, 184
 - at zero frequency, 184
 - circular shifting of time series, 238–9
 - scaling of time series, 238
 - time indexing of time series, 184
- examples of
 - AR(2) data, 172–3, 182, 208, 216, 246, 249, 294, 305, 310
 - AR(4) data, 172–3, 182, 199, 508
 - atmospheric CO₂ data, 335
 - atomic clock data, 327–8
 - chaotic beam data, 228
 - ice profile data, 323
 - ocean noise data, 231
 - ocean wave data, 226–7, 497
 - rotation of earth data, 224
 - harmonic process plus white noise, 533–4, 536, 551
 - ship altitude data, 331
 - three sinusoids (Kay and Marple), 531
 - white noise data, 207, 210
 - Willamette River data, 568, 571, 573
- expected value of, 174–5, 203, 233, 240
 - complex-valued harmonic process plus white noise, 524, 586
 - real-valued harmonic process plus white noise, 524, 536, 585
 - conditional on phase value, 537–8
 - verification via simulation, 242
- in circulant embedding simulation method, 615
- inconsistent estimator of SDF, 163–4, 207, 235
- Laplace, 529
- log likelihood, 308
- Lomb–Scargle, 528–9, 543–4, 588
- modified, 186
- power-law exponent estimation, 408
- reflecting boundary conditions, 244
- rescaled, 541, 581
- spurious peaks, 527
- sum over
 - Fourier frequencies, 237, 540
 - non-Fourier frequencies, 233, 237
- symmetry, 170
- three-point example to verify properties, 238
- truncated, 343, 504
- utility compared to multitaper spectral estimator, 403–12
- variance of, 203, 235, 241
 - for AR(1) process, 243
 - verification via simulation, 242
- phase function (phase shift function), 136
- phase-shifted Fourier transform, 383
- phases (*see* random phases)
- pilot analysis (filters for), 160
- pointwise convergence, 50, 53, 84
- Poisson's formula, 84
- polynomial regression model, 522
- positive
 - definite, 28, 170, 375, 380, 422, 450–2, 454, 457, 461, 464, 480
 - semidefinite, 28, 30, 167, 170, 374–5, 457
- positive-part function, 541, 636, 640
- postcoloring, 198–201, 234, 496, 615
 - AR model approach, 491–2, 510
 - atomic clock data, 329, 339–40
 - ocean wave data, 497–500
- power
 - of a nonperiodic function, 55
 - of a periodic function, 50
- power spectral density function (PSDF), 113, 116
- power-law exponent, 403, 407–12
 - estimation using linear regression and multitaper estimator, 409
 - atomic clock data, 430
 - periodogram, 408
 - periodogram versus multitaper estimation, 410–11
- power-law process, 327–30, 336–7, 403, 407, 430–1
 - ACVS, 391, 434, 605
 - spectral density function (SDF), 391, 434, 605, 610
- predictability of a time series, 17, 403–4, 471–2
- prediction error
 - backward, 455
 - forward, 453, 596
- prediction error filter, 198
- prewhitening, 179, 197–201, 447, 491–2, 503, 506, 615
 - versus multitaper spectral estimator, 424
 - ocean wave data, 496–8, 510
- prewhitening filter, 197, 234, 241, 447, 482, 491–3, 496, 510, 615
- principle of maximum entropy, 447, 471, 473
- probability density function (PDF), 24
 - bivariate standard Gaussian, 634–5
 - chi-square with two degrees of freedom, 37
 - exponential (scaled chi-square with two degrees of freedom), 37, 209
 - gamma (scaled chi-square with arbitrary degrees of freedom), 297

- generalized Gaussian (GG), 633
- Gaussian (normal) with zero mean, 55
- modeling, 631
- spectral window as an example of, 192
- uniform
 - over interval $(-\pi, \pi]$, 37, 45, 598
 - over interval $(-\sqrt{3}, \sqrt{3}]$, 634
- probability integral transform, 634
- probability of an event, 23
- process
 - band-pass, 299, 347
 - high-pass, 299, 347
 - low-pass, 299, 347
 - residual, 550
 - with stationary differences of order d , 340
- profile likelihood, 482–4
- projections, 384, 439
- prolate spheroidal wave functions (PSWFs), 64
- proper
 - complex-valued Gaussian or normal RV, 546–7
 - complex-valued white noise, 517–8, 564
 - process, 30, 32
- PSDF (*see* power spectral density function)
- pseudo-AR process, 553
 - complex-valued, 563
 - finding the coefficients, 591
 - plus white noise is an ARMA process, 554
 - solution to defining equation, 554, 591
 - stationarity condition, 557
- SVD
 - parameter estimator, 566
 - sinusoidal frequency estimator, 567
- purely
 - acausal, 447–8
 - continuous spectrum, 120, 163, 165, 167, 511
 - continuous stationary process, 457, 532
 - discrete spectrum, 120, 511, 513
 - nondeterministic process, 404, 471
- QQ plot
 - Gaussian stationary process assumption, 621–2
 - IID Gaussian assumption, 621
- quadratic lag window (*see* Bartlett–Priestley lag window)
- quadratic spectral estimator, 374–82, 439
 - bias of
 - broad-band, 378
 - indicator, 379–81
 - minimizing indicator of, 380
 - for white noise, 377–8
 - local, 378
 - magnitude indicator, 378–9, 381
- deconstructed multitaper spectral estimator, 376
- examples of
 - direct spectral estimator, 375
 - lag window spectral estimator, 375
 - weighted multitaper spectral estimator, 376
 - justification for, 351
 - WOSA spectral estimator, 423
- expected value of, 376–7, 439
- regularization bandwidth, 377
 - variance of, 380
- quadratic taper, 248
- quadratic trend removal
 - by ordinary least squares (OLS), 332, 443
 - by differencing, 46, 333–4
- quadratic window, 248
 - estimator, 252
- radar clutter, 131
- random amplitudes, 35–7, 111
- random phases, 37–8, 111
 - nonuniformly distributed, 38, 45
 - uniformly distributed, 37, 511, 553, 598, 616
- random variable, 3, 21–3
 - complex-valued, 21, 23
 - vector-valued, 21–2
- random walk, 43
 - frequency noise, 431
- randomly phased sinusoid, 35, 511, 553, 598
 - ACVS for, 553, 591
 - plus white noise, 554
- rank of matrix, 375–6, 380–1, 420, 422–3, 490, 565–6
- Rayleigh
 - distribution, 37, 598
 - quotient, 89, 105
 - theorem, 49
- real part of complex-valued
 - process, 602, 606, 640
 - variable, 61, 486
- real-valued stochastic process
 - continuous parameter/time, 22
 - stationary, 26
 - discrete parameter/time, 22
 - stationary, 26
- realizations, 3, 21–2
- reciprocity relationships, 58–62
 - equivalent width, 58–9
 - fundamental uncertainty relationship, 60–2
 - similarity theorem, 58–60
- rectangular
 - data taper, 188–90, 535
 - kernel, 71, 290
- recursive maximum likelihood autoregressive
 - parameter estimator (RMLE), 485
- reduced (profile) likelihood, 482–4
- reflecting boundary conditions, 181, 183, 216, 219, 244, 424
- reflection coefficient (*see* partial autocorrelation coefficient)
- regularization, 377, 386
 - bandwidth, 357, 377, 425, 428, 434
 - versus spectral window bandwidth, 402
- rejection filtration, 201
- relation function, 31
- relative efficiency, 165
- remodulated filtered series, 592
- repeated lag window spectral estimator, 349–50
- rescaled periodogram, 541, 581
- reshaped
 - lag window, 257, 281–2, 375
 - spectrum, 547, 551, 575, 582, 592
 - Willamette River data, 575

- residual process (EULS), 550
 - periodogram of
 - harmonic process plus white noise, 551
 - Willamette River data, 570
- resistor, 21, 40–1
- resolution
 - loss of, 76, 78, 271, 290, 357, 359, 400, 434, 535
- restoration of power, 195
- Riemann integral, 24, 26, 112–13, 258, 261
 - approximation by a Riemann sum, 174, 254, 258–9, 404, 605–6
- Riemann sum, 174, 254, 258–9, 404, 605–6
- Riemann–Lebesgue lemma, 120, 167
- Riemann–Stieltjes integral
 - ordinary, 24–6, 109, 112, 127
 - stochastic, 110, 112, 127
- ringing, 196, 225, 227, 277, 320, 331
- ripples, 72, 77, 79, 152–3, 189, 253, 275, 290, 294
- robust estimation of SDF via WOSA, 419
- rotation of the earth, 148, 222, 366, 407
- rule of thumb for significance levels, 548, 574, 582

- sample autocorrelation, 3
- sample autocorrelation sequence (sample ACS), 3
 - examples, 4
 - correlation of, 4
 - variance of, 4
- sample mean, 3, 164–6, 184, 195–6, 213
 - variance for AR(1) process, 507
- sampled function, 74
- sampling and aliasing, 122–3, 569
- sampling interval, 84–5, 113, 119, 122–5, 142–3, 163, 207, 232, 352, 374, 432, 445, 505, 511
 - computational details, 220–4, 314–5
 - irregular, 382, 528–9, 543
 - jitter, 124
 - effect on oxygen isotope time series, 124
 - oversampling, 123
 - role in simulation of time series, 594, 601, 609, 613
 - subsampling, 124
 - example, 130–1
- scale preservation (LTI filters), 133, 140
- scatter plot
 - lag 1 and four examples, 2–3
 - lag τ , 2, 5
 - $\tau = 6$ and 9 examples, 6
 - locations of peak frequencies from estimated spectra, 537–8, 556–7, 589, 591
- Schur matrix, 484
- Schuster's test for simple periodicity, 539
- Schwarz inequality, 55, 57, 61, 104, 257
- SDF (*see* spectral density function)
- sea levels, 520
- seasonality/periodicity, 40, 332
 - removal by differencing, 40, 42, 333
- second differencing for quadratic trend removal, 46

- seismic trace of earthquake, 58
- semiperiodogram, 217
- semivariance, 32, 546–7
- sequency (Walsh functions), 138
- sequency-limited and time-limited, 140
- shading sequence, 186
- Shannon number, 66–7, 90, 357, 372–4, 381
- ship altitude data, 330–2, 626–8
 - analysis of longer series, 350
 - direct spectral estimator, 331
 - lag window spectral estimator, 331
 - periodogram, 331
 - plot of, 330, 627
 - simulation of, 626–8
- sidelobe leakage, 77, 185, 386, 388, 390
- sidelobes
 - of smoothing windows, 268–9, 273, 275–8
 - of spectral windows, 281–2
 - Fejér's kernel, 178–80, 328, 527, 532, 535–6
- Siegel's test for compound periodicity, 540–3, 581, 587
 - asymptotic critical value, 542–3, 581
 - exact critical value, 541, 581
 - interpolated critical value, 542, 581
- sign of i convention, 55
- signal-to-noise ratio, 526, 532, 543, 552, 560–1, 567, 578, 587
 - signal in, 526
- similarity theorem, 58–60
- simple periodicity, 540–1, 586
- simulation of
 - atomic clock data, 628–30
 - autoregressive (AR) process, 595
 - zero initial conditions, 595
 - fractionally differenced (FD) process, 440
 - Gaussian AR(1) process
 - via Kay's method, 639
 - Gaussian AR(2) process, 597, 602
 - via circulant embedding, 603
 - via Gaussian spectral synthesis method (GSSM), 608
 - via Kay's method, 597
 - examples, 34
 - Gaussian AR(4) process, 597
 - via circulant embedding, 640–1
 - via Gaussian spectral synthesis method (GSSM), 606, 641
 - via Kay's method, 597, 638
 - examples, 34
 - Gaussian ARMA process, 593–4
 - via Kay's method, 598, 639
 - via McLeod and Hipel's method, 600, 639
 - Gaussian autoregressive (AR) process
 - stationary initial conditions, 595, 599, 624, 637
 - via Kay's method, 597
 - via McLeod and Hipel's method, 599
- harmonic process, 593–4, 637
- moving average (MA) process, 594, 637, 639
 - MA(1) examples, 33
- non-Gaussian time series
 - ARMA, 631, 636, 638
 - with specified ACVS and PDF, 631, 633, 638

- ocean wave data, 619–26
 - uses, 626
- reduced-length time series
 - based on nonparametric SDF estimate, 614
- ship altitude data, 626–8
- time series, 593
 - based on
 - nonparametric SDF estimate, 611, 619–20, 627, 629, 637
 - with non-unity sampling interval, 613, 638
 - method of surrogate time series (MSTS), 615–6
 - parametric SDF estimate, 617, 638
 - periodogram, 615
 - centering, 619, 627–8
 - Gaussianity testing, 621
 - given a portion of its ACVS, 593, 601, 637
 - given its fully specified SDF, 593, 604, 637
 - role of sampling interval, 594, 609, 613
 - summary, 637–8
- white noise, 594, 637
 - examples, 31
- simulation via
 - circulant embedding, 601–4, 619–20, 627, 629, 631
 - ACVS properties, 602, 640
 - computational efficiency, 603
 - issues with ACVS, 615
 - overcoming negative weights, 604, 640
 - weights, 601–2, 605, 629, 640–1
 - Gaussian spectral synthesis method (GSSM), 605–6, 608
 - adjustment for power-law process, 610
 - and adaptive multitaper spectral estimator, 615
 - and prewhitening, 615
 - role of sampling interval, 609
 - weights, 640–1
 - method of surrogate time series (MSTS), 615, 641
 - relation of simulated process to harmonic process, 641
- sinc function, 63, 559
- single-sided versus double-sided SDF, 118
- singular value decomposition (SVD), 425, 564–5
 - and short time series, 567
 - approach to harmonic analysis, 563
 - Willamette River data, 578–80, 592
 - singular value, 565
 - singular vector, 565
- sinusoidal amplitude estimator
 - approximate conditional least squares (ACLS), 515, 525, 549, 569–70, 572–3, 585, 589
 - approximate least squares, 516, 585
 - exact conditional least squares (ECLS), 549, 569–70, 572, 589–90
 - including mean, 552
 - exact least squares
 - complex-valued, 517, 586
 - real-valued, 515, 525, 585
 - exact unconditional least squares (EULS), 549, 569–70, 572
 - including mean, 552
 - Laplace method, 529
 - Lomb–Scargle method, 528
 - multitaper (complex-valued), 546, 551, 574
 - sinusoidal frequency estimator
 - exact unconditional least squares (EULS), 549, 569–70, 572
 - including mean, 552
 - location of maximum (global or local) of AR SDF, 556–8, 591
 - forward and backward least squares, 556
 - direct spectral estimator, 537–8, 556–7, 585, 589
 - rationale for tapering, 535–7
 - periodogram, 526–8, 537–8, 589
 - at Fourier frequencies, 523–4, 585–6
 - at general frequencies, 524–5, 585
 - expected value of estimator, 526
 - convergence of estimator, 527
 - MSE of estimator, 526, 537
 - variance of estimator, 526, 534
 - Thomson's F -test, 574, 582–3
 - parametric approach, 553–8
 - problems with, 558–63
 - three-sinusoids example
 - noise-free (Kay and Marple), 530
 - effect of grid size, 530–2
 - with noise, 532
 - effect of noise variance, 533–4
 - effect of series length, 532, 534
- sinusoidal multitapers, 355, 372, 391, 393–4, 396, 434, 544, 574, 583, 590, 592
 - compared to Slepian multitapers, 393
 - defined, 392
 - equivalent width, 401, 441
 - versus spectral window bandwidth, 402
 - relationship to minimum-bias multitapers, 392
- sinusoidal power estimation (AR SDF), 578
- sinusoidal variations
 - deterministic, 553–4, 558, 590
 - expressible as AR model, 447
 - inexpressible as MA model, 503
 - randomly phased, 35, 511, 553, 561
 - ACVS for, 37, 553, 591
 - examples of realizations, 36
 - plus white noise, 512, 554
 - as ARMA process, 554
 - sum of, 6–7
- skewness, 631
 - ARMA(1,1) process, 632
 - inequality, 633, 641
- Slepian
 - data taper, 190–1, 206, 208, 225, 228, 234, 260, 263, 282, 287, 416, 461, 496, 535–6, 569, 591
 - approximation of zeroth-order, 196
 - lag window, 285, 345
 - multitapers, 355, 360, 362, 372, 391, 434, 544, 574, 582, 590, 592
 - compared to sinusoidal, 393
 - decomposition of energy, 367
 - equivalent width versus spectral window bandwidth, 369
 - sequence, 87, 105, 145, 189, 357
 - double orthogonality, 106
 - zeroth-order as a low-pass filter, 155

- smoothing of
 - direct spectral estimator, 245
 - log direct spectral estimator, 303
 - periodogram estimator, 246, 307
 - smoothing window, 245, 248, 341
 - autocorrelation width of, 251
 - bandwidth, 245, 251, 268, 281, 292, 296, 300–1, 318, 322, 341, 346–7
 - Bartlett–Priestley design, 344
 - choice of, 252
 - finer frequency grid, 255
 - for sea level residuals, 522
 - Grenander, 251, 268, 279, 281
 - lag window spectral estimator, 279
 - Parzen, 251
 - repeated lag window estimator, 349–50
 - bias due to, 256, 290
 - equivalent width of, 251
 - leakage, 268, 287–9, 294, 316, 318, 331, 498
 - parameter m (controls smoothing), 247, 287
 - choice of, 291, 304, 313–4
 - examples of, 318, 322, 328, 330, 339
 - properties of, 250
 - type
 - Bartlett, 269–70
 - Bartlett–Priestley, 274–5
 - Daniell (rectangular), 250, 272–3
 - Gaussian, 277–8
 - modified Daniell, 280
 - Papoulis, 278–9
 - Parzen, 275–6, 280–2, 344
 - rectangular, 271
 - spectral bandwidth, 292, 296, 346
 - spectral density function (SDF), 111
 - after first difference and seasonal filtering, 333–4
 - aliasing effect, 122–3
 - alternative definitions, 114–6
 - and LTI digital filters, 143
 - and subsampling, 130
 - asymmetry for complex-valued process, 118
 - basic properties, 116–7
 - classification examples, 121
 - confidence interval for, 205, 265–7, 342
 - for log of, 205, 227, 229, 266–7, 342
 - double-sided versus single-sided, 118
 - dynamic range, 177, 184, 197, 492, 641
 - for first difference of a process, 158, 326
 - for particular stationary processes
 - AR, 145, 446, 485, 609, 617–8
 - AR(1), 336
 - AR(2), 172–3, 609
 - AR(4), 172–3
 - computation using zero padding, 449
 - ARMA, 145, 159
 - band-limited white noise, 379
 - fractionally differenced (FD), 166, 407–8
 - MA, 130, 144
 - MA(1), 130
 - MA(2), 159
 - power-law, 605, 610
 - white noise, 117
 - Fourier transform of ACVS, 111, 114, 126
 - imputed from noise model, 336
 - linear and log scales, 336–8
 - log of, 301
 - one-sided versus two-sided, 118
 - nonparametric estimators of
 - direct spectral estimator, 186
 - discretely smoothed, 246–7
 - lag window estimator, 248
 - multitaper, 352
 - periodogram, 170
 - discretely smoothed, 307
 - WOSA, 414
 - robust estimation via, 419
 - parametric estimators of
 - AR, 446–7
 - Burg, 466
 - least squares, 475
 - maximum likelihood, 480
 - Yule–Walker, 451
 - MA, 503
 - single-sided versus double-sided, 118
 - symmetry for real-valued process, 111–12, 118, 126
 - two-sided versus one-sided, 118
 - type
 - Gaussian-shaped, 131
 - Lorenzian, 131
 - piecewise constant, 348
 - rectangular, 128
 - triangular, 128
 - unimodal, 297
 - units of (physically meaningful), 374
 - variance-preserving, 337–9
 - versus autocovariance sequence, 124–5
 - weighted sum of uncorrelated processes with SDFs, 130
- spectral distribution function, 117
 - spectral estimator of the Grenander–Rosenblatt type, 252
 - spectral factorization, 636
 - spectral lines, 518, 538–40, 543–4, 548–9
 - splitting of, 558, 560
 - spectral representation
 - interpretation, 110–11
 - of stationary process, 108
 - complex-valued, 114
 - continuous parameter, 113–4, 135
 - discrete parameter, 110, 125
 - nonzero mean, 113
 - with discrete spectra, 518–9
 - with mixed spectra, 518–9
 - theorem, 7, 108–10, 382, 518–9
 - uniqueness, 110
 - spectral reshaping, 547, 551, 575, 582
 - spectral window, 175, 186–8, 255, 353, 414
 - bandwidth, 192, 194, 213–4, 355
 - definitions of
 - autocorrelation width (standard), 194
 - correlation time, 213–4, 355
 - half-power width, 192
 - standard (autocorrelation width), 194
 - variance width, 192, 194
 - definitions (standard) for spectral estimators
 - direct, 194, 235
 - discretely smoothed periodogram, 347

- lag window, 256, 341
- multitaper, 353
- WOSA, 415
- frequency range for statistical results, 206, 213, 244, 268
- standard measure of, 194
- examples of use, 226–7, 229, 294, 317–8, 323, 325, 328, 331, 335, 568, 623–4
- versus regularization bandwidth, 358, 402
- versus smoothing window bandwidth, 296
- versus correlation time, 214, 368–9, 401–2
- bias, 192, 391
- and minimum-bias multitapers, 391–2
- due to central lobe (spectral window bias), 194, 535
- due to sidelobes (leakage), 194, 535
- decay rate, 281, 328
- definitions of for spectral estimators
- direct, 186–8, 234
- lag window, 255, 341
- multitaper, 353
- periodogram, 175
- WOSA, 414
- type
- autocorrelated 100% cosine lag window, 285
- Bartlett lag window, 268, 270
- Bartlett–Priestley lag window, 268, 274
- Bartlett–Priestley taper, 288
- Daniell (rectangular) lag window, 268, 272
- Daniell taper, 288
- Gaussian lag window, 268, 278
- Gaussian taper, 288
- Hanning taper, 191
- modified Daniell window, 280
- 100% cosine lag window, 284
- Papoulis lag window, 268, 279
- Papoulis taper, 288
- Parzen lag window, 268, 276
- $p \times 100\%$ taper, 191
- periodogram, 176
- Slepian lag window, 286
- Slepian multitapers, 358, 364–5
- k th-order taper, 361, 363, 368
- Slepian taper (zeroth-order), 191, 288
- sinusoidal multitapers, 398–401
- k th-order taper, 395, 397
- standard bandwidth approximation, 400–1
- spectrograph estimator, 252
- spectrum, 107–8
- classification of, 120–2
- discrete, 120, 511, 513, 518, 538, 584, 586
- for simple time series model, 5–11
- constant, 11
- estimated
- nonparametric, 11–13
- parametric, 14–5
- high frequency, 9–10
- low frequency, 9–10
- theoretical, 9–11
- integrated, 120
- interpretation via band-pass filtering, 147–8
- line, 120, 511, 522
- mixed, 120, 511, 513, 518, 521–2, 539, 584–5
- purely
- continuous, 120, 165, 167
- discrete, 120, 511
- spectrum analyzers, 419, 444
- spinning rotor, 39
- spontaneous line splitting (*see* spectral lines)
- spurious peaks in
- FBLs pseudo-AR sinusoidal frequency estimator, 567
- periodogram, 527
- square integrable, 48, 53–4, 75
- squared norm, 470
- St. Paul temperature data, 5–6
- standard frequencies (Fourier frequencies), 8, 91, 515
- stationarity
- conditions for
- autoregressive (AR) process, 145, 446–9, 557
- pseudo-AR process, 557
- second-order/weak/covariance, 26–7
- strict/complete/strong, 26, 631, 634
- stationary differences of order d
- process with, 340
- stationary process, 4, 7, 21, 26–31, 107
- and modeling data, 38–41, 45
- examples of
- autoregressive (AR), 33
- autoregressive moving average (ARMA), 35
- bandlimited white noise, 379
- harmonic, 35
- fractionally differenced, 166
- moving average (MA), 32
- white noise, 31
- intrinsically of order d , 340
- nonergodic, 128
- reverse of conjugate of complex-valued process, 502
- reverse of real-valued process, 455, 477, 484
- step function, 24, 26, 120
- step-down Levinson–Durbin recursions, 460–1
- and causality assessment, 453, 509
- and generation of ACVS for an AR process, 460–1, 508
- and simulation of ARMA process, 594–8
- stochastic continuity, 114, 118
- stochastic process, 21–2, 23–5
- jump, 109
- orthogonal, 109
- storm surges, 522
- strict stationarity, 26, 631, 634
- sum of squares (SS), 467, 476, 514
- summability
- absolute, 165
- Cesàro, 78, 165
- superposition (LTI filters), 133, 140
- SVD (*see* singular value decomposition)
- symmetry, 27
- conjugate, 29–30, 48, 54
- Szegő–Kolmogorov theorem, 404
- tables
- AR-based estimates of innovation variance, 500

- Burg's algorithm and sinusoidal frequency estimation, 559
- characteristics of AR SDFs and periodogram for Willamette River data, 578
- comparison of
 - spectral bandwidth measures for direct spectral estimator, 214
 - spectral bandwidth measures for sinusoidal multitaper spectral estimator, 402
 - spectral bandwidth measures for Slepian multitaper spectral estimator, 369
 - window bandwidth measures for discretely smoothed direct spectral estimator, 255
 - Yule-Walker AR(4) and AR(8) coefficient estimates, 459
- exact and approximate critical values for Siegel's test, 542
- harmonic analysis and centering for Willamette River data, 583
- horizontal and vertical components of crisscrosses (ice profile data), 325
- multitaper-based estimates of innovation variance, 432
- NMSE criteria for GSSM-based approximations for AR(4) process, 607
- parameter estimates in harmonic analysis for Willamette River data, 570
- properties of lag windows, 279
- ratio of variances for estimators of power-law exponent, 411
- reduced likelihoods for estimated AR(4) coefficients, 483
- true and estimated sinusoidal frequencies, 533
- variance inflation factor for tapers, 260
- taper, 186
 - compared to lag window, 282–3
 - advantages/disadvantages in use of
 - better performance of Yule-Walker autoregressive spectral estimator, 452, 461
 - decrease in bias due to sidelobes, 179, 185, 189, 234, 535–6
 - decrease in effective sample size, 194–5
 - decrease in pairwise correlation, 207
 - increase in bias due to central lobe, 192, 535
 - converting
 - taper into lag window, 283
 - lag window into taper, 285
 - linear, 186
 - normalization
 - unit energy, 188
 - data-dependent, 195
 - orthogonality, 354, 357, 392, 432
 - plethora of types, 196–7
 - quadratic, 248
 - type
 - Bartlett-Priestley, 287
 - Daniell, 287
 - default, 188, 190
 - Gaussian, 287
 - Hanning, 189–90, 230, 234, 327, 331, 415–7
 - Hanning (alternative definition), 240
 - k th-order minimum-bias, 392–3
 - k th-order sinusoidal, 392–4, 396
 - k th-order Slepian, 357, 360, 362, 368
 - minimum-bias multitapers, 392–3
 - $p \times 100\%$ cosine, 189–90, 260
 - Papoulis, 287
 - Parzen, 287
 - rectangular, 188–90, 535–6
 - sinusoidal multitapers, 355, 392–3, 402–3
 - Slepian, 190, 196, 212, 225, 228, 230, 234, 260, 263, 287, 416, 461, 496, 535–6
 - Slepian multitapers, 355, 357, 372–3
 - variance inflation factor, 260, 262, 342, 417
 - temporal window, 186
 - terrestrial free oscillations, 359, 366
 - test for
 - periodicity
 - Fisher's, 540–1, 586–7
 - in colored noise, 544, 587
 - in white noise, 538, 586–7
 - Schuster's, 539
 - Siegel's, 541, 587
 - Thomson's F -test (multitaper approach), 228, 359, 544–8, 574–5, 582–3, 587, 592
 - with irregular sampling, 543–4
 - white noise (cumulative periodogram), 215–6, 230–1, 573–4
 - textile slivers, 33, 503
 - Thomson's F -test for periodicity, 228, 359, 544–8, 574–5, 582–3, 587, 592
 - tidal
 - analysis, 40, 520–3
 - frequencies, 521
 - time indexing of a series
 - and periodogram, 184
 - time invariance (LTI filters), 133, 140
 - time series simulation, 593
 - based on
 - nonparametric SDF estimate, 611, 619–20, 627, 629, 637
 - with non-unity sampling interval, 613, 638
 - method of surrogate time series (MSTS), 615–6
 - parametric SDF estimate, 617, 638
 - periodogram, 615
 - centering, 619, 627–8
 - examples of,
 - atomic clock data, 628–30
 - ocean wave data, 619–26
 - uses, 626
 - ship altitude data, 626–8
 - for testing of Gaussianity, 621
 - given a portion of its ACVS, 593, 601, 637
 - given its fully specified SDF, 593, 604, 637
 - non-Gaussian
 - ARMA process, 631, 636, 638
 - with specified ACVS and PDF, 631, 633, 638
 - reduced-length
 - based on nonparametric SDF estimate, 614
 - role of sampling interval, 594, 609, 613
 - summary, 637–8
 - time-limited function, 58
 - time reverse of
 - conjugate of complex-valued stationary process
 - is stationary with same ACVS, 502

- real-valued stationary process is stationary with same ACVS, 455, 477, 484
- time-reversibility, 484
- Toeplitz
 - matrix, 29, 156, 457, 462, 480, 484
 - Hermitian, 30
 - system of equations, 452, 461
- trace of a matrix, 376, 378, 439
- trade-off
 - between bias and variance, 80, 296, 309
 - between fidelity and resolution, 80
- transfer function, 136, 142–3, 290
 - and
 - gain function, 136
 - phase function, 136
 - as Fourier transform of impulse response sequence, 143
 - for
 - AR filter, 144–5, 487
 - Hilbert transform, 579, 592
 - MA filter, 144
- transpose of vector, 28
- trend, 40, 151, 332
 - removal, 39, 334, 443
 - as high-pass filtering, 151
- tridiagonal matrix, 88, 95
- trigamma function, 296, 302, 405, 410
- truncated periodogram, 343, 504
- twin peaks, 76–8, 178–9, 192, 200–1, 218–9, 291, 294, 300–1, 306, 337, 382, 458–9, 489
- two-sided versus one-sided SDF, 118

- uncorrelated random variables
 - definition, 31
 - relationship to independent random variables, 31–2
- undersmoothing, 71
- uniform distribution
 - over interval $(-\pi, \pi]$, 37, 45, 598
 - over interval $(-\sqrt{3}, \sqrt{3}]$, 634
- uniqueness of
 - Fourier transform, 55, 82
 - integrated spectrum, 135
 - spectral representation, 110
 - stationary solution to AR equation, 144–5, 446–8
- unit circle, 446, 448–9, 463, 553, 557–8, 566–7, 580, 591–2, 600, 618

- variability of SDF estimators
 - assessment via
 - asymptotic theory, 204–5, 265–7, 342, 354, 485–91
 - simulation, 622–6
- variance
 - definition of
 - complex-valued case, 25
 - real-valued case, 24
- variance inflation factor due to tapering, 260, 262, 342, 417
- variance of
 - linear combination of random variables, 41
 - sample mean for AR(1) process, 507
- sinusoidal frequency estimator using periodogram maximum, 526–7, 534, 592
- spectral estimators
 - basic multitaper, 353–4, 359
 - for Gaussian white noise, 372–4, 402–3
 - direct, 204, 235, 624
 - for complex-valued process, 232
 - verification via simulation, 242
 - lag window, 258–9, 262, 265, 279, 341
 - periodogram, 203, 235, 241
 - for AR(1) process, 243
 - verification via simulation, 242
 - quadratic, 380
 - weighted multitaper, 354
 - WOSA, 415, 435
- stationary process, 27–8
 - bias in estimator (unknown mean), 168, 237
- variance-preserving SDF, 337–9
- variance spectrum for
 - harmonic process, 108
 - simple time series model, 8, 47
- variance-stabilizing transform, 13, 209
- variance width of function, 60, 73, 105, 191–2, 194, 288
- vector transposition, 28

- Walsh functions, 138–40
- weighted covariance estimator, 252
- weighted multitaper spectral estimator, 352–4, 388, 433, 611–13
 - and quadratic spectral estimator, 376
 - bias of, 391
 - covariances of, 437
 - distribution of, 354–5
 - equivalent degrees of freedom (EDOFs) of, 354–5
 - equivalent width, 355, 437
 - expected value of, 352, 389
 - in circulant embedding simulation method, 611–13
 - low-rank approximation, 422–3
 - Parseval's theorem in expectation, 390
 - spectral window for, 352–3, 391
 - bandwidth of, 353
 - variance of, 354
 - via regularization, 382
- weighted overlapped segment averaging (*see* WOSA spectral estimator)
- weights for circulant embedding simulation method (*see* circulant embedding simulation method)
- Welch's overlapped segment averaging spectral estimator (*see* WOSA spectral estimator)
- white noise, 31–2, 120, 169, 184, 197, 236, 381, 386–8, 446, 450, 511, 513, 524, 532, 534, 538, 586
 - and direct spectral estimator, 189, 371
 - and periodogram, 177, 207, 371
 - and quadratic spectral estimator, 377
 - and tapering, 195
 - autocovariance sequence (ACVS), 31
 - band-limited, 379
 - complex-valued, 32

- and proper, 517–8, 564
- continuous parameter, 38
- examples of, 31
- Gaussian, 201, 539
- spectral density function (SDF), 117
- simulation of, 594, 637
- test for (cumulative periodogram), 215, 230–1
- versus colored noise, 120, 126, 241, 372
- white noise variance estimator (harmonic analysis)
 - approximate conditional least squares (ACLS), 549, 570
 - exact conditional least squares (ECLS), 549, 552, 570
 - exact least squares
 - complex-valued, 518
 - real-valued, 516
 - exact unconditional least squares (EULS), 550, 552, 570
 - multitaper, 551
- Whittle likelihood, 307
- Wiener–Khinchine theorem, 118
- Willamette River data, 1–2, 567–8
 - AICC order selection, 575–6
 - analytic series version, 578–81, 592
 - Burg autoregressive spectral estimator, 575–7
 - cumulative periodogram test of residuals, 573–4
 - direct spectral estimator, 569, 591–2
 - FBLs autoregressive spectral estimator, 592
 - fitted values, 571–2
 - harmonic analysis of, 567–81
 - effect of centering, 583
 - integrated autoregressive spectrum, 577
 - multitaper spectral estimator, 575
 - parametric approach to harmonic analysis of, 575–8
 - periodogram, 568
 - of residuals, 570–1, 573
 - plot of
 - complex roots for, 580
 - fitted model, 571–2
 - residuals, 571–2
 - remodulated filtered series, 592
 - reshaped spectrum, 575, 592
 - residuals, 571–2
 - small portion of, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - sample ACVS for, 169
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
 - SVD approach to harmonic analysis of, 578, 592
 - table of characteristics of AR SDFs and periodogram, 578
 - Thomson's F -test for periodicity, 575, 592
 - Yule–Walker autoregressive spectral estimator, 592
- wind speed data, 1–2
 - ACS for
 - sample, 4
 - theoretical, 10
 - lag 1 scatter plot for, 3
 - sample ACVS for, 169
 - simulation of, 11
 - spectrum for
 - estimated, 12, 14
 - theoretical, 10, 12, 14
- window
 - data, 186
 - design, 247
 - for Bartlett–Priestley, 273–4, 319, 344
 - for Daniell, 272, 319
 - lag (*see entry for lag window*)
 - linear, 186
 - quadratic, 248
 - smoothing (*see entry for smoothing window*)
 - spectral (*see entry for spectral window*)
 - temporal, 186
- window closing, 293
 - and WOSA, 416
 - examples, 293–5
- windowed periodogram estimator, 252
- windows and convergence factors, 80
- Wold's decomposition theorem, 404, 471
- Wold's theorem, 117–8, 126, 185
- WOSA spectral estimator, 351, 412–19, 432, 434–6
 - ACVS estimator for, 420, 435
 - AR(2) data, 416–7
 - AR(4) data, 417–8
 - at zero frequency, 420
 - boundary conditions
 - circular, 424
 - reflecting, 424
 - centering for unknown mean, 419–20
 - computational efficiency, 419
 - connection to Bartlett lag window spectral estimator, 270
 - deconstructed multitaper spectral estimator, 420–1, 423
 - EDOFs of, 415–6, 423–4, 442–3
 - expected value of, 414, 435
 - for approximating a multitaper spectral estimator, 424–5
 - in circulant embedding simulation method, 611, 613
 - ocean wave data, 425, 429–30
 - robust SDF estimation, 419
 - shift factor restriction, 420
 - variance of, 415, 435
 - versus multitaper spectral estimator, 414, 417–9, 424
- Yule–Walker autoregressive estimator, 449–52, 505
 - and ACVS for direct spectral estimator, 452, 461–2, 505
 - AR(1) process, 510
 - AR(2) numerical example, 507–8
 - AR(4) data, 458–9, 461–2, 508–9
 - assessing variability (SDF), 485
 - causality, 448, 505

- complex-valued process, 502
- corresponding ACVS, 451, 504, 508
- FPE criterion, 494
- ocean wave data, 496–9
 - order selection via PACS, 498–9
 - order of, 458–9
- peak shifting and splitting, 560–2, 591
- relation to
 - autocorrelation method, 480
 - Burg's algorithm, 471
 - forward least squares, 479
 - Willamette River data, 592
- Yule–Walker equations, 450, 561
 - arising from two related problems, 454–5
 - augmented, 452
 - Levinson–Durbin recursions and, 457

- zero padding, 93–4
 - use in computing AR SDF, 449
 - finer frequency grid, 179, 527, 558, 568
 - for noncyclic convolutions, 106, 169–70, 220
- zero-memory nonlinearity (ZMNL), 634
 - correlation distortion, 635–6