

Index

- “adjustment for covariance” approach, continuous
  - outcome variables 83
  - software packages 282–91
- alternative methods, missing data analysis 234–5
- alternative models
  - autoregressive model 107–8
  - comparison with other alternative models 116–17
  - comparison to standard model 108–14
  - dichotomous outcome variables 138–9
  - experimental study 117–18
  - model of changes 105–7
  - overview 108
  - time-lag model 103–5
- Amsterdam Growth and Health Longitudinal Study 117–18, 153–60
- analysis of covariance 167–8
  - experimental studies 170–2, 176, 181–2
  - longitudinal 188–92, 195–6, 200–1
  - multivariate (MANCOVA) 184–5
- ANOVA (analysis of variance) 24, 33–4
  - see also* univariate analysis of variance
- area under the curve (AUC) 182–3
- autoregressive correlation structure 58–9
- autoregressive model 107–8, 138–9
  - comparison with longitudinal analysis of covariance 188–92
  - comparison with time-lag model 116–17
  - GEE analysis with 109–12
  - mixed model analysis using 113–14
- Barthel index 294–6
- categorical outcome variables
  - example 143–6
  - imputation methods 224–5
  - modeling of time 91–3
  - more than two measurements 142–3
  - relationships with other variables 146–53
  - software 281–2
  - two measurements 141–2
- causal relationships 103
- causality criteria 1–2
- “ceiling” effects, continuous outcome variables 166, 292–4
- changes, modeling of *see* model of changes
- Chi-square test 74
  - see also* McNemar test
- classification of subjects with different developmental trajectories 301–4
- clinical trials *see* experimental studies
- Cochran’s Q test 121, 123–4
- cohort effects 8–10
- “combination” approach 193–6
- comparison of groups *see* group comparisons
- compound symmetry assumption *see* sphericity assumption
- conditional model *see* autoregressive model
- confounding effects 7–13, 15, 102
- continuous outcome variables
  - comments 42–3
  - comparing groups 35–7
  - example dataset 38–41
  - “univariate” approach, a numerical example 37–8
- different contrasts 45–6
  - example dataset 46–8
- experimental studies
  - with more than one follow-up measurement 179–201
  - with only one follow-up measurement 165–76
- more than two measurements 20–3
  - example dataset 29–34
  - numerical example 27–9
  - shape of relationship between an outcome variable and time 26–7
  - univariate approach, a numerical example 23–6
- non-parametric equivalent of MANOVA for repeated measurements 48–50
  - example dataset 50
- non-parametric equivalent of paired *t*-test 18–19
  - example dataset 19–20

317 Index

- post-hoc procedures 44
  - example dataset 44–5
- relationships with other variables 51
  - comparison between GEE and mixed model analysis 81–5
  - example dataset 53–5
  - GEE analysis 57–68
  - longitudinal methods 55–7
  - mixed model analysis 69–81
  - “traditional” methods 51–3
- sample size calculation 237–8
- time modeling 86–93
- two measurements 16–17
  - example dataset 17–18
- univariate vs. multivariate approach 34–5
- contrasts (within-subjects) 45–6
- examples of different 46–8
- “one-within” design 32–3
- post-hoc procedures 48
- correlation structures
  - choice of working 57–60, 61–3
    - for “count” outcome variable 154–8
    - for dichotomous outcome variables 128–33
  - different structures, results with 66–8
  - exchangeable, results of using 63–6
  - software extensions for mixed model analysis 84
- “count” outcome variables 153
  - comparison between GEE and mixed model analysis 160–1
- example datasets 153–4
  - GEE analysis 154–8
  - mixed model analysis 158–60
- software packages 281–2
- Cox proportional hazards regression for recurrent events 208–10
- cross-over trials 14–15
- cross-sectional imputation methods 222
- cross-sectional (traditional) analysis 125–6
  - categorical outcome variables 146–7
  - continuous outcome variables 51–5
  - “count” outcome variables 153
  - dichotomous outcome variables 125–6
  - in experimental studies 210–11
  - summary statistics 182–3
- data augmentation (DA) 226–7
- data structures 4
- development over time *see* time modeling
- developmental trajectories, classification of 301–4
- dichotomous outcome variables
  - comparing groups 121
  - example dataset 122
    - comparing groups 124–5
    - development over time 122–4
  - experimental studies 201
  - other approaches 208–10
  - simple analysis 202–3
  - sophisticated analysis 203–7
  - imputation methods 224–5
  - more than two measurements 121
  - relationships with other variables
    - alternative models 138–9
    - comparison between GEE and mixed model analysis 136–8
    - example dataset using GEE and mixed model analysis 128–36
    - sophisticated methods 126–8
    - “traditional” methods 125–6
  - sample size calculations 238, 239
  - two measurements 119–20
- “difference” contrast 48
- dispersion (scale) parameter, GEE analysis 64, 129, 156
- drop-outs *see* missing data
- epsilon (sphericity coefficient) 31–2
- error sum of squares *see* sum of squares
- estimation procedures
  - GEE and mixed model comparisons 82
  - maximum likelihood 80–1
  - penalized quasi-likelihood (PQL) 279–80
  - restricted maximum likelihood (REML) 73–4, 254–6
- “eta squared”, effect magnitude, MANOVA 33
- example datasets 2–4
- exchangeable correlation structure 58
  - GEE analysis with 92, 99
  - results of linear GEE analysis using 63–6
  - results of Poisson GEE analyses 156–8
- exchangeable covariance structure 282
- example dataset 283–8
- experimental studies 6–7, 13–15, 163–5
  - comments 210–11
  - continuous outcome variables
    - more than one follow-up measurement 179–200
    - only one follow-up measurement 165–76
  - dichotomous outcome variables 201–10
- explained variance 13, 33, 41, 64–5
- F-statistic 22–3
  - ANOVA example 23–6
  - MANOVA for repeated measurements 29–34
- “fit” of a model 116–17
  - indicators 255–6, 262, 270–1
  - and log (restricted) likelihood value 74
- “floor” effects, continuous outcome variables 166, 292–4
- example dataset 294–8
- follow-up measurements 163–5
  - studies with more than one 179–80
  - dichotomous outcome variables 201–10
  - MANOVA for repeated measurements 183–5
  - sample size calculations 238–9
  - simple analysis 180–2

**318 Index**

- follow-up measurements (*cont.*)
  - sophisticated analysis 187–200
  - summary statistics 182–3
  - studies with only one 165–70
    - example dataset 170–6
- Friedman test statistic 48–50
- GEE (generalized estimating equations) 57
  - alternative models 109–12
  - continuous outcome variables
    - comparison with mixed model analysis 81–5
    - different correlation structures 66–8
    - regression coefficients for covariates 60–1
    - results of GEE analysis 63–6
    - selection of correlation structure 61–3
    - software packages 243–9
  - correlation structure, choice of correct 57–60
  - “count” outcome variables 154–8
    - comparison with mixed model analysis 160–1
  - dichotomous outcome variables 128–33
    - comparison with mixed model analysis 136–8
    - intervention effects 203–7
    - software packages 249–53
  - missing data 218–21
    - comparison with mixed model analysis 235–6
  - time modeling 86–93
    - adjustment for time 99–102
    - group comparisons 95–9
- generalized linear model (GLM) *see* MANOVA for repeated measurements
- GENMOD procedure, SAS software 244, 249–50
- GLLAMM procedure, Stata software 148, 281–2, 294
- Greenhouse–Geisser method, sphericity 31–2
- group comparisons, development over time 95–9
  - categorical outcome variables 143
  - continuous outcome variables 35–41
  - dichotomous outcome variables 121
  - “proportion of change” 124–5
- hazard ratio 209–10
- “Helmert” contrast 48
- “hot-deck” imputation method 222
- Huber–White sandwich estimator 59
- imputation methods, missing data
  - alternative approaches 234–5
  - comments 234
  - continuous outcome variables 221–4
  - dichotomous and categorical outcome variables 224–5
  - example dataset 225–34
- independent correlation structure 57–8
  - results of GEE analyses using 129–33
  - results of Poisson GEE analyses 156–8
- independent sample *t*-test 180–1
  - missing data analysis 216–17
- inter-period correlation coefficient (IPC) 12–13
- intervention effects *see* experimental studies
- last value carried forward (LVCF) 222, 225–8
- latent class growth analysis (LCGA) 303
- latent class growth mixture modeling (LCGMM) 303
- learning/test effects 11–12
- likelihood ratio test 74–6, 88–9, 135, 160
- linear GEE analysis *see* GEE, continuous outcome variables
- linear mixed model analysis *see* mixed model analysis, continuous outcome variables
- log (restricted) likelihood value 74
- logistic GEE analysis 128–33
  - comparison with mixed model analysis 136–8
  - “count” outcome variables, software for 282
  - intervention effects 203–7
  - software packages 249–53
- logistic mixed model analysis
  - dichotomous outcome variables 133–6
    - comparison with logistic GEE analysis 136–8
  - missing data 231–4
  - software packages 267–80
  - intervention effects 203–7
- logistic regression analysis, dichotomous outcome variables 126–8
- “long-term exposure” to covariates 51, 54, 126
- longitudinal analysis of covariance 188–92
- longitudinal imputation methods 222
- longitudinal logistic regression 126–8
- longitudinal statistical methods 55–7
- longitudinal studies 1
  - see also* experimental studies
- longitudinal tobit regression 293–7
- LVCF (last value carried forward) 222
  - example illustrating 225–8
- MANCOVA (multivariate analysis of covariance)
  - for repeated measurements 184–5
- MANOVA (multivariate analysis of variance) for repeated measurements
  - contrasts 45–8
  - drawbacks 42–3
  - experimental studies 183–5
  - missing data 219–21, 225
  - “one-within” design 20–3
    - non-parametric equivalent (Friedman test) 48–50
  - “one-within, one-between” design 35–41
  - post-hoc procedures 44–5
  - shape of the relationship between outcome variable and time 26–7
    - numerical example 27–9
  - results from example dataset 29–33
  - results from “naïve” ANOVA analysis 33–4
  - univariate approach within 23–6, 34–5
- Markov model *see* autoregressive model
- maximum likelihood estimation *see* restricted maximum likelihood (REML)

319 Index

- McNemar test 120
  - limitations of 122
  - multivariate extension of 121
- mean, regression to 166–70
  - effect of ignoring, example dataset 170–3
- missing data 212–14
  - analysis of determinants for 216–18
  - analysis performed on datasets with 218–19
    - example dataset 219–21
  - conclusions 236
  - GEE analysis vs. mixed model analysis 235–6
  - generating datasets with 215
  - ignorable or informative 214–15
  - imputation methods
    - alternative approaches 234–5
    - comments 234
    - continuous outcome variables 221–4
    - dichotomous and categorical outcome variables 224–5
    - example dataset 225–34
- mixed model analysis 69–70
  - alternative models 113–14
  - categorical outcome variables 148–53
  - continuous outcome variables 70–2
    - comments 80–1
    - comparison with GEE analysis 81–5
    - example datasets 73–80
    - missing data 226–30
    - software 253–67
  - “count” outcome variables 158–60
    - comparison with GEE analysis 160–1
  - dichotomous outcome variables
    - comparison with GEE analysis 136–8
  - experimental studies 203–7
  - missing data 231–4
  - relationship with several covariates 133–6
  - software 267–80
- missing data 218–19, 220–1, 234
  - vs. GEE analysis 235–6
- MLwiN (multilevel analysis for windows)
  - mixed model analysis
    - continuous outcome variables 264–6
    - dichotomous outcome variables 278–80
- model of changes 105–7, 138–9
  - comparison with other alternative models 116–17
- example dataset 108–9
  - data structure 109
  - GEE analysis 109–12
  - mixed model analysis 113–14
  - experimental study 117–18
- modeling of time *see* time modeling
- multilevel analysis *see* mixed model analysis
- multinomial logistic mixed model analysis 148–53
  - GLLAMM software procedure 281–2
- multinomial logistic regression analysis 147–8
- multiple imputation, missing data 223–4
  - in combination with mixed model analysis 226–9
  - reasons for combining 229
  - dichotomous outcome variables 231–4
    - (in)stability issues 229–30
- multiple longitudinal design 9–10
- multivariate analysis of covariance (MANCOVA)
  - for repeated measurements 184–5
- multivariate analysis of variance *see* MANOVA
- NLMIXED procedure, SAS 268–71
- non-parametric tests 18–19, 48–50
- observational studies 6, 7
  - example dataset 12–13
  - other confounding effects 11–12
  - period and cohort effects 8–10
- odds ratios 126, 203–7
  - interpretation of 129, 147, 149–50
- “one-within” design (MANOVA) 29–34
  - Friedman test 48–50
- “one-within, one-between” design, group comparisons 35–7
  - results from example dataset 38–41
  - “univariate” approach, numerical example 37–8
- outcome variables with upper or lower censoring 292–4
  - example 294–300
  - remarks 300–1
- overdispersion 162
- paired *t*-test 16–18
- penalized quasi-likelihood (PQL), estimation procedure 272–4, 279–80
- period (time of measurement) effects 8
- Poisson regression analysis, “count” outcome variables 153–4
  - GEE analysis 154–8
  - vs. mixed model analysis 160–1
- mixed model analysis 158–60
  - vs. longitudinal two-part model 299–300
- post-hoc procedures, MANOVA 44–5, 48
- posterior predictive distribution, missing data 224
- “proportion of change”
  - categorical outcome variables 142–3
  - example 143–6
- dichotomous outcome variables
  - example 122–5
  - group comparisons 121
  - more than two measurements 121
  - two measurements 120
- proportional hazards regression for recurrent events 208–10
- prospective cohort studies 6
- quadratic development over time 89–91, 96
  - example datasets 294–7
- “quasi-causal” relationships 103
- quasi-likelihood estimation procedures 57, 272–4, 279–80
- R software package
  - GEE analysis
    - continuous outcome variables 245–7

320 Index

- R software package (*cont.*)
  - dichotomous outcome variables package 251
  - mixed model analysis
    - continuous outcome variables 258–9
    - dichotomous outcome variables 271–4
- random coefficient analysis *see* mixed model analysis
- randomized controlled trials (RCTs) 14–15
  - see also* experimental studies
- recurrent events, Cox proportional hazards regression 208–10
- regression to the mean phenomenon 166–70
  - effect of ignoring 170–3
- relative change 168–70
- relative risk 202
  - and hazard ratios 209–10
  - and odds ratios 203
- REML (restricted maximum likelihood) 73–4
  - compared to maximum likelihood estimation 80–1
  - in software packages 254–6, 259, 266–7
- “repeated” contrast 48
- reproducibility of measurements 12–13
- research questions as basis for analysis 2
- “residual change” analysis 168
- restricted maximum likelihood (REML) estimation *see* REML
- robustness, GEE analysis 59, 66
- sample size calculations 237–40
  - example 240–2
- SAS software package
  - GEE analysis
    - continuous outcome variables 244
    - dichotomous outcome variables 249–50
  - mixed model analysis
    - continuous outcome variables 254–6
    - dichotomous outcome variables 268–71
- scale parameter, GEE analysis 64, 129, 156
- semirobust standard error 66
- signed rank sum test (Wilcoxon) 18–20
- simple analytical methods, dichotomous outcome variables
  - comparing groups 121
  - example dataset 122–5
  - experimental studies 202–3
  - more than two measurements 121
  - two measurements 119–20
- “simple” contrast 46–8
- single imputation methods 221–3
- software packages 4, 243
  - “adjustment for covariance” approach 282–3
    - example 283–91
  - categorical and “count” outcome variables 281–2
  - GEE analysis with continuous outcome variables
    - R 245–7
    - SAS 244
    - SPSS 247–8
    - Stata 243–4
    - summary 249
  - GEE analysis with dichotomous outcome variables
    - R 251
    - SAS 249–50
    - SPSS 252–3
    - Stata 249
    - summary 253
  - mixed model analysis with continuous outcome variables
    - MLwiN 264–6
    - R 258–9
    - SAS 254–6
    - SPSS 260–4
    - Stata 253–4
    - summary 266–7
  - mixed model analysis with dichotomous outcome variables 267–8
    - MLwiN 278–80
    - R 271–4
    - SAS 268–71
    - SPSS 274–8
    - Stata 268
    - summary 280
- Solomon four group design 14
- sophisticated analytical methods 126–8, 148, 187–200, 203–7
- sphericity assumption 22–3
  - example dataset 29–34
  - Greenhouse–Geisser adjustment 31–2
- SPIRIT study 297–300
- SPSS software package
  - GEE analysis
    - continuous outcome variables 247–8
    - dichotomous outcome variables 252–3
  - mixed model analysis
    - continuous outcome variables 260–4
    - dichotomous outcome variables 274–8
- Stata software package
  - GEE analysis
    - continuous outcome variables 243–4
    - dichotomous outcome variables 249
  - mixed model analysis
    - continuous outcome variables 253–4
    - dichotomous outcome variables 268
- (stationary) m-dependent correlation structure 58
- statistical notation 4
- statistical prior knowledge 2
- structural equation modeling 301–2
- Stuart–Maxwell test 141–2
  - example dataset 143–4
- sum of squares 24–6
  - individual 28, 34, 70–1
  - “univariate” approach numerical example 37–8
- summary statistics 182–3
- survival approaches 208–10
- t*-tests
  - independent samples 180–1
    - missing data analysis 216–17
  - paired 16–18
    - non-parametric equivalent of 18–20

321 Index

- test/learning effects 11–12
- time of measurement (period) effects 8
  - multiple longitudinal design 9–10
- time-lag model 103–5, 138–9
  - comparison with autoregressive model 116–17
  - GEE analysis 109–12
  - mixed model analysis 113–14
- time modeling
  - adjustment for time 99–102
  - comparing groups 95–9
  - development over time 86–93
  - more than two measurements 20–34
  - shape of relationship 26–7
  - therapy effect at different time points 196–200
  - two measurements 16–20
- tobit regression analysis 293–7
- “traditional” methods 51–3, 125–6, 146
- transformation “factors” 27–9
- trials *see* experimental studies
- two measurements
  - adjustment for covariance between 83
  - categorical outcome variables 141–2
  - continuous outcome variables 16–17
    - definition of change 165–6
    - example dataset 17–18
    - Wilcoxon signed rank sum test 18–20
  - dichotomous outcome variables 119–20
  - see also* MANOVA for repeated measurements
- two-part regression models 293–301
- univariate analysis of variance
  - numerical examples
    - “one-within, one-between” design 37–41
    - simple longitudinal dataset 23–6
  - sphericity assumption 22–3
  - vs. multivariate approach 34–5
- unpaired *t*-test *see* independent sample *t*-test
- unstructured correlation structure 59, 156–8
- unstructured covariance structure 283
  - results from using 283–91
- Wald statistic 126, 134, 265
- Wilcoxon signed rank sum test 18–20
- working correlation structures *see* correlation structures
- zero-inflated Poisson regression 293