

Index

- ACMV
 identifying restriction, 96
 add-on designs, 26
 analysis of variance, 51
 analytic road map, xviii, 112, 113, 130, 151, 152
 ascertainable outcomes, 29
 attrition. *See* dropout
 augmented wGEE
 and double robustness, 88
 autoregressive structure, 50
 auxiliary variables, 56, 81, 117, 135
- baseline observation carried forward (BOCF)
 defined, 20
 Bayesian inference, 39
 Bayesian regression, 98
 bias from missing data, 6
 BOCF, 20, 61, 62, 69, 76, 77, 78, 101, 116
- canonical link, 127
 categorical data, 121, 124, 125, 126, 128
 categorical outcomes
 analysis of, 121
 CCMV
 identifying restrictions, 95, 96, 135, 142
 centralized raters, 25
 classes of MNAR methods, 92
 clustering factors, 133
 Code of Federal Regulations, 35
 Cohen's D, 6
 complete case analyses, 59
 conditional inference, 121, 122, 123
 conditional model, 123
 confidence intervals, 99
 consistent estimators, 88
 defined, 4
 convergence
 ensuring, 54
 Cook's D statistic, 133
 use in influence diagnostics, 133
 correlation, 41
 correlations between repeated measures, 49
- data simulation, 147
de-facto hypotheses, 16, 18, 19, 20, 103, 104, 107, 108, 130, 147
de-jure hypothesis, 16, 17, 18, 19, 104, 106, 130, 147
 descriptive statistics and graphs
 clinical trial example, 136
 design approaches
 to lower rates of missing data, 148
 direct likelihood, 52, 81, 82, 84, 135
 distinction between MAR and MNAR, 11
 dose titration, 25
 double-blind trial, 20

162 **Index**

- doubly robust methods, 87
- dropout, 4
- dropout model, 85, 94, 134, 135
- effectiveness
 - defined, 16
- effectiveness estimand
 - clinical trial example of, 131
- efficacy
 - defined, 16
- efficacy estimand
 - clinical trial example of, 130
- enrichment designs, 23, 24, 148, 149
- estimand
 - defined, 8
 - ITT, 16
 - per protocol, 16
- estimands, 103
 - comparison of, 103
- estimation of estimand 1
 - example of, 104
- estimation of estimand 3
 - example of, 107
- estimation of estimand 4
 - example of, 107
- estimation of estimand 5
 - example of, 108
- estimation of estimand 6
 - example of, 108
- factorization
 - of likelihood function, 67
- Fisher's scoring, 54
- fixed effects, 50, 51, 52, 123, 133, 135
- flexible dosing, 25
- follow-up data, 15, 17, 28, 29, 103, 104, 106, 108, 109, 110, 149
- follow-up periods, 27
- frameworks for inference, 39
- frequentist inference, 39, 47
- full multivariate model, 53
- GEE, 39, 46, 47, 52, 55, 68, 84, 85, 86, 87, 88, 107, 121, 123, 124, 125, 128, 150
- general linear mixed effects model
 - for categorical data, 127
- general linear mixed-effects model, 51
- generalized estimating equations (GEE)
 - defined, 46
- Generalized least squares, 44
- Hamilton Depression Rating Scale, 50, 73, 130
- hierarchical model, 124
- hot deck imputation, 60
- identifying restrictions
 - in pattern mixture models, 95, 118, 135, 142
- identity link, 127, 135
- ignorability of missing data, 68 *See*
- ignorable missing data, 12, 125
 - defined, 12
- imputation model, 63, 82, 98, 104, 135
- incentives for completing treatment, 34
- inclusive modeling
 - clinical trial example, 140
- inclusive models, 56, 65, 89, 117, 134, 135, 143
 - for assessing sensitivity, 117
- independence
 - assumption of, 114
- influence diagnostics for patients
 - clinical trial example, 139
- influence diagnostics for sites
 - clinical trial example, 137
- influential observations
 - influence diagnostics, 115
- informative censoring, 12

163 **Index**

- informed consent, 35
 informed withdrawal of consent, 35
 Intention to Treat (ITT) principle, 10
 intent-to-attend, 34
 intermittent missing data, 4
 investigator payments, 34
 investigator selection, 34
 IPW
 inverse probability weighting, 65, 66, 67, 84, 88

 last observation carried forward (LOCF)
 defined, 37
 latent classes, 94
 latent variables, 94
 least squares estimation
 illustration of, 40
 likelihood function, 45, 46, 67, 68
 likelihood-based analyses
 example with complete data, 74
 example with incomplete data, 78
 likelihood-based methods. *See* maximum likelihood
 linear mixed model, 51, 124, 127, 128, 153
 linear relationships, 40
 linearity, 114
 assumption of, 114
 LOCF, 37, 55, 61, 62, 63, 69, 76, 77, 78, 101, 108, 116, 158
 logistic regression, 66, 84, 93, 115, 134, 135
 logit link, 101, 125, 127, 134
 loss to follow up
 consequences for missing data
 mechanisms, 11
 LSMEANS, 75, 77, 79, 82, 85, 105, 141

 MAR, 11, 12, 56, 64, 65, 66, 67, 68, 69, 78, 81, 84, 85, 88, 91, 96, 101, 110, 112, 113, 116, 117, 125, 128, 131, 134, 135, 136, 140, 142, 143, 144, 145, 147, 149, 151
 marginal and conditional inference
 illustration of, 121
 marginal distributions, 123
 marginal inference, 122, 123
 marginal models, 84, 123, 124, 128
 maximum likelihood, 39, 44, 45, 46, 47, 52, 54, 55, 56, 67, 79, 81, 88, 99, 101, 104, 107, 108, 131, 150
 maximum likelihood-based estimation
 illustration of, 44
 MCAR, 10, 11, 12, 52, 60, 61, 65, 67, 68, 78, 84, 112, 115, 124, 125, 133, 137, 147
 measurement process, 12, 67, 92, 93, 95
 methods of statistical estimation, 39
 minimizing missing data, 23
 missing at random (MAR)
 defined, 10
 missing completely at random (MCAR)
 defined, 10
 missing data mechanisms, 9, 10, 121
 missing not at random (MNAR)
 defined, 10
 missingness process, 12, 67, 68, 84, 92, 93, 95
 mixed-effects models, 51, 52, 126, 128
 MNAR, 11, 12, 57, 64, 65, 66, 68, 69, 91, 92, 93, 112, 115, 116, 117, 134, 135, 143, 144, 145, 147, 149, 151
 bias from, 145
 MNAR analyses
 for assessing sensitivity, 117
 model-based standard errors, 125
 modeling philosophies, 56
 monitoring rates of missing data, 34

164 **Index**

- multiple imputation, 63, 64, 65, 81, 82, 88, 98, 101, 107, 116, 128, 134, 135, 136, 150
 - example with incomplete data, 82
 - for categorical data, 127
- multivariate models, 53
- multivariate normal distribution, 124
- multivariate normal model, 51
- NCMV
 - identifying restriction, 96, 142
- New Drug Application, 129
- Newton-Raphson algorithm, 54
- non-future dependent missing value
 - restrictions (NFD), 96
- non-ignorable methods, 68
- normality, 114
 - assumption of, 114
- Ordinary least squares, 44
- outlier observations, 115
- participant burden, 33
- patient education, 35
- patient ratings, 25
- patient specific factors, 50
- pattern mixture model
 - clinical trial example, 142
 - specification for clinical trial example, 135
- pattern mixture models, 92, 94, 101, 102, 118, 134, 151
- placebo multiple imputation, 96, 118, 134
 - clinical trial example, 143
 - specification for clinical trial example, 136
- pMI, 118, 144
 - effectiveness context, 101
 - efficacy context, 99
 - example with missing data, 99
- poisson link, 127
- power
 - implications of missing data, 30
- primary estimand, 8, 15, 28, 36, 109, 110, 118, 131, 147, 150
- probability of dropout
 - consequences for missing data mechanisms, 11
- PROC GENMOD, 84, 85, 134, 135
- PROC IML, 135
- PROC MI, 98, 135
- PROC MI ANALYZE, 99, 135, 136
- PROC MIXED, 132, 133, 135
- PROC NL MIXED, 136
- property of double robustness, 88
- quasi-likelihood, 125
- random effects, 50, 51, 52, 53, 94, 123, 124, 126, 127, 131
- random variation, 50
- random-effects model, 123
 - for categorical data, 125
- random-effects models, 124
- randomization, 3, 147
- randomized withdrawal design, 24
- region of ignorance, 118, 144
- regression, 40
- rescue medications, 8, 15, 19, 27, 28, 29, 103, 104, 106, 108, 110, 149
- residual diagnostics
 - clinical trial example, 140
- residuals, 114, 115, 140
- restrictive models, 56, 81, 89, 104, 116, 117, 143
- retention. *See* dropout
- reverse factorization, 94
- run-in design, 23, 106

165 **Index**

- run-in designs, 23
- sample size, 30
- sandwich estimator, 55, 125, 132, 133
- selection model
 - clinical trial example, 140
 - specification for clinical trial example, 135
- selection models, 92, 93, 101, 115, 117, 134, 142, 151
 - semi-parametric implementation, 101
- semi-parametric estimators, 87
- sensible analysis, 114
 - defined, 4
- sensitivity analyses, xviii, 8, 65, 102, 111, 112, 116, 117, 118, 119, 130, 131, 134, 144, 150, 151, 152
 - defined, 8
- sensitivity to correlation assumptions
 - clinical trial example, 137
- separability condition, 12
- serial correlation, 50
- shared parameter model, 136
 - clinical trial example, 142
 - specification for clinical trial example, 135
- shared parameter models, 92, 94, 101, 118, 134, 151
- simple imputation strategies, 60, 69
 - example, 76
- software
 - for sensitivity analyses, 151
- standard deviation, 42
- standard error
 - defined, 4
- standardized effect size, 6
- substantive model, 88
 - in multiple imputation, 63, 65
- target population, 24
- time trends, 30, 49, 52, 53, 55
- titration dosing, 26, 52
- training for sites and patients, 34
- treatment changes, 10
- trial conduct
 - to lower rates of missing data, 149
- trimming weights
 - in wGEE, 87
- uncongenial imputation model, 65
- unstructured correlation matrix, 132, 133, 137
- unstructured correlation pattern, 52
- validity of MAR
 - assessing, 116
- validity of MCAR
 - assessing, 115
- variance, 42
- weighted generalized estimating
 - equations (wGEE), 84
- wGEE, 84, 85, 87, 134
 - example with incomplete data, 85
 - in categorical data, 125
- withdrawal. *See* dropout
- within-patient correlations. *See*, 53, 56
- within-patient errors, 52, 53, 56, 131, 132
- working correlation structure, 125