

## Index

- ACMV
  - identifying restriction, 96
- add-on designs, 26
- analysis of variance, 51
- analytic road map, xviii, 112, 113, 130, 151, 152
- ascertainable outcomes, 29
- attrition. *See* dropout
- augmented wGEE
  - and double robustness, 88
- autoregressive structure, 50
- auxiliary variables, 56, 81, 117, 135
  
- baseline observation carried forward (BOCF)
  - defined, 20
- Bayesian inference, 39
- Bayesian regression, 98
- bias from missing data, 6
- BOCF, 20, 61, 62, 69, 76, 77, 78, 101, 116
  
- canonical link, 127
- categorical data, 121, 124, 125, 126, 128
- categorical outcomes
  - analysis of, 121
- CCMV
  - identifying restrictions, 95, 96, 135, 142
- centralized raters, 25
- classes of MNAR methods, 92
  
- clustering factors, 133
- Code of Federal Regulations, 35
- Cohen's D, 6
- complete case analyses, 59
- conditional inference, 121, 122, 123
- conditional model, 123
- confidence intervals, 99
- consistent estimators, 88
  - defined, 4
- convergence
  - ensuring, 54
- Cook's D statistic, 133
  - use in influence diagnostics, 133
- correlation, 41
- correlations between repeated measures, 49
  
- data simulation, 147
- de-facto hypotheses*, 16, 18, 19, 20, 103, 104, 107, 108, 130, 147
- de-jure hypothesis*, 16, 17, 18, 19, 104, 106, 130, 147
- descriptive statistics and graphs
  - clinical trial example, 136
- design approaches
  - to lower rates of missing data, 148
- direct likelihood, 52, 81, 82, 84, 135
- distinction between MAR and MNAR, 11
- dose titration, 25
- double-blind trial, 20

**162**      **Index**

- doubly robust methods, 87
- dropout, 4
- dropout model, 85, 94, 134, 135
- effectiveness
  - defined, 16
- effectiveness estimand
  - clinical trial example of, 131
- efficacy
  - defined, 16
- efficacy estimand
  - clinical trial example of, 130
- enrichment designs, 23, 24, 148, 149
- estimand
  - defined, 8
  - ITT, 16
  - per protocol, 16
- estimands, 103
  - comparison of, 103
- estimation of estimand 1
  - example of, 104
- estimation of estimand 3
  - example of, 107
- estimation of estimand 4
  - example of, 107
- estimation of estimand 5
  - example of, 108
- estimation of estimand 6
  - example of, 108
- factorization
  - of likelihood function, 67
- Fisher's scoring, 54
- fixed effects, 50, 51, 52, 123, 133, 135
- flexible dosing, 25
- follow-up data, 15, 17, 28, 29, 103, 104, 106, 108, 109, 110, 149
- follow-up periods, 27
- frameworks for inference, 39
- frequentist inference, 39, 47
- full multivariate model, 53
- GEE, 39, 46, 47, 52, 55, 68, 84, 85, 86, 87, 88, 107, 121, 123, 124, 125, 128, 150
- general linear mixed effects model
  - for categorical data, 127
- general linear mixed-effects model, 51
- generalized estimating equations (GEE)
  - defined, 46
- Generalized least squares, 44
- Hamilton Depression Rating Scale, 50, 73, 130
- hierarchical model, 124
- hot deck imputation, 60
- identifying restrictions
  - in pattern mixture models, 95, 118, 135, 142
- identity link, 127, 135
- ignorability of missing data, 68 *See*
- ignorable missing data, 12, 125
  - defined, 12
- imputation model, 63, 82, 98, 104, 135
- incentives for completing treatment, 34
- inclusive modeling
  - clinical trial example, 140
- inclusive models, 56, 65, 89, 117, 134, 135, 143
  - for assessing sensitivity, 117
- independence
  - assumption of, 114
- influence diagnostics for patients
  - clinical trial example, 139
- influence diagnostics for sites
  - clinical trial example, 137
- influential observations
  - influence diagnostics, 115
- informative censoring, 12

## 163 Index

- informed consent, 35
- informed withdrawal of consent, 35
- Intention to Treat (ITT) principle, 10
- intent-to-attend, 34
- intermittent missing data, 4
- investigator payments, 34
- investigator selection, 34
- IPW
  - inverse probability weighting, 65, 66, 67, 84, 88
- last observation carried forward (LOCF)
  - defined, 37
- latent classes, 94
- latent variables, 94
- least squares estimation
  - illustration of, 40
- likelihood function, 45, 46, 67, 68
- likelihood-based analyses
  - example with complete data, 74
  - example with incomplete data, 78
- likelihood-based methods. *See* maximum likelihood
- linear mixed model, 51, 124, 127, 128, 153
- linear relationships, 40
- linearity, 114
  - assumption of, 114
- LOCF, 37, 55, 61, 62, 63, 69, 76, 77, 78, 101, 108, 116, 158
- logistic regression, 66, 84, 93, 115, 134, 135
- logit link, 101, 125, 127, 134
- loss to follow up
  - consequences for missing data mechanisms, 11
- LSMEANS, 75, 77, 79, 82, 85, 105, 141
- MAR, 11, 12, 56, 64, 65, 66, 67, 68, 69, 78, 81, 84, 85, 88, 91, 96, 101, 110, 112, 113, 116, 117, 125, 128, 131, 134, 135, 136, 140, 142, 143, 144, 145, 147, 149, 151
- marginal and conditional inference
  - illustration of, 121
- marginal distributions, 123
- marginal inference, 122, 123
- marginal models, 84, 123, 124, 128
- maximum likelihood, 39, 44, 45, 46, 47, 52, 54, 55, 56, 67, 79, 81, 88, 99, 101, 104, 107, 108, 131, 150
- maximum likelihood-based estimation
  - illustration of, 44
- MCAR, 10, 11, 12, 52, 60, 61, 65, 67, 68, 78, 84, 112, 115, 124, 125, 133, 137, 147
- measurement process, 12, 67, 92, 93, 95
- methods of statistical estimation, 39
- minimizing missing data, 23
- missing at random (MAR)
  - defined, 10
- missing completely at random (MCAR)
  - defined, 10
- missing data mechanisms, 9, 10, 121
- missing not at random (MNAR)
  - defined, 10
- missingness process, 12, 67, 68, 84, 92, 93, 95
- mixed-effects models, 51, 52, 126, 128
- MNAR, 11, 12, 57, 64, 65, 66, 68, 69, 91, 92, 93, 112, 115, 116, 117, 134, 135, 143, 144, 145, 147, 149, 151
  - bias from, 145
- MNAR analyses
  - for assessing sensitivity, 117
- model-based standard errors, 125
- modeling philosophies, 56
- monitoring rates of missing data, 34

**164**      **Index**

- multiple imputation, 63, 64, 65, 81, 82, 88, 98, 101, 107, 116, 128, 134, 135, 136, 150
  - example with incomplete data, 82
  - for categorical data, 127
- multivariate models, 53
- multivariate normal distribution, 124
- multivariate normal model, 51
- NCMV
  - identifying restriction, 96, 142
- New Drug Application, 129
- Newton-Raphson algorithm, 54
- non-future dependent missing value
  - restrictions (NFD), 96
- non-ignorable methods, 68
- normality, 114
  - assumption of, 114
- Ordinary least squares, 44
- outlier observations, 115
- participant burden, 33
- patient education, 35
- patient ratings, 25
- patient specific factors, 50
- pattern mixture model
  - clinical trial example, 142
  - specification for clinical trial example, 135
- pattern mixture models, 92, 94, 101, 102, 118, 134, 151
- placebo multiple imputation, 96, 118, 134
  - clinical trial example, 143
  - specification for clinical trial example, 136
- pMI, 118, 144
  - effectiveness context, 101
  - efficacy context, 99
  - example with missing data, 99
- poisson link, 127
- power
  - implications of missing data, 30
- primary estimand, 8, 15, 28, 36, 109, 110, 118, 131, 147, 150
- probability of dropout
  - consequences for missing data mechanisms, 11
- PROC GENMOD, 84, 85, 134, 135
- PROC IML, 135
- PROC MI, 98, 135
- PROC MI ANALYZE, 99, 135, 136
- PROC MIXED, 132, 133, 135
- PROC NL MIXED, 136
- property of double robustness, 88
- quasi-likelihood, 125
- random effects, 50, 51, 52, 53, 94, 123, 124, 126, 127, 131
- random variation, 50
- random-effects model, 123
  - for categorical data, 125
- random-effects models, 124
- randomization, 3, 147
- randomized withdrawal design, 24
- region of ignorance, 118, 144
- regression, 40
- rescue medications, 8, 15, 19, 27, 28, 29, 103, 104, 106, 108, 110, 149
- residual diagnostics
  - clinical trial example, 140
- residuals, 114, 115, 140
- restrictive models, 56, 81, 89, 104, 116, 117, 143
- retention. *See* dropout
- reverse factorization, 94
- run-in design, 23, 106

**165**      **Index**

- run-in designs, 23
- sample size, 30
- sandwich estimator, 55, 125, 132, 133
- selection model
  - clinical trial example, 140
  - specification for clinical trial example, 135
- selection models, 92, 93, 101, 115, 117, 134, 142, 151
  - semi-parametric implementation, 101
- semi-parametric estimators, 87
- sensible analysis, 114
  - defined, 4
- sensitivity analyses, xviii, 8, 65, 102, 111, 112, 116, 117, 118, 119, 130, 131, 134, 144, 150, 151, 152
  - defined, 8
- sensitivity to correlation assumptions
  - clinical trial example, 137
- separability condition, 12
- serial correlation, 50
- shared parameter model, 136
  - clinical trial example, 142
  - specification for clinical trial example, 135
- shared parameter models, 92, 94, 101, 118, 134, 151
- simple imputation strategies, 60, 69
  - example, 76
- software
  - for sensitivity analyses, 151
- standard deviation, 42
- standard error
  - defined, 4
- standardized effect size, 6
- substantive model, 88
  - in multiple imputation, 63, 65
- target population, 24
- time trends, 30, 49, 52, 53, 55
- titration dosing, 26, 52
- training for sites and patients, 34
- treatment changes, 10
- trial conduct
  - to lower rates of missing data, 149
- trimming weights
  - in wGEE, 87
- uncongenial imputation model, 65
- unstructured correlation matrix, 132, 133, 137
- unstructured correlation pattern, 52
- validity of MAR
  - assessing, 116
- validity of MCAR
  - assessing, 115
- variance, 42
- weighted generalized estimating equations (wGEE), 84
  - wGEE, 84, 85, 87, 134
  - example with incomplete data, 85
  - in categorical data, 125
- withdrawal. *See* dropout
- within-patient correlations. *See*, 53, 56
- within-patient errors, 52, 53, 56, 131, 132
- working correlation structure, 125