

# Index

- accelerated aging, 4, 328  
 accident-prone personality, 43  
 accidents/injuries, 24, 44, 45, 50, 53, 54, 55, 78, 79, 155, 325  
 active memory search, 24, 29, 32  
 active-duty soldiers, 1, 2  
 adoption, 5, 73, 163, 233  
 adult household resident, 19  
 adulthood, 14, 50, 54, 113, 119, 168, 169, 170, 196, 306, 323, 327  
 age, 20, 169, 183, 210  
   association with Maladaptive Family Functioning in childhood adversities (MFF CAs), 199  
   association with PTSD, 4, 95, 96, 100, 181, 196, 197, 199, 210, 226  
   association with trauma, 37, 43, 53, 72, 196  
   association with treatment, 143, 146, 147  
   in sample(s), 14, 113, 187, 199, 216, 230, 311, 328, 330  
   mental disorders classification, 110  
   poststratification weighting, 21  
   risk factor for trauma exposure, 2, 4  
 alcohol abuse/dependence, 27, 110, 115, 120, 199  
 age-inappropriate, 73, 164  
 age of diagnosis, 120  
 age of exposure, 164, 165–8, 178, 184, 255, 294  
   age-of-exposure distribution, 44, 51–2  
   association with PTSD or PTSD recovery, 182, 227, 240, 241–3, 257, 268, 274, 310  
   association with treatment, 140  
   disasters, 226, 227  
   first occurrence, 23, 26  
   motor vehicle collisions (MVC), 194–5, 196  
   sexual assault, 181  
   unexpected death (UD), 25, 26, 210, 211  
 age-of-onset (AOO), 6, 27, 28, 31, 35, 74, 102, 112, 117, 128, 241, 254, 288, 303, 306, 327  
   physical and mental conditions, 119, 120  
   PTSD, 96, 100, 103, 289, 326  
   suicide ideation, 133  
 American Red Cross, 5, 6, 233  
 anxiety disorder, 103, 119, 249  
   association with PTSD, 4, 174–9, 192–3, 199, 202, 213, 243  
   association with trauma, 76–9  
   comorbidity, 4, 110–11, 114–16, 251, 190, 303–6, 331  
   impairments, 128–30, 132–3  
   PTSD classification, 253, 260, 270  
   World Mental Health (WMH) survey, 27, 73, 288  
 arthritis, 119–21, 132, 328  
 asthma, 119–21, 132  
 atomic bomb survivors, 1  
 attention deficit (AD), 27, 74, 110, 111, 115, 116–17, 174, 199, 288, 300, 314  
 attention deficit hyperactivity disorder (ADHD), 76, 79, 110, 111, 192, 213, 248, 251, 288, 300  
 autobiographical memory, 31  
 Balanced Repeated Replication (BRR), 22  
 Bayesian Information Criterion, 115  
   behavioral and lifestyle factors, 121  
 behavioral disorders, 73, 111, 131, 192, 224  
 biological parents, 73, 163  
 bipolar disorder (BPD), 2, 27, 72–3, 128–30  
   comorbidity, 76–8, 114, 117, 139, 290, 303  
 bulimia nervosa, 120  
 cancer, 4, 119, 129  
 car bombings, 50  
 cardiovascular disease, 4, 118–19, 121, 329  
 caregiver, 24, 44, 50, 53, 73, 163, 243, 288, 323  
 caregiving relationship, 73, 163  
 causing/witnessing bodily harm, 44, 51, 55  
 Chernobyl disaster, 50  
 child welfare, 73, 164  
 childhood abuse, 43, 54, 72, 78, 119, 163, 169, 170, 181, 300, 304  
 childhood adversities (CAs), 5, 119, 173, 187, 240, 327, 330  
   assessment, 163–4  
   association with PTSD, 5, 164–5, 168–70, 187, 198–9, 201–2, 213, 226–30, 287–91, 294, 309, 325  
   association with recovery, 243, 251  
   association with trauma, 72–4, 78–9, 165–8  
   prevalence, 164  
   witnessing, 73–4, 79, 305  
   types, 160, 163, 323  
 chronic disorder, 4  
 chronic pain, 4, 119–21, 132  
 civilian, 23, 44–53, 79, 95, 128, 154–60  
 clinical contact person (CCP), 35  
 clinical interviews, 117, 230, 257, 321  
 cognitive ability, 4  
 cognitive deficits, 78, 80  
 cognitive-behavioral therapy (CBT), 5, 309  
 cohabitating, 55, 146, 302  
 collective violence, 44, 45, 46, 50, 53, 54, 55  
 combat experience, 24, 44, 50, 79, 243  
 combination therapy, 5  
 comorbidity, 1, 4, 130, 131, 133, 179, 240, 260, 270, 275, 278, 327–9  
   association with PTSD, 251, 287–8, 290  
   development of, 111  
   latent variables, 111–13, 115–18  
   with other mental disorders, 110–18, 251, 255, 280  
   with physical disabilities, 118–22  
   trauma types, 303–4, 306, 326  
 complementary and alternative medicine (CAM), 138, 141, 143  
 component stressors, 25  
 Composite International Diagnostic Interview (CIDI), 2, 13–14, 254–5, 259, 264, 270, 288, 306  
 conservative bias, 170, 281, 321  
 content, 73–4, 137, 163–4, 173, 273–4, 326–7, 329  
   design, 6–7, 17–20, 27–35, 38  
 conduct disorder (CD), 27, 76–7, 79, 110, 111–12, 114–18, 304  
 Conflict Tactics Scale (CTS), 73, 163, 164  
 contact crime, 55  
 core personality dimensions, 72  
 crime victims, 55  
 cross-loading trauma, 51, 74, 76  
 cross-national studies, 7, 33, 43–4  
 cultural differences, 54, 102  
 data collection organization, 17, 18  
 data mining, 5  
 death, 243, 318, 321, 328. *See also* unexpected death (UD)  
   accidentally caused, 76  
   association with PTSD, 228, 230, 255, 257, 303, 314  
   defining trauma, 1, 23, 263  
   motor vehicle collision (MVC) related, 25, 195, 196, 201  
   recovery from, 250–1  
 debt, 7  
 degenerative neurological diseases, 4  
 dementia, 4, 328  
 Detroit Area Survey of Trauma, 2, 4, 187, 200  
 developed countries, 43, 54, 287, 293, 315  
 developing countries, 2, 5, 43, 54, 79  
 developmental delays, 80  
 diabetes, 119, 129, 132, 329

## Index

- Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5), 179, 199, 216–18, 253–4, 256–260, 263, 280, 326  
 dissociative PTSD subtype, 287–8, 293  
 removal of A2 criteria, 8, 253, 263, 269  
 subthreshold PTSD, 273–5, 278–9, 281  
 use of, 1, 5, 6
- Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV), 5–8, 217, 253–7, 259–60, 263–8, 269–71, 280–1, 287, 309, 326  
 definition of trauma, 1, 23, 253  
 in Composite International Diagnostic Interview (CIDI), 6, 13, 27, 96, 174, 240, 273–4  
 PTSD classification, 128
- Diagnostic and Statistical Manual of Mental Disorders, Third Edition (DSM-III), 1–2
- disability, 6, 129, 194, 260  
 work, 129
- disabled, 129
- discrete-time approach, 37
- discrimination, 148
- dispositional negativity, 72
- divorce, 128, 210, 289
- drug use, 27, 78, 112, 116–17, 288. *See also* substance use disorder  
 association with PTSD, 110, 294, 304
- drunk driving, 55, 323
- dysthymia, 111, 120, 199, 248, 254, 274, 288, 300
- early adolescence, 50
- early attachment, 78
- early intervention, 5, 309
- eating disorder, 120
- economic adversity, 73, 74, 163, 164, 169, 227, 288  
 family, 210
- economic development, 36, 43, 148, 311
- economic status, 43, 72
- education, 20  
 association with PTSD, 96, 100, 102, 103, 181, 182, 194–5, 196, 201, 210, 211, 226, 227, 310, 325  
 association with trauma, 36, 53, 55, 196, 302  
 association with treatment, 140, 143, 146  
 high, 55, 140, 216, 230  
 in sample(s), 183, 199, 216, 230  
 low, 55, 100, 102, 103, 146, 181, 184, 201, 230, 325  
 mental disorders classification, 110  
 poststratification, 21  
 termination of, 128
- elderly, 103, 160, 169
- elevated risk, 55, 72, 79, 128, 210, 269, 329
- emotional regulation, 6, 169, 306
- English language, 32, 33
- environmental effects, 173
- epidemiological studies, 2, 7–8, 26, 72, 281, 305, 327  
 past, 1, 3–5, 128, 173, 210, 232, 249, 251, 259, 300, 323  
 episodic memory, 31
- evidence-based treatment, 5
- explicit trauma, 24
- externalizing disorders, 110, 111, 112, 116, 118
- family history, 6
- Family History Research Diagnostic Criteria, 73, 164
- fatality rates, 54, 194
- fibromyalgia, 119
- first episode, 31
- first-responders, 2
- focal trauma, 2, 7, 37, 181, 184, 210, 255, 274, 275, 309, 310, 314, 315
- gastrointestinal distress, 121
- generalized anxiety disorder (GAD), 111, 115, 117, 300, 303
- general interviewing training (GIT), 34
- general population, 1–5, 25–6, 34, 137, 153, 226, 232, 271, 323
- geographic regions, 17, 21
- global public health, 181
- hazards analysis approach, 37
- headache (chronic), 119, 120, 132
- Health and Work Performance questionnaire (HPQ), 131
- healthcare, 138, 147, 288, 306, 315, 332
- health-related surveys, 22
- heart disease, 118, 119, 120, 121, 129, 328
- high blood pressure, 119, 120
- high-income countries, 5, 54, 96, 119, 129, 137, 157  
 barriers to treatment, 137, 139–43, 148  
 conducting surveys in, 18, 36  
 differences from other countries, 43, 102–3, 129, 139–43, 147, 194, 199, 269  
 mental disorders, 113–15, 128  
 PTSD prevalence, 137, 139–43, 148  
 sex differences, 4, 293–4  
 trauma exposure, 2, 182, 195, 325  
 treatment, 147, 329–30
- high-risk, 2, 79, 233, 270, 311, 315, 327
- historical events, 43
- Holocaust victims, 1
- homelessness, 117
- homicide, 26, 50, 210
- human-made disasters, 24, 43, 44, 50, 51, 53, 79, 137, 213, 226, 227, 232, 233
- hyperactivity disorder (HD), 27, 74, 110–11, 115–17, 174, 199, 288, 300, 314
- hyperarousal symptoms, 240, 253, 304, 306
- hypertension, 118, 132, 270
- illness, 7, 26, 45–6, 73, 218, 243, 318  
 association with PTSD, 5, 120–2, 319–23  
 child, 50–5  
 schemas, 31
- immigrants, 54
- impairment, 128–9, 134, 280, 287, 291–4, 301–2, 305, 326, 327, 328  
 PTSD criteria, 253–5, 260, 264, 267, 273–4, 278  
 role, 32, 128, 287, 291
- impulse-control disorder, 111
- impulsivity/extraversion, 72
- industrialized countries, 55
- initial trauma, 43, 55
- institutionalization, 117
- intelligence, 72
- intermittent explosive disorder (IED), 27, 76–7, 112, 114–18, 120, 132
- internalizing disorders, 110, 112, 114, 116, 118, 128–9, 131
- International Statistical Classification of Diseases and Related Health Problems, Eleventh Edition (ICD-11), 6, 253–4, 256–7, 259–60, 326
- International Statistical Classification of Diseases and Related Health Problems, Tenth Edition (ICD-10), 1, 5–6, 13, 27, 253–4, 256–7, 259–60, 326
- interpersonal partner violence, 43
- interpersonal relationships, 306
- interpersonal violence, 44, 50–5, 305  
 association with PTSD, 153, 160–1, 178–9, 201, 213, 228, 257, 311, 314
- intimate partner victimization (IPV), 72
- intimate partner violence, 44, 50–5, 128, 306, 318, 323  
 association with PTSD, 181, 201, 278, 321, 326
- intrinsic factors, 72
- irritable/fearful temperament, 72
- Jackknife Repeated Replication (JRR), 22, 113
- kidnapped, 4, 76, 78, 155, 165, 243, 250, 318
- lifetime core disorder, 19
- lifetime mental disorders, 19, 27–8, 31, 72, 73, 76, 111, 118–19, 181, 310, 315
- low- and middle-income countries (LMICs), 184, 199, 216, 227, 332  
 dissociation, 289, 293–4  
 fatalities, 43, 194  
 in sample(s), 115, 157  
 mental disorders, 113–14  
 PTSD, 182, 211, 187, 195, 227, 325  
 trauma, 157, 182, 195, 232, 325  
 treatment, 139, 141–3, 147–8, 329  
 World Mental Health (WMH) surveys, 111, 119, 181, 194, 227, 287
- lower household income, 100, 102–3
- lower socio-economic status, 2, 4, 103
- low-income countries, 4–5, 18, 54, 269, 325, 328, 332
- machine learning, 202, 233, 309–11, 314–15
- major depressive disorder (MDD), 79, 129, 169  
 association with PTSD, 76, 175–6, 199, 248–9, 274, 305  
 comorbidity, 4, 110, 114–18, 119, 270, 331  
 in World Mental Health (WMH) surveys, 27, 32, 73, 74, 254  
 prevalence, 2, 95

- maladaptive family functioning (MFF), 73,  
 74–5, 163, 243  
 association with PTSD, 164–9, 198–9,  
 213, 230
- maltreatment, 73, 78, 163, 168–70,  
 288, 327
- marital status, 19, 20, 196  
 association with PTSD, 4, 100, 103, 130,  
 182, 197, 211, 302, 311  
 association with trauma, 53, 55, 78, 325  
 association with treatment, 140, 146
- marriage  
 early, 128  
 late, 128  
 on-time, 128
- married, *See* marital status
- maximum-likelihood (ML), 113
- medical diagnoses, 4
- medical treatment, 14, 122, 138
- memory schema, 31
- mental disorders, 26–8, 36, 131, 178  
 association with childhood adversities  
 (CAs), 160, 163, 168–9, 173, 323  
 association with socio-demographic  
 factors, 95  
 association with PTSD, 178, 181–3, 187,  
 199, 202, 269–70, 315  
 association with trauma, 72, 76, 78  
 child- and adolescent-onset, 173  
 comorbidity with PTSD, 4, 110–13,  
 117–18, 120, 288, 303–4, 328–9  
 cost of, 128–30, 132  
 factor analysis of, 113–16  
 postdisaster, 233  
 surveys of, 6, 13, 17–19, 26–8, 31–2, 119  
 treatment, 122, 137, 139, 147–8, 331–2
- mental health  
 literacy, 148  
 needs, 13, 137  
 professional, 5, 138  
 research, 7  
 sector, 5  
 services, 5, 8, 233  
 treatment, 5, 79, 138, 147–8, 331–2
- mental illness, 130
- meta-analysis, 4, 5, 79, 95, 201, 202,  
 249, 309
- methodological challenges, 27–32
- minority, 5, 43, 95, 269
- mood disorder, 4, 27, 76–7, 114, 132, 163,  
 243, 248–9, 305–6
- morbid risk, 35
- motor vehicle collision (MVC), 2, 8, 25, 43,  
 194, 195, 196, 197, 198, 199, 200,  
 201, 202  
 characteristics, 194, 195, 197  
 life-threatening, 195, 197  
 MVC-related PTSD, 194, 195, 199, 200,  
 201, 202  
 survivors, 194, 201
- mugging, 2, 24, 43, 44, 46, 50, 51, 53, 54, 74,  
 243, 321
- National Comorbidity Survey (NCS), 2, 4,  
 226, 327
- national survey, 5, 13, 17, 181
- natural disaster, 4, 24–5, 43, 50–3, 137, 165,  
 243, 251  
 association with PTSD, 79, 155–60,  
 226–33, 315  
 negative emotionality, 72  
 negative health beliefs, 137  
 negative public attitude, 122, 148  
 neglect, 73–4, 78, 163–5, 168, 210, 227, 243,  
 288, 327  
 neural response, 169  
 neurobiological studies, 5, 111  
 neuroticism, 72  
 nonrelative, 26  
 nonresponse bias, 20  
 nonresponse survey, 19, 20  
 nosological, 5, 7, 8, 326
- obsessive compulsive disorder (OCD), 111,  
 115, 117, 120, 254, 274
- one child policy, 50
- oppositional defiant disorder (ODD), 27,  
 76–7, 79, 114, 116, 130, 199, 290
- organized violence, 23, 76–7, 79, 141, 153–4  
 association with PTSD, 155–8, 174,  
 243–4, 314
- panic attacks, 31
- panic disorder, 2, 27, 76–7, 114–17, 129–30,  
 132–3  
 association with PTSD, 174–5, 290
- parent  
 alcohol abuse, 213  
 criminality, 73, 74, 164, 210, 227  
 death, 73, 74, 163, 210, 227, 288  
 divorce, 73, 74, 163, 164, 227, 288  
 loss, 74, 163, 210, 227  
 maladjustment, 73, 78, 163, 288  
 mental disorder, 210, 226  
 mental illness, 73, 74, 213, 289  
 psychopathology, 73, 163–5, 168–70, 173  
 substance use, 73–5, 164, 210, 226, 289  
 violence, 73, 164, 305
- pathological stress reactions, 72
- personality disorder, 79
- pharmacotherapy, 138
- phenomenological, 5
- phobia, 103, 110, 115, 117, 270. *See also*  
 anxiety disorder  
 agoraphobia, 27, 76, 111, 115–17, 120,  
 133  
 association with PTSD, 174, 274, 290,  
 305  
 social, 27, 76, 111, 116, 120, 132, 133  
 association with PTSD, 174, 199, 249,  
 251, 274, 292, 304  
 specific, 27, 76, 111, 115–17, 120, 132  
 association with PTSD, 174, 199, 274,  
 290, 304, 314
- physical abuse, 4, 73, 163, 164, 210, 227, 302,  
 303, 305, 321, 323  
 childhood, 73, 78, 164  
 physical health, 4, 43, 119, 122, 181, 326,  
 328, 329  
 physical health conditions, 4  
 physical ill-health, 119  
 physically impaired, 14
- policy (health care), 148  
 policy (public health), 129, 325  
 policy (social), 30  
 policy makers, 8, 274  
 population registry, 17  
 posttraumatic stress disorder (PTSD)  
 12-month, 96–103, 130, 131–2, 141–8,  
 288–90, 292, 300–6, 326, 329  
 12-month treatment, 138–9, 143, 147  
 association with, 4–5, 37  
 association with aging, 328  
 association with life course events, 128–9  
 association with physical disorders,  
 118–22  
 association with role performance,  
 129–32  
 association with sexual assaults,  
 182–4, 187  
 association with socio-demographic  
 factors, 102–3  
 association with trauma, 25–6, 161, 257,  
 270–1, 301, 326  
 conditional risk, 200  
 definition, 1  
 development of, 4, 43, 103, 181, 182, 325  
 diagnosis, 79, 253, 260, 269, 271, 273  
 dissociative, 6  
 elevated risk, 329  
 episode, 35, 36, 138  
 evidence-based predictors, 5  
 genes, 5  
 history of, 132, 134, 173, 177–9, 240, 315  
 lifetime, 2, 26, 35, 36, 96, 100, 102, 103,  
 137, 153, 304, 321, 327  
 medical-related symptoms, 4  
 mild-moderate, 143, 148  
 prevalence of, 1, 2, 7, 23–4, 26, 35–6,  
 95–103, 177, 187, 311, 321, 325–6  
 by diagnostic system, 254–60  
 Criterion A2, 263–71  
 disaster-related, 160, 227, 232  
 motor vehicle collision (MVC)-related,  
 194–7, 200–1  
 trauma-specific, 153–5  
 unexpected death (UD)-related,  
 211, 217  
 preventive strategies, 5, 38, 72–3, 179,  
 181, 188, 202, 232–3, 309, 315–16,  
 330–2  
 prospective studies, 119  
 remission, 4, 6, 327–8, 329–40  
 recovery, 240–51  
 resilience, 153, 160–1, 327  
 risk, 183, 187–8, 200–2, 216–18, 243, 257,  
 270, 309–15, 325–7  
 A2 criterion, 269, 271  
 disaster-related, 226, 228, 230–3  
 factor, 4–5, 110–11, 118–22, 131, 153,  
 329  
 lifetime, 2, 38, 96  
 overestimation, 26, 181  
 socio-demographics, 196, 102–3  
 trauma exposure, 2, 4, 25, 35–6, 156,  
 160, 168–70, 173, 178–9, 305,  
 318–23  
 severe, 138, 143, 146, 147, 148

Index

- posttraumatic stress disorder (PTSD) (*cont.*)
  - SMH treatment, 143
  - societal burden, 36, 318, 326
  - spousal abuse, 50, 53, 55
  - substance abuse, 73, 95, 163, 213, 254, 288
  - subthreshold, 6, 273–82, 326
  - symptoms, 8, 138, 169, 218, 253, 255, 269, 274, 278, 321, 329, 331
  - syndromic, 72
  - trauma-specific risk, 26, 36
  - treatment, 4, 79, 130, 134, 137–48, 306, 309
    - barriers, 139, 143–145, 148
    - differential, 251, 270, 287, 291–4, 300, 326
    - lack of, 6, 143
    - patterns, 329–32
    - treatment rates, 5, 139, 141–2, 147
    - vulnerability, 161, 163, 169, 170, 173, 179
- post-disaster risk evaluation, 5, 233
- post-trauma symptoms, 1, 251
- premature mortality, 4
- pre-marital mental disorders, 128
- pre-trauma, 72–3, 173, 178, 188, 202, 251, 315–16, 329
  - predictors, 173, 251, 315
  - risk factors, 73, 188, 202, 316
  - vulnerability, 72
- prior traumas, 2, 4, 53–4, 72, 153, 243, 251
  - association with PTSD, 157–61, 177–9, 182, 184–7, 198–9, 213–16, 228–9, 289, 294
- prisoners of war, 1
- privacy, 18
- probability sampling, 17
- prognosis, 4
- psychiatric disorder, 6, 79, 80, 103, 329
- psychopathology, 4, 5, 78, 110, 160, 325. *See also* mental disorders
  - association with PTSD, 173–9, 202, 232, 260, 270
- psychosis, 79
- psychosocial and cognitive-behavioral techniques, 72
- psychotherapy, 138
- psychotic experiences, 110
- PsySTART system, 5, 233
- public awareness, 148
- random trauma, 37–8, 138, 153–7, 217, 281, 310
  - assessment of, 7, 25–8, 329
  - association with PTSD, 174–9, 240–3
  - recovery, 249–51
  - treatment, 139–41, 147
- rape, 1, 4, 5, 7, 25, 45, 50–3, 54–5, 73–4
  - association with PTSD, 155–8, 174, 177, 179, 182–3, 185, 187, 201, 243–4, 321
  - childhood adversity (CA), 73–4, 76, 163
  - risk, 311, 314, 318
  - rates of treatment, 5
- refugee, 23, 44, 50, 53, 137, 160, 243, 251, 330
- region of terror, 50, 155, 157, 160, 315
- relief worker, 23, 44, 153
- respondent booklet (RB), 27, 29, 32
- re-traumatization, 201
- retrospective reports, 28, 117, 120, 169, 199, 250–1, 270, 309, 315
- re-victimization, 43, 160, 305
- romantic partner, 24, 44, 50–3, 55, 74, 76, 78, 243, 318–21, 325
- sampling procedures, 13–20, 38, 95
- schizophrenia, 110
- school shootings, 137
- selective-serotonin uptake inhibitors (SSRIs), 5, 121
- self-blame, 181, 187, 253
- semantic memory, 31
- separation anxiety disorder, 27, 74, 76, 79, 111, 174, 248, 251, 288, 290, 300, 314
- serious injury, 1, 23, 45, 50, 54, 76, 153, 155, 195, 201, 230, 243, 251, 263, 318, 321
- serotonin-norepinephrine reuptake inhibitors (SNRIs), 5
- severe mental and substance use disorders, 5
- severity of exposure, 5
- sex, 1, 20, 21, 43, 53, 72, 255, 310
- sex differences, 4–5, 36, 37
  - mental disorders, 110
  - PTSD, 4–5, 100–1, 103, 196–201, 211, 214–17, 226, 230–1, 311–12, 325
  - PTSD diagnosis, 257, 259, 268, 274, 280, 326
  - recovery, 241–2, 249
  - trauma exposure, 2, 53, 72, 183
  - treatment, 140–1, 143–4, 146, 329
- sexual abuse, 73, 163–8, 181–8, 213, 243, 327
- sexual assault, 26, 44–5, 53, 128, 326, 330
  - association with mental disorders, 73, 76, 78
  - association with PTSD, 154–60, 179, 181–8, 257, 311, 315, 318–22
  - childhood, 50, 73–4, 160, 305
  - in WMH surveys, 25, 163
  - prevalence, 47
  - recovery, 243–4
  - sex differences, 43, 72
  - treatment, 147
- sexual violence, 24, 44–5, 50–5, 74–5, 77–8, 140–1, 243–6, 318–19, 330
  - association with PTSD, 102, 154–60, 181–7, 228, 259, 266–7, 278, 313–14, 320–3, 325–6
- Sheehan Disability Scales (SDS), 129, 130, 131, 139, 288, 291, 300, 301
- sibling, 26, 210
- simple random sampling (SRS), 22
- socio-demographic variables, 20–1, 37, 141–6, 241–3, 251, 327, 329
  - association with PTSD, 100–3, 196–8, 211, 227, 257, 260, 268, 289, 302
  - association with trauma exposure, 43–4, 54, 72, 323
  - risk factor, 4, 95, 181, 201, 309–11, 314–15, 326
- socio-economic status, 43, 95, 103, 143
- spouse, 19, 26, 59, 74, 181, 187, 210, 211, 217, 303, 305, 325
- stalking, 53, 76, 78, 321
- stroke, 118, 119, 328
- student, 34, 53, 96, 109, 187
- subsequent traumas, 44, 27, 72, 79, 163, 173, 174, 177, 178, 179, 321, 323
- substance use disorder, 20, 27, 76–9, 95, 111, 114, 163, 248–9
  - association with PTSD, 4–5, 176, 213, 257, 259–60, 292, 304–5
- suicidality, 1, 5, 132, 133, 233, 240, 254, 255, 273, 278, 287, 288, 291, 292, 294, 326
  - outcomes, 133
- suicide, 25, 26, 50, 73, 117, 132, 133, 139, 164, 199, 210, 288
  - attempts, 133
  - ideation, 133
  - planned attempts, 133
  - unplanned attempts, 133
- survey population, 14–18, 20–2
- target population, 14–16, 21
- Taylor Series linearization method, 22, 37, 38, 44, 96, 241
- therapy, 306, 309, 332, 333
- time-since-onset, 117
- toxic chemical exposure, 50, 79
  - characteristics, 25–6, 182, 188, 202, 216
  - cross-national prevalence of, 43
- trauma
  - accident-related, 73
  - adult, 72
  - aftermath, 202, 218, 232, 233, 269, 270, 309, 315, 316
  - definition, 23
  - discovering a dead body, 46
  - high-risk, 72
  - history, 26–7, 43, 153, 160–1, 177–9, 181, 210, 251, 288, 331
  - life-threatening, 7, 24, 46, 50, 54, 194, 195, 197, 200, 202, 243, 265, 318, 326
  - lifetime, 2, 25–7, 35–8, 43–5, 53–4, 153–7, 216, 243, 254, 273, 315, 318–21
  - memories, 160
  - other, 1, 44, 46, 53, 155, 318, 321, 326
  - predictors, 36–7, 72, 73
  - prevalence, 2, 4, 35–6, 43–51, 153–5, 318, 321, 322–3, 325–6
  - risk, 43, 54–5, 72–3, 78–9, 325
  - related features, 72
  - severe, 26, 270, 332
  - society-wide, 128
  - stressful life event, 5, 53, 163
  - survivors, 5, 128, 160, 181, 202, 226, 232, 233, 251, 287, 330, 331
  - threatened, 44, 50, 243
  - treatment, 78, 140–1, 270
  - types, 4, 23–7, 44–55, 73, 74, 102, 153–61, 163–9, 240, 254, 257, 264, 281, 294, 300, 301–6, 310, 314, 315, 318–23
  - underestimation of, 54
  - war-related, 160, 318, 321

- witness, 1, 23–5, 44–6, 50–5, 250–1, 263, 318
  - association with PTSD, 155–7, 165, 218, 228, 243, 311, 314
- witnessing death, 45, 46, 50, 54, 228, 251
- worst lifetime, 26, 27, 137, 179
- trauma victims, 5, 309, 317
- treatment adherence, 4
- treatment barriers, 139
- trauma exposure, 54–5, 302–6. *See also*
  - prior trauma
  - association with childhood adversities (CAs), 73–5, 78–80
  - association with mental disorders, 76–80
  - association with PTSD, 4, 95, 155–61, 173–8, 315–16
  - association with socio-demographics, 100–3
- under-reported, 28, 169, 184, 323
- unemployment, 4, 129, 325
- unexpected death (UD), 44, 51, 217, 305, 311, 314, 326
  - association with socio-demographics, 53
- association with prior mental disorders, 78
- association with PTSD, 79, 155, 210–16, 303, 314
- cause of, 26, 210, 217–18
  - in WMH surveys, 25–6, 210
  - reported occurrence, 2, 43, 45–6, 50, 54, 153, 243, 318, 321, 323, 325
- veterans, 1, 2, 3, 118, 130, 294, 326, 328, 330
- combat, 137
- violence
  - family, 72–3, 163–4, 243, 289–91, 303
  - marital, 131
  - physical, 130
  - physical victimization, 24, 44–5, 50, 74–9, 140–1
    - association with PTSD, 154–7, 160, 177, 179, 318–23
  - relationship, 78, 130
  - sectarian, 50, 102, 128, 160, 311, 315
- violence-related traumas, 73
- violent crime victimization, 73
- vulnerable groups, 1, 2, 4, 55
- war, 43, 54
  - exposed to, 128, 160
- war zone, 23, 44, 46, 53, 63, 79, 153, 155, 160, 315
- war-torn countries, 54
- welfare, 73, 164
- World Fertility Survey (WFS), 13
- World Health Organization (WHO), 13, 32, 33, 38, 96, 100, 131, 202, 252, 316, 324
- World Mental Health (WMH), 6, 72, 95–6, 111, 128, 132–3, 233, 327, 330, 332
  - developmental transitions, 128–9
  - Initiative, 1, 6, 7, 13, 32, 34, 72, 263–4
  - PTSD diagnostic criterion, 6, 254, 263
  - recall, 7, 120, 160, 199, 230
  - role performance, 129
  - sample, 187, 216, 240, 329
  - subthreshold PTSD, 273
  - survey analyses, 110, 119, 287, 300, 309
  - survey data, 5, 6, 43, 111, 113, 128, 131, 173, 210, 217, 254, 294, 309, 309, 323, 325, 328, 329
  - survey findings, 1, 2, 122, 131, 160, 163, 173, 183, 194, 210, 232, 251, 270, 323, 325–7, 328, 329
  - survey methods, 13–38, 73, 95, 121, 153, 169, 181, 210, 263, 273, 328, 315, 323
  - survey strengths and limitation, 6–7, 102, 160, 194, 199–200, 216, 218, 230–2, 270, 281, 321–3
  - trauma exposure, 43, 44, 50, 53, 54, 55
  - treatment, 137, 138, 139, 147–8
- World War II, 160
- worst trauma, 2, 7, 26–7, 95, 174, 181, 310, 323, 325, 329