

## Index

- acuity scores, 277
- adaptive point process filtering, 303, 310–311
  - application, 324
- adenosine triphosphate (ATP), 332
- Akaike information criterion (AIC), 83, 99, 104, 178, 247, 248, 315
- $\alpha$ -recursion, 19
- Alzheimer's disease, 115
- amygdala, 322
- anterior hippocampus, 322
- APACHE-IV, 270, 277
- approximate marginal inference, 285
- artefacts, 28
- arterial blood pressure (ABP), 290, 302
  - see also* blood pressure
- assumed density filtering (ADF), 40
  - particle filtering as, 42–43
- assumed density smoothing (ADS), 40
- ATR Human Subject Review Committee, 68
- auditory cortices, 68
- automatic relevance determination (ARD), 56
- autoregressive (AR) models, 2, 16
  - cardiophysiological data, 323
  - conditional heteroskedastic (ARCH), 31, 37
  - hidden Markov model (AR-HMM), 257, 259
    - hierarchical Dirichlet process, 260
    - higher-order scalar and vector, 16–17
    - multivariate (MAR), 54, 66, 92–93
    - nearest-neighbors with exogenous input (NNARX) *see* nearest-neighbors
  - autoregressive with exogenous input, 45
    - parameter estimation, 57
    - switching *see* switching vector autoregressive (SVAR)
- autoregressive conditional heteroskedastic (ARCH) model, 31, 37
- autoregressive integrated moving average (ARIMA) model, 27, 323
  - cardiophysiological data, 314, 319–321
- autoregressive moving average (ARMA) model, 86, 91, 309
- baroflex sensitivity, 302
- basal ganglia, 128
- Bayes factor, 177
- Bayes' rule, 3–4, 14, 43, 171, 187
- Bayesian estimation, 4–5
  - heart rate, 348
  - source reconstruction, 55–57
- Bayesian information criterion (BIC), 83, 99, 104, 178
- belief networks, 15
  - latent variables, 30
- beta process autoregressive hidden Markov model (BP-AR-HMM), 261
  - applications to clinical data, 269
- blood oxygen level dependent (BOLD) signal, 79, 81, 102, 114
- blood pressure (BP), 257, 272, 273, 275, 276, 290
  - arterial, 290, 302
  - cardiovascular models and, 307
- blood samples, 25, 28
- Bonferroni correction, 75
- bootstrap filter, 42
- bradycardia, 25
- brain machine interface, 195, 207
- brainstem, 322
- Broyden-Fletcher-Goldfarb-Shanno (BFGS) algorithm, 84, 100, 289
- burn-in, 166
- burst suppression, 330
  - existing models, 332
  - length, 336
  - onset, 330
  - probabilistic modeling, 333–336, 342
    - simplified model, 333 336, 342
  - probability (BSP), 331, 337, 340
  - ratio (BSR), 330
  - reduced order model, 333
  - state space modeling, 338
- C++, 248
- calcarine sulcus, 69
- calcium imaging, 9
- calculus of variations, 61

- cardiac surgery, 343
- cardiophysiological data, 302–303
  - ARIMA model, 319–321
  - brain and autonomic activity correlation, 321–322
  - model order selection, 315
  - modeling robustness, 323
  - point process filter, 304–306, 310–311
  - time-varying frequency analysis, 311
    - dynamic R-R spectrum, 312
    - feedback loop frequency response, 311
    - feedforward loop frequency response, 312, 313
    - dynamic coherence, 313
- cardiovascular dynamics, 274–278
- center manifold theorem, 126
- cerebral metabolic rate of oxygen (CMRO), 332
- Cerebus acquisition system, 218
- change-point test, 232, 233
- Chapman-Kolmogorov equation, 4
- Cholesky decomposition, 300
- clinical data, 10
  - inference algorithms, 262
  - sampling, 264
- clinical monitoring, 10
- cluster analysis, 79
- computational cost
  - decision-making modeling, 182
  - Markov chain Monte Carlo methods, 267
  - source reconstruction, 57–58, 62, 63
- conditional intensity function (CIF), 211, 221, 306, 311
  - interneuronal interactions, 226
- conductance-based neural models, 117
- continuous variables, 1, 16
- control theory, 90
- controllability, 3
- convolution modeling, 117
- correction smoothing, 20
- cortical local field potentials, 293
- cortico-basal ganglia thalamocortical network, 128
- Cox process, 139
  
- data smoothing, 9
- decision modeling, 160–161
  - diffusion-to-bound model, 166–167
    - inference, 168
  - model comparison, 177–178
  - point process models, 163–178
    - Markov chain Monte Carlo method, 164–166
  - switching model, 173–175
    - computational cost, 182
    - latent state sampling, 174–175
    - parameter sampling, 176
- deep learning, 284, 285
- deterministic processes, 1
- deviance information criterion (DIC), 177, 197, 247
- diagonal state space method (dSSM), 59, 72
  - estimation algorithm, 61
  - individual temporal dynamics estimation, 73
  - probabilistic model, 59–60, 64
    - joint probability distribution, 60
  - real data analysis, 68
  - simulation analysis, 66–68
- diencephalon, 322
- diffusion magnetic resonance imaging (dMRI), 58–59, 66
- diffusion-to-bound model
  - inference, 168
    - latent trajectory sampling, 168
    - parameter sampling, 171–173
- direct pathways, 128
- directed acyclic graph (DAG), 233, 284
- Dirichlet process, 201
- discrete state Markov models, parameter estimation, 44
- discrete variables, 1
- dopamine loss, 128
- dynamic causal modeling (DCM), 74, 119, 130, 132
  - learning, 232
  - model optimization, 131
  - model in time domain, 119–122
  - non-EEG applications, 132
  - Validation, 132
- dynamics reversal, 20
  
- electrocorticogrammetry (ECoG), 74–75
- electrocardiogrammetry (ECG)
  - Bayesian filtering, 355–358
  - signal degradation sources, 346, 347
  - signal quality evaluation, 353–355, 361
  - state space approaches, state of art, 347
  - switching Kalman filter, 356, 357
  - X-factor approach, 356–358
  - see also* heart rate
- electrocortigraphy (ECoG), 118
- electroencephalography (EEG), 9, 53, 284
  - automatic segmentation, 337–338
  - burst suppression, 30, 330
  - ‘ground truths’, 114
  - linear approximations to time series, 123
  - modeling approaches, 116
  - signal features between normal and pathological states, 115
  - variational Bayes estimation, 47
  - see also* neural spike trains
- electromyograms, 6
- entopeduncular nucleus (EPN), 129
- epilepsy, 131
- evidence, 8
- expectation, 4
- expectation maximization (EM), 84, 162, 286
  - general linear model with hidden states (GenLMHS), 215
  - Kalman filter, 209, 213

- Poisson linear dynamic systems (PLDS), 147, 153, 155
- switching vector autoregressive (SVAR) models, 286, 289, 291
- variational Bayes (VB) method, 197
- expectation-correction, 25
- eyeblick detection, 21
- factor analysis, 225
- Fano factors, 151
- field-programmable gate array (FPGA), 201
- filtering, 18–19
- Fisher information, 173
- fMRI time series, nearest-neighbors AR with exogenous input (NNARX), 97–98
- Fokker-Planck formalism, 117
- free energy, 197
- frontal eye field (FEF), 161
- full state space model (fSSM), 64–66
  - estimation algorithm, 65–74
  - experimental validation, 75
  - real data analysis, 71
  - statistical test on source activities and interactions, 74
  - structural connectivity, 66–68
- functional magnetic resonance imaging (fMRI), 9, 53, 58–59, 79, 114
  - blood oxygen level dependent (BOLD) signal, 79, 81, 102, 114
  - cardiophysiological data correlation, 321–322
  - clinical examples, 102–106
  - colored noise, 86
  - dynamic causal modeling (DCM), 132
  - innovation approach, 82–84
  - maximum likelihood estimation, 84
  - multivariate autoregressive (MAR) model, 92–93
  - spatial whitening, 94–95
  - nearest-neighbors AR with exogenous input (NNARX), 93
    - with spatial whitening, 98–99
  - non-Gaussian data, 97–98
  - parameter estimation, 99
  - predictive models, 80
    - general linear model (GenLM), 85–90
  - preprocessing, 82
  - time series structure, 81–82
    - exogenous input function, 82
    - stimulation function, 82
- Gaussian distribution, 2, 305
  - heart rate interval, 304
- Gaussian process factor analysis (GPFA), 225
- Gaussian processes, 2
- Gaussian-observation linear dynamical system (GLDS), 138
  - spectral learning, 150–151
- general anesthesia, 317–319, 324
- general linear model (GenLM), 80, 81, 85, 88–89
  - discrete time formulation, 88–89
  - functional magnetic resonance imaging (fMRI) data, 321
  - with hidden states (GenLMHS), 215
    - conditional intensity function, 221
    - Kalman filtering and, 224
    - model identification, 216–217, 220
  - model fitting, 104
  - motor cortex activity, 209
  - observation noise, 89–90
  - parameter estimation, 100–101
  - state space model, 90–91, 108
- generalized autoregressive conditional heteroskedastic (GARCH) model, 31, 37
- generalized pseudo Bayes, 25
- Gibbs sampling, 38
- goal-directed movement control, 9
- gout, 284
- Granger causal analysis, 132
- graphics processing unit (GPU), 168, 182
- hemodynamic response function (HRF), 85, 91
  - state space model (SSM), 87–91
- Hamiltonian Monte Carlo method, 248
- heart beat interval
  - inverse Gaussian distribution, 324
  - nonlinearity, 313
  - probability models, 304
- heart rate, 27, 291
  - beat detector agreement estimation, 348
  - instantaneous indices, 306–307
  - interval analysis, 302
  - power spectrum subband evaluation, 348
  - premature babies, 26
  - skewness and kurtosis, 349
  - state space fusion model, 346–347
  - state space model, 349–351
    - comparison to other techniques, 352
    - observation noise, 350
    - results, 351
  - variability, 283, 302
    - instantaneous indices, 306–307
- hidden Markov model (HMM), 17–21, 23, 47, 162
  - autoregressive (AR-HMM), 257
    - hierarchical Dirichlet process, 260
  - continuous-state, 22–23
  - decision modeling, 174, 175
    - latent state sampling, 174–175
    - parameter sampling, 176
  - deterministic, 30
    - dynamic synapses, 35–37
    - leaky integrate-and-fire, 37–38
  - discrete state, 20–21
  - electrocardiogrammy, 356

- hidden Markov model (HMM) (cont.)
  - filtering, 19
  - first-order, 15
  - fully-observed, 15–17
  - hierarchical Dirichlet prior, 259–260
  - hippocampal spike activity, 186
  - Hopfield membrane potential as, 34
  - inference, 18
    - approximation 38
  - parameter estimation, 32, 43–44
  - prediction, 20
  - second-order, 15
  - sleep-stage scoring, 6
  - smoothing, 19
  - see also* hidden semi-Markov model; Markov chain Monte Carlo methods
- hidden semi-Markov models (HSMM), 162
- hidden states, 209
- hierarchical Dirichlet (HDP) process, 201, 257, 278
  - autoregressive hidden Markov model (HDP-AR-HMM), 263, 278
  - Gibbs sampling, 263
  - Markov chain Monte Carlo (MCMC) algorithms, 266–267
  - parameter settings, 269
- hippocampal neurons, 9
  - hidden Markov model selection, 187–188, 201
  - hidden semi-Markov models (HSMM), 162
  - overdispersed Poisson model, 202
  - population codes, 195
    - experimental results, 198
    - Markov chain, 196
      - as topology graph, 198
      - Variational Bayes (VB) model, 196
    - spike sorting-free decoding, 201
- Hopfield membrane potential, 34
  - augmented network, 35
- importance sampling, 40
- inference, 7–8
  - approximate
    - assumed density filtering, 39
    - importance sampling, 40–41
    - linear Markov models, 38–43
    - Monte Carlo methods, 38
    - sampling, 38
    - variational, 39
  - computational cost, 24
  - difference-to-bound decision model, 168
    - latent trajectory sampling, 168
  - in linear systems, 23
- inferior fusiform gyrus, 69
- inferior occipital gyrus, 69
- inferior temporal cortex, 293
- inferior temporal (IT) cortex, 293
- inhibitory neurons, 122
- inhomogeneous Poisson process (IPP), 215, 220
- innovations, 80, 82–84
- intensive care unit (ICU), 258, 331
  - data generated, 345
  - population heterogeneity, 345
- inverse Gaussian distribution, 305
- inverse problems, 9, 81
- joint probability distribution
  - diagonal state space method, 60
  - full state space method, 65
- Jordan form, 87
- Kalman filter, 47, 84, 106, 146, 208
  - heart rate estimation, 348, 350
  - with hidden states, 219–220
    - decoding, 222
    - higher-value hidden dimension, 226
  - learning, 232
  - motor cortex, 209–211
    - hidden states, 212
    - switching (SKF), 354, 356, 357
- Kernel density estimation (KDE), 190–192, 194
  - density representation, 191
- Kirchoff's law, 117
- KL divergence, 145
- Kolmogorov-Smirnov plot, 305, 315
- Kullback-Leibler divergence, 61, 142
- Laguerre function, 309
- Langevin algorithm, 173, 176
- Laplace expectation maximization, 147
- Laplace transform, 124
- latent Markov model (IMM) *see* hidden Markov model (HMM)
- lateral interparietal area (LIP), 161, 162
  - primates, 163
- learning (algorithmic), 287
  - error back-propagation, 287, 291
  - global label from hidden state proportions, 287
  - sequential labels from local marginals, 288
- learning (animal and human), 9, 231
  - change-point test, 232, 233
  - curve estimation, 232
  - deep brain stimulation, 244–247
  - dynamic causal modeling (DCM), 232
  - example data set, 237–239
    - as random walk, 234
    - response bias in object-place association, 239, 241–244
    - state space model (SSM), 231–233
- leukemia, 284
- LFP model, 119–122
- light-field calcium imaging, 137
- likelihood, 7, 248
- linear dynamical systems (LDS), 22

- Gaussian-observation (GLDS), 138
- inference, 23
- Poisson *see* Poisson linear dynamical system
- smoothing, 23
- switching (SLDS), 23
- local linearization, 89
- logistic regression, 244–247
- lognormal distribution, 305
- LORETA method, 55
- Lyapunov exponents, 125–126
- M1 neurons, 9
- machine learning, 347
- magnetoencephalography (MEG), 9, 53
  - diagonal state space (dSSM) analysis, 68
  - full state space method (fSSM) analysis, 71
  - simulated, 66
  - source reconstruction, 53–54
- MAGNETOM trio, 57
- marginal likelihood, 8
- marginalization, 4, 285, 287
- Markov, Andrey, 1
- Markov chain, 1
- Markov chain Monte Carlo (MCMC), 56, 161, 225, 233, 262
  - learning models, 234, 245
    - convergence, 236
  - neural spike rates
    - burn-in, 166
    - initialization point, 166
    - model comparison, 180
    - sample generation, 165
    - simulated datasets, 178, 179
- Marr-Hildreth operator, 108
- Massachusetts General Hospital, 317–319
- Massachusetts Institute of Technology General Clinical Research Center, 316
- MATLAB, 109, 292
- matrix-normal-inverse-Wishart (MNIW) prior, 259
- maximum *a posteriori* (MAP) estimation, 44, 164
- maximum likelihood estimation (mle), 7, 44, 147, 164
  - Broyden-Fletcher-Goldfarb-Shanno (BFGS) algorithm, 84, 100, 289
  - fMRI time series, 84, 100
  - overfitting, 8
- mean squared error, 178, 223
- medical trainees, 249
- membrane potential, 34, 37
- mesencephalon, 322
- metabolic state, 339
- Metropolis sampling, 38
- Metropolis-Hastings algorithm, 38, 172
- micro-drive arrays, 198
- microdialysis, 114
- MIMIC II database, 258, 268, 351–354
  - association analysis, 270, 272–274
  - HDP-AR-HSMM, 272–274
  - mortality risk estimation, 271–272
- mixed pair hidden Markov model (MPH), 225
- mobile health, 346
- mode proportion, 270
- model fitting, 81
- modulation transfer functions, 124–125
- Monte Carlo techniques, 24, 139
  - Hamiltonian, 248
  - see also* Markov chain Monte Carlo methods
- Morris-Lecar model, 117
- mortality risk estimation, MIMIC II database, 271–273
- motor cortex, 207
  - experimental data, 218
    - collection, 218
  - factor analysis, 225
  - general linear model (GenLM), 211–212
  - hidden state framework, 209
  - Kalman filter, 209–211
    - with hidden states (KFHS), 212
  - primary, 5
- motor neuroprosthetics, 5, 6
- multi-input-multi-output (MIMO) model, 132
- multimove algorithm, 248
- multiple linear regression, 208
- multivariate autoregressive (MAR) model, 54, 66, 73
  - fMRI time series, 92–93
    - parameter numbers, 93
- myocardial infarction, 283
- nearest-neighbors AR with exogenous input (NNARX), 80
  - fMRI time series, 93
    - local (voxel-level) model, 94
    - non-Gaussian data, 97–98
- model fitting tools, 81
- parameter estimation, 99
- spatial whitening, 96, 98–99
- state space variant, 81, 96–99
  - observation noise, 106
  - parameter estimation, 101–102
- negative binomial (NB) distribution, 202
- negentropy, 287, 288
- Nelder-Mead simplex algorithm, 84, 100
- neocortex, 186
- neural decoding, 210
- neural field models, 118
- neural mass models, 116
  - applications, 127–130
  - conductance models, 117
  - convolution modeling, 117, 118
- excitatory cells
  - pyramidal, 123
  - spiny, 122

- neural mass models (cont.)
  - firing rate, 116
  - firing rate curve, 116
  - Fokker-Planck, 117
  - inhibitory cells in supergranular layers, 122
  - linear approximations, 123
  - Lyapunov exponents, 125–126
  - modulation transfer functions, 124–125
  - state space formulation, 122
- neural models, 33
- neural networks, 284
- neural prosthetics, 9, 207
- neural spike trains, 9, 156
  - decision-making and, 160
    - switching model, 173–175
  - linear state space models, 139–141
  - motor cortex
    - latent variable models, 225
    - prior methods, 208
  - as point process, 189
  - Poisson linear dynamic systems, 138–139
  - rat hippocampus
    - Bayesian decoding, 187–188
    - decoding, 186
    - experimental data, 193
    - Kernel density estimation, 190–192, 194
    - sleep states, 202
    - sorting algorithms, 187
    - state estimation from, 141
      - Laplace approximation, 142–144
- neuronal ensembles, 9
- neuronal plasticity, 9
- noise
  - diagonal state space method, 60
  - NNARX-SS method,
- non-Poisson process (NPP), 215, 222
- normalization, 4
- normalized log-likelihood ratio (NLLR), 219, 220
- normalized root mean squared error, 70
- observability, 3
- Othahara syndrome, 330
- overfitting, 8
- parameter estimation, 99
  - autoregressive (AR) models, 45–57
  - discrete state Markov models, 44
  - functional magnetic resonance imaging (fMRI), 99, 108
    - computational expense, 109
  - general linear model (GenLM), 100–101
  - latent Markov model (LMM), 32, 43–44
  - nearest-neighbors AR with exogenous input (NNARX), 99
  - nearest-neighbors AR with exogenous input (NNARX-SS), 101–102
  - Poisson linear dynamic systems (PLDS), 153, 147–155
  - Parkinson's disease, 128
  - particle filtering, 40–42, 168
    - as assumed density filtering (ADF), 42–43
      - neural models of decision-making, 168
  - partition function, 8
  - peri-stimulus time histogram (PSTH), 141
  - phenotypic dynamics, 283
  - physiological cohort time series, 283
  - physiological monitoring
    - artefacts, 25–29
  - PLDSID algorithm, 152
  - point process filter, 208, 217, 324
    - adaptive, 303, 310–311
    - cardiophysiological data, 304–306, 310–311
  - point process models
    - decision-making, 163–178
    - heartbeat intervals, 304–306
    - neural spike trains, 189
  - Poisson distribution, 189
  - Poisson linear dynamic systems (PLDS), 138–139
    - behavior on synthetic data
      - model generation, 152
    - parameter estimation, 153, 147–155
    - subspace identification, 151–152
  - poles, 125
  - polysomnography (PSG), 6
  - population vectors, 208
  - posterior-predictive likelihood, 177
  - pre-whitening, 86
  - premature babies, 25
    - heart rate, 26
  - premature ventricular contractions (PVC), 355
  - preprocessing
    - functional magnetic resonance imaging (fMRI), 82, 95, 102
    - tilt-table experiments, 268
      - see also* filtering; smoothing
  - primary motor cortex (M1), 5, 9
  - primate models
    - learning, 239, 241–244
      - deep brain stimulation, 244–247
    - M1 neurons, 9
    - motor cortex, 207, 218
  - principal components analysis (PCA), 46–47, 79, 353, 354
  - probabilistic time series models, 14–15
    - graphical depiction, 14–15
  - probability density function (pdf), 1
  - probability mass function (pmf), 1
  - posterior predictive distribution, 8
  - Q-Q plot, 305
  - random variables, 1
    - observed and unobserved, 1–2

- rapid eye movement (REM) sleep, 6, 202  
 rat models, 9
  - hippocampal neurons, 186
  - learning, 239–242
 Rauch-Tung-Striebel (RTS) pass, 40, 100  
 recurrent neural network (RNN), 284, 292  
 regularization, 8  
 respiratory effort, 302  
 representation learning, 284  
 reset models, 29–30  
 respiratory pressure, 307  
 respiratory sinus arrhythmia (RSA), 302  
 response-bias models, 239, 241–244  
 risk minimization, 285  
 rwSSM, 68
- SAPS, 277  
 schizophrenia, 115, 131  
 seizure, 123  
 semi-definite programs, 145  
 sequence disambiguation problem, 34  
 sequential importance sampling (SIR), 169  
 signal quality index (SQI), 353, 354, 361  
 silicon multielectrode array recording, 137  
 since-cell recordings, 114  
 sleep-stage scoring, 5–6  
 slow-wake sleep (SWS), 202  
 smoothing, 18, 19, 84
  - assumed density (ADS), 40
  - correction, 19–20
  - Gaussian sum, 25
  - linear dynamical systems (LDS), 23
  - parallel, 19
  - Rauch-Tung-Striebel (RTS) pass, 40
- source reconstruction, 54–56, 59
  - Bayesian, 55–57
  - challenges, 57
  - computational cost, 57–58
  - diagonal state space method, 59–64
  - full state space model (fSSM), 64–66, 73
  - future directions, 74–75
  - nonlinear and nonstationary dynamics, 74
  - norm regularization, 55
  - parameter numbers, 58
  - reduction, 58–59
  - state space methods, 55, 57
- sparseness (of a matrix), 93  
 spatial whitening, 96, 98–99  
 spatiotemporal Poisson process (STPP), 188  
 spectral learning, 149–150  
 spike trains *see* neural spike trains  
 spikes, 137  
 SPM software, 119, 120  
 SPM5, 102  
 state equations, 2  
 state receptive fields, 200  
 state space models (SSM), 2, 54, 186
  - applications, 9
  - burst suppression, 338
  - conditional probability, 2
  - diagonal *see* diagonal state space method
  - full (fSSM), 66
    - source reconstruction, 66, 73
  - general linear model (GenLM), 85–90
  - hemodynamic response function (HRF), 87–91
  - heart rate, 349–351
  - learning, 231–233
    - T-maze, 239
  - linear
    - likelihood function, 7
    - neural spike train modeling, 139–141
    - for neural spike trains, 138
    - state transition probability, 2
- stochastic processes, 1  
 striatum, 128  
 structured output classification, 285  
 subspace identification, 156
  - Gaussian-observation linear dynamical system (GLDS), 150–151
- support vector machines (SVM), 349
  - heart rate, 357
- switching Kalman filters (SKF), 354
  - electrocardiogrammetry (ECG), 356, 357
- switching linear dynamical systems (SLDS), 23, 30, 38, 258
  - clinical applications, 25–29
  - computational tractability, 24
  - independence structure, 24
- switching Markov processes, 258  
 switching vector autoregressive (SVAR) models, 21, 258, 285–289
  - expectation maximization, 286, 289, 291
- gradients
  - inference and hidden state marginal, 297
  - model parameters, 299
  - regression layer, 297
- physiological cohort time series, 283  
 simulated time series, 289–290  
 tilt-table experiments, 290–293  
 time series classification, 285–289  
 synaptic modeling, 35–37
  - depression factors, 36
- tachogrammetry, 303  
 tank model, 87  
 telemedicine, 346  
 temperature probes, 25  
 temporal marked point process, 189  
 temporal smoothing, 86  
 tetraplegia, 9  
 thinning, 235  
 tilt-table experiments, 268, 271, 316–317
  - cardiophysiological data, 316–317
  - patient risk stratification, 270

- tilt-table experiments (cont.)
  - switching autoregressive (SVAR) models, 290–293
- time series, 14
- unobserved variables, 3
- uric acid, 284
- variance of conditional expectation (VarCE), 181
- variational Bayes (VB) technique, 46–47, 61
  - fMRI time series analysis, 84
  - free energy, 61
  - rat hippocampal neurons, 196
- variational estimation, 24, 46–47
- variational Kalman filtering, 62
- vector of covariates (COV), 308
- vegetative state, 131
- Volterra-Wiener series, 302
- weak limit approximation, 263
- Weibull hazard function, 340
- white-matter pathways, 66
- whitening, 80
  - fMRI time series data, 94–95
- wide-sense stationary (WSS) processes, 1
- Wiener-Volterra expansions, 307–309
- WinBUGS, 233, 235, 245, 248
- X-factor, 358
- XClust, 199
- Yokogawa MEG system, 68
- zeroes, 125–126