

Index

- abelian ordering, 223, 261
- admissible
 - Hilbert function, 740
 - labelled polynomial, 782
 - ordering, 7
- algebra, 176
 - almost commutative, 291
 - Boolean, 450
 - Clifford, 280
 - exterior, 280
 - finitely presented, 192
 - standard, 291
 - graded, 192
 - Grassmann, 280
 - Lie, 258
 - magma, 185
 - monomial, 193, 207
 - path, 164, 335
 - S-invariant, 332
 - strictly ordered, 291
 - universal enveloping, 259
 - Weyl, 264, 349, 358, 402–405
- algorithm
 - Buchberger's, 35, 106, 114, 136–137, 165, 171, 485, 565, 587
 - F_4 , 779–781
 - F_5 , 781–792
 - Schreier's, 588, 663
- allgemeine
 - coordinate, 594
- alternating group
 - (SAGBI basis), 519–527
- alternative comprehensive Gröbner basis, 465
- analytic branch, 624
- ancestor, 776
- Anick's resolution, 214
- annihilator, 46, 393
- antichain, 546
- anticommutative, 258
- antigraded ordering, 773
- antisymmetry, 258
- appliance, 185
- approximate quotient, 480
- approximation
 - initial, 626, 628
 - Taylor, 576, 601, 656
- archimedean ordering, 224
- ARGH decomposition, 445
- associated graded
 - module, 353
 - in Hironaka theory, 573
 - Ore extension, 88, 321
 - ring, 267, 318, 336, 351, 353, 567, 592
 - of a local ring, 593
 - vector space, 498
- automaton, 199
 - determined, 199
- Bézout, E.
 - domain, 15, 78–81
 - ring, 15, 109–117
- barrier orderings, 243
- basis
 - L -involution, 775
 - factor-SAGBI, 556
 - Göbel, 539
 - Gröbner, 11, 88
 - reductive, 179
 - of symmetric ideals, 545
 - involution, 761
 - J-marked, 751
 - SAGBI, 514, 552
 - SAGBI–Gröbner, 553
 - shift, 153
 - standard, 568
 - Szekeres-like, 15, 90
- Bergman's Diamond Lemma, 169, 261–264, 266
- Betti numbers, 740
- binomial skew polynomial ring, 298
- Boolean
 - algebra, 450
 - ideal, 451
- Borel, É.
 - escalier, 745–748
 - ideal, 665, 670, 675, 679, 692, 741, 749–750, 752
 - property, 682–690

- Borel, É. (cont.)
 relation, 523, 665, 687, 689, 691, 708–716, 724
 set, 637, 679, 691, 716–720, 744
- Buchberger's
 algorithm, 35, 106, 114, 136–137, 165, 171,
 485, 565, 587
 First Criterion, 51, 94, 128, 142–144, 189, 382,
 386, 608, 777
 reduction, 18, 21, 24, 37, 164, 349, 480, 515,
 540, 565, 581, 609, 614, 619, 689, 779
 Second Criterion, 12, 776
- canonical
 comprehensive Gröbner basis, 437, 439
 form, 20, 23, 398
 Janet, 760
 Gröbner
 cover, 446
 system, 438
 tree, 438
 representatives, 22
- Cartan, E., 664, 758
 characters, 634, 664, 692, 718, 758
 test, 635, 664, 665, 673, 691, 737, 760, 762, 771
- Cauchy, A. L.
 sequences, 581, 579, 593
- center, 69, 268, 368
- chain, 212, 548
- characteristic, 71
- class, 713
 (Janet), 641, 669
- coherent, 163
- complementary valuation ring, 476
- complete, 641
- completion, 54, 93, 579, 592, 643, 766
- comprehensive
 Gröbner basis, 426
 alternative, 465
 canonical, 437, 439
 SAGBI basis, 528
- condition, 426
 quasi-canonical, 430
- conjugate (Ore), 69, 314
- constant
 ring, 69
- contraction, 556
- convex, 232
 cone, 529
- cote, 655, 683
- criteria
 F_5 , 783
 extended F_5 , 785
 Buchberger's First, 51, 94, 128, 142–144, 189,
 382, 386, 608, 778
 Buchberger's Second, 12, 776
 Gebauer–Möller, 12, 129, 135, 157, 332, 553
- critical set, 179
- cycle
 intermediate, 197
 path, 193
 sink, 197
 source, 197
- cyclic
 condition, 293, 301
 group, 9
 path, 193
- deglex ordering, 102, 222, 309
- degree extension, 224
- degree reverse ordering, 309
- Delassus, E., 602, 634, 637, 664, 665, 678,
 684–686, 691, 698, 699, 713, 729, 739
- derivation, 69, 258
- derivative (Ore), 69
- derived, 693, 741
- descending term, 531
- dialytic equations, 651, 656, 665, 698
- Diamond lemma, 169, 261, 264, 266
- Dickson's lemma, 9, 137, 296, 312, 317, 349, 395,
 509, 583, 637, 729
- differential
 operators, 265, 281
- dimension
 Gel'fand–Kirillov, 193
- direct product, 449
- direction ordering, 245
- distinguished, 611
- division
 \succ_{alex} , 773, 778
 involutive, 764
 constructive, 773
 continuous, 773
 pairwise, 771
 Janet, 760, 766, 778
 Pommaret, 760, 766
 Thomas, 766
- domain, 78
 Bezout, 15, 78–81
 well-endowed, 797
- Duval, D.
 field, 626
 splitting, 424
- écart, 582
- echelon set, 109, 126, 188, 397, 752, 785
- effectively given, 337, 341
- elementary symmetric product, 531
- endlichvielen Schritten, 337
- escalier, 20, 664, 745
 Borel, 745–748
 generic, 602, 620, 634, 691
- examples
 Gaber–Kashiwara, 602
 Green–Ufnarowski, 204, 239
 Janet, 638–642, 654
 $k[X, Y, Z]/\mathbb{I}(XZ - YZ)$, 393
 Pan, 11, 14, 16, 26, 50, 346
 \mathbb{Q} as \mathbb{Z} -module, 340

- Riquier, 636–637, 641, 643–648, 766–770
- Sweedler, 490, 491, 516
- Ufnarovski, 138, 198
- Weispfenning ring, 79, 83, 88, 90, 91, 124, 314–324, 336, 349, 350, 357, 410–417
- Weyl algebra, 358
- explicitly given, 337
- extension, 7, 163
 - natural, 261, 264, 267
- exterior algebra, 280
- Factor-SAGBI basis, 556
- fair strategy, 106, 121, 136–142, 385, 487, 519
- faithful
 - canonical Gröbner system, 438
 - canonical Gröbner tree, 438
 - Gröbner system, 429
 - preimage (minimal), 438
- filtration
 - (nested), 492, 498
 - full union, 492
 - intersects to zero, 492
 - \mathbb{K} -filtration, 495, 500, 507, 511
 - normalized, 507
 - equivalence, 492
 - order, 492
- final subductum, 515, 552
- finite segment, 546
- form
 - canonical, 20, 398
 - L -canonical form, 775
 - leading, 43, 89, 103, 116, 118, 267, 272, 363, 368, 567, 568, 593
 - normal, 16, 361
 - Weierstrass, 618
- formal term, 12, 125
- function ring, 176, 336
- \mathcal{G} -invariant
 - orbit, 520
 - special, 533
 - polynomial, 520
- G -presentation, 160, 169, 219, 389
 - symmetric group, 220
- Göbel, M.
 - basis, 539
 - ordering, 532
 - representation, 539
- Gaber–Kashiwara example, 602
- Gateva’s monoid, 87, 288–291, 378
- Gebauer–Möller
 - criteria, 12, 129, 135, 157, 332, 553
 - set, 43, 89, 104, 116, 118, 126–136, 145, 272, 366, 369, 569, 779
 - non-trivial, 120, 382, 384, 386
 - Staggered Linear Bases, 792
- Gel’fand–Kirillov dimension, 193
- generic
 - escalier, 602, 620, 634, 691
 - Gröbner basis, 437
 - ideal, 437
 - initial ideal, 634, 678–681, 685–690
 - Gianni’s proposition, 430
 - Gianni–Kalkbrener theorem, 423, 444
 - Gordan, P., 689
 - Gröbner
 - (strong) representation
 - relative to a complementary valuation ring, 480
 - basis, 11, 88, 359
 - alternative comprehensive, 465
 - canonical comprehensive, 437, 439
 - comprehensive, 426
 - generic, 437
 - minimal, 12
 - reduced, 20, 30, 96
 - regular, 200
 - relative to a complementary valuation ring, 480
 - strong, 11, 88, 359
 - of symmetric ideals, 545
 - reductive basis, 179
 - representation
 - (strong), 359
 - (weak), 359
 - quasi, 118, 272, 366, 369
 - reductive, 179
 - relative to a complementary valuation ring, 480
 - restricted, 323
 - representation (of a polynomial), 11, 88
 - quasi, 12, 104, 116
 - strong, 12
 - weak, 11
 - representation (of an algebra), 445
 - system, 428
 - canonical, 438
 - faithful, 429
 - graded
 - module, 351
 - morphism, 352
 - ring, 351
 - grafting, 185
 - greatest common right divisor, 71
 - group
 - alternating
 - (SAGBI basis), 519–527
 - cyclic, 9
 - ordered abelian, 472
 - reflection, 528
 - ring, 157–160, 335, 349, 405–407, 542
 - symmetric, 301, 520
 - (representation), 220
 - value, 474
 - growth, 192
 - exponential, 192
 - polynomial, 192

- Gunther, N., 522, 634, 637, 664, 678, 686, 687, 691, 739
 GUPI monoid, 473
 Hilbert, D., 634
 function, 692, 700–702
 (admissible), 740
 series, 192
 Hironaka's
 box condition, 620
 Henselian Weierstrass theorem, 602, 620–622
 theorem, 397, 577, 617
 l-regular function, 443
 ideal
 (of monoid), 473
 bilateral, 6
 Boolean, 451
 generic, 437
 homogeneous, 142
 invariant, 543
 left, 6
 letterplace, 151
 lex segment (lexicographical), 694, 713
 of magma algebras, 186
 order, 7, 692
 partial difference, 334
 right, 6
 S-invariant, 332
 semigroup, 7
 shift-invariant, 150
 symmetric, 543
 Szekeres, 15, 266, 269, 288, 340
 of vanishing polynomials, 96
 idempotent, 451
 incident matrix, 193
 index of nilpotency, 59
 index of Puiseux series, 622
 index of regularity, 675, 701, 730, 755
 inf-limited, 352, 495, 565, 773
 initial
 approximation, 626, 628
 exponent, 626
 interreduced set, 187
 invariant ideal, 543
 invariant ring, 69
 inverse system, 651
 involutive, 673, 676, 678, 758, 766
 basis, 621, 761
 cone, 766
 division, 614, 764
 constructive, 773
 continuous, 773
 pairwise, 771
 iterative Ore extension, 85
 J-marked family, 751
 J-reduced, 751
 Jacobi, C. G. I.
 identity, 258
 jacobian, 597, 624, 626, 630
 Janet, M., 338, 573, 633–691, 698, 711, 713, 718, 739, 758, 780
 canonical form, 760
 division, 760, 766, 778
 normal form, 760
 Janet-reduced, 760
 Kandri-Rody–Weispfenning completion, 257, 275–278, 313, 412
 Kleene's Theorem, 200
 Kredel's conditions, 268, 397
 L-canonical form, 775
 L-completion, 766
 L-involutive, 766
 L-involutive basis, 775
 L-normal form, 775
 L-representation, 774
 labelled polynomial, 782
 admissible, 782
 extended normalized, 785
 normalized, 783
 Lagrange, J. L.
 interpolation formula, 341
 Lazard, D., 469, 588
 theorem, 56–63, 83–85, 98
 leading
 coefficient, 7, 87, 175, 318, 351, 652
 derivate, 652
 form, 43, 89, 103, 116, 118, 267, 272, 363, 368, 567, 568, 593
 term, 652
 leitideal, 576, 592, 593
 leitmodul, 574
 length, 102, 221
 function, 797
 letterplace
 ideal, 151
 monomial, 149
 ring, 149
 lex segment ideal, 694, 713
 lexicographical
 ideal, 694, 713
 ordering, 102, 221
 Lie, S.
 bracket, 258
 algebra, 258
 field, 260
 lift, 43, 104, 116, 118, 272, 365, 369, 569, 735
 lifting
 polynomial, 436
 theorem, 44, 89, 104, 246, 321, 335, 588, 662, 735, 737
 Liouville, J., 603
 local smooth system, 598
 standard, 607

- localization, 183, 426, 593
 locally closed, 441
 LSS, 598
- Macaulay, F. S., 740
 matrix, 757, 778
 resultant, 778
 magma, 184
 match, 126, 160
 matrix
 Macaulay's, 757
 Sylvester, 760
 maximal
 monomial, 7, 87, 175, 318, 351
 term, 7, 87, 175, 318, 351
 mesh step, 333
 module
 order, 6, 350
 S-invariant, 325
 semigroup, 6
 Möller's
 lemma, 54, 129–131, 135, 144, 553
 monoid
 G-presentation, 160, 169, 219, 389
 Gateva's, 87, 288–291, 378
 GUPI, 473
 ideal, 473
 Pritchard, 19, 144
 ring, 6, 160–162, 407–410
 free, 336
 skew, 325
 monomes complémentaires, 648
 monomial set, 7
 monotone ordering, 7
 multiplicative
 basis, 163, 176, 390
 system, 39, 143
 variable, 638, 641, 669, 737, 780
- Newton polygon, 632
 Noetherianity, 278, 312, 315, 334, 349, 394–397,
 402, 411, 415, 455, 485, 488–492,
 546–547, 656
 non-trivial Gebauer–Möller set, 120, 382, 384,
 386
 normé, 665, 679, 687, 688
 normal
 (a), 713, 717
 (b), 713, 717, 728, 734, 744
 form
 (weak), 16, 361
 Janet, 760
 L-normal form, 775
 relative to a complementary valuation ring,
 480
 for SG-bases, 553
 strong, 16, 361
 for symideals, 539
 with standard representation, 568
 monomial, word, 193
 normalized, 709
 admissible labelled polynomial, 783
 pair, 784, 786
 normé, 709
 Nullstellensatz bounds, 796
 numération, 710
 (a), 710
 (b), 711
- obstruction, 126
 orbit (G -invariant), 520
 special, 533
 order
 of analytic branch, 624
 ideal, 692
 of irregularity, 68, 81
 module, 6, 350
 of Puiseux series, 622
 ordered abelian group, 472
 ordering
 abelian, 223, 261
 admissible, 7
 antigraded, 773
 archimedean, 224
 barrier, 243
 block, 217, 341, 605
 centerleft, 228, 379
 centerright, 228, 379
 deglex, 102, 222, 309
 degree reverse, 309
 direction, 245
 elimination, 217, 229
 left, 228, 341, 379
 lexicographical, 102, 221
 monomial, 7
 position-over-term, 248, 711, 713
 reverse, 222
 right, 228, 379
 right reductive restriction, 178
 S-termordering, 334
 semigroup, 7, 349, 755
 sequential, 142, 144, 171
 strongly compatible, 231
 symmetric
 cancellation partial, 543
 term, 7
 term-over-position, 711
 weight compatible, 224
- Ore
 algebra, 268, 314
 condition, 260
 extension
 graded, 314, 325, 327
 multivariate, 123
 Ore, E.
 algebra, 85, 335, 418
 condition, 64, 67
 extension, 16, 69, 418–422

Ore, E. (cont.)
 graded, 70, 88
 iterative, 85
 iterative (with commuting variables), 85,
 335, 418
 multivariate, 85, 335, 418
 with zero derivations, 88

pair
 extended rewritable, 786
 normalized, 784, 786
 not extended rewritable, 786

parametrization, 624

path, 193
 algebra, 164, 335
 cycle, 193
 cyclic, 193

permutation
 shift, 549

Poincaré–Birkhoff–Witt theorem, 259, 314

polynomial extension, 28, 178, 276, 436, 454

Pommaret, J. F.
 division, 760, 766

positive set, 472

postfix, 109

prechain, 212

prefix, 158, 179, 212
 reduction, 158

preimage (minimal), 438

preterrace, 459

Pritchard
 monoid, 19

Pritchard, F. L.
 module, 121–124, 144, 397
 monoid, 144

proper endings, 193

pseudovaluation, 352

Puiseux series, 622

quantum plane, 265

quasi-embedding, 546

quasi-Gröbner representation, 43, 104

quasi-inverse, 451

quasi-order, 492
 ~-lex, 232
 reverse ~-antilexicographical, 232

quasi-ordering, 546

quotient
 approximate, 480
 field, 64, 260, 274
 ring, 39, 143

Quotientenkörper, 64

Rabinowitch trick, 250, 445

ramification index, 622

rank
 of an ideal, 594

reduction, 533

redundant
 basis element, 131
 S-element/pair, 52, 135

reflection group, 528

regular
 function, 425, 441–444
 Gröbner basis, 200
 language, 199
 ring, 64, 67, 78

representation
 Gröbner
 (of a polynomial), 11, 12, 88
 (of an algebra), 445
 reductive, 179
 restricted, 323
 Göbel, 539
 of algebraic power series, 609
 standard, 568
 Y-standard, 323

residue class field, 474, 593

resolution, 212, 588

resultant
 Macaulay's, 778
 Sylvester, 562–564

revlex segment, 713, 728, 734, 744

ring
 artinian chain, 59
 Bezout, 15, 109–117
 constant, 69
 function, 176, 336
 group, 157–160, 335, 349, 405–407, 542
 invariant, 69
 letterplace, 149
 monoid, 6, 160–162, 407–410
 free, 336
 quotient, 143
 skew monoid, 325
 skew polynomial, 264
 solvable polynomial, 264, 268, 271–278, 335,
 349
 strong, 23, 96
 twisted polynomial, 279
 valuation, 471
 complementary, 476
 Von Neumann regular, 450
 Zacharias, 19, 37–47, 91–95, 144, 271, 338

Riquier, C., 338, 633, 634, 651, 655, 682–684,
 698, 711, 735, 758

Ritt, J. F., 758

Robinson, L. B., 634, 637, 664, 686, 687, 698, 739

S-element, 51, 125, 139
 redundant, 52, 135
 trivial, 128, 142, 330
 useful, 52, 95, 779
 useless, 50, 51, 126–136, 182–184, 757

S-pair, 51, 125, 139

S-polynomial, 44, 89, 104, 116, 125, 153, 180,
 188, 329, 366, 776

- for symideals, 540
- SAGBI
 - (strong) representation, 515, 552
 - basis, 514, 552
 - comprehensive, 528
 - representation, 515, 552
- SAGBI–Gröbner basis, 553
- saturated
 - weakly, 183
- saturating set, 158
- saturation, 158, 160, 179, 335, 457
- saturator, 183
- Schreier’s algorithm, 588
- segment
 - finite, 546
 - generalized revlex, 749
 - lex, 694, 713
 - revlex, 713, 728, 734, 744
- semi-Thue system, 160, 169, 218, 219, 389
- semigroup
 - module, 6
 - ordering, 7, 349, 755
 - extension wrt LSSs, 605, 608
- sequential, 142, 144, 171
- set
 - critical, 179
 - echelon, 109, 126, 188, 397, 752, 785
 - monomial, 7
 - positive, 472
 - saturating, 158
- SG (SAGBI–Gröbner)
 - (strong) representation, 553
 - basis, 553
 - normal form, 553
 - representation, 553
- SG-basis, 553
- shift, 149
- shift permutation, 549
- shift-basis, 153
- shift-invariant ideal, 150
- shift-uniform, 149
- sig-redundant, 788
- skew
 - monoid ring, 325
 - polynomial ring, 264
 - binomial, 298
- SLSS, 607
- solvable polynomial ring, 264, 268, 271–278, 335, 349
- Spear’s theorem, 336
- Spear’s theorem, 27, 98, 171, 350, 372
 - for standard bases, 610
- special principal ideal ring, 59
- specialization, 426
 - epi, 427
 - essential, 436
- Squier’s theorem, 137, 138, 296
- stable, 178
 - localization, 183
- saturator, 183
- standard
 - basis, 43, 89, 104, 116, 118, 272, 354, 568
 - local smooth system, 607
 - representation, 43, 89, 104, 116, 118, 272, 354, 568
- Steinitz, E.
 - notation, 71
- strategy, fair, 106, 121, 136–142, 385, 487
- strong
 - Gröbner basis, 11, 88
 - ring, 23, 96
- support, 5
 - (of a preterrace), 459
 - (of a terrace), 463
- Sylvester, J. J.
 - matrix, 760
 - resultant, 562–564
- symideal, 537
- symmetric
 - cancellation partial ordering, 543
 - group, 301, 520
 - representation, 220
 - ideal, 543
- symmetry
 - division wrt, 538
 - l.c.d. wrt, 538
- syzygies, 246, 482, 662, 663
 - relative to a complementary valuation ring, 483
 - trivial, 757
- syzygy basis truncated, 145
- syzygy family
 - relative to a complementary valuation ring, 483
- Szekeres
 - ideal, 15, 266, 269, 288, 340, 547
 - level, 15
 - principal generator, 15, 340
 - semigroup, 15
 - theory, 14–16, 90–91, 109, 338, 340, 341, 346
- Szekeres-like basis, 15, 90
- tag variable technique, 219, 254
- Taylor, B.
 - approximation, 576, 601, 656
- term
 - descending, 531
 - multilinear, 531
 - ordering, 7
 - for module, 711
 - special, 533
 - total reduced, 534
- term-ordering in suitable variables
 - filtration, 512
 - valuation, 509
- terrace, 463
- tête-à-tête, 517, 552, 556, 557

820

Thomas, J. M., 758
 division, 766
 trivial (S-element/pair), 128, 142, 330
 Turing machine, 797
 twisted polynomial ring, 279
 type, 82

Ufnarovski, V.
 example, 138, 198
 Graph, 193
 uniform, 165
 useful S-element/pair, 52, 95, 329, 779
 useless (S-element/pair), 50, 51, 126–136,
 182–184, 757

valuation, 474
 valuation ring, 471
 associated to a (nested) filtration, 503
 complementary, 476
 value
 group, 474
 monoid, 476
 vanishing polynomials, 96
 Von Neumann regular ring, 450

Index

weak representation, 11
 Weierstrass, K. T. W.
 division theorem, 619
 form, 618
 preparation theorem, 611
 weight, 102, 224, 572, 601, 659
 extension, 224, 605
 well-endowed domain, 797
 well-founded, 546
 well quasi-ordered, 546
 well quasi-ordering, 546
 Weyl algebra, 358
 witness, 543
 word, 102
 tree, 207
 Wu, W.-t., 758

Zacharias
 canonical representation, 288, 556
 ring, 144, 271
 Zacharias, G.
 canonical representation, 22, 340–349
 ring, 19, 37–47, 91–95, 338
 zariskification, 592, 597