

1

Hyperbolic space and its isometries

In this chapter we gather together basic information about the geometry of two- and three-dimensional hyperbolic spaces and their isometries. This will set the stage for our study of quotient manifolds and orbifolds which begins in the next chapter.

1.1 Möbius transformations

A *Möbius transformation* in the unit sphere \mathbb{S}^n of dimension n is, by definition, the result of a composition of reflections in $(n-1)$ -dimensional spheres in \mathbb{S}^n . It will be orientation-preserving if it is the composition of an even number of reflections. A defining property is that Möbius transformations send $(n-1)$ -dimensional spheres onto $(n-1)$ -dimensional spheres. Automatically, a symmetric pair of points (with respect to reflection) about one sphere gets sent to a symmetric pair about the other.

From now on, the unqualified term *Möbius transformation* will be reserved for those that preserve orientation. The orientation-reversing kind will be called anti-Möbius transformations. For a discussion of the latter, see Exercise (1-31) at the end of the chapter.

The study of hyperbolic 3-manifolds is intimately connected with the study of Möbius transformations on the two-dimensional sphere \mathbb{S}^2 . Via stereographic projection (Figure 1.1), \mathbb{S}^2 is homeomorphic to the extended plane $\mathbb{C} \cup \infty$, and we will freely use this fact to change points of view between the extended plane and the 2-sphere. Under stereographic projection, the collection of circles and straight lines in \mathbb{C} corresponds to the collection of circles on \mathbb{S}^2 ; a straight line in \mathbb{C} corresponds to a circle on \mathbb{S}^2 through the north pole. With this correspondence in mind, we can refer to the collection of circles and lines in \mathbb{C} simply as “circles”. Moreover stereographic projection is a conformal map, that is, it preserves angles between intersecting arcs—in particular, angles of intersection between circles.

Möbius transformations in two dimensions are *fractional linear transformations* of the extended plane. That is, a Möbius transformation acting on $\mathbb{C} \cup \infty$ has the form

$$z \mapsto A(z) = \frac{az + b}{cz + d}, \quad \text{with } a, b, c, d \in \mathbb{C} \text{ such that } ad - bc \neq 0. \quad (1.1)$$




Fig. 1.1. Stereographic projection.

(When $ad - bc = 0$ the expression on the right is a constant, so the map is not a Möbius transformation.) As we will see shortly, a map of this form can indeed be expressed as the composition of an even number of reflections in circles (in fact, two or four circles: see Exercise (1-7)). The symmetry properties of such maps are established in Exercise (1-2).

Möbius transformations are conformal maps. In fact, the only conformal homeomorphisms of $\mathbb{C} \cup \infty$ are Möbius transformations.

We will generally assume that the representation in (1.1) is *normalized*, meaning that $ad - bc = 1$. Then we can identify the group of Möbius transformations with the quotient $\text{PSL}(2, \mathbb{C}) := \text{SL}(2, \mathbb{C}) / \pm I$, where $\text{SL}(2, \mathbb{C})$ is the group of 2×2 matrices of determinant one and I is the identity matrix:

$$A(z) = \frac{az + b}{cz + d} \longleftrightarrow \pm \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad A^{-1}(z) \longleftrightarrow \pm \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$$

The \pm ambiguity cannot be avoided. We will not keep inserting it, unless it plays an essential role. In any case the value of changing from transformations to matrices lies mainly in the algebra of composition. If A, B are Möbius transformations, the Möbius transformation resulting from the application of A followed by B is written BA ; the corresponding matrix is just the usual product BA of the component matrices, in the order written. The \pm ambiguity follows along. We will hop from one to the other, the representation as a transformation to the representation as a matrix, depending on which best suits the situation, without changing the labeling.

Two Möbius transformations A, B are *conjugate* if there is a Möbius transformation U such that $B = UAU^{-1}$. Conjugate transformations have the same geometry: U effects transfer of the geometry of A to that of B .

The expression $ABA^{-1}B^{-1}$ is called the *commutator* of A and B and written as $[A, B]$. Two elements commute if and only if their commutator is the identity.¹

The *trace* of a Möbius transformation A is, by definition, the trace of the normalized matrix of A :

¹ The alternative conventions $[A, B] = B^{-1}A^{-1}BA$ or $A^{-1}B^{-1}AB$ are preferred by some authors; they do the same job, but the formulas come out differently.

1.1 Möbius transformations

3

$$\tau_A = \operatorname{tr} A = \pm(a + d).$$

It is invariant under conjugation. The \pm ambiguity can be avoided either by using τ_A^2 or by specifying $0 \leq \arg \tau_A < \pi$.

By solving the equation $A(z) = z$, we find that a Möbius transformation $\neq \operatorname{id}$ has one or two fixed points in \mathbb{S}^2 , namely $(a - d \pm \sqrt{\tau_A^2 - 4})/2c$, when $c \neq 0$, or otherwise the points ∞ and $b/(d - a) = ab/(1 - a^2)$. Here $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $ad - bc = 1$. Only the identity can have three fixed points.

Given three distinct points $(p_2, p_3, p_4) \in \mathbb{S}^2$, there exists a unique Möbius transformation T sending p_2 to 0, p_3 to 1, p_4 to ∞ . It is given by

$$T(z) = \frac{(z - p_2)(p_3 - p_4)}{(z - p_4)(p_3 - p_2)} = (z, p_2, p_3, p_4),$$

when none of the points p_i is ∞ . By taking the limit as some $p_i \rightarrow \infty$, we obtain the correct expression for $p_i = \infty$. The expression (z, p_2, p_3, p_4) is called the *cross ratio* of the four points.²

- Given distinct triples (p_2, p_3, p_4) , (p'_2, p'_3, p'_4) there is a unique Möbius transformation taking one triple onto the other.
- Cross ratios are invariant under Möbius transformations:

$$(Az, Ap_2, Ap_3, Ap_4) = (z, p_2, p_3, p_4) \quad \text{for any } A.$$

To establish the second item, note that both Möbius transformations $S(z) = (z, Ap_2, Ap_3, Ap_4)$ and $T \circ A^{-1}(z)$ take (Ap_2, Ap_3, Ap_4) to $(0, 1, \infty)$ so they are identical. Replace z by $A(z)$.

Apart from the identity, Möbius transformations fall into one of three types:

A is *parabolic* if the following equivalent properties hold.

- A is conjugate to $z \mapsto z + 1$.
- A has exactly one fixed point in \mathbb{S}^2 .
- $\tau_A = \pm 2$ and $A \neq \operatorname{id}$.

A is *elliptic* if the following equivalent properties hold.

- A is conjugate to $z \mapsto e^{2i\theta}z$, with $2\theta \not\equiv 2\pi$.
- $\tau_A \in (-2, +2)$.
- A has exactly two fixed points, and the derivative of A has absolute value 1 at each of them.

A is *loxodromic* if the following equivalent properties hold.

- A is conjugate to $z \mapsto \lambda^2z$, with $|\lambda| > 1$.
- $\tau_A \in \mathbb{C} \setminus [-2, +2]$.
- A has two fixed points; at one $|A'| < 1$ (attracting), at the other $|A'| > 1$ (repelling).

² In the induced direction on \mathbb{R} , UHP lies to the left; also $(z, 0, 1, \infty) = z$. A common alternate definition results in $(z, 1, 0, \infty) = z$.

We will use the term *standard forms* for the conjugates just listed. The geometry of a general normalized Möbius transformation A is most easily read off from the conjugate standard form. Note that the elliptic $z \mapsto 1/z$ is conjugate to $z \mapsto -z$.

A loxodromic Möbius transformation A has a collection of *loxodromic curves* or *invariant spirals* in \mathbb{S}^2 . (In navigation, a *loxodromic curve* or *rumb line* is a path of constant bearing: it makes equal oblique angles with all meridians, and so coils around the poles without ever reaching them.) For the standard form $z \mapsto \lambda^2 z$, one such spiral is given by

$$z(t) = \lambda^{2t}, \quad -\infty < t < \infty.$$

If σ denotes the segment $0 \leq t < 1$ of the spiral, the various images $\{A^n(\sigma)\}$ cover the spiral without overlap. See Figure 1.2.

For additional structure in special cases see Wright [2006].

The term *hyperbolic transformation* has historically been used to designate a loxodromic transformation whose trace is real. Such a transformation is conjugate to $z \mapsto \lambda^2 z$ with $\lambda > 1$. Nowadays the term “hyperbolic” is also used for a loxodromic element acting in hyperbolic 3-space.

The classification is proved by first conjugating A so that one fixed point lies at ∞ and the other, if there is one, at 0. The further conjugation $z \mapsto 1/z$ that interchanges 0 and ∞ may be needed to put the attracting fixed point at ∞ .

If $p \in \mathbb{C}$ is a fixed point of $A \neq \text{id}$, p is attracting if and only if $|A'(p)| < 1$ and repelling if and only if $|A'(p)| > 1$. The transformation A is parabolic if and only if $A'(p) = 1$; A is elliptic if and only if $|A'(p)| = 1$ but $A'(p) \neq 1$.

Upon referring to the normalized matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, we find that the *eigenvalues* are $\lambda, \lambda^{-1} = \frac{1}{2}(\text{tr } A \pm \sqrt{\text{tr}^2 A - 4})$. The corresponding eigenvectors $\begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ satisfy

$$\frac{\alpha}{\beta} = \frac{\lambda - d}{c} = p \quad \text{and} \quad \frac{\alpha}{\beta} = \frac{\lambda^{-1} - d}{c} = q,$$

where p, q are the fixed points. Like the trace, the eigenvalues are invariant under conjugation. The eigenvalues of an elliptic transformation have the form $e^{\pm i\theta}$ and the




Fig. 1.2. Invariant spiral of a loxodromic with trace $\lambda + \lambda^{-1} = 1.976 + 0.005i$.

trace is $2 \cos \theta$. A loxodromic transformation has eigenvalues $\lambda^{\pm 1}$ and trace $\lambda + \lambda^{-1}$. We can choose λ so that $|\lambda| > 1$, that is, so that λ is the expanding eigenvalue.

The expanding eigenvalue of a loxodromic element A can be expressed as a cross ratio by the formula

$$\lambda^2 = (z, p_+, A(z), p_-),$$

where p_+, p_- are the attracting and repelling fixed points. (It is enough to confirm this when $p_+ = \infty$ and $p_- = 0$.)

We can write $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ as

$$Az = \frac{1}{-c^2(z + d/c)} + \frac{a}{c} \quad \text{if } c \neq 0, \quad Az = \frac{a}{d} \left(z + \frac{b}{a} \right) \quad \text{if } c = 0. \quad (1.2)$$

This expresses A in terms of simple building blocks: maps in standard form, plus the map $z \mapsto 1/z$. Each of these has the property of preserving (generalized) circles. Therefore any Möbius transformation preserves circles, as mentioned earlier. Likewise each building block is easily seen to be a composition of two reflections, so a Möbius transformation is the composition of an even number of reflections.

Three distinct points p_2, p_3, p_4 uniquely determine a circle C , with an orientation determined by their order. When C is a proper circle or any closed curve, the *positive orientation* is determined by a sequence of points p_2, p_3, p_4 in counterclockwise direction, that is when the interior of the curve lies to the left. Let q_2, q_3, q_4 be another set of distinct points, and C' the circle through them. The Möbius transformation T that sends $p_i \rightarrow q_i$ automatically sends C onto C' . If both are proper circles, T sends the interior of C to the interior of C' if and only if the triples give both circles the same, positive or negative, orientations.

The transformation $T : z \rightarrow w$ can be expressed in terms of cross ratios as

$$(w, q_2, q_3, q_4) = (z, p_2, p_3, p_4).$$

But if we focus simply on sending C to C' , and a designated side of C to a designated side of C' , it is easier to find T by cross ratio using the symmetry property: A Möbius transformation sends points symmetric with respect to reflection in one circle, to a pair of points symmetric in its image (Exercise (1-2)). For a proper circle, the most conspicuous symmetric points are its center and ∞ . If the circle is $\{z : |z - z_0| < r\}$, the points $|a| < 1$ and $|a^*| > 1$ are symmetric if and only if $a^* - z_0 = r^2 / \overline{(a - z_0)}$.

A cross ratio (p, p_2, p_3, p_4) is real if and only if the four points lie on a circle in \mathbb{S}^2 . The cross ratio is positive if and only if (p, p_3, p_4) gives the circle the same orientation as (p_2, p_3, p_4) .

We are now ready to show that Möbius transformations in $\mathbb{C} \cup \infty$ can be extended to Möbius transformations acting in upper half-space (UHS) $\{\vec{x} = (z, t) : z \in \mathbb{C}, t > 0\}$ (and, correspondingly, lower half-space (LHS)). The simplest way to see this is by applying the following observation. Each Möbius transformation is the composition of an even number of reflections in circles or lines in \mathbb{C} . A reflection in a circle extends

naturally to the reflection in the upper hemisphere bounded by that circle. Likewise the reflection in a straight line extends to the reflection in the vertical half-plane bounded by that line. (The same argument shows that Möbius transformations on $S^n = \mathbb{R}^n \cup \{\infty\}$ extend to upper half $(n+1)$ -space.)

A Möbius transformation acting on $\mathbb{C} \cup \infty$ sends a given circle to another circle or line. Its extension to upper half-space will therefore map the hemisphere bounded by the circle to the hemisphere or half-plane bounded by the image of the circle. We conclude that the extension to upper half-space maps the totality of hemispheres and vertical half-planes onto itself.

If two hemispheres intersect, or a hemisphere and a vertical half-plane intersect, the intersection is a semicircle which is orthogonal to \mathbb{C} . If two vertical half-planes intersect, they intersect in a vertical half-line orthogonal to \mathbb{C} . The extension of a Möbius transformation thus maps the totality of half-lines and semicircles orthogonal to \mathbb{C} onto itself. The dihedral angles between intersecting hemispheres is the same as the angle of intersection between their bounding circles in \mathbb{C} .

It is useful to explicitly work out the formula for extension to upper half-space $\{\vec{x} = (z, t) : z \in \mathbb{C}, t > 0\}$. We first extend the building blocks. First,

$$\begin{aligned} z \mapsto az & \text{ becomes } (z, t) \mapsto (az, |a|t); \\ z \mapsto z + b & \text{ becomes } (z, t) \mapsto (z + b, t). \end{aligned}$$

The inversion $z \mapsto z^{-1}$ is most easily dealt with as the composition of two anti-Möbius transformations: $z \mapsto \bar{z}$ (reflection in a line) and $z \mapsto z/|z|^2 = \bar{z}^{-1}$ (reflection in the unit circle). Extending to reflections in a vertical plane and the unit hemisphere, we get respectively $(z, t) \mapsto (\bar{z}, t)$ and

$$\vec{x} \mapsto \frac{\vec{x}}{|\vec{x}|^2} \quad \text{or} \quad (z, t) \mapsto \left(\frac{z}{|z|^2 + t^2}, \frac{t}{|z|^2 + t^2} \right).$$

Therefore,

$$z \mapsto \frac{1}{z} \quad \text{becomes} \quad (z, t) \mapsto \left(\frac{\bar{z}}{|z|^2 + t^2}, \frac{t}{|z|^2 + t^2} \right).$$

Composing the building blocks we find that the extension of $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is

$$(z, t) \mapsto \left(-\frac{\overline{z + d/c}}{c^2(|z + d/c|^2 + t^2)} + \frac{a}{c}, \frac{t}{|c|^2(|z + d/c|^2 + t^2)} \right) \quad \text{when } c \neq 0.$$

$$(z, t) \mapsto \left(\frac{a}{d}(z + b/a), \left| \frac{a}{d} \right| t \right) \quad \text{when } c = 0.$$

A Möbius transformation acting in UHS acts symmetrically in LHS under the correspondence $(z, t) \leftrightarrow (z, -t)$.

1.2 Hyperbolic geometry

In the euclidean plane, there is exactly one line through a given point and not meeting a given line disjoint from the point; this is the famous fifth postulate of Euclid. It

gradually became clear in the nineteenth century that one can have a self-consistent and interesting geometry where this postulate is not valid—where “parallel” lines are not unique and indeed exist in uncountable abundance. This became known as *hyperbolic geometry*. Though the name was bestowed in connection with conics and projective geometry [Klein 1871, p. 72], it is a doubly felicitous choice, because the Greeks had named the hyperbola after the word for excess (compare “hyperbole”, from the same Greek word). Hyperbolic geometry certainly has an excess of lines—and of “room”—compared to euclidean geometry!

Here are some of the salient features that distinguish hyperbolic geometry from the familiar euclidean and spherical geometry.

- (i) The angle sum Σ of a hyperbolic triangle Δ satisfies $0 < \Sigma < \pi$; in fact, Σ equals $\pi - \text{area } \Delta$. The limiting case $\Sigma = 0$ is achieved by *ideal triangles* whose vertices are “at infinity”: we will have more to say about such *ideal vertices* soon (p. 14). At the other extreme, the case $\Sigma = \pi$ is the limiting case of hyperbolic triangles of very small area. Indeed, on the infinitesimal scale, hyperbolic geometry is euclidean.
- (ii) There are no similarities in hyperbolic space—one cannot scale a figure up or down without changing its angles and shape. It follows, for instance, that all hyperbolic triangles with the same angles are isometric (hyperbolic triangles are “rigid”), and also that the choice of a unit of length is not arbitrary, as in euclidean space; one can privilege a unit having some special property, say the side length of an equilateral triangle whose vertex angles are $\pi/4$.
- (iii) For any $0 \leq \theta < \pi/(n-2)$ there is a regular n -sided hyperbolic polygon with vertex angles θ . More generally, a necessary and sufficient condition for the existence of an n -sided convex polygon with vertex angles θ_i (with $0 \leq \theta_i < \pi$) in clockwise order is that $\sum \theta_i < (n-2)\pi$. The polygon is uniquely determined up to isometry and its area is $(n-2)\pi - \sum \theta_i$.

We now discuss the most commonly used *conformal models* of the hyperbolic plane and of hyperbolic space with their metrics.




Fig. 1.3. Disk and upper half-plane models of \mathbb{H}^2 showing the same geodesics.

1.2.1 The hyperbolic plane

The *upper half-plane model* is $\{z \in \mathbb{C} : \text{Im } z > 0\}$ with the metric

$$ds = \frac{|dz|}{\text{Im } z}.$$

Here $\text{Im } z$ is the notation for the imaginary part. The *unit disk model* is $\{z \in \mathbb{C} : |z| < 1\}$ with the metric

$$ds = \frac{2|dz|}{1 - |z|^2}.$$

The two models are equivalent under any Möbius transformation that maps the upper half-plane onto the unit disk. We will denote either one of these models by \mathbb{H}^2 , the notation for the *hyperbolic plane*. These models have the following properties.

- (i) The metrics are *infinitesimally euclidean*; at each point they equal a rescaled euclidean metric. Thus the angle between two curves in the disk or upper half-plane is the same whether measured in the hyperbolic or the euclidean geometry; as a result these models are often called *conformal*. (For other models see Exercise (1-25) and following.)
- (ii) \mathbb{H}^2 is *complete* in its metric. Every arc tending to the “boundary” has infinite length.
- (iii) The hyperbolic metric is invariant under any Möbius transformation that maps the upper half-space or disk model onto itself. These transformations comprise the full group of orientation-preserving isometries.
- (iv) The hyperbolic lines (geodesics) in the upper half-plane model are semicircles orthogonal to \mathbb{R} and vertical half lines. In the disk model they are diameters and circular arcs orthogonal to $\{|z| = 1\}$.
- (v) Two hyperbolic triangles with the same vertex angles are isometric.

Proof of Property (v). Applying an isometry, we may assume that the base of the two triangles Δ, Δ' lie on the same geodesic ℓ , and that the left side of the two triangles also lie on the same geodesic ℓ^* . Denote the right edges by δ, δ' . If $\delta = \delta'$ then $\Delta = \Delta'$.

Now two distinct geodesics, which intersect a third geodesic at the same angle cannot themselves intersect. For then the angle sum would exceed π .

In view of this fact, assume $\delta \cap \ell$ separates $\delta' \cap \ell$ from the vertex $\ell^* \cap \ell$. δ' cannot intersect δ . Therefore δ' cannot intersect ℓ^* , in contradiction to the assumption that Δ' is a triangle. \square

1.2.2 Hyperbolic space

The *upper half-space model* is $\{(z, t) : z \in \mathbb{C}, t > 0\}$ with the metric

$$ds = \frac{|d\vec{x}|}{t}, \quad |d\vec{x}|^2 = |dz|^2 + dt^2.$$

1.2 Hyperbolic geometry

9




Fig. 1.4. Ball and upper half-space model of \mathbb{H}^3 showing geodesic planes.

The *ball model* is $\{\vec{x} \in \mathbb{R}^3 : |\vec{x}| < 1\}$ with the metric

$$ds = \frac{2|d\vec{x}|}{1 - |\vec{x}|^2}.$$

The two models are equivalent by a Möbius transformation that maps one to the other. Stereographic projection extends to such a Möbius transformation (Exercise (1-11)). We will refer to either of these models with its metric as *hyperbolic space* and denote it by \mathbb{H}^3 .

Here is a further list of properties:

- (i) The hyperbolic metric is infinitesimally euclidean and correctly represents the angles in \mathbb{H}^3 .
- (ii) \mathbb{H}^3 is complete in its metric: an arc tending to the “boundary” has infinite length.
- (iii) The hyperbolic planes in the upper half-space model are hemispheres orthogonal to \mathbb{C} and vertical euclidean half-planes. The lines (geodesics) are semi-circles orthogonal to \mathbb{C} and vertical euclidean half-lines. In the ball model the hyperbolic planes are spherical caps orthogonal to the unit sphere, and equatorial planes. The lines are circular arcs orthogonal to the unit sphere, and euclidean diameters.
- (iv) The hyperbolic metric is invariant under any Möbius transformation that maps the upper half-space or the ball model onto itself. These transformations form the full group of orientation-preserving isometries.
- (v) The volume V of a hyperbolic ball and the surface area S of its bounding sphere grow exponentially with the hyperbolic radius ρ . The ratio of the surface area to the volume approaches 2 as $\rho \rightarrow \infty$.
- (vi) Two convex hyperbolic polyhedra that are combinatorially the same with the same dihedral angles and valence 3 at all vertices are isometric [Rivin 1996; Bobenko and Springborn 2004].
- (vii) **Andreev-Thurston Theorem** [Roeder et al. 2007] and [Bowers and Stephenson 1996]. Also see Exercise (4-17). Let \mathbb{T} be a triangulation of \mathbb{S}^2 and $E = \{e_i\}$ the set of edges. Assume that \mathbb{T} is not equivalent to the edge set of a tetrahedron. Assign an angle $\Phi : E \rightarrow [0, \frac{\pi}{2}]$ to each edge e_i . The assignments must satisfy two conditions:

- (a) If the sequence of edges (e_1, e_2, e_3) form a closed loop on \mathbb{S}^2 satisfying $\sum_{i=1}^3 \Phi(e_i) \geq \pi$, then these edges bound a triangle in \mathbb{T} .
- (b) If the sequence (e_1, e_2, e_3, e_4) form a closed loop on \mathbb{S}^2 satisfying $\sum_{i=1}^4 \Phi(e_i) = 2\pi$ then these edges form the boundary of two adjacent triangles.

Then there is a realization of \mathbb{T} , (1) as a geodesic triangulation (in the spherical metric) $\mathbb{T}_{geod} \subset \mathbb{S}^2$, unique up to Möbius equivalence, together with (2) a family $C = \{C_i\}$ of circles, centered at the vertices of \mathbb{T}_{geod} , with the property that the two circles in C centered at the endpoints of an edge e_i intersect at angle $\Phi(e_i)$.

- (viii) Erect the hyperbolic planes $\{P_{C_i}\} \subset \mathbb{H}^3$ rising from the circles $\{C_i\}$. Assume, say in the ball model, that 0 is in the “exterior” $Ext(P_i)$ of each plane. Then $\cap_i Ext(P_i)$ is a hyperbolic polyhedron with dihedral angles $\{\Phi(e_i)\}$ coming from the intersecting circles centered at the endpoints of the edges $\{e_i\}$.

For analysis of properties of ideal polyhedrons (all vertices at “infinity” in \mathbb{H}^3) see Rivin [1996].

Restricting the hyperbolic metric to a hyperbolic plane in the model yields the two-dimensional hyperbolic metric on that plane. Particular cases are the vertical half-plane rising from \mathbb{R} in the upper half-space model and the equatorial plane in the ball model.

Proof of Property (iii). For the proof that the Möbius transformations are orientation-preserving isometries of the models, see Exercise 1-7. Here we show that there are no other such isometries, concentrating on the hyperbolic plane.

Given three positive distances d_1, d_2, d_3 satisfying the triangle inequality, and a point z on an oriented line $\ell \in \mathbb{H}^2$, there is a triangle Δ with a vertex at z , a side of length d_1 lying on the positive side of ℓ , a side of length d_2 lying in the right side of ℓ and sharing the vertex z , and a third side of length d_3 . There is one other triangle with sides of these lengths: the reflection of Δ in ℓ . One of the two is uniquely determined if an ordering of the vertices is given and required to give the positive orientation of the triangle they bound.

Given an orientation-preserving isometry T , the T -images of three points not on a line are not on a hyperbolic line either. There is a Möbius transformation A such that $A \circ T$ fixes the three points. It then pointwise fixes the sides of the triangle they determine, and then fixes the whole triangle, say Δ . That is, $T(z) = A^{-1}(z)$, for $z \in \Delta$. If Δ' is a triangle sharing an edge with Δ , there is Möbius transformation A_1 such that $T(z) = A_1^{-1}(z)$ on Δ' . Necessarily $A_1 = A$. Continuing on, building up the whole plane \mathbb{H}^2 by adding in succession adjacent triangles, we conclude that $T \equiv A$. \square

Proof of Property (iv). Working in the upper half-space model, in view of (iii), we need only prove that the vertical axis ℓ , rising from $z = 0$ is itself a geodesic. Fix a pair of points $t_1, t_2 \in \ell$. Let $\vec{x} = \vec{x}(s)$ be a smooth parameterized arc α between $t_1 < t_2 \in \ell$. Differentiating with respect to s ,